




BAYLOR
UNIVERSITY

MLP Algebraic Expansion and Simplification:

Reducing the computational error and increasing computational efficiency,

Blake Edward Schultze

June 13, 2015



$$\begin{aligned}
\frac{1}{\beta^2(u)p^2(u)} &= a_0 + a_1u + a_2u^2 + a_3u^3 + a_4u^4 + a_5u^5 \\
\int_{u_0}^{u_1} \frac{(u_1 - u)^2}{\beta^2(u)p^2(u)} \frac{du}{X_0} &= \frac{1}{X_0} \int_{u_0}^{u_1} (u_1 - u)^2 (a_0 + a_1u + a_2u^2 + a_3u^3 + a_4u^4 + a_5u^5) du \\
&= \frac{1}{X_0} \int_{u_0}^{u_1} (u_1^2 - 2u_1u + u^2) (a_0 + a_1u + a_2u^2 + a_3u^3 + a_4u^4 + a_5u^5) du \\
&= \frac{1}{X_0} \int_{u_0}^{u_1} \left[u_1^2 (a_0 + a_1u + a_2u^2 + a_3u^3 + a_4u^4 + a_5u^5) - 2u_1u (a_0 + a_1u + a_2u^2 + a_3u^3 + a_4u^4 + a_5u^5) + u^2 (a_0 + a_1u + a_2u^2 + a_3u^3 + a_4u^4 + a_5u^5) \right] du \\
&= \frac{1}{X_0} \int_{u_0}^{u_1} \left[u_1^2 (a_0 + a_1u + a_2u^2 + a_3u^3 + a_4u^4 + a_5u^5) - 2u_1 (a_0u + a_1u^2 + a_2u^3 + a_3u^4 + a_4u^5 + a_5u^6) + (a_0u^2 + a_1u^3 + a_2u^4 + a_3u^5 + a_4u^6 + a_5u^7) \right] du \\
&= \frac{1}{X_0} \left[u_1^2 \left(a_0u + \frac{a_1u^2}{2} + \frac{a_2u^3}{3} + \frac{a_3u^4}{4} + \frac{a_4u^5}{5} + \frac{a_5u^6}{6} \right) - 2u_1 \left(\frac{a_0u^2}{2} + \frac{a_1u^3}{3} + \frac{a_2u^4}{4} + \frac{a_3u^5}{5} + \frac{a_4u^6}{6} + \frac{a_5u^7}{7} \right) + \left(\frac{a_0u^3}{3} + \frac{a_1u^4}{4} + \frac{a_2u^5}{5} + \frac{a_3u^6}{6} + \frac{a_4u^7}{7} + \frac{a_5u^8}{8} \right) \right] \Big|_{u=u_0}^{u_1} \\
&= \frac{1}{X_0} \left[\left(a_0u_1^3 + \frac{a_1u_1^4}{2} + \frac{a_2u_1^5}{3} + \frac{a_3u_1^6}{4} + \frac{a_4u_1^7}{5} + \frac{a_5u_1^8}{6} \right) - u_1^2 \left(a_0u_0 + \frac{a_1u_0^2}{2} + \frac{a_2u_0^3}{3} + \frac{a_3u_0^4}{4} + \frac{a_4u_0^5}{5} + \frac{a_5u_0^6}{6} \right) - 2 \left(\frac{a_0u_1^3}{2} + \frac{a_1u_1^4}{3} + \frac{a_2u_1^5}{4} + \frac{a_3u_1^6}{5} + \frac{a_4u_1^7}{6} + \frac{a_5u_1^8}{7} \right) \right. \\
&\quad \left. + 2u_1 \left(\frac{a_0u_0^2}{2} + \frac{a_1u_0^3}{3} + \frac{a_2u_0^4}{4} + \frac{a_3u_0^5}{5} + \frac{a_4u_0^6}{6} + \frac{a_5u_0^7}{7} \right) + \left(\frac{a_0u_1^3}{3} + \frac{a_1u_1^4}{4} + \frac{a_2u_1^5}{5} + \frac{a_3u_1^6}{6} + \frac{a_4u_1^7}{7} + \frac{a_5u_1^8}{8} \right) - \left(\frac{a_0u_0^3}{3} + \frac{a_1u_0^4}{4} + \frac{a_2u_0^5}{5} + \frac{a_3u_0^6}{6} + \frac{a_4u_0^7}{7} + \frac{a_5u_0^8}{8} \right) \right] \\
&= \frac{1}{X_0} \left\{ a_0u_1^3 \left(1 - 1 + \frac{1}{3} \right) + a_1u_1^4 \left(\frac{1}{2} - \frac{2}{3} + \frac{1}{4} \right) + a_2u_1^5 \left(\frac{1}{3} - \frac{2}{4} + \frac{1}{5} \right) + a_3u_1^6 \left(\frac{1}{4} - \frac{2}{5} + \frac{1}{6} \right) + a_4u_1^7 \left(\frac{1}{5} - \frac{2}{6} + \frac{1}{7} \right) + a_5u_1^8 \left(\frac{1}{6} - \frac{2}{7} + \frac{1}{8} \right) \right. \\
&\quad \left. - u_1^2 \left(a_0u_0 + \frac{a_1u_0^2}{2} + \frac{a_2u_0^3}{3} + \frac{a_3u_0^4}{4} + \frac{a_4u_0^5}{5} + \frac{a_5u_0^6}{6} \right) + 2u_1 \left(\frac{a_0u_0^2}{2} + \frac{a_1u_0^3}{3} + \frac{a_2u_0^4}{4} + \frac{a_3u_0^5}{5} + \frac{a_4u_0^6}{6} + \frac{a_5u_0^7}{7} \right) - \left(\frac{a_0u_0^3}{3} + \frac{a_1u_0^4}{4} + \frac{a_2u_0^5}{5} + \frac{a_3u_0^6}{6} + \frac{a_4u_0^7}{7} + \frac{a_5u_0^8}{8} \right) \right\}
\end{aligned}$$

$$\int_{u_0}^{u_1} \frac{(u_1 - u)^2}{\beta^2(u)p^2(u)} \frac{du}{X_0} = \frac{1}{X_0} \left\{ \frac{a_0u_1^3}{3} + \frac{a_1u_1^4}{12} + \frac{a_2u_1^5}{30} + \frac{a_3u_1^6}{60} + \frac{a_4u_1^7}{105} + \frac{a_5u_1^8}{168} - u_1^2 \left(a_0u_0 + \frac{a_1u_0^2}{2} + \frac{a_2u_0^3}{3} + \frac{a_3u_0^4}{4} + \frac{a_4u_0^5}{5} + \frac{a_5u_0^6}{6} \right) \right. \\
\left. + 2u_1 \left(\frac{a_0u_0^2}{2} + \frac{a_1u_0^3}{3} + \frac{a_2u_0^4}{4} + \frac{a_3u_0^5}{5} + \frac{a_4u_0^6}{6} + \frac{a_5u_0^7}{7} \right) - \left(\frac{a_0u_0^3}{3} + \frac{a_1u_0^4}{4} + \frac{a_2u_0^5}{5} + \frac{a_3u_0^6}{6} + \frac{a_4u_0^7}{7} + \frac{a_5u_0^8}{8} \right) \right\}$$

$$\begin{aligned}
\int_{u_0}^{u_1} \frac{u_1 - u}{\beta^2(u)p^2(u)} \frac{du}{X_0} &= \frac{1}{X_0} \int_{u_0}^{u_1} (u_1 - u) (a_0 + a_1 u + a_2 u^2 + a_3 u^3 + a_4 u^4 + a_5 u^5) du \\
&= \frac{1}{X_0} \left[u_1 \left(a_0 u + \frac{a_1 u^2}{2} + \frac{a_2 u^3}{3} + \frac{a_3 u^4}{4} + \frac{a_4 u^5}{5} + \frac{a_5 u^6}{6} \right) - \left(\frac{a_0 u^2}{2} + \frac{a_1 u^3}{3} + \frac{a_2 u^4}{4} + \frac{a_3 u^5}{5} + \frac{a_4 u^6}{6} + \frac{a_5 u^7}{7} \right) \right]_{u=u_0}^{u_1} \\
&= \frac{1}{X_0} \left[\left(a_0 u_1^2 + \frac{a_1 u_1^3}{2} + \frac{a_2 u_1^4}{3} + \frac{a_3 u_1^5}{4} + \frac{a_4 u_1^6}{5} + \frac{a_5 u_1^7}{6} \right) - u_1 \left(a_0 u_0 + \frac{a_1 u_0^2}{2} + \frac{a_2 u_0^3}{3} + \frac{a_3 u_0^4}{4} + \frac{a_4 u_0^5}{5} + \frac{a_5 u_0^6}{6} \right) \right. \\
&\quad \left. - \left(\frac{a_0 u_1^2}{2} + \frac{a_1 u_1^3}{3} + \frac{a_2 u_1^4}{4} + \frac{a_3 u_1^5}{5} + \frac{a_4 u_1^6}{6} + \frac{a_5 u_1^7}{7} \right) + \left(\frac{a_0 u_0^2}{2} + \frac{a_1 u_0^3}{3} + \frac{a_2 u_0^4}{4} + \frac{a_3 u_0^5}{5} + \frac{a_4 u_0^6}{6} + \frac{a_5 u_0^7}{7} \right) \right] \\
&= \frac{1}{X_0} \left\{ \left[a_0 u_1^2 \left(\frac{1}{1} - \frac{1}{2} \right) + a_1 u_1^3 \left(\frac{1}{2} - \frac{1}{3} \right) + a_2 u_1^4 \left(\frac{1}{3} - \frac{1}{4} \right) + a_3 u_1^5 \left(\frac{1}{4} - \frac{1}{5} \right) + a_4 u_1^6 \left(\frac{1}{5} - \frac{1}{6} \right) + a_5 u_1^7 \left(\frac{1}{6} - \frac{1}{7} \right) \right] \right. \\
&\quad \left. - u_1 \left(a_0 u_0 + \frac{a_1 u_0^2}{2} + \frac{a_2 u_0^3}{3} + \frac{a_3 u_0^4}{4} + \frac{a_4 u_0^5}{5} + \frac{a_5 u_0^6}{6} \right) + \left(\frac{a_0 u_0^2}{2} + \frac{a_1 u_0^3}{3} + \frac{a_2 u_0^4}{4} + \frac{a_3 u_0^5}{5} + \frac{a_4 u_0^6}{6} + \frac{a_5 u_0^7}{7} \right) \right\}
\end{aligned}$$

$$\int_{u_0}^{u_1} \frac{u_1 - u}{\beta^2(u)p^2(u)} \frac{du}{X_0} = \frac{1}{X_0} \left[\left(\frac{a_0 u_1^2}{2} + \frac{a_1 u_1^3}{6} + \frac{a_2 u_1^4}{12} + \frac{a_3 u_1^5}{20} + \frac{a_4 u_1^6}{30} + \frac{a_5 u_1^7}{42} \right) - u_1 \left(a_0 u_0 + \frac{a_1 u_0^2}{2} + \frac{a_2 u_0^3}{3} + \frac{a_3 u_0^4}{4} + \frac{a_4 u_0^5}{5} + \frac{a_5 u_0^6}{6} \right) + \left(\frac{a_0 u_0^2}{2} + \frac{a_1 u_0^3}{3} + \frac{a_2 u_0^4}{4} + \frac{a_3 u_0^5}{5} + \frac{a_4 u_0^6}{6} + \frac{a_5 u_0^7}{7} \right) \right]$$

$$\begin{aligned}
\int_{u_0}^{u_1} \frac{1}{\beta^2(u)p^2(u)} \frac{du}{X_0} &= \frac{1}{X_0} \int_{u_0}^{u_1} (a_0 + a_1 u + a_2 u^2 + a_3 u^3 + a_4 u^4 + a_5 u^5) du \\
&= \frac{1}{X_0} \left[a_0 u + \frac{a_1 u^2}{2} + \frac{a_2 u^3}{3} + \frac{a_3 u^4}{4} + \frac{a_4 u^5}{5} + \frac{a_5 u^6}{6} \right]_{u=u_0}^{u_1}
\end{aligned}$$

$$\int_{u_0}^{u_1} \frac{1}{\beta^2(u)p^2(u)} \frac{du}{X_0} = \frac{1}{X_0} \left[\left(a_0 u_1 + \frac{a_1 u_1^2}{2} + \frac{a_2 u_1^3}{3} + \frac{a_3 u_1^4}{4} + \frac{a_4 u_1^5}{5} + \frac{a_5 u_1^6}{6} \right) - \left(a_0 u_0 + \frac{a_1 u_0^2}{2} + \frac{a_2 u_0^3}{3} + \frac{a_3 u_0^4}{4} + \frac{a_4 u_0^5}{5} + \frac{a_5 u_0^6}{6} \right) \right]$$

$$\begin{aligned}
\int_{u_1}^{u_2} \frac{(u_2 - u)^2}{\beta^2(u)p^2(u)} \frac{du}{X_0} &= \frac{1}{X_0} \int_{u_1}^{u_2} (u_2 - u)^2 (a_0 + a_1u + a_2u^2 + a_3u^3 + a_4u^4 + a_5u^5) du = \frac{1}{X_0} \int_{u_1}^{u_2} (u_2^2 - 2u_2u + u^2) (a_0 + a_1u + a_2u^2 + a_3u^3 + a_4u^4 + a_5u^5) du \\
&= \frac{1}{X_0} \left[u_2^2 \left(a_0u + \frac{a_1u^2}{2} + \frac{a_2u^3}{3} + \frac{a_3u^4}{4} + \frac{a_4u^5}{5} + \frac{a_5u^6}{6} \right) - 2u_2 \left(\frac{a_0u^2}{2} + \frac{a_1u^3}{3} + \frac{a_2u^4}{4} + \frac{a_3u^5}{5} + \frac{a_4u^6}{6} + \frac{a_5u^7}{7} \right) + \left(\frac{a_0u^3}{3} + \frac{a_1u^4}{4} + \frac{a_2u^5}{5} + \frac{a_3u^6}{6} + \frac{a_4u^7}{7} + \frac{a_5u^8}{8} \right) \right]_{u=u_1}^{u_2} \\
&= \frac{1}{X_0} \left[\left(a_0u_2^3 + \frac{a_1u_2^4}{2} + \frac{a_2u_2^5}{3} + \frac{a_3u_2^6}{4} + \frac{a_4u_2^7}{5} + \frac{a_5u_2^8}{6} \right) - u_2^2 \left(a_0u_1 + \frac{a_1u_1^2}{2} + \frac{a_2u_1^3}{3} + \frac{a_3u_1^4}{4} + \frac{a_4u_1^5}{5} + \frac{a_5u_1^6}{6} \right) - 2 \left(\frac{a_0u_2^3}{2} + \frac{a_1u_2^4}{3} + \frac{a_2u_2^5}{4} + \frac{a_3u_2^6}{5} + \frac{a_4u_2^7}{6} + \frac{a_5u_2^8}{7} \right) \right. \\
&\quad \left. + 2u_2 \left(\frac{a_0u_1^2}{2} + \frac{a_1u_1^3}{3} + \frac{a_2u_1^4}{4} + \frac{a_3u_1^5}{5} + \frac{a_4u_1^6}{6} + \frac{a_5u_1^7}{7} \right) + \left(\frac{a_0u_2^3}{3} + \frac{a_1u_2^4}{4} + \frac{a_2u_2^5}{5} + \frac{a_3u_2^6}{6} + \frac{a_4u_2^7}{7} + \frac{a_5u_2^8}{8} \right) - \left(\frac{a_0u_1^3}{3} + \frac{a_1u_1^4}{4} + \frac{a_2u_1^5}{5} + \frac{a_3u_1^6}{6} + \frac{a_4u_1^7}{7} + \frac{a_5u_1^8}{8} \right) \right] \\
&= \frac{1}{X_0} \left\{ a_0u_2^3 \left(1 - 1 + \frac{1}{3} \right) + a_1u_2^4 \left(\frac{1}{2} - \frac{2}{3} + \frac{1}{4} \right) + a_2u_2^5 \left(\frac{1}{3} - \frac{2}{4} + \frac{1}{5} \right) + a_3u_2^6 \left(\frac{1}{4} - \frac{2}{5} + \frac{1}{6} \right) + a_4u_2^7 \left(\frac{1}{5} - \frac{2}{6} + \frac{1}{7} \right) + a_5u_2^8 \left(\frac{1}{6} - \frac{2}{7} + \frac{1}{8} \right) \right. \\
&\quad \left. - u_2^2 \left(a_0u_1 + \frac{a_1u_1^2}{2} + \frac{a_2u_1^3}{3} + \frac{a_3u_1^4}{4} + \frac{a_4u_1^5}{5} + \frac{a_5u_1^6}{6} \right) + 2u_2 \left(\frac{a_0u_1^2}{2} + \frac{a_1u_1^3}{3} + \frac{a_2u_1^4}{4} + \frac{a_3u_1^5}{5} + \frac{a_4u_1^6}{6} + \frac{a_5u_1^7}{7} \right) - \left(\frac{a_0u_1^3}{3} + \frac{a_1u_1^4}{4} + \frac{a_2u_1^5}{5} + \frac{a_3u_1^6}{6} + \frac{a_4u_1^7}{7} + \frac{a_5u_1^8}{8} \right) \right\}
\end{aligned}$$

$$\begin{aligned}
\int_{u_1}^{u_2} \frac{(u_2 - u)^2}{\beta^2(u)p^2(u)} \frac{du}{X_0} &= \frac{1}{X_0} \left\{ \left[\frac{a_0u_2^3}{3} + \frac{a_1u_2^4}{12} + \frac{a_2u_2^5}{30} + \frac{a_3u_2^6}{60} + \frac{a_4u_2^7}{105} + \frac{a_5u_2^8}{168} \right] - u_2^2 \left(a_0u_1 + \frac{a_1u_1^2}{2} + \frac{a_2u_1^3}{3} + \frac{a_3u_1^4}{4} + \frac{a_4u_1^5}{5} + \frac{a_5u_1^6}{6} \right) \right. \\
&\quad \left. + 2u_2 \left(\frac{a_0u_1^2}{2} + \frac{a_1u_1^3}{3} + \frac{a_2u_1^4}{4} + \frac{a_3u_1^5}{5} + \frac{a_4u_1^6}{6} + \frac{a_5u_1^7}{7} \right) - \left(\frac{a_0u_1^3}{3} + \frac{a_1u_1^4}{4} + \frac{a_2u_1^5}{5} + \frac{a_3u_1^6}{6} + \frac{a_4u_1^7}{7} + \frac{a_5u_1^8}{8} \right) \right\} \\
&= \frac{1}{X_0} \left\{ \left(\frac{a_0u_2^3}{3} + \frac{a_1u_2^4}{12} + \frac{a_2u_2^5}{30} + \frac{a_3u_2^6}{60} + \frac{a_4u_2^7}{105} + \frac{a_5u_2^8}{168} \right) + u_1 (-a_0u_2^2) + u_1^2 \left(\frac{-a_1u_2^2}{2} + \frac{2a_0u_2}{2} \right) + u_1^3 \left(\frac{-a_2u_2^2}{3} + \frac{2a_1u_2}{3} - \frac{a_0}{3} \right) + u_1^4 \left(\frac{-a_3u_2^2}{4} + \frac{2a_2u_2}{4} - \frac{a_1}{4} \right) \right. \\
&\quad \left. + u_1^5 \left(\frac{-a_4u_2^2}{5} + \frac{2a_3u_2}{5} - \frac{a_2}{5} \right) + u_1^6 \left(\frac{-a_5u_2^2}{6} + \frac{2a_4u_2}{6} - \frac{a_3}{6} \right) + u_1^7 \left(\frac{2a_5u_2}{7} - \frac{a_4}{7} \right) + u_1^8 \left(\frac{a_5}{8} \right) \right\} \\
&= \frac{1}{X_0} \left\{ \left(\frac{a_0u_2^3}{3} + \frac{a_1u_2^4}{12} + \frac{a_2u_2^5}{30} + \frac{a_3u_2^6}{60} + \frac{a_4u_2^7}{105} + \frac{a_5u_2^8}{168} \right) + u_1 (-a_0u_2^2) + u_1^2 \left(\frac{-a_1u_2^2 + 2a_0u_2}{2} \right) + u_1^3 \left(\frac{-a_2u_2^2 + 2a_1u_2 - a_0}{3} \right) + u_1^4 \left(\frac{-a_3u_2^2 + 2a_2u_2 - a_1}{4} \right) \right. \\
&\quad \left. + u_1^5 \left(\frac{-a_4u_2^2 + 2a_3u_2 - a_2}{5} \right) + u_1^6 \left(\frac{-a_5u_2^2 + 2a_4u_2 - a_3}{6} \right) + u_1^7 \left(\frac{2a_5u_2 - a_4}{7} \right) + u_1^8 \left(\frac{-a_5}{8} \right) \right\}
\end{aligned}$$

$$\begin{aligned}
\int_{u_1}^{u_2} \frac{(u_2 - u)^2}{\beta^2(u)p^2(u)} \frac{du}{X_0} &= \frac{1}{X_0} \left\{ \left(\frac{a_0u_2^3}{3} + \frac{a_1u_2^4}{12} + \frac{a_2u_2^5}{30} + \frac{a_3u_2^6}{60} + \frac{a_4u_2^7}{105} + \frac{a_5u_2^8}{168} \right) + u_1 (-a_0u_2^2) + u_1^2 \left[-\left(\frac{a_1}{2} \right) u_2^2 + 2 \left(\frac{a_0}{2} \right) u_2 \right] + u_1^3 \left[-\left(\frac{a_2}{3} \right) u_2^2 + 2 \left(\frac{a_1}{3} \right) u_2 - \frac{a_0}{3} \right] \right. \\
&\quad \left. + u_1^4 \left[-\left(\frac{a_3}{4} \right) u_2^2 + 2 \left(\frac{a_2}{4} \right) u_2 - \frac{a_1}{4} \right] + u_1^5 \left[-\left(\frac{a_4}{5} \right) u_2^2 + 2 \left(\frac{a_3}{5} \right) u_2 - \frac{a_2}{5} \right] + u_1^6 \left[-\left(\frac{a_5}{6} \right) u_2^2 + 2 \left(\frac{a_4}{6} \right) u_2 - \frac{a_3}{6} \right] + u_1^7 \left[2 \left(\frac{a_5}{7} \right) u_2 - \frac{a_4}{7} \right] + u_1^8 \left[-\frac{a_5}{8} \right] \right\}
\end{aligned}$$

$$\begin{aligned}
\int_{u_1}^{u_2} \frac{u_2 - u}{\beta^2(u)p^2(u)} \frac{du}{X_0} &= \frac{1}{X_0} \int_{u_1}^{u_2} (u_2 - u) (a_0 + a_1 u + a_2 u^2 + a_3 u^3 + a_4 u^4 + a_5 u^5) du \\
&= \frac{1}{X_0} \left[u_2 \left(a_0 u + \frac{a_1 u^2}{2} + \frac{a_2 u^3}{3} + \frac{a_3 u^4}{4} + \frac{a_4 u^5}{5} + \frac{a_5 u^6}{6} \right) - \left(\frac{a_0 u^2}{2} + \frac{a_1 u^3}{3} + \frac{a_2 u^4}{4} + \frac{a_3 u^5}{5} + \frac{a_4 u^6}{6} + \frac{a_5 u^7}{7} \right) \right]_{u=u_1}^{u_2} \\
&= \frac{1}{X_0} \left[\left(a_0 u_2^2 + \frac{a_1 u_2^3}{2} + \frac{a_2 u_2^4}{3} + \frac{a_3 u_2^5}{4} + \frac{a_4 u_2^6}{5} + \frac{a_5 u_2^7}{6} \right) - u_2 \left(a_0 u_1 + \frac{a_1 u_1^2}{2} + \frac{a_2 u_1^3}{3} + \frac{a_3 u_1^4}{4} + \frac{a_4 u_1^5}{5} + \frac{a_5 u_1^6}{6} \right) \right. \\
&\quad \left. - \left(\frac{a_0 u_2^2}{2} + \frac{a_1 u_2^3}{3} + \frac{a_2 u_2^4}{4} + \frac{a_3 u_2^5}{5} + \frac{a_4 u_2^6}{6} + \frac{a_5 u_2^7}{7} \right) + \left(\frac{a_0 u_1^2}{2} + \frac{a_1 u_1^3}{3} + \frac{a_2 u_1^4}{4} + \frac{a_3 u_1^5}{5} + \frac{a_4 u_1^6}{6} + \frac{a_5 u_1^7}{7} \right) \right] \\
&= \frac{1}{X_0} \left\{ \left[a_0 u_2^2 \left(\frac{1}{1} - \frac{1}{2} \right) + a_1 u_2^3 \left(\frac{1}{2} - \frac{1}{3} \right) + a_2 u_2^4 \left(\frac{1}{3} - \frac{1}{4} \right) + a_3 u_2^5 \left(\frac{1}{4} - \frac{1}{5} \right) + a_4 u_2^6 \left(\frac{1}{5} - \frac{1}{6} \right) + a_5 u_2^7 \left(\frac{1}{6} - \frac{1}{7} \right) \right] \right. \\
&\quad \left. - u_2 \left(a_0 u_1 + \frac{a_1 u_1^2}{2} + \frac{a_2 u_1^3}{3} + \frac{a_3 u_1^4}{4} + \frac{a_4 u_1^5}{5} + \frac{a_5 u_1^6}{6} \right) + \left(\frac{a_0 u_1^2}{2} + \frac{a_1 u_1^3}{3} + \frac{a_2 u_1^4}{4} + \frac{a_3 u_1^5}{5} + \frac{a_4 u_1^6}{6} + \frac{a_5 u_1^7}{7} \right) \right\}
\end{aligned}$$

$$\int_{u_1}^{u_2} \frac{u_2 - u}{\beta^2(u)p^2(u)} \frac{du}{X_0} = \frac{1}{X_0} \left[\left(\frac{a_0 u_2^2}{2} + \frac{a_1 u_2^3}{6} + \frac{a_2 u_2^4}{12} + \frac{a_3 u_2^5}{20} + \frac{a_4 u_2^6}{30} + \frac{a_5 u_2^7}{42} \right) - u_2 \left(a_0 u_1 + \frac{a_1 u_1^2}{2} + \frac{a_2 u_1^3}{3} + \frac{a_3 u_1^4}{4} + \frac{a_4 u_1^5}{5} + \frac{a_5 u_1^6}{6} \right) + \left(\frac{a_0 u_1^2}{2} + \frac{a_1 u_1^3}{3} + \frac{a_2 u_1^4}{4} + \frac{a_3 u_1^5}{5} + \frac{a_4 u_1^6}{6} + \frac{a_5 u_1^7}{7} \right) \right]$$

$$\begin{aligned}
\int_{u_1}^{u_2} \frac{1}{\beta^2(u)p^2(u)} \frac{du}{X_0} &= \frac{1}{X_0} \int_{u_1}^{u_2} (a_0 + a_1 u + a_2 u^2 + a_3 u^3 + a_4 u^4 + a_5 u^5) du \\
&= \frac{1}{X_0} \left[a_0 u + \frac{a_1 u^2}{2} + \frac{a_2 u^3}{3} + \frac{a_3 u^4}{4} + \frac{a_4 u^5}{5} + \frac{a_5 u^6}{6} \right]_{u=u_1}^{u_2}
\end{aligned}$$

$$\int_{u_1}^{u_2} \frac{1}{\beta^2(u)p^2(u)} \frac{du}{X_0} = \frac{1}{X_0} \left[\left(a_0 u_2 + \frac{a_1 u_2^2}{2} + \frac{a_2 u_2^3}{3} + \frac{a_3 u_2^4}{4} + \frac{a_4 u_2^5}{5} + \frac{a_5 u_2^6}{6} \right) - \left(a_0 u_1 + \frac{a_1 u_1^2}{2} + \frac{a_2 u_1^3}{3} + \frac{a_3 u_1^4}{4} + \frac{a_4 u_1^5}{5} + \frac{a_5 u_1^6}{6} \right) \right]$$

$$\int_{u_0}^{u_1} \frac{(u_1 - u)^2}{\beta^2(u)p^2(u)} \frac{du}{X_0} = \frac{1}{X_0} \left\{ \left(\frac{a_0}{3} \right) u_1^3 + \left(\frac{a_1}{12} \right) u_1^4 + \left(\frac{a_2}{30} \right) u_1^5 + \left(\frac{a_3}{60} \right) u_1^6 + \left(\frac{a_4}{105} \right) u_1^7 + \left(\frac{a_5}{168} \right) u_1^8 - u_1^2 \left[(a_0)u_0 + \left(\frac{a_1}{2} \right) u_0^2 + \left(\frac{a_2}{3} \right) u_0^3 + \left(\frac{a_3}{4} \right) u_0^4 + \left(\frac{a_4}{5} \right) u_0^5 + \left(\frac{a_5}{6} \right) u_0^6 \right] \right. \\ \left. + 2u_1 \left[\left(\frac{a_0}{2} \right) u_0^2 + \left(\frac{a_1}{3} \right) u_0^3 + \left(\frac{a_2}{4} \right) u_0^4 + \left(\frac{a_3}{5} \right) u_0^5 + \left(\frac{a_4}{6} \right) u_0^6 + \left(\frac{a_5}{7} \right) u_0^7 \right] - \left[\left(\frac{a_0}{3} \right) u_0^3 + \left(\frac{a_1}{4} \right) u_0^4 + \left(\frac{a_2}{5} \right) u_0^5 + \left(\frac{a_3}{6} \right) u_0^6 + \left(\frac{a_4}{7} \right) u_0^7 + \left(\frac{a_5}{8} \right) u_0^8 \right] \right\}$$

$$\int_{u_0}^{u_1} \frac{u_1 - u}{\beta^2(u)p^2(u)} \frac{du}{X_0} = \frac{1}{X_0} \left\{ \left[\left(\frac{a_0}{2} \right) u_1^2 + \left(\frac{a_1}{6} \right) u_1^3 + \left(\frac{a_2}{12} \right) u_1^4 + \left(\frac{a_3}{20} \right) u_1^5 + \left(\frac{a_4}{30} \right) u_1^6 + \left(\frac{a_5}{42} \right) u_1^7 \right] - u_1 \left[(a_0)u_0 + \left(\frac{a_1}{2} \right) u_0^2 + \left(\frac{a_2}{3} \right) u_0^3 + \left(\frac{a_3}{4} \right) u_0^4 + \left(\frac{a_4}{5} \right) u_0^5 + \left(\frac{a_5}{6} \right) u_0^6 \right] \right. \\ \left. + \left[\left(\frac{a_0}{2} \right) u_0^2 + \left(\frac{a_1}{3} \right) u_0^3 + \left(\frac{a_2}{4} \right) u_0^4 + \left(\frac{a_3}{5} \right) u_0^5 + \left(\frac{a_4}{6} \right) u_0^6 + \left(\frac{a_5}{7} \right) u_0^7 \right] \right\}$$

$$\int_{u_0}^{u_1} \frac{1}{\beta^2(u)p^2(u)} \frac{du}{X_0} = \frac{1}{X_0} \left\{ \left[(a_0)u_1 + \left(\frac{a_1}{2} \right) u_1^2 + \left(\frac{a_2}{3} \right) u_1^3 + \left(\frac{a_3}{4} \right) u_1^4 + \left(\frac{a_4}{5} \right) u_1^5 + \left(\frac{a_5}{6} \right) u_1^6 \right] - \left[(a_0)u_0 + \left(\frac{a_1}{2} \right) u_0^2 + \left(\frac{a_2}{3} \right) u_0^3 + \left(\frac{a_3}{4} \right) u_0^4 + \left(\frac{a_4}{5} \right) u_0^5 + \left(\frac{a_5}{6} \right) u_0^6 \right] \right\}$$

$$\int_{u_1}^{u_2} \frac{(u_2 - u)^2}{\beta^2(u)p^2(u)} \frac{du}{X_0} = \frac{1}{X_0} \left\{ \left[\left(\frac{a_0}{3} \right) u_2^3 + \left(\frac{a_1}{12} \right) u_2^4 + \left(\frac{a_2}{30} \right) u_2^5 + \left(\frac{a_3}{60} \right) u_2^6 + \left(\frac{a_4}{105} \right) u_2^7 + \left(\frac{a_5}{168} \right) u_2^8 \right] - u_2^2 \left[(a_0)u_1 + \left(\frac{a_1}{2} \right) u_1^2 + \left(\frac{a_2}{3} \right) u_1^3 + \left(\frac{a_3}{4} \right) u_1^4 + \left(\frac{a_4}{5} \right) u_1^5 + \left(\frac{a_5}{6} \right) u_1^6 \right] \right. \\ \left. + 2u_2 \left[\left(\frac{a_0}{2} \right) u_1^2 + \left(\frac{a_1}{3} \right) u_1^3 + \left(\frac{a_2}{4} \right) u_1^4 + \left(\frac{a_3}{5} \right) u_1^5 + \left(\frac{a_4}{6} \right) u_1^6 + \left(\frac{a_5}{7} \right) u_1^7 \right] - \left[\left(\frac{a_0}{3} \right) u_1^3 + \left(\frac{a_1}{4} \right) u_1^4 + \left(\frac{a_2}{5} \right) u_1^5 + \left(\frac{a_3}{6} \right) u_1^6 + \left(\frac{a_4}{7} \right) u_1^7 + \left(\frac{a_5}{8} \right) u_1^8 \right] \right\}$$

$$\int_{u_1}^{u_2} \frac{u_2 - u}{\beta^2(u)p^2(u)} \frac{du}{X_0} = \frac{1}{X_0} \left\{ \left[\left(\frac{a_0}{2} \right) u_2^2 + \left(\frac{a_1}{6} \right) u_2^3 + \left(\frac{a_2}{12} \right) u_2^4 + \left(\frac{a_3}{20} \right) u_2^5 + \left(\frac{a_4}{30} \right) u_2^6 + \left(\frac{a_5}{42} \right) u_2^7 \right] - u_2 \left[(a_0)u_1 + \left(\frac{a_1}{2} \right) u_1^2 + \left(\frac{a_2}{3} \right) u_1^3 + \left(\frac{a_3}{4} \right) u_1^4 + \left(\frac{a_4}{5} \right) u_1^5 + \left(\frac{a_5}{6} \right) u_1^6 \right] \right. \\ \left. + \left[\left(\frac{a_0}{2} \right) u_1^2 + \left(\frac{a_1}{3} \right) u_1^3 + \left(\frac{a_2}{4} \right) u_1^4 + \left(\frac{a_3}{5} \right) u_1^5 + \left(\frac{a_4}{6} \right) u_1^6 + \left(\frac{a_5}{7} \right) u_1^7 \right] \right\}$$

$$\int_{u_1}^{u_2} \frac{1}{\beta^2(u)p^2(u)} \frac{du}{X_0} = \frac{1}{X_0} \left\{ \left[(a_0)u_2 + \left(\frac{a_1}{2} \right) u_2^2 + \left(\frac{a_2}{3} \right) u_2^3 + \left(\frac{a_3}{4} \right) u_2^4 + \left(\frac{a_4}{5} \right) u_2^5 + \left(\frac{a_5}{6} \right) u_2^6 \right] - \left[(a_0)u_1 + \left(\frac{a_1}{2} \right) u_1^2 + \left(\frac{a_2}{3} \right) u_1^3 + \left(\frac{a_3}{4} \right) u_1^4 + \left(\frac{a_4}{5} \right) u_1^5 + \left(\frac{a_5}{6} \right) u_1^6 \right] \right\}$$

Note that in the above equations, terms in bold denote those that can be precalculated, either before compilation (as is the case for the a_i/c_j terms, which are constants) or once per iteration (as is the case with the u_0/u_2 terms). Note also that the terms in green, blue, cyan, and brown highlight the terms that are exactly the same and appear in multiple expressions.

Since the only thing that really matters in our calculations is the depth $u_1 - u_0$ and the remaining depth $u_2 - u_1$, we see that these are relative distances, which are unaffected by a translation applied to each term (i.e. u_0 , u_1 , and u_2). Thus, we can simplify our calculations considerably by simply defining the entry coordinate $u_0 = 0$

so that the depth inside the object is $u_1 - u_0 = u_1$. Therefore, the above calculations reduce to

$$\int_{u_0}^{u_1} \frac{(u_1 - u)^2}{\beta^2(u)p^2(u)} \frac{du}{X_0} = \frac{1}{X_0} \left[\left(\frac{a_0}{3}\right)u_1^3 + \left(\frac{a_1}{12}\right)u_1^4 + \left(\frac{a_2}{30}\right)u_1^5 + \left(\frac{a_3}{60}\right)u_1^6 + \left(\frac{a_4}{105}\right)u_1^7 + \left(\frac{a_5}{168}\right)u_1^8 \right]$$

$$\int_{u_0}^{u_1} \frac{u_1 - u}{\beta^2(u)p^2(u)} \frac{du}{X_0} = \frac{1}{X_0} \left[\left(\frac{a_0}{2}\right)u_1^2 + \left(\frac{a_1}{6}\right)u_1^3 + \left(\frac{a_2}{12}\right)u_1^4 + \left(\frac{a_3}{20}\right)u_1^5 + \left(\frac{a_4}{30}\right)u_1^6 + \left(\frac{a_5}{42}\right)u_1^7 \right]$$

$$\int_{u_0}^{u_1} \frac{1}{\beta^2(u)p^2(u)} \frac{du}{X_0} = \frac{1}{X_0} \left[(a_0)u_1 + \left(\frac{a_1}{2}\right)u_1^2 + \left(\frac{a_2}{3}\right)u_1^3 + \left(\frac{a_3}{4}\right)u_1^4 + \left(\frac{a_4}{5}\right)u_1^5 + \left(\frac{a_5}{6}\right)u_1^6 \right]$$

$$\int_{u_1}^{u_2} \frac{(u_2 - u)^2}{\beta^2(u)p^2(u)} \frac{du}{X_0} = \frac{1}{X_0} \left\{ \left[\left(\frac{a_0}{3}\right)u_2^3 + \left(\frac{a_1}{12}\right)u_2^4 + \left(\frac{a_2}{30}\right)u_2^5 + \left(\frac{a_3}{60}\right)u_2^6 + \left(\frac{a_4}{105}\right)u_2^7 + \left(\frac{a_5}{168}\right)u_2^8 \right] - u_2^2 \left[(a_0)u_1 + \left(\frac{a_1}{2}\right)u_1^2 + \left(\frac{a_2}{3}\right)u_1^3 + \left(\frac{a_3}{4}\right)u_1^4 + \left(\frac{a_4}{5}\right)u_1^5 + \left(\frac{a_5}{6}\right)u_1^6 \right] \right. \\ \left. + 2u_2 \left[\left(\frac{a_0}{2}\right)u_1^2 + \left(\frac{a_1}{3}\right)u_1^3 + \left(\frac{a_2}{4}\right)u_1^4 + \left(\frac{a_3}{5}\right)u_1^5 + \left(\frac{a_4}{6}\right)u_1^6 + \left(\frac{a_5}{7}\right)u_1^7 \right] - \left[\left(\frac{a_0}{3}\right)u_1^3 + \left(\frac{a_1}{4}\right)u_1^4 + \left(\frac{a_2}{5}\right)u_1^5 + \left(\frac{a_3}{6}\right)u_1^6 + \left(\frac{a_4}{7}\right)u_1^7 + \left(\frac{a_5}{8}\right)u_1^8 \right] \right\}$$

$$\int_{u_1}^{u_2} \frac{u_2 - u}{\beta^2(u)p^2(u)} \frac{du}{X_0} = \frac{1}{X_0} \left\{ \left[\left(\frac{a_0}{2}\right)u_2^2 + \left(\frac{a_1}{6}\right)u_2^3 + \left(\frac{a_2}{12}\right)u_2^4 + \left(\frac{a_3}{20}\right)u_2^5 + \left(\frac{a_4}{30}\right)u_2^6 + \left(\frac{a_5}{42}\right)u_2^7 \right] - u_2 \left[(a_0)u_1 + \left(\frac{a_1}{2}\right)u_1^2 + \left(\frac{a_2}{3}\right)u_1^3 + \left(\frac{a_3}{4}\right)u_1^4 + \left(\frac{a_4}{5}\right)u_1^5 + \left(\frac{a_5}{6}\right)u_1^6 \right] \right. \\ \left. + \left[\left(\frac{a_0}{2}\right)u_1^2 + \left(\frac{a_1}{3}\right)u_1^3 + \left(\frac{a_2}{4}\right)u_1^4 + \left(\frac{a_3}{5}\right)u_1^5 + \left(\frac{a_4}{6}\right)u_1^6 + \left(\frac{a_5}{7}\right)u_1^7 \right] \right\}$$

$$\int_{u_1}^{u_2} \frac{1}{\beta^2(u)p^2(u)} \frac{du}{X_0} = \frac{1}{X_0} \left\{ \left[(a_0)u_2 + \left(\frac{a_1}{2}\right)u_2^2 + \left(\frac{a_2}{3}\right)u_2^3 + \left(\frac{a_3}{4}\right)u_2^4 + \left(\frac{a_4}{5}\right)u_2^5 + \left(\frac{a_5}{6}\right)u_2^6 \right] - \left[(a_0)u_1 + \left(\frac{a_1}{2}\right)u_1^2 + \left(\frac{a_2}{3}\right)u_1^3 + \left(\frac{a_3}{4}\right)u_1^4 + \left(\frac{a_4}{5}\right)u_1^5 + \left(\frac{a_5}{6}\right)u_1^6 \right] \right\}$$

Notice that several other terms involve the coordinate u_2 , which remains constant throughout the MLP calculations, so this can be calculated once and stored so it does not need to be calculated again. Also note that several polynomial terms (those in green and blue) are common to several terms and need only be calculated once each iteration. Then since there are only 3 polynomial terms in u_1 that are not common to other terms, there are only a total of 5 polynomial terms in u_1 that need to be calculated each iteration. However, notice that the coefficients on many of these terms, and indeed on many of the polynomial terms in general, involve only a small number of different ratios of the a_i coefficients. Therefore, not only do we define the a_i coefficients prior to compilation, but we also calculate the a_i/c_j terms prior to compilation since these remain constant through the entire MLP process, not just for a single history. For each a_i , there are only a handful of corresponding c_j that arise in these polynomial expansions (there are a total of 26 total values of a_i/c_j that appear in these expressions) and precalculating these and storing them as constant saves a considerable number of computations and time. Hence, there are 3 levels of computational simplification that can be exploited here: (1) the various a_i/c_j are calculated before compilation and stored as constant, (2) the terms involving u_0 and u_2 are calculated once per history, where the definition $u_0 = 0$ reduces this to just those terms involving u_2 , and (3) the polynomial terms in u_1 which appear in multiple expressions are calculated once per iteration (i.e. for each depth) and inserted into each expression where it appears. We then see that inserting the 5th order approximation of the $1/\beta^2(x)p^2(x)$ term into the integrals, performing the integration, manipulating the result algebraically to gather like terms, and simplifying the final expression has made it possible to identify unnecessary and redundant calculations and allowed us

to reduce the computational load significantly (by perhaps an order of magnitude or more). In addition, the term

$$\int_{u_0}^{u_1} \frac{(u_1 - u)^2}{\beta^2(u)p^2(u)} \frac{du}{X_0} = \frac{1}{X_0} \left[\left(\frac{a_0}{3}\right)u_1^3 + \left(\frac{a_1}{12}\right)u_1^4 + \left(\frac{a_2}{30}\right)u_1^5 + \left(\frac{a_3}{60}\right)u_1^6 + \left(\frac{a_4}{105}\right)u_1^7 + \left(\frac{a_5}{168}\right)u_1^8 \right]$$

$$\int_{u_0}^{u_1} \frac{u_1 - u}{\beta^2(u)p^2(u)} \frac{du}{X_0} = \frac{1}{X_0} \left[\left(\frac{a_0}{2}\right)u_1^2 + \left(\frac{a_1}{6}\right)u_1^3 + \left(\frac{a_2}{12}\right)u_1^4 + \left(\frac{a_3}{20}\right)u_1^5 + \left(\frac{a_4}{30}\right)u_1^6 + \left(\frac{a_5}{42}\right)u_1^7 \right]$$

$$\int_{u_0}^{u_1} \frac{1}{\beta^2(u)p^2(u)} \frac{du}{X_0} = \frac{1}{X_0} \left[(a_0)u_1 + \left(\frac{a_1}{2}\right)u_1^2 + \left(\frac{a_2}{3}\right)u_1^3 + \left(\frac{a_3}{4}\right)u_1^4 + \left(\frac{a_4}{5}\right)u_1^5 + \left(\frac{a_5}{6}\right)u_1^6 \right]$$

$$\int_{u_1}^{u_2} \frac{(u_2 - u)^2}{\beta^2(u)p^2(u)} \frac{du}{X_0} = \frac{1}{X_0} \left\{ \left[\left(\frac{a_0}{3}\right)u_2^3 + \left(\frac{a_1}{12}\right)u_2^4 + \left(\frac{a_2}{30}\right)u_2^5 + \left(\frac{a_3}{60}\right)u_2^6 + \left(\frac{a_4}{105}\right)u_2^7 + \left(\frac{a_5}{168}\right)u_2^8 \right] - u_2^2 \left[(a_0)u_1 + \left(\frac{a_1}{2}\right)u_1^2 + \left(\frac{a_2}{3}\right)u_1^3 + \left(\frac{a_3}{4}\right)u_1^4 + \left(\frac{a_4}{5}\right)u_1^5 + \left(\frac{a_5}{6}\right)u_1^6 \right] \right. \\ \left. + 2u_2 \left[\left(\frac{a_0}{2}\right)u_1^2 + \left(\frac{a_1}{3}\right)u_1^3 + \left(\frac{a_2}{4}\right)u_1^4 + \left(\frac{a_3}{5}\right)u_1^5 + \left(\frac{a_4}{6}\right)u_1^6 + \left(\frac{a_5}{7}\right)u_1^7 \right] - \left[\left(\frac{a_0}{3}\right)u_1^3 + \left(\frac{a_1}{4}\right)u_1^4 + \left(\frac{a_2}{5}\right)u_1^5 + \left(\frac{a_3}{6}\right)u_1^6 + \left(\frac{a_4}{7}\right)u_1^7 + \left(\frac{a_5}{8}\right)u_1^8 \right] \right\}$$

$$\int_{u_1}^{u_2} \frac{u_2 - u}{\beta^2(u)p^2(u)} \frac{du}{X_0} = \frac{1}{X_0} \left\{ \left[\left(\frac{a_0}{2}\right)u_2^2 + \left(\frac{a_1}{6}\right)u_2^3 + \left(\frac{a_2}{12}\right)u_2^4 + \left(\frac{a_3}{20}\right)u_2^5 + \left(\frac{a_4}{30}\right)u_2^6 + \left(\frac{a_5}{42}\right)u_2^7 \right] - u_2 \left[(a_0)u_1 + \left(\frac{a_1}{2}\right)u_1^2 + \left(\frac{a_2}{3}\right)u_1^3 + \left(\frac{a_3}{4}\right)u_1^4 + \left(\frac{a_4}{5}\right)u_1^5 + \left(\frac{a_5}{6}\right)u_1^6 \right] \right. \\ \left. + \left[\left(\frac{a_0}{2}\right)u_1^2 + \left(\frac{a_1}{3}\right)u_1^3 + \left(\frac{a_2}{4}\right)u_1^4 + \left(\frac{a_3}{5}\right)u_1^5 + \left(\frac{a_4}{6}\right)u_1^6 + \left(\frac{a_5}{7}\right)u_1^7 \right] \right\}$$

$$\int_{u_1}^{u_2} \frac{1}{\beta^2(u)p^2(u)} \frac{du}{X_0} = \frac{1}{X_0} \left\{ \left[(a_0)u_2 + \left(\frac{a_1}{2}\right)u_2^2 + \left(\frac{a_2}{3}\right)u_2^3 + \left(\frac{a_3}{4}\right)u_2^4 + \left(\frac{a_4}{5}\right)u_2^5 + \left(\frac{a_5}{6}\right)u_2^6 \right] - \left[(a_0)u_1 + \left(\frac{a_1}{2}\right)u_1^2 + \left(\frac{a_2}{3}\right)u_1^3 + \left(\frac{a_3}{4}\right)u_1^4 + \left(\frac{a_4}{5}\right)u_1^5 + \left(\frac{a_5}{6}\right)u_1^6 \right] \right\}$$

1 MLP Calculations

$$\begin{aligned}
y_{MLP} &= \begin{bmatrix} t_1 \\ \theta_1 \end{bmatrix} = (\Sigma_1^{-1} + R_1^T \Sigma_2^{-1} R_1)^{-1} (\Sigma_1^{-1} R_0 y_0 + R_1^T \Sigma_2^{-1} y_2) \\
R_0 y_0 &= \begin{bmatrix} 1 & u_1 - u_0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} t_0 \\ \theta_0 \end{bmatrix} = \begin{bmatrix} t_0 + (u_1 - u_0) \theta_0 \\ \theta_0 \end{bmatrix} \\
&= (\Sigma_1^{-1} + R_1^T \Sigma_2^{-1} R_1)^{-1} \\
&= \left(\begin{bmatrix} \sigma_{t_1} & \sigma_{t_1 \theta_1} \\ \sigma_{t_1 \theta_1} & \sigma_{\theta_1} \end{bmatrix}^{-1} + \begin{bmatrix} 1 & 0 \\ u_2 - u_1 & 1 \end{bmatrix} \begin{bmatrix} \sigma_{t_2} & \sigma_{t_2 \theta_2} \\ \sigma_{t_2 \theta_2} & \sigma_{\theta_2} \end{bmatrix}^{-1} \begin{bmatrix} 1 & u_2 - u_1 \\ 0 & 1 \end{bmatrix} \right)^{-1} \\
&= \left(\frac{1}{\sigma_{t_1} \sigma_{\theta_1} - (\sigma_{t_1 \theta_1})^2} \begin{bmatrix} \sigma_{\theta_1} & -\sigma_{t_1 \theta_1} \\ -\sigma_{t_1 \theta_1} & \sigma_{t_1} \end{bmatrix} + \frac{1}{\sigma_{t_2} \sigma_{\theta_2} - (\sigma_{t_2 \theta_2})^2} \begin{bmatrix} 1 & 0 \\ u_2 - u_1 & 1 \end{bmatrix} \begin{bmatrix} \sigma_{\theta_2} & -\sigma_{t_2 \theta_2} \\ -\sigma_{t_2 \theta_2} & \sigma_{t_2} \end{bmatrix} \begin{bmatrix} 1 & u_2 - u_1 \\ 0 & 1 \end{bmatrix} \right)^{-1} \\
&= \left(\begin{bmatrix} \sigma_{\theta_1}' & -\sigma_{t_1 \theta_1}' \\ -\sigma_{t_1 \theta_1}' & \sigma_{t_1}' \end{bmatrix} + \begin{bmatrix} 1 & 0 \\ u_2 - u_1 & 1 \end{bmatrix} \begin{bmatrix} \sigma_{\theta_2}' & -\sigma_{t_2 \theta_2}' \\ -\sigma_{t_2 \theta_2}' & \sigma_{t_2}' \end{bmatrix} \begin{bmatrix} 1 & u_2 - u_1 \\ 0 & 1 \end{bmatrix} \right)^{-1} \\
&= \left(\begin{bmatrix} \sigma_{\theta_1}' & -\sigma_{t_1 \theta_1}' \\ -\sigma_{t_1 \theta_1}' & \sigma_{t_1}' \end{bmatrix} + \begin{bmatrix} 1 & 0 \\ u_2 - u_1 & 1 \end{bmatrix} \begin{bmatrix} \sigma_{\theta_2}' & (u_2 - u_1) \sigma_{\theta_2}' - \sigma_{t_2 \theta_2}' \\ -\sigma_{t_2 \theta_2}' & \sigma_{t_2}' - (u_2 - u_1) \sigma_{t_2 \theta_2}' \end{bmatrix} \right)^{-1} \\
&= \left(\begin{bmatrix} \sigma_{\theta_1}' & -\sigma_{t_1 \theta_1}' \\ -\sigma_{t_1 \theta_1}' & \sigma_{t_1}' \end{bmatrix} + \begin{bmatrix} \sigma_{\theta_2}' & (u_2 - u_1) \sigma_{\theta_2}' - \sigma_{t_2 \theta_2}' \\ (u_2 - u_1) \sigma_{\theta_2}' - \sigma_{t_2 \theta_2}' & (u_2 - u_1)^2 \sigma_{\theta_2}' - 2(u_2 - u_1) \sigma_{t_2 \theta_2}' + \sigma_{t_2}' \end{bmatrix} \right)^{-1} \\
&= \begin{bmatrix} \sigma_{\theta_1}' + \sigma_{\theta_2}' & (u_2 - u_1) \sigma_{\theta_2}' - \sigma_{t_1 \theta_1}' - \sigma_{t_2 \theta_2}' \\ (u_2 - u_1) \sigma_{\theta_2}' - \sigma_{t_1 \theta_1}' - \sigma_{t_2 \theta_2}' & (u_2 - u_1)^2 \sigma_{\theta_2}' - 2(u_2 - u_1) \sigma_{t_2 \theta_2}' + \sigma_{t_1}' + \sigma_{t_2}' \end{bmatrix}^{-1} \\
&= \frac{\begin{bmatrix} (u_2 - u_1)^2 \sigma_{\theta_2}' - 2(u_2 - u_1) \sigma_{t_2 \theta_2}' + \sigma_{t_1}' + \sigma_{t_2}' & (u_1 - u_2) \sigma_{\theta_2}' + \sigma_{t_1 \theta_1}' + \sigma_{t_2 \theta_2}' \\ (u_1 - u_2) \sigma_{\theta_2}' + \sigma_{t_1 \theta_1}' + \sigma_{t_2 \theta_2}' & \sigma_{\theta_1}' + \sigma_{\theta_2}' \end{bmatrix}}{(\sigma_{\theta_1}' + \sigma_{\theta_2}') [(u_2 - u_1)^2 \sigma_{\theta_2}' - 2(u_2 - u_1) \sigma_{t_2 \theta_2}' + \sigma_{t_1}' + \sigma_{t_2}] - [(u_2 - u_1) \sigma_{\theta_2}' - \sigma_{t_1 \theta_1}' - \sigma_{t_2 \theta_2}']^2} \\
&= \frac{\begin{bmatrix} (u_2 - u_1)^2 \sigma_{\theta_2}' - 2(u_2 - u_1) \sigma_{t_2 \theta_2}' + \sigma_{t_1}' + \sigma_{t_2}' & (u_1 - u_2) \sigma_{\theta_2}' + \sigma_{t_1 \theta_1}' + \sigma_{t_2 \theta_2}' \\ (u_1 - u_2) \sigma_{\theta_2}' + \sigma_{t_1 \theta_1}' + \sigma_{t_2 \theta_2}' & \sigma_{\theta_1}' + \sigma_{\theta_2}' \end{bmatrix}}{\sigma_{\theta_1}' \sigma_{\theta_2}' (u_2 - u_1)^2 + 2(\sigma_{\theta_2}' \sigma_{t_1 \theta_1}' - \sigma_{\theta_1}' \sigma_{t_2 \theta_2}') (u_2 - u_1) - (\sigma_{t_1 \theta_1}' + \sigma_{t_2 \theta_2}')^2 + (\sigma_{\theta_1}' + \sigma_{\theta_2}') (\sigma_{t_1}' + \sigma_{t_2}')}
\end{aligned}$$

$$\begin{aligned}
& \left(\Sigma_1^{-1} R_0 y_0 + R_1^T \Sigma_2^{-1} y_2 \right) \\
&= \begin{bmatrix} \sigma_{t_1} & \sigma_{t_1 \theta_1} \\ \sigma_{t_1 \theta_1} & \sigma_{\theta_1} \end{bmatrix}^{-1} \begin{bmatrix} 1 & u_1 - u_0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} t_0 \\ \theta_0 \end{bmatrix} + \begin{bmatrix} 1 & 0 \\ u_2 - u_1 & 1 \end{bmatrix} \begin{bmatrix} \sigma_{t_2} & \sigma_{t_2 \theta_2} \\ \sigma_{t_2 \theta_2} & \sigma_{\theta_2} \end{bmatrix}^{-1} \begin{bmatrix} t_2 \\ \theta_2 \end{bmatrix} \\
&= \frac{1}{\sigma_{t_1} \sigma_{\theta_1} - \sigma_{t_1 \theta_1}^2} \begin{bmatrix} \sigma_{\theta_1} & -\sigma_{t_1 \theta_1} \\ -\sigma_{t_1 \theta_1} & \sigma_{t_1} \end{bmatrix} \begin{bmatrix} 1 & u_1 - u_0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} t_0 \\ \theta_0 \end{bmatrix} + \frac{1}{\sigma_{t_2} \sigma_{\theta_2} - \sigma_{t_2 \theta_2}^2} \begin{bmatrix} 1 & 0 \\ u_2 - u_1 & 1 \end{bmatrix} \begin{bmatrix} \sigma_{\theta_2} & -\sigma_{t_2 \theta_2} \\ -\sigma_{t_2 \theta_2} & \sigma_{t_2} \end{bmatrix} \begin{bmatrix} t_2 \\ \theta_2 \end{bmatrix} \\
&= \begin{bmatrix} \sigma_{\theta_1}' & -\sigma_{t_1 \theta_1}' \\ -\sigma_{t_1 \theta_1}' & \sigma_{t_1}' \end{bmatrix} \begin{bmatrix} 1 & u_1 - u_0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} t_0 \\ \theta_0 \end{bmatrix} + \begin{bmatrix} 1 & 0 \\ u_2 - u_1 & 1 \end{bmatrix} \begin{bmatrix} \sigma_{\theta_2}' & -\sigma_{t_2 \theta_2}' \\ -\sigma_{t_2 \theta_2}' & \sigma_{t_2}' \end{bmatrix} \begin{bmatrix} t_2 \\ \theta_2 \end{bmatrix} \\
&= \begin{bmatrix} \sigma_{\theta_1}' & -\sigma_{t_1 \theta_1}' \\ -\sigma_{t_1 \theta_1}' & \sigma_{t_1}' \end{bmatrix} \begin{bmatrix} t_0 + (u_1 - u_0) \theta_0 \\ \theta_0 \end{bmatrix} + \begin{bmatrix} 1 & 0 \\ u_2 - u_1 & 1 \end{bmatrix} \begin{bmatrix} \sigma_{\theta_2}' t_2 - \sigma_{t_2 \theta_2}' \theta_2 \\ \sigma_{t_2}' \theta_2 - \sigma_{t_2 \theta_2}' t_2 \end{bmatrix} \\
&= \begin{bmatrix} \sigma_{\theta_1}' [t_0 + (u_1 - u_0) \theta_0] - \sigma_{t_1 \theta_1}' \theta_0 \\ \sigma_{t_1}' \theta_0 - \sigma_{t_1 \theta_1}' [t_0 + (u_1 - u_0) \theta_0] \end{bmatrix} + \begin{bmatrix} \sigma_{\theta_2}' t_2 - \sigma_{t_2 \theta_2}' \theta_2 \\ (u_2 - u_1) (\sigma_{\theta_2}' t_2 - \sigma_{t_2 \theta_2}' \theta_2) + \sigma_{t_2}' \theta_2 - \sigma_{t_2 \theta_2}' t_2 \end{bmatrix}
\end{aligned}$$

$$\begin{aligned}
&= \begin{bmatrix} \sigma_{\theta_1}' [t_0 + (u_1 - u_0)\theta_0] - \sigma_{t_1\theta_1}' \theta_0 + \sigma_{\theta_2}' t_2 - \sigma_{t_2\theta_2}' \theta_2 \\ \sigma_{t_1}' \theta_0 - \sigma_{t_1\theta_1}' [t_0 + (u_1 - u_0)\theta_0] + (u_2 - u_1) (\sigma_{\theta_2}' t_2 - \sigma_{t_2\theta_2}' \theta_2) + \sigma_{t_2}' \theta_2 - \sigma_{t_2\theta_2}' t_2 \end{bmatrix} \\
&= \begin{bmatrix} \sigma_{\theta_1}' t_0 + \sigma_{\theta_1}' (u_1 - u_0)\theta_0 - \sigma_{t_1\theta_1}' \theta_0 + \sigma_{\theta_2}' t_2 - \sigma_{t_2\theta_2}' \theta_2 \\ \sigma_{t_1}' \theta_0 - \sigma_{t_1\theta_1}' [t_0 + (u_1 - u_0)\theta_0] + (u_2 - u_1) (\sigma_{\theta_2}' t_2 - \sigma_{t_2\theta_2}' \theta_2) + \sigma_{t_2}' \theta_2 - \sigma_{t_2\theta_2}' t_2 \end{bmatrix} \\
&= \begin{bmatrix} \sigma_{\theta_1}' t_0 + [\sigma_{\theta_1}' (u_1 - u_0) - \sigma_{t_1\theta_1}'] \theta_0 + \sigma_{\theta_2}' t_2 - \sigma_{t_2\theta_2}' \theta_2 \\ \sigma_{t_1}' \theta_0 - \sigma_{t_1\theta_1}' [t_0 + (u_1 - u_0)\theta_0] + (u_2 - u_1) (\sigma_{\theta_2}' t_2 - \sigma_{t_2\theta_2}' \theta_2) + \sigma_{t_2}' \theta_2 - \sigma_{t_2\theta_2}' t_2 \end{bmatrix}
\end{aligned}$$

Now we need to matrix multiply the 2×2 matrix from the left hand term by the 2×1 vector from the right hand term:

$$\begin{aligned}
& (\Sigma_1^{-1} + R_1^T \Sigma_2^{-1} R_1)^{-1} \\
&= \begin{pmatrix} \Sigma_1^{-1}[0] & \Sigma_1^{-1}[1] \\ \Sigma_1^{-1}[2] & \Sigma_1^{-1}[3] \end{pmatrix} + \begin{pmatrix} R_1^T[0] & R_1^T[1] \\ R_1^T[2] & R_1^T[3] \end{pmatrix} \begin{pmatrix} \Sigma_2^{-1}[0] & \Sigma_2^{-1}[1] \\ \Sigma_2^{-1}[2] & \Sigma_2^{-1}[3] \end{pmatrix} \begin{pmatrix} R_1[0] & R_1[1] \\ R_1[2] & R_1[3] \end{pmatrix} \\
&= \begin{pmatrix} \Sigma_1^{-1}[0] & \Sigma_1^{-1}[1] \\ \Sigma_1^{-1}[2] & \Sigma_1^{-1}[3] \end{pmatrix} + \begin{pmatrix} R_1^T[0] & R_1^T[1] \\ R_1^T[2] & R_1^T[3] \end{pmatrix} \begin{pmatrix} \Sigma_2^{-1}[0]R_1[0] + \Sigma_2^{-1}[1]R_1[2] & \Sigma_2^{-1}[0]R_1[1] + \Sigma_2^{-1}[1]R_1[3] \\ \Sigma_2^{-1}[2]R_1[0] + \Sigma_2^{-1}[3]R_1[2] & \Sigma_2^{-1}[2]R_1[1] + \Sigma_2^{-1}[3]R_1[3] \end{pmatrix} \\
&= \left\{ \begin{pmatrix} \Sigma_1^{-1}[0] & \Sigma_1^{-1}[1] \\ \Sigma_1^{-1}[2] & \Sigma_1^{-1}[3] \end{pmatrix} + \begin{pmatrix} R_1^T[0] & R_1^T[1] \\ R_1^T[2] & R_1^T[3] \end{pmatrix} \begin{pmatrix} \Sigma_2^{-1}[0]R_1[0] + \Sigma_2^{-1}[1]R_1[2] & \Sigma_2^{-1}[0]R_1[1] + \Sigma_2^{-1}[1]R_1[3] \\ \Sigma_2^{-1}[2]R_1[0] + \Sigma_2^{-1}[3]R_1[2] & \Sigma_2^{-1}[2]R_1[1] + \Sigma_2^{-1}[3]R_1[3] \end{pmatrix} \right\}^{-1} \\
&= \left\{ \begin{pmatrix} \Sigma_1^{-1}[0] + R_1^T[0](\Sigma_2^{-1}[0]R_1[0] + \Sigma_2^{-1}[1]R_1[2]) + R_1^T[1](\Sigma_2^{-1}[2]R_1[0] + \Sigma_2^{-1}[3]R_1[2]) & \Sigma_1^{-1}[1] + R_1^T[0](\Sigma_2^{-1}[0]R_1[1] + \Sigma_2^{-1}[1]R_1[3]) + R_1^T[1](\Sigma_2^{-1}[2]R_1[1] + \Sigma_2^{-1}[3]R_1[3]) \\ \Sigma_1^{-1}[2] + R_1^T[2](\Sigma_2^{-1}[0]R_1[0] + \Sigma_2^{-1}[1]R_1[2]) + R_1^T[3](\Sigma_2^{-1}[2]R_1[0] + \Sigma_2^{-1}[3]R_1[2]) & \Sigma_1^{-1}[3] + R_1^T[2](\Sigma_2^{-1}[0]R_1[1] + \Sigma_2^{-1}[1]R_1[3]) + R_1^T[3](\Sigma_2^{-1}[2]R_1[1] + \Sigma_2^{-1}[3]R_1[3]) \end{pmatrix} \right\}^{-1} \\
&= (\Sigma_1^{-1} R_0 y_0 + R_1^T \Sigma_2^{-1} y_2) \\
&= \begin{pmatrix} \Sigma_1^{-1}[0] & \Sigma_1^{-1}[1] \\ \Sigma_1^{-1}[2] & \Sigma_1^{-1}[3] \end{pmatrix} \begin{pmatrix} R_0[0] & R_0[1] \\ R_0[2] & R_0[3] \end{pmatrix} \begin{pmatrix} y_0[0] \\ y_0[1] \end{pmatrix} + \begin{pmatrix} R_1^T[0] & R_1^T[1] \\ R_1^T[2] & R_1^T[3] \end{pmatrix} \begin{pmatrix} \Sigma_2^{-1}[0] & \Sigma_2^{-1}[1] \\ \Sigma_2^{-1}[2] & \Sigma_2^{-1}[3] \end{pmatrix} \begin{pmatrix} y_2[0] \\ y_2[1] \end{pmatrix} \\
&= \begin{pmatrix} \Sigma_1^{-1}[0] & \Sigma_1^{-1}[1] \\ \Sigma_1^{-1}[2] & \Sigma_1^{-1}[3] \end{pmatrix} \begin{pmatrix} R_0[0]y_0[0] + R_0[1]y_0[1] \\ R_0[2]y_0[0] + R_0[3]y_0[1] \end{pmatrix} + \begin{pmatrix} R_1^T[0] & R_1^T[1] \\ R_1^T[2] & R_1^T[3] \end{pmatrix} \begin{pmatrix} \Sigma_2^{-1}[0]y_2[0] + \Sigma_2^{-1}[1]y_2[1] \\ \Sigma_2^{-1}[2]y_2[0] + \Sigma_2^{-1}[3]y_2[1] \end{pmatrix} \\
&= \begin{pmatrix} \Sigma_1^{-1}[0] (R_0[0]y_0[0] + R_0[1]y_0[1]) + \Sigma_1^{-1}[1] (R_0[2]y_0[0] + R_0[3]y_0[1]) \\ \Sigma_1^{-1}[2] (R_0[0]y_0[0] + R_0[1]y_0[1]) + \Sigma_1^{-1}[3] (R_0[2]y_0[0] + R_0[3]y_0[1]) \end{pmatrix} + \begin{pmatrix} R_1^T[0] (\Sigma_2^{-1}[0]y_2[0] + \Sigma_2^{-1}[1]y_2[1]) + R_1^T[1] (\Sigma_2^{-1}[2]y_2[0] + \Sigma_2^{-1}[3]y_2[1]) \\ R_1^T[2] (\Sigma_2^{-1}[0]y_2[0] + \Sigma_2^{-1}[1]y_2[1]) + R_1^T[3] (\Sigma_2^{-1}[2]y_2[0] + \Sigma_2^{-1}[3]y_2[1]) \end{pmatrix} \\
&= \begin{pmatrix} \Sigma_1^{-1}[0] (R_0[0]y_0[0] + R_0[1]y_0[1]) + \Sigma_1^{-1}[1] (R_0[2]y_0[0] + R_0[3]y_0[1]) \\ \Sigma_1^{-1}[2] (R_0[0]y_0[0] + R_0[1]y_0[1]) + \Sigma_1^{-1}[3] (R_0[2]y_0[0] + R_0[3]y_0[1]) \end{pmatrix} + \begin{pmatrix} R_1^T[0] (\Sigma_2^{-1}[0]y_2[0] + \Sigma_2^{-1}[1]y_2[1]) + R_1^T[1] (\Sigma_2^{-1}[2]y_2[0] + \Sigma_2^{-1}[3]y_2[1]) \\ R_1^T[2] (\Sigma_2^{-1}[0]y_2[0] + \Sigma_2^{-1}[1]y_2[1]) + R_1^T[3] (\Sigma_2^{-1}[2]y_2[0] + \Sigma_2^{-1}[3]y_2[1]) \end{pmatrix} \\
&= \begin{pmatrix} \Sigma_1^{-1}[0] (R_0[0]y_0[0] + R_0[1]y_0[1]) + \Sigma_1^{-1}[1] (R_0[2]y_0[0] + R_0[3]y_0[1]) + R_1^T[0] (\Sigma_2^{-1}[0]y_2[0] + \Sigma_2^{-1}[1]y_2[1]) + R_1^T[1] (\Sigma_2^{-1}[2]y_2[0] + \Sigma_2^{-1}[3]y_2[1]) \\ \Sigma_1^{-1}[2] (R_0[0]y_0[0] + R_0[1]y_0[1]) + \Sigma_1^{-1}[3] (R_0[2]y_0[0] + R_0[3]y_0[1]) + R_1^T[2] (\Sigma_2^{-1}[0]y_2[0] + \Sigma_2^{-1}[1]y_2[1]) + R_1^T[3] (\Sigma_2^{-1}[2]y_2[0] + \Sigma_2^{-1}[3]y_2[1]) \end{pmatrix}
\end{aligned}$$