

Description

Background

Carbapenem-resistant *Acinetobacter baumannii* (CRAB), globally recognized as a “superbug”, has emerged as a leading cause of life-threatening nosocomial infections. Possessing extraordinary environmental adaptability, CRAB can survive on dry surfaces for weeks and resist most common disinfectants, making it highly prone to nosocomial transmission in ICUs and high-risk areas^[1]. It invades hosts via damaged physical barriers or medical devices, requiring only a low inoculum ($10^3\text{--}10^5$ CFU) for stable colonization. Its infection spectrum covers pulmonary infections (VAP, 30–40%), bloodstream infections (CRBSI, 25–35%), and wound/soft tissue infections (15–20%), accompanied by staggering mortality rates—up to 50–70% for bloodstream infections and 70–80% for meningitis^[2].

The clinical threat stems from the synergy of multidrug resistance mechanisms (e.g., OXA-type carbapenemase production, outer membrane porin loss) and biofilm formation, which increases bacterial drug resistance by 10–1000 times. Current antibiotic therapies are plagued by high costs (15,000–20,000 RMB per course), severe adverse effects (e.g., nephrotoxicity), and an inability to specifically target biofilms, leading to frequent recurrence.

While targeting the Quorum Sensing (QS) system is a promising strategy, existing methods face critical challenges in targeting efficiency, signal amplification, and therapeutic monitoring, limiting their clinical translation.

Brief Introduction

Diagnosis Module

We utilize a colloidal gold immunochromatography test strip combined with a visible light analysis device. The strip uses glass fiber pads to filter impurities^[3], and the nitrocellulose membrane provides strong dipole interactions for protein immobilization^[4]. By analyzing signal molecule ratios through grayscale

algorithms^[5], the system enables rapid CRAB screening and distinguishes between “colonization” and “infection” states.

Delivery Module

The core circuit implements a “sensing-amplification-execution” architecture to detect and disarm CRAB biofilms^{[6][7]}, specifically targeting 3-OH-C12-HSL^[8], the primary quorum-sensing^[9] signal, and utilizing PvdQ enzyme for efficient long-chain AHL degradation^{[10][11]}. The system incorporates engineered chemotaxis receptors for active targeting and a positive feedback loop for sustained efficacy. Tailored strategies include: pH-sensitive liposomes encapsulating PvdQ for site-specific release in acidic pulmonary foci^{[12][13]}; engineered OMVs with lipid raft structures^[14] and dephosphorylated protamine targeting^[15] for bloodstream delivery^[16]; and a thermosensitive double-layer hydrogel system for wounds. The hydrogel utilizes Pluronic F127, QCS, and CMC for thermosensitive gelling and antimicrobial action^{[17][18][19][20][21][22][23]}, with core-shell micro-carriers^{[24][25][26]} and a dual-chamber prefilled syringe^[27] ensuring stability and biosafety.

Visualization Module

Real-time efficacy monitoring is achieved through specific mechanisms: a genetic memory switch triggers BpsA to produce blue pigment in wound dressings^[30]; an enzyme-probe competition mechanism generates water-soluble indigo carmine to indicate pulmonary therapeutic endpoints^[29]; and PEG-shielded prodrug probes release red signals^[30] upon specific protease CpaA cleavage^[31] in bloodstream infections, excreted via urine.

Kill Switch

A dual biocontainment system using the MazEF toxin-antitoxin module ensures safety. It features an arabinose-dependent “timed elimination” mechanism and a doxycycline-inducible “manual emergency kill” switch to guarantee complete clearance of the engineered bacteria after treatment.

Reference

- [1] Congcong Liu,Kaichao Chen,Yuchen Wu,Ling Huang,Yinfei Fang,Jiayue Lu. Epidemiological and genetic characteristics of clinical carbapenem-resistant

Acinetobacter baumannii strains collected countrywide from hospital intensive care units (ICUs) in China[J].Emerging Microbes & Infections , 17 Feb 2022 , <https://doi.org/10.1080/22221751.2022.2093134>

- [2] Carmi Bartal, Kenneth V. I. Rolston & Lior Nesher. Carbapenem-resistant Acinetobacter baumannii: Colonization, Infection and Current Treatment Options. Israel Infect Dis Ther (2022) 11:683–694 , <https://doi.org/10.1007/s40121-022-00597-w>
- [3] Amadeo Sena-Torralba, Ruslán Álvarez-Diduk, Claudio Parolo, Andrew Pipe, A. Merkoçi. Toward Next Generation Lateral Flow Assays: Integration of Nanomaterials [J]. Chemical Reviews, 6 September 2022, 14881–14905 DOI: 10.1021/acs.chemrev.1c01012
- [4] Rishabh Gandotra (a), Feng-Chih Kuo (c), Mel S. Lee (d), Gwo-Bin Lee (a) (b). A paper-based aptamer-sandwich assay for detection of HNP 1 as a biomarker for periprosthetic joint infections on an integrated microfluidic platform [J]. Analytica Chimica Acta, 15 November 2023, 341879
- [5] Junqi Shen, Zhengyi Cai, Cheng Zhang, Xinyue Feng, Chenzhi Zhang, Huan Zhao, Chuanlin Yin, Bo Wang, Xiaoping Yu, Biao Zhang. Dual-Mode Quantitative Immunochromatographic Assay for Highly Sensitive On-Site Detection of Ciprofloxacin in Fish Products [J]. Foods, 2025 Mar 25;14(7):1132. DOI: 10.3390/foods14071132.
- [6] Rumbo-Feal S, Gómez MJ, Gayoso C, Álvarez-Fraga L, Cabral MP, Aransay AM, Rodríguez-Ezpeleta N, Fullaondo A, Valle J, Tomás M, Bou G, Poza M. Whole transcriptome analysis of *Acinetobacter baumannii* assessed by RNA-sequencing reveals different mRNA expression profiles in biofilm compared to planktonic cells. PLoS One. 2013 Aug 30;8(8):e72968. doi: 10.1371/journal.pone.0072968. PMID: 24023660; PMCID: PMC3758355.
- [7] Upmanyu K, Haq QMR, Singh R. Factors mediating *Acinetobacter baumannii* biofilm formation: Opportunities for developing therapeutics. Curr Res Microbiol Sci. 2022 Mar 28;3:100131. doi: 10.1016/j.crmicr.2022.100131. PMID: 35909621; PMCID: PMC9325880.
- [8] Cui B, Peng G, Wang LE, Deng Y. Signaling in *Acinetobacter baumannii*: Quorum sensing and nucleotide second messengers. Comput Struct Biotechnol J. 2025 May 22;27:2168–2175. doi: 10.1016/j.csbj.2025.05.032. PMID: 40510767; PMCID: PMC12162036.
- [9] Zhong S, He S. Quorum Sensing Inhibition or Quenching in *Acinetobacter baumannii*: The Novel Therapeutic Strategies for New Drug Development. Front Microbiol. 2021 Feb 1;12:558003. doi: 10.3389/fmicb.2021.558003. PMID: 33597937; PMCID: PMC7882596.
- [10] Bokhove M, Nadal Jimenez P, Quax WJ, Dijkstra BW. The quorum-quenching N-acyl homoserine lactone acylase PvdQ is an Ntn-hydrolase with an unusual substrate-binding pocket. Proc Natl Acad Sci U S A. 2010 Jan 12;107(2):731–736. doi: 10.1073/pnas.0909320107. PMID: 20078900; PMCID: PMC2818030.

07(2):686-91. doi: 10.1073/pnas.0911839107. Epub 2009 Dec 22. PMID: 20080736; PMCID: PMC2818923.

- [11] Vogel J, Jansen L, Setroikromo R, Cavallo FM, van Dijl JM, Quax WJ. Fighting *Acinetobacter baumannii* infections with the acylase PvdQ. *Microbes Infect.* 2022 Jun;24(4):104951. doi: 10.1016/j.micinf.2022.104951. Epub 2022 Feb 10. PMID: 35151875.
- [12] Tsirogianni AG, Chountoulesi M, Pippa N. Recent Advances in pH-Responsive Liposomes: Lessons Learnt and New Directions in Nanomedicine Development. *Materials (Basel)*. 2025 Sep 13;18(18):4295. doi: 10.3390/ma18184295. PMID: 41010139; PMCID: PMC12471921.
- [13] Mahmoudzadeh M, Magarkar A, Koivuniemi A, Rög T, Bunker A. Mechanistic Insight into How PEGylation Reduces the Efficacy of pH-Sensitive Liposomes from Molecular Dynamics Simulations. *Mol Pharm.* 2021 Jul 5;18(7):2612-2621. doi: 10.1021/acs.molpharmaceut.1c00122. Epub 2021 Jun 6. PMID: 34096310; PMCID: PMC8289284.
- [14] Li J, Yu J, Song J, Zhang Y, Li N, Wang Z, Qin M, Zhao M, Zhang B, Huang R, Zhou S, Liu Y, He Z, Liu H, Dan Liu, Wang Y. Galloylated liposomes enable targeted drug delivery by overcoming protein corona shielding. *Nat Commun.* 2025 Aug 25;16(1):7926. doi: 10.1038/s41467-025-63198-4. PMID: 40854890; PMCID: PMC12378351.
- [15] Chhetri KB, Jang YH, Lansac Y, Maiti PK. Effect of phosphorylation of protamine-like cationic peptide on the binding affinity to DNA. *Biophys J.* 2022 Dec 20;121(24):4830-4839. doi: 10.1016/j.bpj.2022.09.025. Epub 2022 Sep 26. PMID: 36168289; PMCID: PMC9808561.
- [16] Peulen TO, Wilkinson KJ. Diffusion of nanoparticles in a biofilm. *Environ Sci Technol.* 2011 Apr 15;45(8):3367-73. doi: 10.1021/es103450g. Epub 2011 Mar 24. PMID: 21434601.
- [17] Qu J, Zhao X, Liang Y, Zhang T, Ma PX, Guo B. Antibacterial adhesive injectable hydrogels with rapid self-healing, extensibility and compressibility as wound dressing for joints skin wound healing. *Biomaterials.* 2018 Nov;183:185-199. doi: 10.1016/j.biomaterials.2018.08.044. Epub 2018 Aug 24. PMID: 30172244.
- [18] Li S, Yang C, Li J, et al. Progress in Pluronic F127 Derivatives for Application in Wound Healing and Repair. *Int J Nanomedicine.* 2023;18:4485-4505. Published 2023 Aug 7. doi:10.2147/IJN.S418534
- [19] Pelegrino MT, De Araujo Lima B, Do Nascimento MHM, Lombello CB, Brochi M, Seabra AB. Biocompatible and Antibacterial Nitric Oxide-Releasing Pluronic F-127/Chitosan Hydrogel for Topical Applications. *Polymers (Basel)*. 2018 Apr 18;10(4):452. doi: 10.3390/polym10040452. PMID: 30966487; PMCID: PMC6415216.
- [20] Li S, Yang C, Li J, Zhang C, Zhu L, Song Y, Guo Y, Wang R, Gan D, Shi J,

- Ma P, Gao F, Su H. Progress in Pluronic F127 Derivatives for Application in Wound Healing and Repair. *Int J Nanomedicine*. 2023 Aug 7;18:4485-4505. doi: 10.2147/IJN.S418534. PMID: 37576462; PMCID: PMC10416793.
- [21] Kaiting Cheng, Yiming Fang, Liang Bai, Feng Gui, Junchi Ma, Huimin Gao, Yadong Zhao, Xingtao Xu, Quaternized chitosan / Pluronic F127 thermosensitive hydrogel with high antibacterial properties for wound dressing, *Progress in Natural Science: Materials International*, Volume 33, Issue 5, 2023, Pages 581-592, ISSN 1002-0071, <https://doi.org/10.1016/j.pnsc.2023.11.002>.
- [22] Huang A, Chen Y, Wu C. Wound Dressing Double-Crosslinked Quick Self-Healing Hydrogel Based on Carboxymethyl Chitosan and Modified Nanocellulose. *Polymers (Basel)*. 2023 Aug 13;15(16):3389. doi: 10.3390/polym15163389. PMID: 37631446; PMCID: PMC10459649.
- [23] Alven S, Aderibigbe BA. Chitosan and Cellulose-Based Hydrogels for Wound Management. *Int J Mol Sci*. 2020 Dec 18;21(24):9656. doi: 10.3390/ijms21249656. PMID: 33352826; PMCID: PMC7767230.
- [24] Łętocha, A., et al., Effect of Encapsulation of *Lactobacillus casei* in Alginat e-Tapioca Flour Microspheres Coated with Different Biopolymers on the Viability of Probiotic Bacteria. *ACS Applied Materials & Interfaces*, 2024. 16(39): p. 52878–52893.
- [25] Allen ME, Kamilova E, Monck C, et al. Engineered Bacteria as Living Biosensors in Dermal Tattoos. *Adv Sci (Weinh)*. 2024;11(30):e2309509. doi:10.1002/advs.202309509
- [26] Li S, Ji Y, Xue X, et al. Hierarchical biofilm models using sodium alginate beads containing bacteria embedded in a cellulose-chitosan hydrogel matrix. *J Mater Chem B*. 2025;13(12):3952-3958. Published 2025 Mar 20. doi:10.1039/d4tb02015d
- [27] Anupam Chanda. Double chamber Devices and their advantages[J]. *Journal of Addictive Disorders and Mental Health*, 2022, 2(2): 1-9.
- [28] Brown AS, Calcott MJ, Collins VM, Owen JG, Ackerley DF. The indigoidine synthetase BpsA provides a colorimetric ATP assay that can be adapted to quantify the substrate preferences of other NRPS enzymes. *Biotechnol Lett*. 2020 Dec;42(12):2665-2671. doi: 10.1007/s10529-020-02972-4 IF: 2.1 Q3 . Epub 2020 Jul 17. PMID: 32681380.
- [29] Li, Z.; Lu, R.; Zhang, J.; Zhu, Y.; Mu, W. Recent Advances on Natural Pigment Indigoidine: Occurrence, Biosynthetic Approaches, and Applications. *J. Agric. Food Chem.* 2025, 73 (30), 18545-18560. DOI: 10.1021/acs.jafc.5c05724.
- [30] O'Leary MK, Chen SS, Westblade LF, Alabi CA. Design of a PEGylated Antimicrobial Prodrug with Species-Specific Activation. *Biomacromolecules*. 2021 Feb 8;22(2):984-992. doi: 10.1021/acs.biomac.0c01695 . Epub 2021 Jan 11. PMID: 33428376; PMCID: PMC8270352.

[31] Haurat MF, Scott NE, Di Venanzio G, Lopez J, Pluvinage B, Boraston AB, Ferracane MJ, Feldman MF. 2020. The Glycoprotease CpaA Secreted by Medically Relevant Acinetobacter Species Targets Multiple O-Linked Host Glycoproteins. *mBio* 11:10.1128/mbio.02033-20. <https://doi.org/10.1128/mbio.02033-20>