

A Negative Fire Feedback in a Transitional Forest of Southeastern Amazonia



Jennifer Balch, Yale University

Dan Nepstad, Woods Hole Research Center

Paulo Brando, University of Florida

Lisa Curran, Yale University

Osvaldo Portela, Instituto de Pesquisa Ambiental da Amazonia

Oswaldo de Carvalho Jr., Kent University

Paul LeFebvre, Woods Hole Research Center

Predominant paradigms...

Fire begets more fire in Amazon forests.

&

Moisture dynamics control fire behavior.

Objectives & Hypotheses

- Compare fuel loads & moisture, microclimate, and fire behavior in an annually burned plot after 1, 2, and 3 prescribed burns

Expect an **increase in fuels** and **drier understory** leads to increased burned area and fire intensity

Objectives & Hypotheses

- After two burns test the effects of fuel additions, in canopy gaps and non-gaps, on fire intensity, fire spread rate, and burn extent

Expect **microclimate** rather than fuels controlling fire behavior



Transitional forest ~400,000 km²

Rainfall: 1740 mm yr⁻¹

Severe 5-mo dry season:

(< 10 mm 3 mo, < 50 mm 2 mo)

Soils: Oxisols (Haplustox)

Leaf litter production: 4.3 Mg ha⁻¹yr⁻¹


Biomass: 191 ± 6 Mg ha⁻¹

Canopy height: 20 ± 1 m

LAI max: ~ 5 m² m⁻²

Lower diversity: ~100 tree species

Dominance: Lauraceae






Experimental Design:

- Two fire regimes
- 1st 100-ha burn Aug. 2004
- 2nd 50-ha burn Sept. 2005
- 3rd 50-ha burn Aug/Sept. 2006
- 4th 100-ha burn Aug. 2007



Fazenda Tanguro: Plano de Amostragem

1500 m




Inventários:


Area Total	> 40cm DAP
	> 20cm DAP
	> 10cm DAP

Sub-Parcelas:




Forest flammability:






1st Burn - 2004




80% burned

2nd Burn - 2005




80% burned

3rd Burn - 2006



40% burned!

Flame Height (cm)



1st, 2nd, and 3rd burn

CARBON CONSEQUENCES:




Dead biomass combustion (\pm SE)

44.5 (\pm 21.2) Mg ha⁻¹

23.2 (\pm 9.1) Mg ha⁻¹

15.3 (\pm 9.3) Mg ha⁻¹


(during 1, 2, and 3
consecutive annual burns)



“Shouldn’t you rake those up first and *then* set them on fire?”



The New Yorker



3 tons of leaves later....

Fire behavior properties (s.e.)

	Fuel added	No fuel added
	$n = 18$	$n = 18$
Proportion of area burned	0.85 (7)	0.16 (6)
Fire spread rate (m/min)	0.18 (0.01)	0.10 (0.01)
Flame height (cm)	42 (4)	23 (4)
Flame width (cm)	27 (3)	16 (2)

Conclusions...

- Increasing fire frequency causes a negative feedback on fire in transitional forests - underscores the importance of considering the **rates of fuel inputs/outputs**
- **Fine fuels** appear to be controlling fire behavior in transitional forests - shift away from importance of microclimate & moisture witnessed in more humid forests

2007 Burn Season

















Packard



Heinz
Foundation



thanks...