



# ***Spatial variability of the deforestation along the Amazon River main stem floodplain***

*Evlyn Novo, INPE*

*Adriana G. Affonso, INPE*

*Laura L. Hess, UCSB*

*John M. Melack, UCSB*






## ***Why focus in land use changes in the wetland?***

- *wetland represents 20 % of the Amazon region (Melack et al. 2004).*
- *it may affect the role of wetlands and rivers in carbon transport and release by changing the carbon pools*
- *it may affect water quality and biodiversity not only in the wetland*

# *Amazon Basin Wetland*





***Covers around  
20 % of the entire  
basin (Melack et al. 2004)***

***Source: Hess et al. 2006***

- ***Is a globally significant source of atmospheric methane (Melack et al. 2004)***
- ***Length of inundation controls vegetation composition (Piedade et al. 2001)***

***Vegetative cover varies as a function of hydrologic regime, solute and sediment content of waters (Klinge et al. 1990; Junk, 1997)***


## ***Growing human pressure on the Amazon River floodplain – the richest varzea type wetland in the Amazon.***



Manaus-Boa Vista



Belém-Brasília

Cuiabá-Santarém





## *Increased presence of humans reflected in the increase in cattle raising*



Source: IBGE, 2000






# Background

- *It is well documented that the floodplain is being intensively used for cattle in both Pará and Amazonas states,*
- *It is still missing information on the spatial variability of this occupation.*
- *The impact of cattle ranching on floodplain ecology may increase as the proportion of wetland within a given region increases.*
- *There are striking differences in the baseline algae primary productivity expressed as chlorophyll concentration from measurements reviewed in literature (Maleck and Forsberg, 2001) and measurements made in Curuai Lake (Barbosa, 2005) and extrapolated to the Parintins- Almeirim reaches (Novo et al. 2006).*


## ***How to tell differences between natural and human induced patterns in the land cover?***




Water



Bare or herbaceous



Forest



Shrub



# What is the question?

- *Are those differences in primary productivity natural?*
- *Are they connected to cattle ranching in the floodplain?*






# Methodology

- *In order to answer those questions the following data were quantified:*
  - *proportion of wetland for each municipality in the Amazonas (37) and Para (18) States using the non-validated Amazon wetland mask;*
  - *the deforested area for each municipality using the digital deforestation data base provided by PRODES project;*
  - *the deforested area within a 2 km buffer from the Amazon River bank, assuming that the floodplain is mainly used for owners living nearby the river banks.*
  - *Herd size (number of heads) per municipality (IBGE, 2002);*
  - *Population (inhabitants) per municipality (2004);*
  - *Distance from the market centers (Manaus and Belém);*
  - *Eutrophic Lake Area (index derived from MODIS images - 2002)*




# Divisão de Sensoriamento Remoto

## ***STUDY AREA***




## Steps to obtain the data


**1- Overlay the wetland mask on the municipality map**




**2- build a 2 km buffer-wetland mask**




**3 -build a mosaic of 60 digital deforestation maps**




**4- build the land cover/deforestation map for the 55 municipalities**



## *wetland deforestation*



## *2 buffer-wetland deforestation*



***2km Buffer deforestation: deforestation occurring in Terra Firme at the edge of the wetland***



## ***Summary – Pará State***

- ***18 municipalities occupying wetland in the Amazon River main stem***
- ***Total Wetland Area = 43,381 km<sup>2</sup>***
- ***Total Deforestation in the wetland= 2,773km<sup>2</sup>***
- ***Percentage of deforestation=6.4 %***
- ***Total 2km buffer deforestation= 8,410 km<sup>2</sup>***
- ***Percentage of buffer deforestation=7.2 %***
- ***Total herd size = 894,369***
- ***Total Population = 6,189,550***







### ***Summary – Amazonas State***


- ***37 municipalities occupying wetland in the Amazon River main stem***
- ***Total Wetland Area = 107,248***
- ***Total Deforestation in the wetland= 5,012***
- ***Percentage of deforestation= 4.6 %***
- ***Total 2km buffer deforestation= 6,471***
- ***Percentage of buffer deforestation= 4.0 %***
- ***Total herd size = 590,262***
- ***Total Population = 2,582,737***
- ***Manaus Population= 1,592,555***
- ***70 % of the Population in the Amazon state is in those 37 municipalities***




# Divisão de Sensoriamento Remoto




















## ***55 municipalities along the Amazon River main stem floodplain***












- *What are the consequences of deforestation, cattle ranching, increased human pressure on:*
- *algae primary productivity – chl concentration in lake*
- *Water quality – lake eutrophication resulting from huge inputs of nutrients*





# Divisão de Sensoriamento Remoto






# Divisão de Sensoriamento Remoto












# Divisão de Sensoriamento Remoto











## *Final considerations*

- *Deforestation area in the Amazon River floodplain presents distinct patterns.*
- *Pará state- the deforestation is clearly underestimated because PRODES started after considerable deforestation had took place.*
- *Amazonia state - deforestation is more closely connected to the presence of cattle ranching in the floodplain.*
- *In spite of the limited samples on chlorophyll concentration, the results suggests a clear connection between cattle ranching and the occurrence of large areas of eutrophic water*