


Human exposure to Mercury as a consequence of deforestation by slash and burn agriculture in the Amazon

- Marc Lucotte, Robert Davidson, Nicolina Farella & Nicolas Mainville
- Institut des sciences de l'environnement
- Université du Québec à Montréal, Canada
- LBA-ECO 9th science meeting, Sao Paulo, Nov. 11, 2005

Human exposure to Mercury as a consequence of deforestation by slash and burn agriculture in the Amazon

- Marc Lucotte, Robert Davidson, Nicolina Farella & Nicolas Mainville
- Institut des sciences de l'environnement
- Université du Québec à Montréal, Canada
- **Projects mainly funded by IDRC, NSERC and CNPq**




Preconceived ideas:

- Presence of mercury attributable to small scale gold mines
 - Communities exposed to Hg mostly in regions of gold mines

No decreasing gradient in Hg Conc. in SPM along the Tapajós River away from gold mines

Hg, ng/g d.w.




Area nearest to gold mines


Area furthest to gold mines

(Roulet et al., 1998)


Recent Hg pollution (less than 40 years) as evidenced in lake sediments records



Half carnivorous fish have Hg concentrations over the recommended limit of 0.5 ppm



Relationship between fish consumption and human exposure to Hg as seen by Hg levels in hair




(Passos et al., 2003)

Now, two questions:

- Do the observed Hg levels affect the health of local communities?
- If this Hg does not come from goldmining, what is its origin?

Donna Mergler's famous diagram on Human Health & Mercury



Dexterity

Vision



Other health problems related to chronic exposure to low levels of Hg:

- Cardiovascular problems

(Filion et al., 2005)

- Auto-immune system problems


(Silbergeld et al. 2005)

...


Two questions:

- Do the observed Hg levels affect the health of local communities?
- If this Hg does not come from goldmining, what is its origin?


High Hg Concentrations in river bank soils: natural origin (Roulet et al., 1999)




Rapid progression of colonization




Important loss of Hg in soils recently (6 years)
deforested for agriculture (Roulet et al., 1998)



PCA Clay Soils at 25 cm




PCA Clay Soils at 50 cm



(Farella, 2005)

PCA Sandy-clay Soils at 25 cm



Any solutions to the problem?

- Short term: limit exposure
- Long term: limit deforestation
of sensitive areas


COMUNIDADE DE SÃO LUIS DO TAPAJÓS, SECA 2004

Para diminuir o mercúrio que está no seu corpo,
coma de preferência os peixes com pouco mercúrio.


A COR VERMELHA INDICA PEIXES COM ALTO NÍVEL DE MERCÚRIO




A COR AMARELA INDICA PEIXES COM RAZOÁVEL NÍVEL DE MERCÚRIO



A COR VERDE INDICA PEIXES COM BAIXO NÍVEL DE MERCÚRIO



01 PORÇÃO DE BICO-DE-PATO CONTÉM A MESMA QUANTIDADE DE MERCÚRIO
QUE 06 PORÇÕES DE ARACÚS E QUE 10 PORÇÕES DE PACÚS-PIRANHA



ESTA TABELA FOI ELABORADA ATRAVÉS DA ANÁLISE DOS TEORES DE MERCÚRIO EM
PEIXES DE TAMANHO MÉDIO, CAPTURADOS NO PARANÁ, SÃO LUIZ DO TAPAJÓS

PROJETO CARUSO: Universidade Federal do Rio de Janeiro (UFRJ), Universidade do Québec em Montreal (UQAM)
e Centro de Pesquisa para Desenvolvimento Internacional do Canadá (CRDI).

Participatory research:

Red: high mercury levels

Yellow: moderate mercury levels

Green: low mercury levels



Fish consumption (1995-2000)

(n = 47) (Mergler et al., 2005)

	1995	2000
	123 meals per week	122 meals per week
Carnivorous	59 (48%)	21 (18%)
Herbivorous	20 (16%)	64 (52%)

Hair Hg levels (1995-2000)

(n = 47) (Mergler et al., 2005)

1995	2000	reduction
17.9 µg/g	10.9 µg/g***	39.1%

*** p < 0.001

Any solutions to the problem?

- Short term: limit exposure
- Long term: Reconcile Fishing and farming by developing sustainable regional agroforestry

Developing sustainable regional agroforestry

- Understand social background of communities


Activities on family lots in relation to origin

		PA	NE
Bananas	kg/y	14,600	22,700
Rice	kg/y	700	1,900
Cattle	beef	7.9	12.5
Fishing	kg/week	16	8
Hunting	%	64	55


Developping sustainable regional agroforestry

- Understand social background
 - Understand usual farming practices

After 19 years of land colonization




Average lot size of 70 ha (350 m x 2 km)



The number of sequences (culture - fallow) varies a lot among farmers.

The sequence always begins with culture and ends with pasture.



40% farmers prefer
Only one sequence

Developping sustainable regional agroforestry


- Understand social background
 - Understand usual practices
- Elaborate adapted agroforestry models

Developping sustainable regional agroforestry

- Understand social background
 - Understand usual practices
- Elaborate agroforestry models
 - Divulgate through adapted environmental education

SEQÜÊNCIA DOS CULTIVOS E MERCURIO


1 Mata



2 Primeira queimada




3 Primeira roça



4 Roça e gado depois alguns anos




5 Capoeira



6 Segunda queimada




Regionalization of the project: multiple workshops including representatives of communities, NGO's, local and national governmental institutions



Information web about mercury between inhabitants of Brasília Legal

Relations **between groups** / **with health people** / **intern within groups**



Sustainable regional agroforestry practices

Ribeirinhos will not change their behaviors for the sake of the Carbon cycle, neither for fears of climate changes nor biodiversity loss.

But, they will act for a better quality of life, a better health of their family!

Sustainable regional agroforestry practices

Ribeirinhos will not change their behaviors for the sake of the Carbon cycle, neither for fears of climate changes nor biodiversity loss.


But, they will act for a better quality of life, a better health of their family!

In that sense, Hg lixiviation from newly deforested soils is a good pretext to change their practices!

Obrigado!

<http://www.unites.uqam.ca/saam>

SAAM - Sistemas Agrícolas Amazônicos



Contaminação mercurial dos ecossistemas aquáticos do Rio Tapajós na região amazônica:

Uma alavanca para a implementação de modelos agrícolas ribeirinhos baseados no Desenvolvimento sustentável

Objetivos

Área de estudo

Instituições e membros

Links

Notícias

Intranet

Uma proposta de networking - pesquisa - intervenção

Banco Mundial
Biodôme de Montréal
CIAT
EMBRAPA
FCAP
IEC
IBAMA

INCRA
MPEG
PSA
SUDAM
UFPA
UFRJ
UQAM

Ecosystem representation of Agroforestry practices

Agricultural practices

Cultivated crops


Plant cover on slopes

Tree plantation


Pastures

Burning frequency


Ecosystem representation of Agroforestry practices



Ecosystem representation of Agroforestry practices



Ecosystem representation of Agroforestry practices



Ecosystem representation of Agroforestry practices

