

# The origin of stream flow in small Amazon forest and pasture watersheds: an end-member mixing analysis approach

Joaquín Chaves<sup>1</sup>, Christopher Neill<sup>1</sup>, Helmut Elsenbeer<sup>2</sup>, Alex Krusche<sup>3</sup>, Sonja Germer<sup>2</sup>, and Sérgio Gouveia Neto<sup>3</sup>

<sup>1</sup>The Ecosystems Center, Marine Biological Laboratory, Woods Hole, MA 02543

<sup>2</sup> Universität Potsdam, Institut für Geoökologie, Potsdam, Germany

<sup>3</sup>Laboratório de Ecologia Isotópica, CENA –USP, Piracicaba, SP, Brasil



Biological Discovery in Woods Hole

*Founded in 1888 as the Marine Biological Laboratory*



**USP** Universidade de São Paulo  
Brasil

# Flowpaths:

Precipitation falling in watersheds may take a number of pathways to streams



Biological Discovery in Woods Hole

Founded in 1888 as the Marine Biological Laboratory

# Flowpaths:

Precipitation falling in watersheds may take a number of pathways to streams

- Fast

- ◎ *Overland flow*
- ◎ *Shallow subsurface flow*



Biological Discovery in Woods Hole

Founded in 1888 as the Marine Biological Laboratory

# Flowpaths:

Precipitation falling in watersheds may take a number of pathways to streams

- Fast

- Overland flow*

- Shallow subsurface flow*

- Slow

- *Groundwater*

- Soil water*



Biological Discovery in Woods Hole  
Founded in 1888 as the Marine Biological Laboratory

# Flowpaths:

*The conditions encountered by water along flowpaths control biogeochemical transformations that determine stream water chemistry*



Biological Discovery in Woods Hole

Founded in 1888 as the Marine Biological Laboratory

# Flowpaths:

## Early thinking:

- © High surface infiltrability in undisturbed tropical soils led to predominantly vertical flowpaths
- © Precluding fast lateral flows such as overland flow except on steep hillslopes



Biological Discovery in Woods Hole

Founded in 1888 as the Marine Biological Laboratory



# Flowpaths:

Recent work :

Rainfall intensities can exceed soil infiltration capacities, leading to lateral flows




Biological Discovery in Woods Hole

Founded in 1888 as the Marine Biological Laboratory

# Flowpaths:

Question :

What is the importance of vertical vs. horizontal flowpaths to stream flow in lowland Amazon forest?



Biological Discovery in Woods Hole

Founded in 1888 as the Marine Biological Laboratory




# Flowpaths:



Question :

How does the contribution of these flowpaths change as forest is converted to pasture?



Biological Discovery in Woods Hole

Founded in 1888 as the Marine Biological Laboratory

# End-Member Mixing Analysis (EMMA)

*Journal of Hydrology*, 116 (1990) 321–343  
Elsevier Science Publishers B.V., Amsterdam · Printed in The Netherlands

321

[3]

## MODELLING STREAMWATER CHEMISTRY AS A MIXTURE OF SOILWATER END-MEMBERS — AN APPLICATION TO THE PANOLA MOUNTAIN CATCHMENT, GEORGIA, U.S.A.

RICHARD P. HOOPER<sup>1</sup>, NILS CHRISTOPHERSEN<sup>2</sup> and NORMAN E. PETERS

<sup>1</sup>U.S. Geological Survey, Water Resources Division, Doraville, GA 30360 (U.S.A.)

<sup>2</sup>Center for Industrial Research, Environmental Chemistry Division, Oslo (Norway)

328





Fig. 3. Dissolved silica and magnesium EMMA plot.



# Investigating the applicability of end-member mixing analysis (EMMA) across scale: A study of eight small, nested catchments in a temperate forested watershed

April L. James<sup>1,2</sup> and Nigel T. Roulet<sup>1,3</sup>




**Figure 4.** Bivariate solute plots of Lk catchment (147 ha) stream water chemistry, located at the outflow of the Westcreek watershed ( $n = 123$ ). Ions and ALK are in units of  $\mu\text{eq/L}$ ; DOC is in units of  $\log(\mu\text{mol/L})$ ; EC is in units of  $\mu\text{S/cm}$ .



## Investigating the applicability of end-member mixing analysis (EMMA) across scale: A study of eight small, nested catchments in a temperate forested watershed

April L. James<sup>1,2</sup> and Nigel T. Roulet<sup>1,3</sup>



**Figure 10.** Lk (147 ha) 2-D mixing space or U-space: (a) Lk stream water chemistry only and (b) stream water chemistry from all catchments. Projected values of end-members represent average observed values from individual instruments (e.g., wells, piezometers, lysimeters). These data are also used in the calculation of the median value of each end-member represented in Table 5.

# End-Member Mixing Analysis (EMMA)

## ❖ Principal Component Analysis (PCA)

*“The main purpose of PCA is to find a lower-dimensional space,  $U$ , where most of the observations can be assumed to lie within a specified accuracy”* Christophersen & Hooper, 1992

## References:

Christophersen, N & Hooper RP **Multivariate Analysis of Stream Water Chemical Data: The Use of Principal Components Analysis for the End-Member Mixing Problem.** WATER RESOURCES RESEARCH, VOL. 28, NO. 1, PAGES 99–107, 1992

Hooper RP **Diagnostic tools for mixing models of stream water chemistry.** Water Resources Research, Volume 39, Issue 3, pp. HWC 2-1, 2003




Biological Discovery in Woods Hole

Founded in 1888 as the Marine Biological Laboratory



# End-Member Mixing Analysis (EMMA)

## ❖ Principal Component Analysis (PCA)...*in a nutshell*






Biological Discovery in Woods Hole

Founded in 1888 as the Marine Biological Laboratory

# Study site:

Fazenda Rancho Grande, Rondônia



▲ Throughfall collector

x Overland flow collector

▼ H flume

○ Lysimeter nest

• Groundwater well

⊕ Precipitation collector



Biological Discovery in Woods Hole

Founded in 1888 as the Marine Biological Laboratory

Surface

12.5 cm

20 cm




Fig. 3. Infiltrability and Ksat as a function of land use (at surface, at a depth of 12.5 and 20 cm: panels A–C, respectively). The crossbar within the box shows the median, the length of the box reflects the interquartile range, the fences are marked by the extremes if there are no outliers, or else by the largest and smallest observation that does not qualify for an outlier. Outliers are defined as data points more than 1.5 times the interquartile range away from the upper or lower quartile. The notches represent the 95% confidence interval for the median, and overlapping notches from two box plots indicate that there is no significant difference between the medians.



Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

SCIENCE @ DIRECT®

Forest Ecology and Management 222 (2006) 23–34

Forest Ecology  
and  
Management

[www.elsevier.com/locate/foreco](http://www.elsevier.com/locate/foreco)

## The influence of land-use changes on soil hydraulic properties: Implications for runoff generation

Beate Zimmermann<sup>a,\*</sup>, Helmut Elsenbeer<sup>a</sup>, Jorge M. De Moraes<sup>b</sup>

<sup>a</sup>Institute of Geography, University of Potsdam, P.O. Box 601553, 14415 Potsdam, Germany

<sup>b</sup>CEMEX, Universidade de São Paulo, C.P. 95, 13400-970 Piracicaba, SP, Brazil

Received 7 December 2004; received in revised form 4 June 2005; accepted 3 October 2005



Biological Disc

Founded in 1888 as the Marine Biological Laboratory



# Study site:

Fazenda Rancho Grande, Rondônia




Biological Discovery in Woods Hole

Founded in 1888 as the Marine Biological Laboratory

## Study site:

Fazenda Rancho Grande, Rondônia




Biological Discovery in Woods Hole

Founded in 1888 as the Marine Biological Laboratory




# Water yields



Biological Discovery in Woods Hole

Founded in 1888 as the Marine Biological Laboratory

# Mixing Diagrams:




Biological Discovery in Woods Hole

*Founded in 1888 as the Marine Biological Laboratory*

# Mixing Diagrams:




## Forest



Biological Discovery in Woods Hole

*Founded in 1888 as the Marine Biological Laboratory*


## A photograph of a grassy field with a wooden fence and a haystack in the foreground, and a dense forest in the background. The field is covered in tall, green grass. A wooden fence runs across the middle ground, and a haystack is visible in the center. The background is a dense forest of tall trees under a blue sky with some clouds.



# Mixing Diagrams:



## Pasture




**MBL**

Biological Discovery in Woods Hole

Founded in 1888 as the Marine Biological Laboratory



# Overall Results:



Biological Discovery in Woods Hole

*Founded in 1888 as the Marine Biological Laboratory*



Throughfall



Groundwater




Soil water



Overland flow

# Overall Results:




Biological Discovery in Woods Hole

Founded in 1888 as the Marine Biological Laboratory


# Overall Results:



Entire rainy season



Pasture



Throughfall



Groundwater



Soil water



Overland flow



Biological Discovery in Woods Hole

Founded in 1888 as the Marine Biological Laboratory

# Conclusions:

- ◎ Surface flow increased dramatically after the establishment of pasture.
- ◎ Sources to storm flow to small streams changed considerably following deforestation
- ◎ Transition to pasture caused a shift towards predominantly faster flowpaths
- ◎ Flowpath hydrology in pasture is “simpler” than in forest



# We are we going with this:

- ◎ Use these results to tests our knowledge and observations about the biogeochemistry of these landscapes.
- ◎ Test EMMA at larger scales (1<sup>st</sup>, 2<sup>nd</sup> order catchments).
- ◎ Apply these approach in areas of the Amazon where mechanized, large scale agriculture is growing rapidly.





## **Acknowledgements**

We thank Alexandra A. Montebelo and Gustavo G. Baldi at the Centro de Energia Nuclear na Agricultura in Piracicaba for conducting the chemical analyses. Shelby Hayhoe, Sonya Remington, Mathew Shamey, Tobias Vetter, and Lisa Werther helped during the field work phase of this study. Support for this study was provided by the US National Science Foundation grant DEB-0315656 and by grants from Brazilian agencies FAPESP # 03/13172-2 and CNPq # 420199/2005-5. Special thanks to the Schmitz family for allowing us to work on their land.



Biological Discovery in Woods Hole

Founded in 1888 as the Marine Biological Laboratory





Photo: Lisa Werther



Biological Discovery in Woods Hole

Founded in 1888 as the Marine Biological Laboratory

Table 1: Precipitation and runoff at the research watersheds during the period of study.

		Early <sup>a</sup>	Late <sup>b</sup>	Total
Precipitation events, count		73	90	163
Precipitation, mm		1020	1164	2184
Runoff, mm (# events)	Forest	2.2 (15)	14.5 (29)	16.7 (44)
	Pasture	101 (20)	277 (35)	378 (55)
Sampled events, count	Forest	10	12	22
	Pasture	11	16	27
Precipitation during sampled events, mm	Forest	332	352	684
	Pasture	313	311	624
Runoff during sampled events, mm	Forest	0.6	2.1	2.6
	Pasture	29.1	104.4	133.5

<sup>a</sup> Early rainy season, from August 4<sup>th</sup> to December 31<sup>st</sup>, 2004.

<sup>b</sup> Late rainy season, from January 1<sup>st</sup> 2005 to April 25<sup>th</sup>, 2005



Biological Discovery in Woods Hole

*Founded in 1888 as the Marine Biological Laboratory*