




# Water and Solute Balances in Small Amazonian Forest and Pasture Watersheds

Chris Neill, Sonja Germer, Alex Krusche, Joaquín Chaves,  
Helmut Elsenbeer, Sergio Gouveia Neto, Adriana Castellanos Bonilla

Marine Biological Laboratory  
University of São Paulo  
University of Potsdam

# Fazenda Rancho Grande, Rondônia



▲ Throughfall collector

x Overland flow collector

▽ H flume

○ Lysimeter nest


• Groundwater well

⊕ Precipitation collector

# Fazenda Rancho Grande, Rondônia



# Infiltration and hydraulic conductivity



Zimmermann et al. 2006


# Annual water yield

Rain=2286 mm

176 rainfall events

70 generated flow  
in pasture

36 generated flow  
in forest



# Quickflow and Baseflow



		No. Quickflow	Baseflow	Total	
		events	----- mm-----		
Forest	TDWS	11	2	0	2
	WS	25	23	0	23
	Total	26	25	0	25
Pasture	TDWS	12	48	0	48
	WS	58	360	9	369
	Total	70	408	9	417

# Water balance

	Forest		Pasture	
	mm (% of precip.)			
Precipitation	2286			
Interception <sup>a</sup>	50	(2.2)	-	
Streamflow	25	(1.1)	417	(18)
Evapotranspiration	1387	(61)	1024	(45)
Groundwater recharge	824	(36)	846	(37)



# Partitioning water sources with end member modeling



Throughfall



Groundwater




Soil water



Overland flow




# Solute inputs in TF and rain are highly seasonal




# Solute inputs in TF and rain are highly seasonal

	Total Flux		Ratio TDW / W	
	---kg ha <sup>-1</sup> ---			
	Rain	TF	Rain	TF
K	9	78	1.0	0.9
Ca	18	19	0.9	0.6
NO <sub>3</sub> -N	1	2	7	5
NH <sub>4</sub> -N	4	5	3	1.3


# Solute concentrations vary within rain events




# Solute concentrations vary within flowpaths



# Solute concentrations vary within flowpaths



# Pathways of solute export





# Relative solute balances in forest and pasture

	Forest / Pasture
$K^+$	2.2
$Ca^{2+}$	0.7
$NO_3^-$ -N	0.3
$NH_4^+$ -N	1.8

# Conclusions

- 1) Large decreases to soil infiltrability and permeability lead to 18 x increase in streamflow from pasture
- 2) Quickflow dominates surface water output in forest and pasture
- 3) Large seasonal and within-rain event concentrations of most solutes likely related to fragmented regional landscape
- 4) Large amount of  $\text{NO}_3^-$  removal between soils and groundwater apparently occurs at soil-groundwater interface

# Conclusions

- 5) Biogeochemical “homogenization” of flowpath chemistry in pasture leads to low DIN export
- 6) Despite importance of surficial flowpaths to stream flow, most solute export is via groundwater
- 7) Despite large shifts in runoff generation and differences in flowpath chemistry, differences in total export are surprisingly modest

# Acknowledgments

Reynaldo Victoria  
Linda Deegan  
Vicky Ballester  
Tobia Vetter  
Lisa Werther  
Shelby Hayhoe  
Mathew Shamey  
Xandra Montebelo  
Gustavo Baldi  
Sonia Remington  
Schmitz family

NASA-LBA

FAPESP

CNPq

NSF

DAAD



Biological Discovery in Woods Hole

*Founded in 1888 as the Marine Biological Laboratory*

**USP** Universidade de São Paulo  
Brasil

