

## ABSTRACT

We report the first ever genomic characterization (of a RPTP type IIB gene) of an RNA-Protein Terminus complex Recombinant Prototyp Tyrosine Triphosphate (RPT-3) segment. Alternatively, different variants may be produced from these RTGAs (recombinant Protease Terminal Proteins) which are thought to have originated as separate modular proteins that fusion into

## INTRODUCTION

The RPTP<sub>p</sub> gene encodes a protein that is expressed in the nucleus of the cell. The gene encodes a protein that is expressed in the nucleus of the cell. This gene has been identified in multiple human tissues, such as the brain and pancreas, and has also been found in a number of other organisms, including yeast, rodents, and flies.

The RPTP<sub>p</sub> gene encodes a protein that is expressed in the nucleus of the cell. The gene encodes a protein that is expressed in the nucleus of the cell. The gene has been identified in multiple human tissues, such as the brain and pancreas, and has also been found in a number of other organisms, including yeast, rodents, and flies. The RPTP<sub>p</sub> gene encodes a protein that is expressed in the The regulation of various cellular functions, including signal transduction, growth, differentiation, cell adhesion, and axon guidance, is facilitated by protein tyrosine phosphorylation. This regulatory mechanism relies on the equilibrium between protein-tyrosines and proteins involved in these functions. All receptor-like protein (RPTPs) are classified into several classes based on their extracellular region, transmembrane segment(s), and at least one intracellular catalytic domain. After cleavage, N-terminal segments remain non-covalently There have been reports of type IIB phosphatases (PTP, PTP and RPTP+) that are located on human chromosomes 1, 6, and 18; the hPCP-2 is on chromatin 1; The php-CD[n] and its RNA (cytosine peptide 7), which is believed to be the same as PCP2, with several more human Rpt proteins (like Peptetapis), including mPPmam4 (#Npappapidae 8 Recent studies have revealed that RPTP, the most recently isolated member of the IIB family, is mostly restricted to the central nervous system. In the CNS, expression is developmentally controlled and a unique boundary region is formed in what is now known as the cerebellar cortex. The presence of cytosolic proteins such as serotonin and glutathione indicate that this molecule is involved in signal transmission through the extracellular segment of its RNA. A sequence of events has been conducted to identify the human RPTP gene, which is mapped to chromosome 20q12-13.1; it is flanked by the phospholipase C gamma 1 and splicing factor SRp55-2 genes, and the mouse gene maps to an area on the left side of the genome that is closely related to Pltp, as well as two other markers, D2Mit22 and D2-Mit52. The mouse region containing this gene has not been fully assembled.

## CONCLUSION

Remarkable conclusions This paper presents details on cloning (for the first time), the genomic structure and alternative splicing of mouse and human genes, as well as an 8 kb 3'UTR in human RPTP. Unlike other RNAs from ESCs, this gene has a significant length: approximately 1 megabase pairs of genomic DNA forms extracellular modules encoded by protein domains (typically around phase 0, and otherwise insignificant) That segment is the largest known PCR enzyme, called the ectodomain1, by the MARN2, by which we know