

## Intestinal parasitic infections in Thai HIV-infected patients with different immunity status

### ABSTRACT

In case of a tropical epidemic, it is important to focus on the non-opportunistic intestinal parasite infections that are prevalent among HIV-infected patients who present with diarrhea.

### INTRODUCTION

The interaction between the MS2 coat protein and its translational operator is a well-established example of RNA-protein recognition, utilizing genetic, biochemical, and structural methods. Figure 1 displays the primary and secondary structures of the recombinant rRNA hairpin that establish contacts with both subunits of each coat proteins dimer. The coat protein complex with its RNA target is highly intricate, as two unpaired adenosines are inserted into equivalent pockets on different subunits of the coat dimer (Figure 2). The interactions between A-4 and A-10 with coat proteins involve non-identical contacts with the same five amino acid residues, Val29, Thr45, Ser47, Finally, and Lys61. The use of X-ray crystallographic analysis indicates specific amino acid-nucleotide interactions, but fails to provide a clear explanation of their respective roles in RNA-binding and translational repression. In the experiments described here, we used amino acid substitutions of A-pocket amino acids in single-chain coat protein heterodimers to determine the significance of each residue's interaction with A-4 and A-10.

### CONCLUSION

After taking antipsychotics, peripheral lymphocytes' molecular biologically-determined dopamine receptors are reactive, as demonstrated in this study.