

- RNMC: kinetic Monte Carlo implementations for
- 2 complex reaction networks
- Laura Zichi 10 1,2*, Daniel Barter 10 3*, Eric Sivonxay 10 3*, Evan Walter Clark
- Spotte-Smith 1,4, Rohith Srinivaas Mohanakrishnan^{1,4}, Emory M. Chan⁵,
- Kristin Aslaug Persson^{4,5¶}, and Samuel M. Blau^{3¶}
- 1 Materials Science Division, Lawrence Berkeley National Laboratory, Berkeley, CA, USA 94720 2
- 7 Department of Physics, University of Michigan Ann Arbor, Ann Arbor, MI, USA 48109 3 Energy
- 8 Storage and Distributed Resources, Lawrence Berkeley National Laboratory, Berkeley, CA USA 94720 4
- 9 Department of Materials Science and Engineering, University of California Berkeley, CA, USA 94720 5
- Molecular Foundry, Lawrence Berkeley National Laboratory, Berkeley, CA, USA 94720 ¶ Corresponding
- 11 author * These authors contributed equally.

DOI: 10.xxxxx/draft

Software

- Review 🗗
- Repository 🖸
- Archive 🗗

Editor: Open Journals ♂ Reviewers:

@openjournals

Submitted: 01 January 1970 **Published:** unpublished

License

Authors of papers retain copyright and release the work under a 24 Creative Commons Attribution 4.0 International License (CC BY 4.0)

Summary

17

18

Macroscopic chemical and physical phenomena are driven by microscopic interactions at the atomic and molecular scales. In order to capture complex processes with high fidelity, simulation methods that bridge disparate time and length scales are needed. While techniques like molecular dynamics and *ab initio* simulations capture dynamics and reactivity at high resolution, they cannot be used beyond relatively small length (hundreds to thousands of atoms) and time scales (picoseconds to microseconds). Kinetic Monte Carlo (kMC) approaches overcome these limitations to bridge length and time scales across several orders of magnitude while retaining relevant microscopic resolution, making it a powerful and flexible tool.

Here, we present RNMC, an easy-to-use, modular, high-performance kMC simulation framework that enables modeling of complex systems. RNMC consists of a core module defining the common features of kMC algorithms, including an implementation of the Gillespie algorithm (Gillespie, 1977), input/output operations leveraging SQLite databases, random number sampling, threading logic for parallel execution, and dependency graphs for efficient event propensity updates. In addition, there are currently three modules defining kMC implementations for different types of applications. The GMC (Gillespie Monte Carlo) module enables simulations of reaction networks in a homogeneous (well-mixed) environment. GMC is a basic tool that is appropriate for general simulations of solution-phase chemistry. The NPMC (NanoParticle Monte Carlo) module enables simulation of dynamics in nanoparticles with 3D statistical field theory and supports one- and two-site interactions. Finally, the LGMC (Lattice Gillespie Monte Carlo) module is designed for simulations of multi-phase systems (especially at solid-fluid interfaces) where chemical and electrochemical reactions can occur between a lattice region and a homogeneous region. We have designed RNMC to be easily extensible, enabling users to add additional kMC modules for other diverse chemical and physical systems.

Statement of need

Three are many existing kMC implementations, including several open source examples (e.g. the Stochastic Parallel PARticle Kinetic Simulator or SPPARKS (Garcia Cardona et al., 2009) and kmos (Hoffmann et al., 2014)). RNMC began as a fork of SPPARKS but differs in several important ways. First, because RNMC uses the widely supported SQLite database engine for simulation inputs and outputs, it facilitates the automation of simulations. Second, RNMC



has a focus on modularity. All simulators leverage the small core library, which serves as a common interface through the use of templating. As long as they can operate through this shared core, different simulation implementations are totally independent. This means that new developers need only read and understand the core library to be able to add new capabilities to RNMC, lowering the barrier to entry, and further reduces the likelihood that new additions will adversely affect pre-existing code.

The simulation modules already implemented in RNMC provide unique capabilities that are not widely available in other open source codes. NPMC is specifically designed for 3D simulations of the complex photophysical interaction networks in nanocrystals (Teitelboim et al., 2019), 50 particularly multi-domain heterostructures whose optical properties cannot be calculated 51 deterministically (Skripka et al., 2023). NPMC can be used to simulate energy transfer interactions between dopants in nanoparticles, their radiative transitions, and nonlinear processes such 53 as upconversion (Chan, 2015) and photon avalanching (Skripka et al., 2023). LGMC is also somewhat unique in that it can simulate multi-phase systems and electrochemical processes. Simulations using LGMC can include a lattice region and a homogeneous solution region which can interact via interfacial reactions. Electrochemcial reactions can be treated using Marcus 57 theory (Marcus, 1965) or Butler-Volmer kinetics (Newman & Balsara, 2021). Because it allows for a dynamic lattice region, LGMC is also appropriate for simulations of nucleation and growth, 59 dissolution, precipitation, and related phenomena.

We have already used the GMC module in a number of prior works in applications related to Li-ion and Mg-ion batteries (Barter et al., 2023; Spotte-Smith et al., 2022, 2023). We note that these simulations included tens of millions of reactions, demonstrating that RNMC is able to scale to large and complex reaction networks. In addition, we have used NPMC to perform Bayesian optimization of upconverting nanoparticles (Xia et al., 2023).

Acknowledgements

This project was intellectually led by the Laboratory Directed Research and Development 67 Program of Lawrence Berkeley National Laboratory under U.S. Department of Energy Contract No. DE-AC02-05CH11231. L.Z. was supported in part by the U.S. Department of Energy, Office of Science, Office of Workforce Development for Teachers and Scientists (WDTS) under the Science Undergraduate Laboratory Internships Program (SULI). E.W.C.S.-S. was supported by the Kavli Energy NanoScience Institute Philomathia Graduate Student Fellowship. Work at the Molecular Foundry (E.M.C., K.A.P) was supported by the Office of Science, Office of Basic Energy Sciences, of the U.S. Department of Energy under Contract No. DE-AC02-05CH11231. Additional support came from the Joint Center for Energy Storage Research (JCESR), an Energy Innovation Hub funded by the U.S. Department of Energy, Office of Science, Basic Energy Sciences. This code was developed and tested using computational resources provided by the National Energy Research Scientific Computing Center (NERSC), a U.S. Department of Energy Office of Science User Facility under Contract No. DE-AC02-05CH11231, the Eagle and Swift HPC systems at the National Renewable Energy Laboratory (NREL), and the Lawrencium HPC cluster at Lawrence Berkeley National Laboratory.

References

Barter, D., Spotte-Smith, E. W. C., Redkar, N. S., Khanwale, A., Dwaraknath, S., Persson, K. A., & Blau, S. M. (2023). Predictive stochastic analysis of massive filter-based electrochemical reaction networks. *Digital Discovery*, 2(1), 123–137. https://doi.org/10.1039/D2DD00117A

Chan, E. M. (2015). Combinatorial approaches for developing upconverting nanomaterials:

High-throughput screening, modeling, and applications. *Chemical Society Reviews*, *44*(6),

1653–1679. https://doi.org/10.1039/C4CS00205A



- Garcia Cardona, C., Wagner, G. J., Tikare, V., Holm, E. A., Plimpton, S. J., Thompson, A.
 P., Slepoy, A., Zhou, X. W., Battaile, C. C., & Chandross, M. E. (2009). Crossing the mesoscale no-mans land via parallel kinetic monte carlo. Sandia National Laboratories (SNL), Albuquerque, NM,; Livermore, CA.
- Gillespie, D. T. (1977). Exact stochastic simulation of coupled chemical reactions. The Journal
 of Physical Chemistry, 81(25), 2340–2361.
- Hoffmann, M. J., Matera, S., & Reuter, K. (2014). Kmos: A lattice kinetic monte carlo
 framework. Computer Physics Communications, 185(7), 2138–2150. https://doi.org/10.
 1016/j.cpc.2014.04.003
- Marcus, R. A. (1965). On the theory of electron-transfer reactions. VI. Unified treatment for homogeneous and electrode reactions. *The Journal of Chemical Physics*, 43(2), 679–701.
 https://doi.org/10.1063/1.1696792
- Newman, J., & Balsara, N. P. (2021). Electrochemical systems. John Wiley & Sons.
- Skripka, A., Lee, M., Qi, X., Pan, J.-A., Yang, H., Lee, C., Schuck, P. J., Cohen, B. E., Jaque, D., & Chan, E. M. (2023). A generalized approach to photon avalanche upconversion in luminescent nanocrystals. *Nano Letters*, 23(15), 7100–7106. https://doi.org/10.1021/acs.nanolett.3c01955
- Spotte-Smith, E. W. C., Blau, S. M., Barter, D., Leon, N. J., Hahn, N. T., Redkar, N. S., Zavadil, K. R., Liao, C., & Persson, K. A. (2023). Chemical reaction networks explain gas evolution mechanisms in Mg-ion batteries. *Journal of the American Chemical Society*, 145(22), 12181–12192. https://doi.org/10.1021/jacs.3c02222
- Spotte-Smith, E. W. C., Kam, R. L., Barter, D., Xie, X., Hou, T., Dwaraknath, S., Blau, S. M., & Persson, K. A. (2022). Toward a mechanistic model of solid–electrolyte interphase formation and evolution in lithium-ion batteries. *ACS Energy Letters*, 7(4), 1446–1453. https://doi.org/10.1021/acsenergylett.2c00517
- Teitelboim, A., Tian, B., Garfield, D. J., Fernandez-Bravo, A., Gotlin, A. C., Schuck, P. J., Cohen, B. E., & Chan, E. M. (2019). Energy transfer networks within upconverting nanoparticles are complex systems with collective, robust, and history-dependent dynamics.
 The Journal of Physical Chemistry C, 123(4), 2678–2689. https://doi.org/10.1021/acs.jpcc.9b00161
- Xia, X., Sivonxay, E., Helms, B. A., Blau, S. M., & Chan, E. M. (2023). Accelerating the design of multishell upconverting nanoparticles through bayesian optimization. *Nano Letters*, 23(23), 11129–11136. https://doi.org/10.1021/acs.nanolett.3c03568