

# A simulation framework for exploring spatio-temporal dynamics in mixed fisheries

Paul J. Dolder<sup>a,b,\*</sup>, C  il  n Minto<sup>a</sup>, Jean-Marc Guarini<sup>c</sup>, Jan-Jaap Poos<sup>d</sup>

<sup>a</sup>*Galway-Mayo Institute of Technology (GMIT), Dublin Road, Galway, Ireland*

<sup>b</sup>*Centre for Environment, Fisheries and Aquaculture Science (Cefas), Pakefield Road, Lowestoft, UK*

<sup>c</sup>*Universit   Pierre et Marie Curie, 4 Place Jussieu, 75005 Paris, France*

<sup>d</sup>*Wageningen Marine Research, Haringkade 1 1976 CP IJmuiden, Netherlands*

---

## Abstract

[A concise and factual abstract is required. The abstract should state briefly the purpose of the research, the principal results and major conclusions. An abstract is often presented separately from the article, so it must be able to stand alone. For this reason, References should be avoided, but if essential, then cite the author(s) and year(s). Also, non-standard or uncommon abbreviations should be avoided, but if essential they must be defined at their first mention in the abstract itself. Graphical abstract: Although a graphical abstract is optional, its use is encouraged as it draws more attention to the online article. The graphical abstract should summarize the contents of the article in a concise, pictorial form designed to capture the attention of a wide readership. Graphical abstracts should be submitted as a separate file in the online submission system. Image size: Please provide an image with a minimum of 531 X 1328 pixels (h X w) or proportionally more. The image should be readable at a size of 5 x 13 cm using a regular screen resolution of 96 dpi. Preferred file types: TIFF, EPS, PDF or MS Office files.]

*Keywords:* Some, keywords, here. Max 6, American "spelling"

*2010 MSC:* 00-01, 99-00

---

\*Corresponding author

Email address: [paul.dolder@gmit.ie](mailto:paul.dolder@gmit.ie) (Paul J. Dolder)

## 1. Introduction

[Guidance:: State the objectives of the work and provide an adequate background, avoiding a detailed literature survey or a summary of the results.]

5      Fishers exploit fish populations that are heterogenously distributed in space and time without prior knowledge of species distributions and non-selective fishing gear. Fisheries managed by single-species quotas catch an assemblage of species, known as mixed fisheries, leading to discarding of overquota catch. Reducing discarding is crucial to ensure biological and economic sustainability  
10 of fisheries. As such there is increasing interest in technical solutions such as gear, spatial closures as a way of avoiding discarding unwanted catch.

Use of spatial management as a tool is hampered by lack of knowledge of fish and fishery spatiotemporal dyanmics and understanding of the scale at  
15 which processes are important for management. Understanding of the correct scale for spatial management is crucial in order to implement measures at a resolution that ensures effective management[1] while minimising economic impact; for example a scale which promote species avoidance for vulnerable or low quota species while allowing continance of sustainable fisheries for available  
20 quota species.

Ensuring measures are implemented at an appropriate scale has been a challenge in the a past that has led to ineffectual measures with unitended consequences such as limited impact towards the management objective or increased  
25 benthic impact on previously unexploited areas (e.g. the cod closure in the North Sea[2, 3]). Since then more refined spatial information has become available through the combination of logbook and Vessel Monitoring System (VMS) data[4, 5, 6, 7], though such information is patchy and derived from an inherently biased sampling programme (i.e. targeted fishing). Further, generally fishers  
30 only recorded landings (not catch) on a daily basis. This leads to questions

about the validity of inferences that can be drawn from landings data assigned to VMS activity pings.

In order to test the assumption that VMS associated landings data can be  
35 used to draw inference on the underlying population structures we develop a  
simulation model where population dynamics are known rather than inferred  
from sampling or commercial catches. Population movement is driven by a  
random (diffusive) and directed (advective) process and we incorporate charac-  
terisation of a number of different fisheries exploiting four fish populations with  
40 different spatial and population demographics.

Using our model we simulation 20 years of exploitation of the fish populations  
and use the results from the fishing model to draw inference on the underlying  
population structures. We compare this inference to i) a simulated [stratified]  
45 fixed-site sampling design commonly used for fisheries monitoring purposes, ii)  
the underlying population structures input to the simulation.

[Could fit a geostatistical model (e.g. VAST) to the fisheries-dependent and  
fisheries-independent data, though may be overkill...]

50

We simulate a fishery closure to protect one species based on the fishery-  
dependent inferred distributions at a spatial and temporal scale typical in fish-  
eries management, and assess a theoretical "benefit" to the population, and  
effect on the other three populations. Further, we extend our analysis to a  
55 range of spatial and temporal scales to assess the impact of these processes on  
the success of the management measure.

## 2. Materials and Methods

[Guidance: Provide sufficient details to allow the work to be reproduced  
60 by an independent researcher. Methods that are already published should be  
summarized, and indicated by a reference. If quoting directly from a previously  
published method, use quotation marks and also cite the source. Any modifi-  
cations to existing methods should also be described.]

65 We develop a simulation model with a modular event-based approach, where  
modules occur on independent time-scales appropriate to capture the character-  
istic of the process modelled. The fishing model operated on a tow-by-tow basis,  
while population dynamics (fishing and natural mortality, growth) operate on a  
daily time-step. Population movement occurs on a two-weekly time-step, while  
70 recruitment occurs periodically each year for a set time period (e.g. 3 weeks) at  
at specified point individual to a species. The model structure is summarised in  
Figure 1.

In the following section, we describe each of the model components; 1) Popu-  
75 lation dynamics, 2) Recruitment dynamics, 3) Population movement, 4) fishery  
dynamics.

### 2.1. Population dynamics

The basic population level processes are simulated using a modified two-  
80 stage Deriso-Schnute delay difference model [8, 9, 10] occurring at a daily time-  
step. Here, population biomass growth and depletion for pre-recruits and fish  
recruited to the fishery are modelled separately as a function of previous re-  
cruited biomass, intrinsic population growth and recruitment:




Figure 1: Overview Schematic of simulation model

$$\begin{aligned}
 B_{y,w+1} = & \\
 & (1 + \rho)B_{y,w} \cdot e^{-Z_{y,w}} - \rho \cdot e^{-Z_{y,w}} \quad \times \\
 & (B_{y,w-1} \cdot e^{-Z_{y,w-1}} + Wt_{R-1} \cdot \alpha_{w-1} \cdot R_{\tilde{y}(y,w-1)}) \quad + \\
 & Wt_R \cdot \alpha_w \cdot R_{\tilde{y}(y,w)}
 \end{aligned}$$

where  $\rho$  is Brody's coefficient, shown to be approximately equal to  $\exp(-K)$ ,  
85 where  $K$  is the growth rate from a von bertalanffy logistic growth model [9].  
 $Wt_{R-1}$  is the weight of fish prior to recruitment, while  $Wt_R$  is the recruited  
weight.  $\alpha_w$  represents the proportion of fish recruited during the week, while  
 $R_y$  is the annual recruits.

90 Mortality  $Z$  can be decomposed to natural mortality,  $M$ , and fishing mor-  
tality,  $F$ , where both  $M$  and  $F$  are instantaneous rates with  $M$  fixed and  $F$   
calculated by solving the Baranov catch equation [11] for  $F$ :

$$C_w = \frac{F_w}{F_w + M_w} * (1 - e^{-(F_w + M_w)}) * B$$

where  $C$  is the summed catch from the fishing model across all fleets and  
vessels for the population during the day, and  $B$  the daily biomass for the species.

95

## 2.2. Recruitment dynamics

Recruitment is modelled through a function relating the biomass at time of  
recruitment to recruits, either as a stochastic Beverton-Holt stock-recruit form  
([12]):

$$\bar{R} = \frac{(\alpha * B)}{(\beta + B)}$$

$$R \sim N[(\bar{R}, \sigma^2)]$$

100 Where  $\alpha$  is the maximum recruitment rate,  $\beta$  the spawning stock biomass  
(SSB) required to produce half the maximum, and  $B$  current SSB;

or a stochastic Ricker form [13]

$$\bar{R} = B * e^{(\alpha - \beta * B)}$$

$$R \sim N[(\bar{R}, \sigma^2)]$$

where  $\alpha$  is the maximum productivity per spawner and  $\beta$  the density dependent reduction in productivity as the SSB increases.

### 2.3. Population movement

In order to simulate how fish populations might be distributed in space and time, we employed a Gaussian spatial process to model habitat suitability for each of the populations, with an advection-diffusion process to control how the populations moved in time.

For the habitat we define a Gaussian random field process,  $\{S(x) : x \in \mathbb{R}^2\}$ , that is a stochastic process where any collection of locations  $x_1, \dots, x_n$  where for each  $x_i \in \mathbb{R}^2$ , the joint distribution of  $S = \{S(x_1), \dots, S(x_n)\}$  is multivariate Gaussian. The distribution is specified by its *mean function*,  $\mu(x) = E[S(x)]$  and its *covariance function*,  $\gamma(x, x') = \text{Cov}\{S(x), S(x')\}$  [14].

The specification of the covariance structure directly affects the smoothness of the surfaces which the process generates. We used the *Matérn* family of covariance structures, one where the correlation strength weakens the further the distance apart (i.e. the correlation between  $S(x)$  and  $S(x')$  decreases as the distance  $u = \|x - x'\|$  increases). The *Matérn* correlation is a two-parameter family where:

$$\rho(u) = \{2^{\kappa-1} \Gamma \kappa\}^{-1} (u/\phi)^\kappa K_\kappa(u/\phi)$$

$K_\kappa(\cdot)$  is a modified Bessel function of order  $\kappa$ ,  $\phi > 0$  is a scale parameter with the dimensions of distance, and  $\kappa > 0$ , called the order, is a shape parameter which determines the smoothness of the underlying process.

The population is initialised at a single location, and subsequently moves according to a probabilistic distribution based on habitat suitability and distance

from current cell.

$$Pr(B|A) = \frac{e^{-\lambda * d_{AB}} \cdot Hab_B^2}{\sum_{c=1}^C e^{-\lambda * d} \cdot Hab^2} \quad (1)$$

Where  $d_{AB}$  is the euclidean distance between cell  $A$  and cell  $B$ , and  $\lambda$  is some rate of decay.

135

During specified weeks of the year, the habitat quality is modified for spawning habitats, meaning each population has a concentrated area where spawning takes place and the population moves towards this in the weeks prior to spawning.

140

#### 2.4. Fleet dynamics

The fleet dynamics can be broadly categorised into three components; fleet targeting - which determines the fleet catch efficiency and preference towards a particular species; trip-level decisions, which determine the initial location  
145 to be fished at the beginning of a trip; and within-trip decisions, determining movement from one fishing spot to another within a trip. A further sub-model controls local depletion of fish populations at fished sites.

##### 2.4.1. Fleet targeting

Each fleet of  $n$  vessels is characterised by both a general efficiency,  $Q$ , and a  
150 species specific efficiency,  $Q_s$ . Thus, the product of these parameters affects the overall catch rates for the fleet and the preferential targeting of one species over another. This, in combination with the parameter choice for the step-function (as well as some randomness from the exploratory fishing process) determines the preference of fishing locations for the fleet. All species prices are kept the  
155 same, across fleets, though can be made to vary seasonally.



#### 2.4.2. Trip-level decisions

Several studies (e.g.[15, 16, 17]) have confirmed past activity and past catch rates are strong predictors of fishing location choice. For this reason, the fleet dynamics sub-model includes a learning component, where a vessels initial fishing location in a trip is based on selecting from previously successful fishing locations. This is achieved by sorting all previous fishing events in the previous trip as well as the previous time periods in past years, and choosing randomly from the top x % of fishing events in value. Simulation testing indicating this increased the mean value of catches for the vessels, over just relying on the correlated random walk function.

#### 2.4.3. Within-trip decisions

Fishing locations within a trip are determined by a Random Walk process. A Random walk was chosen as it is commonly used in ecology to describe animal movement which searching for homogeneously distributed prey about which there is uncertain knowledge. In a random walk, movement is a stochastic process through a series of steps that can either be equal in length or take some other functional form. The direction of the random walk can be correlated, a characteristic known as 'persistence', providing some overall location directional movement [18] or uncorrelated.

A *lévy walk* is a particular form of random walk characterised by a heavy-tailed distribution of step-length and has received a lot of attention in ecological theory in recent years as having shown to have very similar characteristics as those observed by animals in nature, and being a near optimum searching strategy for predators pursuing patchily distributed prey [19, 20]. [21] showed that Peruvian anchovy fishermen have a stochastic search pattern similar to that observed with a lévy walk. However, it remains a subject of debate, with the contention that search patterns may be more simply characterised as random walks [22] with specific patterns related to the characteristics of the prey field [23].

We use a modified random walk where directional change is based on a correlated circular distribution where a favourable fishing ground is likely to be "fished back over" by the vessel returning in the direction it came from and step length (i.e. the distance travelled from the current to the next fishing location) is determined by relating recent fishing success, measured as the summed value of fish caught,

$$Rev = \sum_{s=1}^{\infty} C_s \cdot Pr_s$$

where  $C_s$  is catch of a species, and  $Pr_s$  price of a species, to step distance. Here, when fishing is successful vessels remain in a similar location and continue to exploit the local fishing grounds. When unsuccessful, they move some distance  
190 away from the current fishing location. The movement distance retains some degree of stochasticity, which can be controlled separately.

The step function takes the form:

$$StepL = e^{\log(\beta_1) + \log(\beta_2) - (\log(\frac{\beta_1}{\beta_3}))} * Rev$$

So that, a step from (x1,y1) to (x2, y2) is defined by:

$$\begin{aligned} (x2, y2) &= x1 + StepL \cdot \cos(\frac{\pi \cdot Br}{180}), \\ &y1 + StepL \cdot \sin(\frac{\pi \cdot Br}{180}) \\ \text{where } Br &\sim U[(0, 360)] \end{aligned}$$

#### 195 2.4.4. Local population depletion

Where several fishing vessels are exploiting the same fish population competition is known to play an important role in local distribution of fishing effort [24]. If several vessels are fishing on the same patch of fish, local depletion and interference will affect fishing location choice of the fleet as a whole [25, 26]. In  
200 order to account for this behaviour, the fishing sub-model operates spatially on

a daily time-step so that for future days the biomass available to the fishery is reduced in the areas fished. The cumulative effect is to make heavily fished areas less attractive as future fishing opportunities.

## 205 2.5. Fisheries independent survey

A fisheries-independent survey is simulated where fishing on a regular grid begins each year at the same time for a given number of stations (a fixed station survey design). Catches of the populations present are recorded but not removed from the population. This provides a fishery independent snap shot of  
210 the populations at a regular spatial distribution each year, similar to scientific surveys undertaken by fisheries research agencies.

## 3. Theory/calculation

[Guidance: A Theory section should extend, not repeat, the background to  
215 the article already dealt with in the Introduction and lay the foundation for further work. In contrast, a Calculation section represents a practical development from a theoretical basis.]

HERE DESCRIBE THE PARAMETERISATION, AND ANY COMPARISON AGAINST 'REAL' DATA

## 220 4. Results

Present simulated closures in terms of % change in population biomass and fishery.

[Guidance: Results should be clear and concise.]

## 5. Discussion

225 [Guidance: This should explore the significance of the results of the work, not repeat them. A combined Results and Discussion section is often appropriate. Avoid extensive citations and discussion of published literature.]

## 6. Conclusions

[Guidance: The main conclusions of the study may be presented in a short  
230 Conclusions section, which may stand alone or form a subsection of a Discussion  
or Results and Discussion section.]

## Appendices

[Guidance: If there is more than one appendix, they should be identified  
as A, B, etc. Formulae and equations in appendices should be given separate  
235 numbering: Eq. (A.1), Eq. (A.2), etc.; in a subsequent appendix, Eq. (B.1)  
and so on. Similarly for tables and figures: Table A.1; Fig. A.1, etc.]




Figure 2: habitat

## Abbreviations

Detail any unusual ones used.

## Acknowledgements

240 those providing help during the research..




Figure 3: Annual Summary

## Funding

This work was supported by the MARES doctoral training program; and the Centre for Environment, Fisheries and Aquaculture Science seedcorn program.

## References

- 245 [1] D. C. Dunn, S. M. Maxwell, A. M. Boustany, P. N. Halpin, Dynamic ocean management increases the efficiency and efficacy of fisheries management, Proceedings of the National Academy of Sciences (2016) 201513626doi:




Figure 4: f dynamics

10.1073/pnas.1513626113.

URL <http://www.pnas.org/content/early/2016/01/02/1513626113>.

**abstract**

- [2] A. D. Rijnsdorp, G. J. Piet, J. J. Poos, Effort allocation of the Dutch beam trawl fleet in response to a temporarily closed area in the North Sea, *Ices Cm* 2001/N: 01 (2001) 1–17.

- [3] T. A. Dinmore, D. E. Duplisea, B. D. Rackham, D. L. Maxwell, S. Jennings, Impact of a large-scale area closure on patterns of fishing disturbance and




Figure 5: vessel movement

the consequences for benthic communities, *ICES Journal of Marine Science* 60 (2) (2003) 371–380. doi:10.1016/S1054-3139(03)00010-9.

- [4] J. Lee, A. B. South, S. Jennings, Developing reliable, repeatable, and accessible methods to provide high-resolution estimates of fishing-effort distributions from vessel monitoring system (VMS) data, *ICES Journal of Marine Science* 67 (6) (2010) 1260–1271. doi:10.1093/icesjms/fsq010.

- [5] F. Bastardie, J. R. Nielsen, C. Ulrich, J. Egekvist, H. Degel, Detailed mapping of fishing effort and landings by coupling fishing logbooks with






Figure 6: vessel movement multi

satellite-recorded vessel geo-location, Fisheries Research 106 (1) (2010) 41–

53. doi:10.1016/j.fishres.2010.06.016.

URL <http://dx.doi.org/10.1016/j.fishres.2010.06.016>

- [6] H. D. Gerritsen, C. Lordan, C. Minto, S. B. M. Kraak, Spatial patterns in the retained catch composition of Irish demersal otter trawlers: High-resolution fisheries data as a management tool, Fisheries Research 129-130 (2012) 127–136. doi:10.1016/j.fishres.2012.06.019.

URL <http://dx.doi.org/10.1016/j.fishres.2012.06.019>




Figure 7: vessel movement value

- [7] M. Mateo, L. Pawlowski, M. Robert, Highly mixed fisheries: fine-scale spatial patterns in retained catches of French fisheries in the Celtic Sea, ICES Journal of Marine Science: Journal du Conseil (2016) fsw129doi:10.1093/icesjms/fsw129.

275

URL <http://icesjms.oxfordjournals.org/lookup/doi/10.1093/icesjms/fsw129>

- [8] R. B. Deriso, Harvesting Strategies and Parameter Estimation for an Age-Structured Model, Canadian Journal of Fisheries and Aquatic Sci-




Figure 8: fleet movement




Figure 9: spatial catch composition

280 ences 37 (2) (1980) 268–282. [arXiv:arXiv:1410.7455v3](https://arxiv.org/abs/1410.7455v3), doi:10.1139/f80-034.




Figure 10: spatial catch composition with clustering

- [9] J. Schnute, A genera theory for analysis of catch and effort data, Canadian Journal of Fisheries and Aquatic Sciences 42 (1970) (1985) 414–429. doi: 10.1139/f85-057.
- 285 [10] C. M. Dichmont, A. E. Punt, A. Deng, Q. Dell, W. Venables, Application of a weekly delay-difference model to commercial catch and effort data for tiger prawns in Australia ’ s Northern Prawn Fishery, Fisheries Research 65 (2003) 335–350. doi:10.1016/j.fishres.2003.09.024.
- [11] R. Hilborn, C. Walters, Quantitative fisheries stock assessment: Choice, dynamics and uncertainty, Vol. 2, 1992. arXiv:arXiv:1011.1669v3, doi: 10.1007/BF00042883.
- 290 [12] R. J. Beverton, S. J. Holt, On the Dynamics of Exploited Fish Populations (1957) 533doi:10.1007/978-94-011-2106-4.
- [13] W. E. Ricker, Stock and recruitment, Journal of the Fisheries Research Board of Canada 11 (5) (1954) 559 – 623.
- 295 [14] P. J. Diggle, P. J. Ribeiro, Model-based Geostatistics (Springer Series in Statistics), Vol. 1, 2007.




Figure 11: survey index

URL <http://www.amazon.es/Model-based-Geostatistics-Springer-Series-Statistics/dp/0387329072>

- 300 [15] T. Hutton, S. Mardle, S. Pascoe, R. a. Clark, Modelling fishing location choice within mixed fisheries: English North Sea beam trawlers in 2000 and 2001, ICES Journal of Marine Science 61 (8) (2004) 1443–1452. doi: 10.1016/j.icesjms.2004.08.016.
- [16] A. N. Tidd, T. Hutton, L. T. Kell, J. L. Blanchard, Dynamic prediction  
305 of effort reallocation in mixed fisheries, Fisheries Research 125-126 (2012)




Figure 12: survey spatial abundance

243–253. doi:10.1016/j.fishres.2012.03.004.

URL <http://dx.doi.org/10.1016/j.fishres.2012.03.004>

- [17] R. Girardin, Y. Vermard, O. Thébaud, A. Tidd, P. Marchal, Predicting fisher response to competition for space and resources in a mixed demersal fishery, *Ocean & Coastal Management* 106 (2015) 124–135. doi:10.1016/j.ocecoaman.2015.01.017.

URL <http://linkinghub.elsevier.com/retrieve/pii/S0964569115000265>




Figure 13: Realised step function

- [18] E. A. Codling, M. J. Plank, S. Benhamou, J. R. S. Interface, Random walk  
 315 models in biology., *Journal of the Royal Society, Interface / the Royal Society* 5 (25) (2008) 813–34. doi:10.1098/rsif.2008.0014.  
 URL [http://www.pubmedcentral.nih.gov/articlerender.fcgi?](http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2504494&tool=pmcentrez&rendertype=abstract)  
[artid=2504494&tool=pmcentrez&rendertype=abstract](http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2504494&tool=pmcentrez&rendertype=abstract)
- [19] F. Bartumeus, M. G. E. Da Luz, G. M. Viswanatham, J. Catalan, Ani-  
 320 mal Search Strategies: A Quantitative Random Walk Analysis, *Ecological Society of America* 86 (11) (2005) 3078–3087. doi:10.1890/04-1806.
- [20] D. W. Sims, E. J. Southall, N. E. Humphries, G. C. Hays, C. J. A. Brad-  
 shaw, J. W. Pitchford, A. James, M. Z. Ahmed, A. S. Brierley, M. A.  
 Hindell, D. Morritt, M. K. Musyl, D. Righton, E. L. C. Shepard, V. J.  
 325 Wearmouth, R. P. Wilson, M. J. Witt, J. D. Metcalfe, Scaling laws of  
 marine predator search behaviour, *Nature* 451 (7182) (2008) 1098–U5.  
 doi:10.1038/nature06518.
- [21] S. Bertrand, A. Bertrand, R. Guevara-Carrasco, F. Gerlotto, Scale-  
 invariant movements of fishermen: The same foraging strategy as nat-  
 330 ural predators, *Ecological Applications* 17 (2) (2007) 331–337. doi:

10.1890/06-0303.

URL <http://www.ncbi.nlm.nih.gov/pubmed/17489242>

- 335 [22] T. Sakiyama, Y.-P. Gunji, Emergence of an optimal search strategy from a simple random walk., *Journal of the Royal Society, Interface* 10 (86) (2013) 20130486. doi:10.1098/rsif.2013.0486.  
URL <http://www.ncbi.nlm.nih.gov/pubmed/23804445>
- [23] D. W. Sims, N. E. Humphries, R. W. Bradford, B. D. Bruce, Lévy flight and Brownian search patterns of a free-ranging predator reflect different prey field characteristics, *Journal of Animal Ecology* 81 (2) (2012) 432–  
340 442. doi:10.1111/j.1365-2656.2011.01914.x.
- [24] D. M. Gillis, R. M. Peterman, Implications of interference among fishing vessels and the ideal free distribution to the interpretation of CPUE, *Canadian Journal of Fisheries and Aquatic Sciences* 55 (1) (1998) 37–46. doi:10.1139/f97-206.
- 345 [25] A. Rijnsdorp, Competitive interactions among beam trawlers exploiting local patches of flatfish in the North Sea, *ICES Journal of Marine Science* 57 (4) (2000) 894–902. doi:10.1006/jmsc.2000.0576.  
URL <http://icesjms.oxfordjournals.org/cgi/doi/10.1006/jmsc.2000.0576>
- 350 [26] J.-J. Poos, A. D. Rijnsdorp, An "experiment" on effort allocation of fishing vessels: the role of interference competition and area specialization, *Canadian Journal of Fisheries and Aquatic Sciences* 64 (2) (2007) 304–313. doi:10.1139/f06-177.