

Jessup, B., and J. Stamp. 2020. Development of indices of biotic integrity for assessing macroinvertebrate assemblages in Massachusetts freshwater Wadeable streams. Prepared for the Massachusetts Department of Environmental Protection.

Jessup, B., Block, B. and J. Stamp. 2021. Development of an Index of Biotic Integrity for macroinvertebrates in freshwater low gradient Wadeable streams in Massachusetts. Prepared for the Massachusetts Department of Environmental Protection.

Johnson, Z., S. Leibowitz and R. Hill. 2018. Revising the index of watershed integrity national maps. *Science of the Total Environment*. 10.1016/j.scitotenv.2018.10.112.

Karr, J.R. 1981. Assessment of biotic integrity using fish communities. *Fisheries*, 6(6):21-27.

Leppo, E.W., J. Stamp, and J. van Sickle. 2021. BioMonTools: Tools for Biomonitoring and Bioassessment. R package version 0.5.0.9039. <https://github.com/leppott/BioMonTools>.

Lussier, S.M., S.N. da Silva, M. Charpentier, J.F. Heltshe, S.M. Cormier, D.J. Klemm, M. Chintala, and S. Jayaraman. 2008. The influence of suburban land use on habitat and biotic integrity of coastal Rhode Island streams. *Environmental Monitoring and Assessment* 139:119 – 136.

MassDEP. 2004. CN 187.1. QAPP for 2004 Biological Monitoring and Habitat Assessment. Massachusetts Department of Environmental Protection, Division of Watershed Management. Worcester, MA. 99 p.

McKay, L., T. Bondelid, T. Dewald, J. Johnston, R. Moore, and A. Reah. 2012. NHDPlus Version 2: User Guide. Washington, DC: U.S. Environmental Protection Agency, Office of Water.

Merritt, R.W. and K.W. Cummins (editors). 1996. An introduction to the aquatic insects of North America, 3rd ed. Kendall/Hunt Publishing Company, Dubuque, Iowa.

Nuzzo, R. 2003. CN 32.2. Standard Operating Procedures: Water Quality Monitoring in Streams Using Aquatic Macroinvertebrates. Massachusetts Department of Environmental Protection, Division of Watershed Management. Worcester, MA. 35 p.

Poff, N. L.; J. D. Olden, N. K. M. Vieira, D. S. Finn, M. P. Simmons, and B. C. Kondratieff. 2006. Functional trait niches of North American lotic insects: traits-based ecological applications in light of phylogenetic relationships. *Journal of the North American Benthological Society* 25(4):730–755.

R Core Team. 2020. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/>

Roth, N. E., M. T. Southerland, J. C. Chaillou, J. H. Vølstad, S. B. Weisberg, H. T. Wilson, D. G. Heimbuch, and J. C. Seibel. 1997. Maryland Biological Stream Survey: Ecological status of non-tidal streams in six basins sampled in 1995. Maryland Department of Natural Resources, Chesapeake Bay and Watershed Programs, Monitoring and Non-tidal Assessment, Annapolis, Maryland. CBWP-MANTAEA-97-2.

Stoddard, J. L., A. T. Herlihy, D. V. Peck, R. M. Hughes, T. R. Whittier, and E. Tarquinio. 2008. A process for creating multimetric indices for large-scale aquatic surveys. *Journal of the North American Benthological Society*, 2008, 27(4):878–891

Tetra Tech. 2019. Sampling and Analysis Plan - Data Collection for Development of an Index of Biotic Integrity for Freshwater Low-Gradient Wadeable Streams in Southern New England. Prepared for the NEIWPCC, Lowell, MA. Prepared by Tetra Tech, Montpelier, VT.

Thornbrugh, D. J., S. G. Leibowitz, R. A. Hill, M. H. Weber, Z. C. Johnson, A. R. Olsen, J. E. Flotemersch, J. L. Stoddard, and D. V. Peck. 2018. Mapping watershed integrity for the conterminous United States. *Ecological Indicators*, 85:1133-1148.

U.S. Environmental Protection Agency (U.S. EPA). 2011. Level III and IV ecoregions of the continental United States. U.S. EPA, National Health and Environmental Effects Research Laboratory, Corvallis, Oregon, Map scale 1:3,000,000. Available online at: <https://www.epa.gov/eco-research/level-iii-and-iv-ecoregions-continental-united-states>.

U.S. Environmental Protection Agency (U.S. EPA). 2012. Implications of Climate Change for Bioassessment Programs and Approaches to Account for Effects. Global Change Research Program, National Center for Environmental Assessment, Washington, DC; EPA/600/R-11/036F. Available online: <https://cfpub.epa.gov/ncea/global/recordisplay.cfm?deid=239585>

U.S. Environmental Protection Agency (U.S. EPA). 2013. Biological Assessment Program Review: Assessing Level of Technical Rigor to Support Water Quality Management. EPA 820-R-13-001. U.S. Environmental Protection Agency, Washington, DC.

U.S. Environmental Protection Agency (U.S. EPA). 2016. Regional Monitoring Networks (RMNs) to detect changing baselines in freshwater wadeable stream. (EPA/600/R-15/280). Washington, DC: Office of Research and Development, Washington. Authorship: Jen Stamp, Anna Hamilton, Britta G. Bierwagen, Debbie Arnwine, Margaret Passmore and Jonathan Witt. Available online: <https://cfpub.epa.gov/ncea/global/recordisplay.cfm?deid=307973>

Yoder, C. O., and E. T. Rankin. 1995. Biological criteria program development and implementation in Ohio. In W. S. Davis & T. P. Simon (Eds.), *Biological assessment and criteria: tools for water resource planning and decision making* (pp. 109–144). Boca Raton: Lewis Publishers.

Yuan, L. 2006. Estimation and Application of Macroinvertebrate Tolerance Values. Report No. EPA/600/P-04/116F. National Center for Environmental Assessment, Office of Research and Development, U.S. Environmental Protection Agency, Washington, D.C.