Section 8.5: Congruence Modulo n

Juan Patricio Carrizales Torres

May 30, 2022

This chapter discusses the previously seen topic of **Congruence Modulo n**, but now with the lens of **Equivalence relations**. Basically, the author proved that every relation on \mathbb{Z} defined by the congruence modulo of some $n \geq 2$ is an equivalence relation with n equivalence classes. This follows from the **Division Algorithm**, namely in the case for $n \geq 2$, any integer m can be expressed uniquely as m = kn + r, where $k \in \mathbb{Z}$ and $0 \leq r < n$.

Another interesting idea is the logical equivalence between coditions that define equivalence relations. For example, let R_1 and R_2 be relations on some nonempty set defined by $a R_1 b$ if P(a,b) and $a R_2 b$ if Q(a,b). The fact that $P(a,b) \iff Q(a,b)$ for some other condition Q(n), implies that $R_1 = R_2$. Hence, one can show that two relations have the same distinct equivalence classes by just showing that there is a biconditional relation between the conditions that define them.

Problem 47. The relation R on \mathbb{Z} defined by a R b if $a^2 \equiv b^2 \pmod{4}$ is known to be an equivalence relation. Determine the distinct equivalence classes.

Solution 47. Let's first consider [0]. We know that

$$[0] = (x \in \mathbb{Z} : x R 0)$$

$$= (x \in \mathbb{Z} : x^2 = 4k, k \in \mathbb{Z})$$

$$= (x \in \mathbb{Z} : 4 \mid x^2) = (x \in \mathbb{Z} : 2 \mid x^2)$$

$$= (x \in \mathbb{Z} : 2 \mid x).$$

Hence, [0] is the set of all even integers. Now we are left with the odd ones, so let's check what are the elements of [1]. We know that

[1] =
$$(x \in \mathbb{Z} : x R 1)$$

= $(x \in \mathbb{Z} : x^2 - 1 = 4k, k \in \mathbb{Z}).$

Note that $x^2 - 1 = (x - 1)(x + 1)$ is an even integer. Thus, it is a necessary and sufficient condition that either $2 \mid (x-1)$ or $2 \mid (x+1)$. Therefore, x = 2a+1 or x = 2b-1 = 2(b-1)+1 for $a, b \in \mathbb{Z}$, and so [1] is the set of odd integers.