

# $K^+$ Pathway for Code version 2.0

Tim David

January 17, 2018

# Todo list

	I have kept this in to make code easier to understand. It has no explicit parameters	
	in it for the $K^+$ pathway . . . . .	9
	what about diffusion into the ECS for both $Na^+$ and $K^+$ ? . . . . .	11
5	what are the definitions and values of $K_{Na_k}$ , $K_{K_s}$ . . . . .	13
	why do we have $\frac{F_{KIR_i}}{\gamma_i}$ when they have both the same dimensions but one value	
	$F_{KIR_i}$ is 750 and the other $\gamma_i$ is 1970 ? . . . . .	15
	We should note here that the membrane potential coupling $V_{coupling_i}^{SMC-EC}$ is an approx-	
10	imation that assumes the gradient of concentrations is negligible and hence	
	only the membrane potential diffusion term is non-zero determined from the	
	electro-diffusion theory. . . . .	16
	diffusion, NaK and BK fluxes defined here . . . . .	17

# Chapter 1

## Notes for reading

15 The following notes, definitions and equations provide the reader with a comprehensive guide to version 1.2 of NVU. The document is set out in sections where each section contains the equations for each compartment, namely neuron, synaptic cleft, astrocyte, perivascular space, smooth muscle cell, endothelial cell, extracellular space and finally the lumen. The reader will find multiple definitions and equations but by dividing the  
20 document into sections corresponding to compartments it is hoped that a more clear understanding is obtained. Concentrations as written on the left-hand-side of the o.d.e. are given by the notation of  $N_j$  where  $j$  can be any species such as  $\text{Na}^+$  or  $\text{Ca}^{2+}$ . True concentrations are written with square brackets as in  $[\text{Ca}^{2+}]_n$ . In point of fact they are equivalent. Subscripts on variable such as concentrations denote the compartment,  
25  $n$ =neuron,  $k$ =astrocyte,  $s$ =synaptic cleft,  $i$ =smooth muscle cell,  $j$ =endothelial cell,  $e$  = extracellular space. Concentrations with "hats" denote those in the ER/SR stores.

### 1.0.1 Version 1.2 /2.0 difference

The basic difference between version 1.2 and version 2.0 is the new neuron model. This is based on the work of Chang et al [1] and that of Kager et al [9]. For this version the neuron  
30 model has 4 compartments; i) soma/axon, ii) dendrite, iii) post synaptic terminal and extracellular space (ECS). Ion channels for  $\text{Na}^+$ , and  $\text{K}^+$  efflux into the ECS.  $\text{K}^+$  is buffered in the ECS and a portion of the  $\text{K}^+$  flux is passed into the synaptic cleft compartment. On reaching a certain concentration of  $\text{K}^+$  in the synaptic cleft glutamate is pumped into the synaptic cleft. This glutamate is taken up by both the post-synaptic neuron and the  
35 astrocyte. The neuron is stimulated by injection of a current of specified value into the soma/axon compartment.

Schematic of the full set of pathways is shown in Figure ??

## 1.1 Global Constants

$F$	Faraday's constant	96500 C mole <sup>-1</sup>
$T$	Temperature	300 K
$R_{gas}$	Gas constant	8.315 J mole K <sup>-1</sup>

## Chapter 2

# State variables, initial values and parameter values

In the actual Matlab code the state variables are defined as follows

- $v_{sa}$  : membrane potential of soma/axon, mV
- $v_d$  : membrane potential of dendrite, mV
- $K_{sa}$  : K+ concentration of soma/axon, mM
- $K_d$  : K+ concentration of dendrite, mM
- $Na_d$  : Na+ concentration of dendrite, mM
- $K_e$  : K+ concentration of ECS, mM
- $Na_e$  : Na+ concentration of ECS, mM
- $Buf f_e$  : Buffer concentration for K+ buffering in ECS, mM

### Gating variables

- m1 : Activation gating variable, soma/axon NaP channel (Na+) m2 : Activation gating variable, soma/axon KDR channel (K+) m3 : Activation gating variable, soma/axon KA channel (K+) m4 : Activation gating variable, dendrite NaP channel (Na+) m5 : Activation gating variable, dendrite NMDA channel (Na+) m6 : Activation gating variable, dendrite KDR channel (K+) m7 : Activation gating variable, dendrite KA channel (K+) m8 : Activation gating variable, soma/axon NaT channel (Na+)
- h1 : Inactivation gating variable, soma/axon NaP channel (Na+) h2 : Inactivation gating variable, soma/axon KA channel (K+) h3 : Inactivation gating variable, dendrite NaP channel (Na+) h4 : Inactivation gating variable, dendrite NMDA channel (Na+) h5 : Inactivation gating variable, dendrite KA channel (K+) h6 : Inactivation gating variable, soma/axon NaT channel (Na+)

## NO pathway

65

- $Ca_n$  :  $Ca^{2+}$  in the post-synaptic neuron  $nNOS_{actn}$  : activated NOS in the post-synaptic neuron  $NO_n$  : Nitric Oxide in the post-synaptic neuron

Table 2.1: Initial resting values and other parameter values of the neuron model, from Chang et al[1]

Parameters	Values	Units	Description
$v_m$	-70	$mV$	membrane potential
$[K^+]_e$	3.5	$mM$	extracellular space potassium ion concentration
$[K^+]_i$	133.5	$mM$	intracellular potassium ion concentration of neuron
$[Na^+]_e$	140	$mM$	extracellular space sodium ion concentration
$[Na^+]_i$	10	$mM$	intracellular sodium ion concentration of neuron
$[O_2]_0$	$2 \times 10^{-2}$	$mM$	baseline concentration of oxygen in the tissue
$B_0$	0.9	$ml/100mg/s$	baseline cerebral blood flow
$[O_2]_b$	$4 \times 10^{-2}$	$mM$	blood oxygen concentration
$J_0$	$2.5 \times 10^{-2}$	$mM/s$	steady state change in oxygen concentration due to cerebral blood flow
$R_a$	$1.83 \times 10^5$	$\Omega$	input resistance of dendritic tree
$\delta_d$	$4.5 \times 10^{-2}$	$cm$	half-length of dendrite
$A_s$	$1.586 \times 10^{-5}$	$cm^2$	surface area of soma
$A_d$	$2.6732 \times 10^{-4}$	$cm^2$	surface area of dendrite
$V_s$	$2.160 \times 10^{-9}$	$cm^3$	volume of soma
$V_d$	$5.614 \times 10^{-9}$	$cm^3$	volume of dendrite
$S_e$	$4.1179 \times 10^{-6}$	$cm$	volume to surface area ratio of the extracellular space
$C_m$	$7.5 \times 10^{-5}$	$s/\Omega cm^2$	membrane capacitance
$I_{max}$	$1.48 \times 10^{-3}$	$mA/cm^2$	$Na^+/K^+$ -ATPase rate
$D_{Na^+}$	$1.33 \times 10^{-5}$	$cm^2/s$	Sodium diffusion coefficient
$D_{K^+}$	$1.96 \times 10^{-5}$	$cm^2/s$	Potassium diffusion coefficient
$D_{Cl^-}$	$2.03 \times 10^{-5}$	$cm^2/s$	Chlorine diffusion coefficient

Table 2.2: Rate expressions and parameter values used in the voltage dependent channel currents of the neuron model, from Chang et al[1]

<b>Currents</b> $mA/cm^2$	$g_{Ion,GHK}$ $mAcm$	<b>Gates</b> $m^p h^q$	<b>Voltage dependent rate functions</b>
$I_{Na,P}$	$2 \times 10^{-6}$	$m^2 h$	$\alpha_m = \frac{1}{6(1+\exp[-(0.143E_m+5.67)])}$ $\beta_m = \frac{\exp[-(0.143E_m+5.67)]}{6(1+\exp[-(0.143E_m+5.67)])}$ $\alpha_h = 5.12 \times 10^{-8} \exp[-(0.056E_m + 2.94)]$ $\beta_h = \frac{1.6 \times 10^{-6}}{1+\exp[-(0.2E_m+1.25)]}$
$I_{K,DR}$	$10 \times 10^{-5}$	$m^2$	$\alpha_m = 0.016 \frac{E_m+34.9}{1-\exp[-(0.2E_m+6.98)]}$ $\beta_m = 0.25 \exp[-(0.25E_m + 1.25)]$
$I_{K,A}$	$1 \times 10^{-5}$	$m^2 h$	$\alpha_m = 0.02 \frac{E_m+56.9}{1-\exp[-(0.1E_m+5.69)]}$ $\beta_m = 0.0175 \frac{E_m+29.9}{\exp(0.1E_m+2.99)-1}$ $\alpha_h = 0.016 \exp[-(0.056E_m + 4.61)]$ $\beta_h = \frac{0.5}{1+\exp[-(0.2E_m+11.98)]}$
$I_{NMDA}$	$1 \times 10^{-5}$	$mh$	$\alpha_m = \frac{0.5}{1+\exp\left(\frac{13.5-[K^+]}{1.42}\right)}$ $\beta_m = 0.5 - \alpha_m$ $\alpha_h = \frac{1}{2000\left(1+\exp\left[\frac{[K^+]e^{-6.75}}{0.71}\right]\right)}$ $\beta_h = 5 \times 10^{-5} - \alpha_h$

## Chapter 3

# Equations for each compartment

### 70 3.1 Neuron

#### 3.1.1 Nernst potential for Na,K ions in soma and dendrite (Cl constant)

$$E_{K_{sa}} = \frac{RT}{F} \ln\left(\frac{K_e}{K_{sa}}\right) \quad (3.1.1)$$

$$E_{K_d} = \frac{RT}{F} \ln\left(\frac{K_e}{K_d}\right) \quad (3.1.2)$$

#### 3.1.2 Leak fluxes of Na,K,Cl in soma and dendrite using HH

$$J_{Kleak_{sa}} = g_{Kleak_{sa}}(v_{sa} - E_{K_{sa}}) \quad (3.1.3)$$

$$J_{Kleak_d} = g_{Kleak_d}(v_d - E_{K_d}) \quad (3.1.4)$$

$$(3.1.5)$$

$$J_{Kleak_{sa}} = p.gKleak_{sa} * (v_{sa} - E_{K_{sa}}); \quad J_{Kleak_d} = p.gKleak_d * (v_d - E_{K_d});$$

#### 3.1.3 Dendrite (with subscript d)

### 75 Na flux through NaP channel in dendrite using GHK



$$m4_{\alpha} = \frac{1}{6(1 + \exp(-((0.143v_d) + 5.67)))} \quad (3.1.6)$$

$$m4_{\beta} = \frac{\exp(-((0.143v_d) + 5.67))}{6(1 + \exp(-((0.143v_d) + 5.67)))} \quad (3.1.7)$$

$$h3_{\alpha} = \frac{5.12e - 8 \exp(-((0.056v_d) + 2.94))}{1.6e - 6} \quad (3.1.8)$$

$$h3_{\beta} = \frac{1.6e - 6}{1 + \exp(-((0.2 * v_d) + 8))} \quad (3.1.9)$$

$$J_{NaPd} = (m4^2 h3 g_{NaP} F v_d \frac{(Na_d - (\exp(\frac{-v_d F}{RT}) Na_e))}{(\frac{RT}{F} (1 - \exp(\frac{-v_d F}{RT})))}) \quad (3.1.10)$$

$$(3.1.11)$$

$$J_{NMDA_{K_d}} = M(v, Mg) ((m5 h4 g_{NMDA} F v_d \frac{(K_d - (\exp(\frac{v_d F}{RT}) K_e))}{(\frac{RT}{F} (1 - \exp(\frac{-v_d F}{RT})))}) \quad (3.1.12)$$

$$M(v, Mg) = \frac{1}{(1 + 0.33)} Mg \exp(-((0.07v_d + 0.7))) \quad (3.1.13)$$

$$J_{NMDA_{K_d}} = ((m5.*h4.*p.g_{NMDA_G} Hk * p.Farad.*v_d.*(K_d - (\exp(-v_d/p.ph).*K_e)))./(p.ph*(1 - \exp(-v_d/p.ph))))$$

**K flux through KDR channel in dendrite using GHK**

$$m6_{\alpha} = \frac{0.016((v_d + 34.9))}{(1 - \exp(-((0.2 * v_d) + 6.98)))} \quad (3.1.14)$$

$$m6_{\beta} = \frac{0.25 \exp(-((0.025 * v_d) + 1.25))}{1} \quad (3.1.15)$$

$$J_{KDR_d} = m6^2 g_{KDR} F v_d \frac{(K_d - (\exp(\frac{-v_d F}{RT}) K_e))}{(\frac{RT}{F} (1 - \frac{v_d F}{RT}))} \quad (3.1.16)$$

**K flux through KA channel in dendrite using GHK**

$$m7_{\alpha} = \frac{0.02((v_d + 56.9))}{(1 - \exp(-((0.1v_d) + 5.69)))} \quad (3.1.17)$$

$$m7_{\beta} = \frac{0.0175((v_d + 29.9))}{(\exp(((0.1 * v_d) + 2.99)) - 1)} \quad (3.1.18)$$

$$h5_{\alpha} = \frac{0.016 \exp(-((0.056v_d) + 4.61))}{1} \quad (3.1.19)$$

$$h5_{\beta} = \frac{0.5}{(1 + \exp(-((0.2 * v_d) + 11.98)))} \quad (3.1.20)$$

$$J_{KA_d} = m7^2 h5 g_{KA} F v_d \frac{(K_d - (\exp(\frac{-v_d F}{RT}) K_e))}{(\frac{RT}{F} (1 - \frac{v_d F}{RT}))} \quad (3.1.21)$$

### 3.1.4 Soma/Axon (with subscript sa)

80 **K flux through KDR channel in soma using GHK**

$$m2_{\alpha} = \frac{0.016((v_{sa} + 34.9))}{(1 - \exp(-((0.2v_{sa}) + 6.98))))} \quad (3.1.22)$$

$$m2_{\beta} = 0.25 \exp(-((0.025v_{sa}) + 1.25)) \quad (3.1.23)$$

$$J_{KDR_{sa}} = m2^2 g_{KDR} F v_{sa} \frac{(K_{sa} - (\exp(\frac{-v_{sa}F}{RT})K_e))}{(\frac{RT}{F}(1 - \frac{-v_{sa}F}{RT}))} \quad (3.1.24)$$

$$(3.1.25)$$

**K flux through KA channel in soma using GHK input current**

$$m3_{\alpha} = \frac{0.02(v_{sa} + 56.9)}{(1 - \exp(-((0.1v_{sa}) + 5.69))))} \quad (3.1.26)$$

$$m3_{\beta} = \frac{0.0175(v_{sa} + 29.9)}{(\exp(0.1v_{sa} + 2.99) - 1))} \quad (3.1.27)$$

$$h2_{\alpha} = 0.016 \exp(-((0.056v_{sa} + 4.61))) \quad (3.1.28)$$

$$h2_{\beta} = \frac{0.5}{1 + \exp(-((0.2v_{sa} + 11.98)))} \quad (3.1.29)$$

$$J_{KA_{sa}} = m3^2 h2 g_{KA} F v_{sa} \frac{(K_{sa} - (\exp(\frac{-v_{sa}F}{RT})K_e))}{(\frac{RT}{F}(1 - \frac{-v_{sa}F}{RT}))} \quad (3.1.30)$$

$$(3.1.31)$$

**flux through the NaK-ATPase pump**

$$J_{pump1_{sa}} = (1 + (\frac{K_{init_e}}{K_e}))^{-2} (1 + (\frac{Na_{init_{sa}}}{Na_{sa}}))^{-3} \quad (3.1.32)$$

$$J_{pump1_{init_{sa}}} = 0.0312 \quad (3.1.33)$$

$$J_{pump1_d} = (1 + (\frac{K_{init_e}}{K_e}))^{-2} (1 + (\frac{Na_{init_d}}{Na_d}))^{-3} \quad (3.1.34)$$

$$J_{pump1_{init_d}} = 0.0132 \quad (3.1.35)$$

$$(3.1.36)$$

### 3.1.5 Total ion fluxes

I have kept this in to make code easier to understand. It has no explicit parameters in it for the  $K^+$  pathway

85 **Total ion fluxes in soma**

$$J_{Na_{tot_{sa}}} = J_{NaP_{sa}} + J_{Na_{leak_{sa}}} + J_{Na_{pump_{sa}}} + J_{NaT_{sa}} \quad (3.1.37)$$

$$J_{K_{tot_{sa}}} = J_{KDR_{sa}} + J_{KA_{sa}} + J_{K_{leak_{sa}}} + J_{K_{pump_{sa}}} \quad (3.1.38)$$

$$J_{leak_{tot_{sa}}} = g_{leak_{sa}}(v_{sa} - E_{Cl_{sa}}) \quad (3.1.39)$$

$$(3.1.40)$$

### Total ion fluxes in dendrite

$$J_{Na_{tot_d}} = J_{NaP_d} + J_{Na_{leak_d}} + J_{Na_{pump_d}} + J_{Na_{NM DA_d}} \quad (3.1.41)$$

$$J_{K_{tot_d}} = J_{KDR_d} + J_{KA_d} + J_{K_{leak_d}} + J_{K_{pump_d}} + J_{K_{NM DA_d}} \quad (3.1.42)$$

$$J_{leak_{tot_d}} = g_{leak_d}(v_d - E_{Cl_d}) \quad (3.1.43)$$

$$(3.1.44)$$

### Total ion fluxes in soma and dendrite

$$J_{tot_{sa}} = J_{Na_{tot_{sa}}} + J_{K_{tot_{sa}}} + J_{leak_{tot_{sa}}} \quad (3.1.45)$$

$$J_{tot_d} = J_{Na_{tot_d}} + J_{K_{tot_d}} + J_{leak_{tot_d}} \quad (3.1.46)$$

$$(3.1.47)$$

### Tissue oxygen

$$J_{pump2_0} = 0.0952 \quad (3.1.48)$$

$$J_{pump2_{O2_0}} = 1 \quad (3.1.49)$$

$$CBF = CBF_{init} \frac{R^4}{R_{init}^4} \quad (3.1.50)$$

$$(3.1.51)$$

**Note** The pump functions could look like this

$$J_{pump2_0} = 2 \left( 1 + \frac{O2_0}{(((1-\alpha_{O2})O2_0) + \alpha_{O2}O2_0))} \right)^{-1} \quad (3.1.52)$$

$$J_{pump2_{O2_0}} = 2 * (1 + O2_0./(((1 - p.alpha_O2) * p.O2_0) + p.alpha_O2 * p.O2_0)).^{-1}$$

## 3.2 Conservation equations for neuron compartment

change in concentration of Na,K in the soma

$$\frac{dK_{sa}}{dt} = \frac{-A_s}{FV_s} J_{K_{tot_{sa}}} + \frac{D_k(V_d + V_s)}{2dhod^2 V_s} (K_d - K_{sa}) \quad (3.2.1)$$

$$(3.2.2)$$

95 change in concentration of Na,K in the dendrite

$$\frac{dK_d}{dt} = \frac{-A_d}{FV_d} J_{K_{tot_{sa}}} + \frac{D_k(V_d + V_s)}{2dhod^2 V_s} (K_{sa} - K_d) \quad (3.2.3)$$

$$(3.2.4)$$

## 3.3 Extra Cellular Space (ECS with subscript e)

change in buffer for K+ in the extracellular space

$$\frac{dBuf f_e}{dt} = \frac{\mu K_e (B_0 - Buf f_e)}{1 + \exp(-((K_e - 5.5)/1.09))} - \mu Buf f_e \quad (3.3.1)$$

$$(3.3.2)$$

change in concentration of Na,K in the extracellular space

$$\frac{dK_e}{dt} = \frac{1}{F \textcolor{red}{f}_e} \left( \frac{\textcolor{red}{A}_s J_{K_{tot_{sa}}}}{V_s} + \frac{\textcolor{red}{A}_d J_{K_{tot_d}}}{V_d} \right) \quad (3.3.3)$$

$$(3.3.4)$$

what about diffusion into the ECS for both Na<sup>+</sup> and K<sup>+</sup>?

K<sup>+</sup> concentration in the SC

$$\frac{dN_{K_s}}{dt} = k_C f(t) - \frac{dN_{K_k}}{dt} + J_{BK_k} + \frac{R_s}{\tau_s} \{ [K^+]_e - [K^+]_s \} \quad (3.3.5)$$

$\tau_s$  is defined in equation 3.8.5 and has a value of 2.8 secs.  $[K^+]_e$  is the potassium concentration in the ECS and  $[K^+]_s$  is the potassium concentration in the synaptic cleft. Na<sup>+</sup> concentration in the SC

$$\frac{dN_{Na_s}}{dt} = -k_C f(t) - \frac{dN_{Na_k}}{dt} \quad (3.3.6)$$

HCO<sub>3</sub> concentration in the SC

$$\frac{dN_{HCO_{3s}}}{dt} = - \frac{dN_{HCO_{3k}}}{dt} \quad (3.3.7)$$

100

Cl concentration in the synaptic cleft is derived by assuming electro-neutrality:

$$[Cl^{-1}]_s = [Na^+]_s + [K^+]_s - [HCO_3^{-1}]_s \quad (3.3.8)$$

### 3.4 Astrocyte (with subscript k)

K<sup>+</sup> concentration in the AC :

$$\frac{dN_{K_k}}{dt} = -J_{K_k} + 2J_{NaK_k} + J_{NKCC1_k} + J_{KCC1_k} - J_{BK_k} \quad (3.4.1)$$

K<sup>+</sup> flux through the Ca<sup>2+</sup> mediated BK channel :

$$J_{BK_k} = \frac{\textcolor{red}{g}_{BK_k}}{F} w_k (v_k - E_{BK_k}) \quad (3.4.2)$$

$g_{BK_k}$  is evaluated in the code using

`g_BK_k = p.G_BK_k*1e-12/p.A_ef_k; the_value = p.G_BK_k`

105

is 225 (as listed in the parameter list in Astrocyte.m)

---

<sup>1</sup>model estimation

$V_{\text{spine}}$	dendritic spine volume	$8 \times 10^{-8}$ nL	[15]
$\kappa_{\text{ex}}$	decay rate constant of internal $\text{Ca}^{2+}$ concentration	$1.6 \times 10^3$ s $^{-1}$	[15]
$[\text{Ca}^{2+}]_{\text{rest}}$	resting internal calcium concentration	0.1 $\mu\text{M}$	[15]
$\lambda_{\text{buf}}$	buffer capacity	20 (dim.less)	[15]
$V_{\text{max,nNOS}}$	maximum nNOS activation rate	$25 \times 10^{-3}$ $\mu\text{M}$	M.E. <sup>1</sup>
$K_{\text{m,nNOS}}$	Michaelis constant	$9.27 \times 10^{-2}$	[8]
$\mu_{\text{deact},n}$	rate constant at which nNOS is deactivated	0.0167 s $^{-1}$	[4]
$K_{\text{m},A}$	Michaelis constant	650 $\mu\text{M}$	[15]
$K_{\text{m},B}$	Michaelis constant	2800 $\mu\text{M}$	[15]
$v_n$	neuronal membrane potential	-0.04 V	M.E. but see Kager et al model maybe -0.05 -or -0.06 V
$G_{\text{M}}$	conductance of NMDA receptor	$4.6 \times 10^4$ fS	[15]
$P_{\text{Ca}}/P_{\text{M}}$	ratio of NMDA receptor permeability to $\text{Ca}^{2+}$ to permeability to monovalent ions	3.6 (dim.less)	[15]
$[\text{Ca}^{2+}]_{\text{ex}}$	external calcium concentration	$2 \times 10^3$ $\mu\text{M}$	[15]
$[\text{M}]$	concentration of monovalent ions	$1.3 \times 10^5$ $\mu\text{M}$	[15]
$\alpha_v$	voltage-dependent $\text{Mg}^{2+}$ block parameter	-80 V $^{-1}$	[15]
$\beta_v$	voltage-dependent $\text{Mg}^{2+}$ block parameter	0.02 V	[15]
$n_{\text{NR2},A}$	average number of NR2A NMDA receptors	0.63 (dim.less)	[15]
$n_{\text{NR2},B}$	average number of NR2A NMDA receptors	11 (dim.less)	[15]
$Q_1$	$\text{Ca}^{2+}$ -CaM binding constant	$1.9 \times 10^5$ $\mu\text{M}^{-1}$	[5]
$Q_2$	$\text{Ca}^{2+}$ -CaM binding constant	$2.1 \times 10^5$ $\mu\text{M}^{-1}$	[5]
$Q_3$	$\text{Ca}^{2+}$ -CaM binding constant	$0.4 \times 10^5$ $\mu\text{M}^{-1}$	[5]
$Q_4$	$\text{Ca}^{2+}$ -CaM binding constant	$0.26 \times 10^5$ $\mu\text{M}^{-1}$	[5]
$V_{\text{max,NO},n}$	maximum catalytic rate of neuronal NO production	4.22 s $^{-1}$	[2]
$[\text{O}_2]_n$	$\text{O}_2$ concentration in the neuron	200 $\mu\text{M}$	M.E.
$K_{\text{m,O}_2,n}$	Michaelis constant for nNOS for $\text{O}_2$	243 $\mu\text{M}$	[3]
$[\text{L-Arg}]_n$	L-Arg concentration in the neuron	100 $\mu\text{M}$	[3]
$K_{\text{m,L-Arg},n}$	Michaelis constant for nNOS for L-Arg	1.5 $\mu\text{M}$	[2]
$k_{\text{O}_2,n}$	$\text{O}_2$ reaction rate constant	$9.6 \times 10^{-6}$ $\mu\text{M}^{-2}$ s $^{-1}$	[10]
$x_{nk}$	distance between centres of NE and AC	25 $\mu\text{m}$	M.E.
$D_{\text{c,NO}}$	NO diffusion coefficient	3300 $\mu\text{m}^2\text{s}^{-1}$	[12]

$k_C$	Input scaling parameter	$7.35 \times 10^{-5} \mu\text{M m s}^{-1}$	[14]
-------	-------------------------	--	------

Open probability of the BK channel ( $\text{s}^{-1}$ ):

$$\frac{dw_k}{dt} = \phi_w (w_\infty - w_k) \quad (3.4.3)$$

$$\phi_w = \psi_n \cosh\left(\frac{v_k - v_3}{2v_4}\right) \quad (3.4.4)$$

$$v_3 = -\frac{v_5}{2} \tanh\left[\frac{[\text{Ca}^{2+}]_k - Ca_3}{Ca_4}\right] + v_6 \quad (3.4.5)$$

$\psi_n$	characteristic time scale for BK channel	$2.664 \text{ s}^{-1}$
$v_4$	measure of the spread of $w_\infty$	8 millivolts
$v_5$	shift in $w_\infty$ as a function of $\text{Ca}^{2+}$	15 millivolts
$v_6$	BK open probability constant	-55 millivolts
$Ca_3$	BK open probability constant	0.4 $\mu\text{M}$
$Ca_4$	BK open probability constant	0.35 $\mu\text{M}$
$EET_{shift}$	EET dependent voltage shift	2 mV $M^{-1}$

**Equilibrium state BK-channel as a function of the concentration of EET in the astrocytic cytosol:**

$$w_\infty = 0.5 \left( 1 + \tanh \left( \frac{v_k EET_{shift} [EET]_k - v_3}{v_4} \right) \right) \quad (3.4.6)$$

### 3.4.1 Fluxes into and out of the astrocyte

$\text{K}^+$  flux

$$J_{K_k} = \frac{gK_k}{F} (v_k - E_{K_k}) \quad (3.4.7)$$

$\text{Cl}$  and  $\text{K}^+$  flux through the KCC1 channel

$$J_{KCC1_k} = C_{input} \frac{gKCC1_k}{F} \frac{R_g T}{F} \ln \left( \frac{K_s Cl_s}{K_k Cl_k} \right) \quad (3.4.8)$$

$\text{Na}^+$ ,  $\text{K}^+$  and  $\text{Cl}$  flux through the NKCC1 channel

$$J_{NKCC1_k} = C_{input} \frac{gNKCC1_k}{F} \frac{R_g T}{F} \ln \left( \frac{Na_s K_s Cl_s^2}{Na_k K_k Cl_k^2} \right) \quad (3.4.9)$$

Flux through the sodium potassium pump

$$J_{NaK_k} = J_{NaK_{max}} \frac{Na_k^{1.5}}{Na_k^{1.5} + K_{Na_k}^{1.5}} \frac{K_s}{K_s + K_{K_s}} \quad (3.4.10)$$

what are the definitions and values of  $K_{Na_k}$ ,  $K_{K_s}$

$F$	Faraday's constant	$9.649 \times 10^4 \text{ C mol}^{-1}$	
$R_g$	Gas constant	$8.315 \text{ J mol}^{-1} \text{ K}^{-1}$	
$T$	Temperature	300 K	
$g_{K_k}$	Specific ion conductance of potassium	$40 \times 10^3 \text{ } \Omega^{-1} \text{ m}^{-2}$	[14]
$g_{Na_k}$	Specific ion conductance of sodium	$1.314 \times 10^3 \text{ } \Omega^{-1} \text{ m}^{-2}$	[14]
$K_{Na_k}$		$40 \times 10^3 \text{ } \Omega^{-1} \text{ m}^{-2}$	[14]
$g_{Na_k}$	Specific ion conductance of sodium	$1.314 \times 10^3 \text{ } \Omega^{-1} \text{ m}^{-2}$	[14]
$g_{NBC_k}$	Specific ion conductance of the NBC cotransporter	$7.57 \times 10^2 \text{ } \Omega^{-1} \text{ m}^{-2}$	[14]
$g_{KCC1_k}$	Specific ion conductance of the KCC1 cotransporter	$10 \text{ } \Omega^{-1} \text{ m}^{-2}$	[14]
$g_{NKCC1_k}$	Specific ion conductance of the NKCC1 cotransporter	$55.4 \text{ } \Omega^{-1} \text{ m}^{-2}$	[14]
$J_{NaK_{max}}$	Maximum flux through the NaKATPase pump	$1.42 \times 10^{-3} \text{ } \mu\text{M ms}^{-1}$	[14]
$g_{BK_k}$	Specific ion conductance of the BK channel	$1.16 \times 10^3 \text{ } \Omega^{-1} \text{ m}^{-2}$	[7]
$C_{input}$	Block function to switch the channel on and off	0 ; 1 [-]	

### 3.5 Perivascular Space (with subscript p)

$\text{K}^+$  concentration in the PVS (in  $\mu\text{M}$ ):

$$\frac{dK_p}{dt} = \frac{J_{BK_k}}{R_k R_{pa}} + \frac{J_{KIR_i}}{R_{ps}} + \frac{J_{TRPV_k}}{R_k R_{pa}} \quad (3.5.1)$$

115 The ODE for the PVS  $\text{Ca}^{2+}$  concentration is

$$\frac{dCa_p}{dt} = -\frac{J_{TRPV_k}}{V R_{pa}} + \frac{J_{VOCC_i}}{V R_{ps}} - Ca_{decay_k} (Ca_p - Ca_{min_k}) \quad (3.5.2)$$

$$(3.5.3)$$

$R_{pa}$	Volume ratio of PVS to AC	$10^{-3} \text{ [-]}$	[13]
$R_{ps}$	Volume ratio of PVS to SMC	$10^{-3} \text{ [-]}$	[13]
$Ca_{decay_k}$	Rate of decay of $\text{Ca}^{2+}$ in the PVS	$0.5 \text{ s}^{-1}$	
$Ca_{min_k}$	steady state value of $\text{Ca}^{2+}$ in PVS	2 mM	

### 3.6 Smooth Muscle Cell

Flux through the sodium potassium pump (in  $\mu\text{M s}^{-1}$ ):

$$J_{NaK_i} = F_{NaK} \quad (3.6.1)$$

$F_{NaK}$	Rate of the potassium influx by the sodium potassium pump	$4.32 \times 10^{-2} \text{ } \mu\text{M s}^{-1}$	[11]
-----------	---	---	------

$G_{Cl_i}$	Whole-cell conductance for $\text{Cl}^-$ current	$1.34 \times 10^{-3} \text{ } \mu\text{M mV}^{-1} \text{s}^{-1}$	[11]
$v_{Cl_i}$	Reversal potential for $\text{Cl}^-$ channels.	-25.0 mV	[11]

Potassium flux through potassium channel (in  $\mu\text{M s}^{-1}$ ):

$$J_{K_i} = G_{K_i} w_i (v_i - E_{K_i}) \quad (3.6.2)$$

$G_{K_i}$	Whole-cell conductance for $\text{K}^+$ efflux.	$4.46 \times 10^{-3} \text{ } \mu\text{M mV}^{-1} \text{s}^{-1}$	[11]
$v_{K_i}$	Nernst potential	-94 mV	[11]

Flux through KIR channels in the SMC (in  $\mu\text{M s}^{-1}$ ):

$$J_{KIR_i} = \frac{F_{KIR_i} g_{KIR_i}}{\gamma_i} (v_i - v_{KIR_i}) \quad (3.6.3)$$

why do we have  $\frac{F_{KIR_i}}{\gamma_i}$  when they have both the same dimensions but one value  $F_{KIR_i}$  is 750 and the other  $\gamma_i$  is 1970 ?

Nernst potential of the KIR channel in the SMC (in mV):

$$v_{KIR_i} = z_1 K_p - z_2 \quad (3.6.4)$$

Conductance of KIR channel (in  $\mu\text{M mV}^{-1} \text{s}^{-1}$ ):

$$g_{KIR_i} = \exp(z_5 v_i + z_3 K_p - z_4) \quad (3.6.5)$$

$c_{wi}$	Translation factor for $\text{Ca}^{2+}$ dependence of $K_{Ca}$ channel activation sigmoidal.	0.0 $\mu\text{M}$	[11]
$\beta_i$	Translation factor for membrane potential dependence of $K_{Ca}$ channel activation sigmoidal.	0.13 $\mu\text{M}^2$	[11]
$v_{Ca_{3i}}$	Half-point for the $K_{Ca}$ channel activation sigmoidal.	-27 mV	[11]
$R_{K_i}$	Maximum slope of the $K_{Ca}$ activation sigmoidal.	12 mV	[11]
$z_1$	Model estimation for membrane voltage KIR channel	$4.5 \times 10^3 \text{ mV } \mu\text{M}^{-1}$	[6]
$z_2$	Model estimation for membrane voltage KIR channel	112 mV	[6]
$z_3$	Model estimation for the KIR channel conductance	$4.2 \times 10^2 \text{ mV}^{-1} \text{s}^{-1}$	[6]
$z_4$	Model estimation for the KIR channel conductance	$12.6 \text{ } \mu\text{M mV}^{-1} \text{s}^{-1}$	[6]
$z_5$	Model estimation for the KIR channel conductance	$-7.4 \times 10^{-2} \text{ } \mu\text{M mV}^{-2} \text{s}^{-1}$	[6]

$F_{KIR_i}$	Scaling factor of potassium efflux through the KIR channel	750 mV $\mu\text{M}^{-1}$	
-------------	--	---------------------------	--



IP<sub>3</sub> degradation (in  $\mu\text{M s}^{-1}$ ):

$$J_{\text{degrad}_i} = k_{di} I_i \quad (3.6.6)$$

$k_{di}$	Rate constant of IP <sub>3</sub> degradation	$0.1 \text{ s}^{-1}$	[11]
----------	--	----------------------	------

120

We should note here that the membrane potential coupling  $V_{\text{coupling}_i}^{\text{SMC-EC}}$  is an approximation that assumes the gradient of concentrations is negligible and hence only the membrane potential diffusion term is non-zero determined from the electro-diffusion theory.

K<sup>+</sup> concentration in the SMC (in  $\mu\text{M}$ ):

$$\frac{d[K_i^+]}{dt} = J_{Na/K_i} - J_{KIR_i} - J_{K_i} \quad (3.6.7)$$

$\gamma_i$	Change in membrane potential by a scaling factor	$1970 \text{ mV } \mu\text{M}^{-1}$	[11]
$\lambda_i$	Rate constant for opening	$45.0 \text{ s}^{-1}$	[11]

### 3.7 Endothelial Cell

Potassium efflux through the  $J_{BK_{Ca_j}}$  channel and the  $J_{SK_{Ca_j}}$  channel (in  $\mu\text{M s}^{-1}$ ):

$$J_{K_j} = G_{\text{tot}_j}(v_j - v_{K_j})(J_{BK_{Ca_j}} + J_{SK_{Ca_j}}) \quad (3.7.1)$$

$G_{\text{tot}_j}$	Total potassium channel conductivity.	$6927 \text{ pS}$	[11]
$v_{K_j}$	K <sup>+</sup> equilibrium potential	$-80.0 \text{ mV}$	[11]

Potassium efflux through the  $J_{BK_{Ca_j}}$  channel (in  $\mu\text{M s}^{-1}$ ):

$$J_{BK_{Ca_j}} = 0.2 \left( 1 + \tanh \left( \frac{(\log_{10}[Ca^{2+}]_j - c)(v_j - b_j) - a_{1j}}{m_{3bj}(v_j + a_{2j}(\log_{10}[Ca^{2+}]_j - c) - b_j)^2 + m_{4bj}} \right) \right) \quad (3.7.2)$$

Potassium efflux through the  $J_{SK_{Ca_j}}$  channel (in  $\mu\text{M s}^{-1}$ ):

$$J_{SK_{Ca_j}} = 0.3 \left( 1 + \tanh \left( \frac{\log_{10}[Ca^{2+}]_j - m_{3sj}}{m_{4sj}} \right) \right) \quad (3.7.3)$$

$c$	Model constant, further explanation see reference	-0.4 $\mu\text{M}$	[11]
$b_j$	Model constant, further explanation see reference	-80.8 mV	[11]
$a_{1j}$	Model constant, further explanation see reference	53.3 $\mu\text{M mV}$	[11]
$a_{2j}$	Model constant, further explanation see reference	53.3 mV $\mu\text{M}^{-1}$	[11]
$m_{3bj}$	Model constant, further explanation see reference	$1.32 \times 10^{-3} \mu\text{M mV}^{-1}$	[11]
$m_{4bj}$	Model constant, further explanation see reference	0.30 $\mu\text{M mV}$	[11]
$m_{3sj}$	Model constant, further explanation see reference	-0.28 $\mu\text{M}$	[11]
$m_{4sj}$	Model constant, further explanation see reference	0.389 $\mu\text{M}$	[11]

### 125 3.8 Extracellular Space

diffusion, NaK and BK fluxes defined here

$$\frac{d[K]_e}{dt} = -VRJ_{diff} + J_K - J_{NaK} \quad (3.8.1)$$

where

$$J_{diff} = \frac{1}{\tau_s}(K_e - K_s) \quad (3.8.2)$$

$$J_K = \textcolor{red}{G}_K w_i (v_i - E_K) \quad (3.8.3)$$

$$J_{NaK} = F_{NaK} \quad (3.8.4)$$

The flux  $J_K$  and the open probability  $w_i$  are defined in equations ?? to ??. and

$$\tau_s = \frac{(\Delta x_s)^2}{2D_K} \quad (3.8.5)$$

$$D_K = \frac{D_{free}}{\lambda_0^2} \quad (3.8.6)$$

130 here  $\Delta x_s$  is the effective diffusion distance and  $D_{free}$  is the diffusion coefficient of potassium in a free medium,  $\lambda_0$  the tortuosity factor since diffusion is hindered by the narrow confines of the extracellular space. **At this time volume ratios are used only for the transfer of potassium from the synaptic cleft to the ECS.**

### 3.9 Lumen

$\Delta x_s$	$10^{-4}$	m	average distance across two adjacent astrocyte arms
$D_{free}$	$4.58 \times 10^{-9}$	$m^2 s^{-1}$	potassium diffusion coefficient in free media
$\lambda_0$	1.6	non-dimensional	tortuosity factor
$G_K$	$4.46 \times 10^{-3}$	$\mu M m V^{-1} s^{-1}$	whole SMC conductance for $K^+$ efflux
$E_K$	-94	mV	Nernst potential for the SMC BK channel
$F_{NaK}$	$4.32 \times 10^{-2}$	$\mu M s^{-1}$	rate of $K^+$ influx by the sodium/potassium pump.

# Bibliography

- [1] **Chang, J. C.; Brennan, K. C.; He, D.; Huang, H.; Miura, R. M.; Wilson, P. L. and Wylie, J. J. (2013):** A Mathematical Model of the Metabolic and Perfusion Effects on Cortical Spreading Depression, PloS one, Vol. 8, No. 8 p. e70469.
- [2] **Chen, K. and Popel, A. S. (2006):** Theoretical analysis of biochemical pathways of nitric oxide release from vascular endothelial cells, Free Radical Biology and Medicine, Vol. 41, No. 4 pp. 668–680.
- [3] **Chen, K. and Popel, A. S. (2007):** Vascular and perivascular nitric oxide release and transport: Biochemical pathways of neuronal nitric oxide synthase (NOS1) and endothelial nitric oxide synthase (NOS3), Free Radical Biology and Medicine, Vol. 42, No. 6 pp. 811–822.
- [4] **Comerford, a.; Plank, M. J. and David, T. (2008):** Endothelial nitric oxide synthase and calcium production in arterial geometries: an integrated fluid mechanics/cell model., Journal of biomechanical engineering, Vol. 130, No. 1 p. 011010.
- [5] **Crouch, T. H. and Klee, C. B. (1980):** Positive cooperative binding of calcium to bovine brain calmodulin., Biochemistry, Vol. 19, No. 16 pp. 3692–3698.
- [6] **Filosa, J. a.; Bonev, A. D.; Straub, S. V.; Meredith, A. L.; Wilkerson, M. K.; Aldrich, R. W. and Nelson, M. T. (2006):** Local potassium signaling couples neuronal activity to vasodilation in the brain., Nature neuroscience, Vol. 9, No. 11 pp. 1397–1403.
- [7] **Gonzalez-Fernandez, B., J.M. Ermentrout (1994):** On the Origin and Dynamics of the Vasomotion of Small Arteries, Mathematical Biosciences, Vol. 167, No. 2 pp. 127–167.
- [8] **Hayashi, Y.; Nishio, M.; Naito, Y.; Yokokura, H.; Nimura, Y.; Hidaka, H. and Watanabe, Y. (1999):** Regulation of neuronal nitric-oxide synthase by calmodulin kinases, Journal of Biological Chemistry, Vol. 274, No. 29 pp. 20597–20602.
- [9] **Kager, H.; Wadman, W. J. and Somjen, G. G. (2000):** Simulated seizures and spreading depression in a neuron model incorporating interstitial space and ion concentrations., Journal of neurophysiology, Vol. 84, No. 1 pp. 495–512.

- 165 [10] **Kavdia, M.; Tsoukias, N. M. and Popel, A. S. (2002)**: Model of nitric oxide  
diffusion in an arteriole: impact of hemoglobin-based blood substitutes., *American  
Journal of Physiology, Heart and Circulatory Physiology*, Vol. 282, No. 6 pp. H2245–  
H2253.
- [11] **Koenigsberger, M.; Sauser, R.; Bény, J.-L. and Meister, J.-J. (2006)**: Ef-  
170 ffects of arterial wall stress on vasomotion., *Biophysical journal*, Vol. 91, No. 5 pp.  
1663–1674.
- [12] **Malinski, T.; Taha, Z.; Grunfeld, S.; Patton, S.; Kapturczak, M. and  
Tomboulia, P. (1993)**: Diffusion of nitric oxide in the aorta wall monitored in  
situ by porphyrinic microsensors., *Biochemical and Biophysical Research Communi-  
175 cations*, Vol. 193, No. 3 pp. 1076–1082.
- [13] **Nagelhus, E.; Horio, Y. and Inanobe, A. (1999)**: Immunogold evidence sug-  
gests that coupling of K<sup>+</sup> siphoning and water transport in rat retinal muller cells is  
mediated by a coenrichment of kir4. 1 and aqp4 in specific membrane domains, *Glia*,  
Vol. 63, No. 1 pp. 47–54.
- 180 [14] **Østby, I.; Øyehaug, L.; Einevoll, G. T.; Nagelhus, E. A.; Plahte, E.;  
Zeuthen, T.; Lloyd, C. M.; Ottersen, O. P. and Omholt, S. W. (2009)**:  
Astrocytic Mechanisms Explaining Neural-Activity-Induced Shrinkage of Extraneu-  
ronal Space, *PLoS Computational Biology*, Vol. 5, No. 1 pp. 1–12.
- [15] **Santucci, D. M. and Raghavachari, S. (2008)**: The effects of NR2 subunit-  
185 dependent NMDA receptor kinetics on synaptic transmission and CaMKII activation,  
*PLoS Computational Biology*, Vol. 4, No. 10.