Supporting Information

Synthesis of Vinylcycloheptadienes by the Nickel-Catalyzed Three-Component [3+2+2] Cocyclization. Application to the Synthesis of Polycyclic Compounds

Shinsuke Komagawa,[†] Kouhei Takeuchi,[†] Ikuo Sotome,[†] Isao Azumaya,[‡] Hyuma Masu,[‡] Ryu Yamasaki,[†] and Shinichi Saito^{*,†}

Department of Chemistry, Faculty of Science, Tokyo University of Science, Kagurazaka, Shinjuku, Tokyo 162-8601, Japan, and Faculty of Pharmaceutical Sciences at Kagawa Campus, Tokushima Bunri University, Kagawa 769-2193, Japan

ssaito@rs.kagu.tus.ac.jp

Table of contents

Experimental details. S2-11
Figure S1, S2 and S3. S12

¹H and ¹³C NMR spectra of **3a-3e**, **4g**, **6a-6c**, **6g**, **8**, **10**, **12a-12d**, **13b**, **14c**, **15b**, **16d**. S13-56

NOESY Spectrum of *endo-***12a**. S57

General Information

Reagents were commercially available and used without further purification unless otherwise noted. Ethyl (2c), $^{2-5}$ $(1).^{1}$ 2-benzyl-1-buten-3-yne 3-decen-1-yne (2d), 6,7 cyclopropylideneacetate (2f), 10 8-hexyl-1,1,2,2,3,3,4,4,5,5,6,6,6-tridecafluoro-(7*E*)-buten-9-yne (2e), 8,9 1-decen-3-yne diisopropoxytitanium(IV) dichloride¹¹ were prepared as reported. Chemical shifts were reported in delta units (δ) relative to chloroform (7.24 ppm for ¹H NMR and 77.0 ppm for ¹³C NMR) or dimethyl sulfoxide (2.49 ppm for ¹H NMR and 39.7 ppm for ¹³C NMR). Multiplicity is indicated by s (singlet), d (doublet), t (triplet), q (quartet) or m (multiplet). Coupling constants, J, are reported in Hz. The crystal structure was solved by direct methods SIR92 (Altomare et al., 1994). For refining the structures, performing a structure analysis, and producing crystallographic illustrations, the program TEXSAN (Rigaku/MSC, 1998) was used. All non-hydrogen atoms were refined anisotropically. All hydrogen atoms were included as their calculated positions.

Nickel(0)-Catalyzed Cycloaddition of Ethyl Cyclopropylideneacetate (1) and Conjugated Enynes (Table 1). A Representative Procedure. To a dark red mixture of Ni(cod)₂ (27.5 mg, 0.1 mmol) and PPh₃ (52.5 mg, 0.2 mmol) in dry toluene (0.5 mL) was added dropwise a solution of 1 (126 mg, 1 mmol) and 2 (3 mmol) in dry toluene (0.5 mL) at room temperature over 3 h under Ar. The progress of the reaction was monitored by TLC and GC-MS, and the mixture was stirred until the starting material 1 disappeared. The mixture was passed through a short silica gel column (ether). Evaporation of the solvent gave an oil, which was further purified by silica gel column chromatography to give 3.

(*E*)-1-Ethoxycarbonylmethylene-3,5-(di-1-methylethenyl)-2,4-cycloheptadiene (3a): purified by silica gel column chromatography (hexane/AcOEt 30:1 and hexane/CH₂Cl₂ 3:1); yellow oil; ¹H NMR (300 MHz, CDCl₃) 6.39 (s, 1H), 6.19 (s, 1H), 5.71 (s, 1H), 5.23 (s, 1H), 5.14 (s, 1H), 5.05 (s, 2H), 4.13 (q, J = 7.2 Hz, 2H), 3.16-3.11 (m, 2H), 2.54-2.50 (m, 2H), 1.97 (s, 3H), 1.96 (s, 3H), 1.25 (t, J = 7.2 Hz, 3H); ¹³C NMR (75 MHz, CDCl₃) 166.8, 158.3, 147.3, 146.0, 143.9, 143.0, 131.0, 122.7, 117.4, 115.1, 114.1, 59.6, 31.8, 26.1, 21.8, 21.1, 14.3; IR (neat) 2976, 1706, 1585, 1444, 1374, 1264, 1213, 1154, 1040, 891 cm⁻¹. HR-MS (EI) Calcd for C₁₇H₂₂O₂: 258.1620. Found: 258.1611.

(*E*)-1-Ethoxycarbonylmethylene-3,5-(di-1-cyclohexenyl)-2,4-cycloheptadiene (3b): purified by silica gel column chromatography (hexane/AcOEt 30:1 and hexane/CH₂Cl₂ 3:1); yellow oil; ¹H NMR (300 MHz, CDCl₃) 6.27 (s, 1H), 6.04 (t, J = 4 Hz, 1H), 6.02 (s, 1H), 5.91 (t, J = 4 Hz, 1H), 5.64 (s, 1H), 4.11 (q, J = 7.2 Hz, 2H), 3.13-3.09 (m, 2H), 2.49-2.45 (m, 2H), 2.22-2.11 (m, 8H), 1.69-1.54 (m, 8H), 1.25 (t, J = 7.2 Hz, 3H); ¹³C NMR (75 MHz, CDCl₃) 167.1, 159.4, 148.4, 145.8, 139.9, 136.6, 128.2, 127.1, 125.9, 120.2, 115.7, 59.4, 32.3, 27.2, 26.3, 26.2, 26.0, 25.9, 23.0, 23.0, 22.1, 22.0, 14.4; IR (neat) 2928, 1704, 1582, 1236, 1150, 1039, 756 cm⁻¹. HR-MS (ESI) Calcd for $(C_{23}H_{30}O_2)[M+Na]^+$: 361.2138. Found: 361.2138.

(*E*)-1-Ethoxycarbonylmethylene-3,5-(di-1-benzylethenyl)-2,4-cycloheptadiene (3c): purified by silica gel column chromatography (hexane/AcOEt 20:1); yellow oil; 1 H NMR (300 MHz, CDCl₃) 7.29-6.99 (m, 10H), 6.27 (s, 1H), 6.08 (s, 1H), 5.55 (s, 1H), 5.40 (s, 1H), 5.06 (s, 1H), 4.94 (s, 1H), 4.85 (s, 1H) 4.10 (q, J = 7.2 Hz, 2H), 3.65 (s, 2H), 3.41 (s, 2H) 3.07-3.04 (m, 2H), 2.45-2.41 (m, 2H), 1.24 (t, J = 7.2 Hz, 3H); 13 C NMR (125 MHz, CDCl₃) 166.7, 157.9, 150.2, 146.6, 146.5, 143.6, 139.8, 139.2, 131.9, 128.7, 128.5, 128.4, 128.3, 126.1, 126.1, 124.4, 117.2, 116.0, 59.6, 41.4, 40.8, 31.6, 26.7, 14.3; IR (neat) 3027, 2979, 1704, 1585, 1495, 1453, 1400, 1266, 1215, 1154, 1039, 895, 735, 699 cm⁻¹. HR-MS (ESI) Calcd for ($C_{29}H_{30}O_{2}$)[M+Na]⁺: 433.2138. Found: 433.2139.

$$CO_2Et$$
 C_6H_{13}

(*E*)-1-Ethoxycarbonylmethylene-3,5-(di-1-octenyl)-2,4-cycloheptadiene (3d): purified by silica gel column chromatography (hexane/ether 30:1); yellow oil; 1 H NMR (300 MHz, CDCl₃) 6.16-5.83 (m, 6H), 5.64 (s, 1H), 4.13 (q, J = 7.2 Hz, 2H), 3.14-3.11 (m, 2H), 2.46-2.43 (m, 2H), 2.13 (quint, J = 6.8 Hz, 4H), 1.40-1.23 (m, 19H), 0.89-0.85 (m, 6H); 13 C NMR (125 MHz, CDCl₃) 167.0, 158.4, 146.8, 140.4, 134.4, 133.1, 133.0, 132.7, 132.2, 122.9, 116.4, 59.6, 33.2, 33.1, 31.7, 29.5, 29.4, 29.3, 28.9, 25.1, 22.6, 14.4, 14.1; IR (neat) 2926, 2854, 1705, 1579, 1465, 1402, 1279, 1230, 1148, 1040, 961, 890 cm $^{-1}$. HR-MS (ESI) Calcd

for $(C_{27}H_{42}O_2)[M+Na]^+$: 421.3077. Found: 421.3079.

$$CO_2Et$$
 C_6H_{13}
 C_6F_{13}
 C_6H_{13}

(*E*)-1-Ethoxycarbonylmethylene-3,5-[di-1-hexyl-(2*E*)-perfluorohexyletenyl]-2,4-cycloheptadiene (3e): purified by silica gel column chromatography (hexane/AcOEt 30:1); yellow oil; ¹H NMR (300 MHz, CDCl₃) 6.31 (s, 1H), 5.88 (s, 1H), 5.77 (s, 1H), 5.48 (t, J = 15.6 Hz, 2H), 4.15 (q, J = 7.2 Hz, 2H), 3.19-3.16 (m, 2H), 2.45-2.42 (m, 6H), 1.40-1.15 (m, 16H), 1.27 (t, J = 7.2 Hz, 3H), 0.87-0.82 (m, 6H); ¹³C NMR (125 MHz, CDCl₃) 166.4, 159.5 (t, ${}^{3}J_{CF} = 5$ Hz), 155.4 (t, ${}^{3}J_{CF} = 5$ Hz), 150.1, 143.1, 134.4, 124.6, 119.9, 114.3 (t, ${}^{2}J_{CF} = 23$ Hz), 112.8 (t, ${}^{2}J_{CF} = 23$ Hz), 119-108 (m), 60.1, 31.5, 31.5, 30.8, 30.1, 29.3, 29.2, 29.0, 28.1, 22.5, 14.2, 13.8; IR (neat) 2933, 2862, 1712, 1240, 1202, 1165, 1146 cm⁻¹. Anal. Calcd for C₃₉H₄₀F₂₆O₂: C, 45.27; H, 3.90. Found: C, 45.08; H, 3.89.

(*E*)-1-Ethoxycarbonylmethylene-3,5-(di-1-oxoethyl)-2,4-cycloheptadiene (4g): purified by silica gel column chromatography (hexane/AcOEt 3:1); yellow oil; 1 H NMR (300 MHz, CDCl₃) 7.57 (s, 1H), 7.31 (s, 1H), 6.03 (s, 1H), 4.19 (q, J = 7.2 Hz, 2H), 3.09-3.05 (m, 2H), 2.64-2.61 (m, 2H), 2.47 (s, 3H), 2.42 (s, 3H), 1.28 (t, J = 7.2 Hz, 3H); 13 C NMR (75 MHz, CDCl₃) 198.4, 198.3, 165.5, 154.0, 146.3, 145.5, 138.6, 131.0, 125.5, 60.6, 28.5, 26.3, 25.7, 22.8, 14.2; IR (neat) 2981, 1711, 1668, 1597, 1435, 1356, 1250, 1166, 1034, 899 cm ${}^{-1}$. HR-MS (ESI) Calcd for ($C_{15}H_{18}O_4$)[M+Na] ${}^{+}$: 285.1097. Found: 285.1056.

Nickel(0)-Catalyzed Three-Component Cycloaddition of Ethyl Cyclopropylideneacetate (1), Conjugated Enynes and (Trimethylsilyl)acetylene (5) (Table 2). A Representative Procedure. To a dark red mixture of Ni(cod)₂ (27.5 mg, 0.1 mmol) and PPh₃ (52.5 mg, 0.2 mmol) in dry toluene (0.5 mL) was added dropwise a solution of 1 (126 mg, 1 mmol), 2 (1 mmol), and (trimethylsilyl)acetylene 5 (4 mmol) in dry toluene (0.5 mL) at room temperature over 3 h under Ar. The progress of the reaction was monitored by TLC and GC-MS, and the mixture was stirred until the starting material 1 disappeared. The mixture was passed through a short silica gel column (ether). Evaporation of the solvent gave an oil, which was further purified by silica gel column chromatography to give 6.

Reaction Condition for the Ni(PPh₃)₂Br₂-Zn-PPh₃ System (entry 2, Table 2). A mixture of Ni(PPh₃)₂Br₂ (372 mg, 0.5 mmol), PPh₃ (262 mg, 1 mmol), and Zn dust (654 mg, 10 mmol) in dry toluene

(2.5 mL) was stirred at room temperature over 1 h and changed color from green to dark red. To this mixture was added dropwise a solution of **1a** (630 mg, 5 mmol), **2a** (0.48 mL, 5 mmol), and **5** (2.1 mL, 15 mmol) in dry toluene (2.5 mL) at room temperature for 5 h under Ar. The mixture was stirred for 14 h and passed through a short silica gel column (ether). Evaporation of the solvent gave an oil, which was further purified by silica gel column chromatography (hexnane/ether 30:1) to give **6a** (905 mg, 62%).

(*E*)-1-Ethoxycarbonylmethylene-3-(1-methylethenyl)-5-trimethylsilyl-2,4-cycloheptadiene (6a): purified by silica gel column chromatography (hexane/CH₂Cl₂ 3:1); pale yellow oil; ¹H NMR (300 MHz, CDCl₃) 6.38 (s, 1H), 6.36 (s, 1H), 5.71 (s, 1H), 5.13 (s, 1H), 5.05 (s, 1H), 4.13 (q, J = 7.2 Hz, 2H), 3.07-3.04 (m, 2H), 2.33-2.29 (m, 2H), 1.95 (s, 3H), 1.25 (t, J = 7.2 Hz, 3H), 0.10 (s, 9H); ¹³C NMR (125 MHz, CDCl₃) 166.8, 158.5, 151.7, 145.6, 143.5, 133.7, 131.6, 117.4, 115.1, 59.6, 32.3, 27.6, 21.7, 14.3, -2.2; IR (neat) 2954, 1709, 1586, 1444, 1403, 1376, 1249, 1214, 1155, 1088, 1067, 1039, 892, 838, 752 cm⁻¹. HR-MS (ESI) Calcd for $(C_{17}H_{26}O_2Si)[M+Na]^+$: 313.1594. Found: 313.1594.

(*E*)-1-Ethoxycarbonylmethylene-3-(1-cyclohexenyl)-5-trimethylsilyl-2,4-cycloheptadiene (6b): purified by silica gel column chromatography (hexane/AcOEt 30:1); pale yellow oil; 1 H NMR (300 MHz, CDCl₃) 6.30 (s, 1H), 5.91 (t, J = 4.0 Hz, 1H), 5.67 (s, 1H), 4.12 (q, J = 7.2 Hz, 2H), 3.07-3.03 (m, 2H), 2.31-2.27 (m, 2H), 2.18-2.14 (m, 4H), 1.70-1.64 (m, 2H), 1.62-1.56 (m, 2H), 1.25 (t, J = 7.2 Hz, 3H), 0.10 (s, 9H); 13 C NMR (125 MHz, CDCl₃) 167.0, 159.2, 151.4, 144.6, 139.0, 134.1, 129.6, 127.3, 116.4, 59.5, 32.8, 27.6, 26.9, 26.0, 22.9, 21.9, 14.3, -2.2; IR (neat) 2930, 1707, 1585, 1248, 1217, 1152, 1087, 1039, 838, 752 cm⁻¹. HR-MS (ESI) Calcd for $(C_{20}H_{30}O_{2}Si)[M+Na]^{+}$: 353.1907. Found: 353.1907.

(E)-1-Ethoxycarbonylmethylene-3-(1-benzylethenyl)-5-trimethylsilyl-2,4-cycloheptadiene (6c): purified

by silica gel column chromatography (hexane/AcOEt 25:1 and hexane/CH₂Cl₂ 3:1); pale yellow oil; 1 H NMR (300 MHz, CDCl₃) 7.28-7.13 (m, 5H), 6.34 (s, 1H), 6.26 (s, 1H), 5.58 (s, 1H), 5.23 (d, J = 1.3 Hz, 1H), 4.98 (d, J = 1.3 Hz, 1H), 4.11 (q, J = 7.2 Hz, 2H), 3.59 (s, 2H), 3.02-2.98 (m, 2H), 2.25-2.22 (m, 2H), 1.24 (t, J = 7.2 Hz, 3H), 0.08 (s, 9H); 13 C NMR (75 MHz, CDCl₃) 166.8, 158.4, 152.2, 149.8, 143.3, 139.3, 134.1, 132.6, 128.8, 128.3, 126.1, 117.1, 116.0, 59.6, 41.5, 32.3, 27.6, 14.3, -2.3; IR (neat) 2954, 1708, 1586, 1453, 1401, 1249, 1214, 1154, 1086, 1039, 895, 838, 751, 699 cm⁻¹. HR-MS (ESI) Calcd for ($C_{22}H_{20}O$)[M+Na]⁺: 389.1907. Found: 389.1907.

(*E*)-1-Ethoxycarbonylmethylene-3-(1-butoxyethenyl)-5-trimethylsilyl-2,4-cycloheptadiene (6g): purified by silica gel column chromatography (hexane/AcOEt 30:1 and hexane/ether 30:1); pale yellow oil; 1 H NMR (300 MHz, $C_{6}D_{6}$) 7.09 (s, 1H), 6.62 (s, 1H), 5.97 (s, 1H), 4.51 (d, J = 2.3 Hz, 1H), 4.13 (d, J = 2.3 Hz, 1H), 4.01 (q, J = 7.2 Hz, 2H), 3.50 (t, J = 6.3 Hz, 1H), 3.39-3.35 (m, 2H), 2.29-2.25 (m, 2H), 1.52 (tt, 7.3 Hz, 6.3 Hz, 2H), 1.31 (qt, 7.3 Hz, 7.3 Hz, 2H), 0.99 (t, J = 7.2 Hz, 3H), 0.81 (t, J = 7.3 Hz, 3H), 0.02 (s, 9H); 13 C NMR (125 MHz, $C_{6}D_{6}$) 166.4, 162.2, 158.0, 152.2, 138.6, 133.3, 132.8, 119.1, 85.1, 67.6, 59.6, 32.0, 31.3, 28.1, 19.7, 14.4, 13.9, -2.4; IR (neat) 2957, 1710, 1594, 1266, 1249, 1214, 1154, 1081, 838 cm ${}^{-1}$. HR-MS (ESI) Calcd for $(C_{20}H_{32}O_{3}Si)[M+Na]^{+}$: 371.2013. Found: 371.2013.

(*E*)-1-Ethoxycarbonylmethylene-3-(1-methylethenyl)-4,5-diphenyl-2,4-cycloheptadiene (8): purified by silica gel column chromatography (hexane/CH₂Cl₂ 2:1); pale yellow solid; mp 86-87 °C; ¹H NMR (300 MHz, CDCl₃) 7.13-6.79 (m, 10H), 6.68 (s, 1H), 5.84 (d, J = 1.2 Hz, 1H), 4.89 (s, 1H), 4.80 (s, 1H), 4.13 (q, J = 7.2 Hz, 2H), 3.26-3.21 (m, 2H), 2.78-2.74 (m, 2H), 1.67 (s, 3H), 1.25 (t, J = 7.2 Hz, 3H); ¹³C NMR (150 MHz, CDCl₃) 166.6, 157.9, 148.7, 145.6, 145.0, 142.3, 140.2, 135.2, 131.3, 130.6, 129.6, 127.7, 127.1, 126.3, 125.9, 117.2, 116.5, 59.6, 37.1, 34.0, 21.6, 14.3; IR (KBr) 2966, 1700, 1587, 1488, 1443, 1396, 1252, 1208, 1149, 1031, 899, 752, 697 cm⁻¹. Anal. Calcd for C₂₆H₂₆O₂: C, 84.29; H, 7.07. Found: C, 84.15; H, 7.13.

(*E*)-1-Ethoxycarbonylmethylene-3-methoxymethyl-5-(1-methylethenyl)-2,4-cycloheptadiene (10): purified by silica gel column chromatography (hexane/AcOEt 10:1); pale yellow oil; 1 H NMR (300 MHz, CDCl₃) 6.29 (s, 1H), 6.00 (s, 1H), 5.66 (s, 1H), 5.23 (s, 1H), 5.06 (s, 1H), 4.13 (q, J = 7.2 Hz, 2H), 3.97 (s, 2H), 3.32 (s, 3H), 3.14-3.11 (m, 2H), 2.56-2.52 (m, 2H), 1.95 (s, 3H), 1.25 (t, J = 7.2 Hz, 3H); 13 C NMR (75 MHz, CDCl₃) 166.9, 157.3, 148.9, 143.3, 140.6, 132.1, 122.4, 117.0, 114.5, 78.4, 59.7, 57.9, 29.3,26.6, 21.1, 14.3; IR (KBr) 2980, 1705, 1592, 1445, 1402, 1269, 1158, 1905, 1042, 879 cm $^{-1}$. HR-MS (ESI) Calcd for $(C_{16}H_{22}O_3)[M+Na]^{+}$: 285.1461. Found: 285.1460.

Diels-Alder Reaction of 6a and 11 (Table 3). A Representative Procedure. A mixture of **6a** (145 mg, 0.5 mmol) and **11** (0.75 mmol) in toluene (2 mL) was stirred at the designated temperature. After the reaction completed, the mixture was purified by silica gel column chromatography to give **12**.

Diels-Alder Reaction of 5a and 6 in the presence of TiCl₂(Oi-Pr)₂. A Representative Procedure. To a mixture of **6a** (145 mg, 0.5 mmol) and **11** (0.75 mmol) in toluene (2 mL) was added a solution of TiCl₂(Oi-Pr)₂ (1.2 mL, 0.5 M in toluene, 0.6 mmol) at rt. After the reaction completed, the mixture was purified by silica gel column chromatography to give **12** or **16**.

Four-Component One-Pot [3+2+2]/[4+2] Cycloaddition of 1, 2, 5, and 11b (eq 9). To a dark red mixture of Ni(cod)₂ (27.5 mg, 0.1 mmol) and PPh₃ (52.5 mg, 0.2 mmol) in dry toluene (0.5 mL) was added dropwise a solution of 1 (127 mg, 1 mmol), 2a (0.095 mL, 1 mmol), and 5 (0.57 mL, 4 mmol) in dry toluene (0.5 mL) at rt over 3 h under Ar. After 16 h, toluene (1 mL) and 11b (128 mg, 1 mmol) was added and the mixture was stirred at rt for 23 h. The crude product was passed through a short silica gel column (ether) and further purified by silica gel column chromatography (hexane/AcOEt 8:1) to give 12b (227.9 mg, 54%).

endo-12a: purified by silica gel column chromatography (hexane/AcOEt 3:1); white solid; mp 67-68 °C; 1 H NMR (500 MHz, CDCl₃) 6.39 (s, 1H), 5.92 (s, 1H), 4.16 (q, J = 7.2 Hz, 2H), 3.61 (dd, J = 9.8, 5.8 Hz, 1H), 3.44 (ddd, J = 9.8, 6.7, 2.4 Hz, 1H), 3.32 (m, 1H), 3.04-2.99(m, 1H), 2.83-2.78 (m, 1H), 2.68 (dd, J = 15.5, 2.3 Hz, 1H), 2.49-2.41 (m, 3H), 1.89 (s, 3H), 1.27 (t, J = 7.2 Hz, 3H), 0.04 (s, 9H); 13 C NMR (125 MHz, CDCl₃) 173.3, 170.5, 165.9, 159.1, 144.6, 134.7, 130.8, 130.4, 116.7, 59.9, 50.2, 44.8, 41.3, 31.8, 31.6, 31.2,

20.4, 14.2, -1.9; IR (KBr) 2955, 1844, 1779, 1704, 1650, 1248, 1179, 992, 937, 837 cm $^{-1}$. HR-MS (EI) Calcd for $C_{21}H_{28}O_5Si$: 388.1706. Found: 388.1714.

12b: purified by silica gel column chromatography (hexane/AcOEt 8:1); colorless solid; mp 156-158 °C; 1 H NMR (300 MHz, CDCl₃) 6.17 (s, 2H), 4.21 (qd, J = 7.2, 3.4 Hz, 2H), 3.89 (s, 1H), 3.59 (dt, J = 11.5, 4.1 Hz, 1H), 3.20 (d, J = 18.4 Hz, 1H), 3.00 (d, J = 18.4 Hz, 1H), 2.62-2.56 (m, 2H), 2.52-2.43 (m, 1H) 1.82 (s, 3H), 1.30 (t, J = 7.2 Hz, 3H), 0.06 (s, 9H); 13 C NMR (75 MHz, CDCl₃) 164.5, 152.7, 150.1, 130.2, 125.7, 125.4, 124.2, 110.8, 110.7, 110.5, 109.0, 60.6, 53.4, 44.2, 38.7, 37.0, 34.1, 27.7, 19.7, 14.1, -2.1; IR (KBr) 2954, 1715, 1658, 1446, 1381, 1248, 1209, 1189, 1154, 1041, 837, 751 cm $^{-1}$. Anal. Calcd for $C_{23}H_{26}N_4O_2Si$: C, 66.00; H, 6.26; N, 13.39. Found: C, 65.97; H, 6.26; N, 13.19.

endo-12c: purified by silica gel column chromatography (CH₂Cl₂/ether 20:1); white amorphous; mp 59-61 °C; ¹H NMR (300 MHz, CDCl₃) 7.45-7.16 (m, 5H), 6.47 (s, 1H), 5.97 (s, 1H), 4.13 (qd, J = 7.2, 1.3 Hz, 2H), 3.54-3.49 (m, 2H), 3.33-3.27 (m, 1H), 3.05-2.99 (m, 1H), 2.93-2.85 (m, 1H), 2.71 (dd, J = 15.8, 3.8 Hz, 1H), 2.52-2.44 (m, 3H), 1.88 (s, 3H), 1.25 (t, J = 7.1 Hz, 3H), 0.05 (s, 9H); ¹³C NMR (125 MHz, CDCl₃) 178.0, 175.9, 166.2, 160.4, 144.0, 134.6, 131.9, 131.3, 130.4, 129.1, 128.6, 126.3, 119.8, 115.5, 59.7, 50.0, 43.6, 40.6, 31.7, 31.3, 20.3, 14.3, -1.8; IR (KBr) 2954, 1712, 1499, 1383, 1249, 1177, 1044, 839, 754, 692 cm⁻¹. Anal. Calcd for $C_{27}H_{33}NO_4Si$: C, 69.94; H, 7.17; N, 3.02. Found: C, 69.71; H, 7.27; N, 3.06.

endo-12d: purified by silica gel column chromatography (hexane/AcOEt 10:1); white solid; mp 141-142 °C; 1 H NMR (500 MHz, CDCl₃) 8.01-7.99 (m, 1H), 7.89-7.88 (m, 1H), 7.70-7.68 (m, 2H), 6.34 (s, 1H), 5.58 (s, 1H), 4.06 (q, J = 7.2 Hz, 2H), 3.80 (t, J = 5.2 Hz, 1H), 3.58-3.49 (m, 2H), 3.30 (tt, J = 5.2, 4.9 Hz, 1H), 2.92 (td, J = 11.9, 3.1 Hz, 1H), 2.58 (d, J = 18.5 Hz, 1H), 2.41-2.21 (m, 3H), 1.70 (s, 3H), 1.20 (t, J = 7.2 Hz, 3H), 0.04 (s, 9H); 13 C NMR (125 MHz, CDCl₃) 198.0, 196.3, 167.2, 166.0, 144.9, 136.0, 134.5, 134.1, 134.0,

132.9, 130.5, 127.5, 126.8, 126.6, 116.3, 59.4, 55.0, 49.9, 48.5, 34.1, 32.3, 30.6, 20.2, 14.3, -1.8; IR (KBr) 2952, 1713, 1692, 1631, 1594, 1449, 1376, 1285, 1251, 1205, 1165, 1047, 967, 836 cm⁻¹. Anal. Calcd for $C_{27}H_{32}O_4Si$: C, 72.28; H, 7.19. Found: C, 72.05; H, 7.17.

13b: purified by silica gel column chromatography (hexane/AcOEt 10:1); colorless solid; mp 133-134 °C; 1 H NMR (300 MHz, CDCl₃) 6.34 (s, 1H), 6.17 (s, 1H), 4.21 (qd, J = 7.2, 3.2 Hz, 2H), 3.97 (s, 1H), 3.82 (m, 2H), 3.64-3.59 (m, 1H), 3.26 (d, J = 18.0 Hz, 1H), 3.17 (d, J = 18.0 Hz, 1H), 2.67-2.46 (m, 3H), 1.62 (tt, J = 7.3, 6.6 Hz, 2H), 1.40 (qt, J = 7.3, 7.3 Hz, 2H), 1.30 (t, J = 7.2 Hz, 3H), 0.92 (t, J = 7.3 Hz, 3H), 0.06 (s, 9H); 13 C NMR (75 MHz, CDCl₃) 164.6, 152.8, 148.4, 144.0, 127.8, 124.1, 113.6, 110.5, 110.4, 110.1, 108.7, 70.8, 60.7, 52.7, 44.1, 39.0, 34.5, 33.3, 31.7, 27.8, 19.0, 14.1, 13.7, -2.1; IR (KBr) 2960, 1718, 1650, 1443, 1379, 1249, 1178, 1039, 835, 748 cm⁻¹. Anal. Calcd for $C_{26}H_{32}N_4O_3Si$: C, 65.52; H, 6.77; N, 11.75. Found: C, 65.64; H, 6.79; N, 11.52.

*endo-***14c:** purified by silica gel column chromatography (hexane/AcOEt 2:1); white solid; mp 181-182 °C; 1 H NMR (300 MHz, CDCl₃ at 50 °C) 7.47-6.83 (m, 15H), 6.33 (bs, 1H), 4.13 (q, J = 7.2 Hz, 2H), 3.68-3.62 (m, 1H), 3.64 (dd, J = 9.4, 6.3 Hz, 1H) 3.38-3.31 (m, 2H), 2.87-2.52 (m, 5H), 1.49 (s, 3H), 1.23 (t, J = 7.2 Hz, 3H); 13 C NMR (75 MHz, CDCl₃ at 50 °C) 177.9, 176.6, 166.2, 142.5, 141.7, 141.3, 137.4, 133.8, 132.4, 130.0, 129.5, 129.0, 128.6, 127.9, 127.6, 126.7, 126.4, 125.9, 59.7, 49.2, 44.4, 40.6, 34.7, 30.0, 21.3, 14.2; 1 H NMR (300 MHz, DMSO-d₆ at 90 °C) 7.52-6.87 (m, 15H), 6.46 (bs, 1H), 4.08 (q, J = 7.2 Hz, 2H), 3.80-3.71 (m, 1H), 3.77 (d, J = 6.0 Hz, 1H), 3.55-3.47 (m, 1H), 3.23-3.11 (m, 1H), 2.80-2.92 (1H, overlap), 2.67-2.53 (m, 4H), 1.43 (s, 3H), 1.18 (t, J = 7.2 Hz, 3H); 13 C NMR (75 MHz, DMSO-d₆ at 90 °C) 177.5, 176.3, 165.3, 141.8, 141.1, 139.8, 136.4, 132.5, 132.2, 129.1, 128.5, 128.2, 127.6, 127.1, 126.6, 126.1, 125.6, 125.1, 119.2, 58.6, 47.7, 43.8, 33.3, 30.1, 28.6, 20.2, 13.4; IR (KBr) 2933, 1707, 1498, 1378, 1234, 1165, 702, 691 cm⁻¹. Anal. Calcd for C₃₆H₃₃NO₄: C, 79.53; H, 6.12; N, 2.58. Found: C, 79.48; H, 6.18; N, 2.54.

*exo-***14c:** purified by silica gel column chromatography (hexane/AcOEt 2:1); white solid; mp 245 °C; 1 H NMR (300 MHz, CDCl₃) 7.53-6.72 (m, 15H), 5.68(s, 1H), 4.21-4.16 (m, 1H), 4.17 (q, J = 7.2 Hz, 2H), 3.64 (dd, J = 9.4, 2.1 Hz, 1H), 3.42-3.33 (m, 2H), 3.19-3.11 (m, 1H), 2.86-2.67 (m, 2H), 2.56-2.54 (m, 2H), 1.34 (s, 3H), 1.29 (t, J = 7.2 Hz, 3H); 13 C NMR (125 MHz, CDCl₃) 190.7, 178.6, 166.2, 160.8, 143.3, 142.1, 141.7, 139.6, 134.5, 133.3, 131.9, 129.8, 129.6, 129.2, 128.8, 127.8, 127.6, 126.4, 126.3, 125.9, 112.7, 60.1, 48.8, 41.7, 38.7, 36.3, 30.7, 29.0, 21.5, 14.3; IR (KBr) 2903, 1711, 1637, 1497, 1283, 1195, 1165, 1138, 1043, 745, 700 cm⁻¹. Anal. Calcd for $C_{36}H_{33}NO_4$: C, 79.53; H, 6.12; N, 2.58. Found: C, 79.63; H, 6.15; N, 2.58.

15b: purified by silica gel column chromatography (hexane/AcOEt 1:1); white solid; mp 247 °C (decomp.); 1 H NMR (300 MHz, CDCl₃) 6.30 (s, 2H), 4.63 (s, 1H), 4.39 (s, 1H), 4.19 (q, J = 7.2 Hz, 2H), 4.08-3.97 (m, 1H), 3.45-3.20 (m, 3H), 2.97-2.84 (m, 2H), 2.65-2.57 (m, 1H), 2.21 (s, 3H), 2.21-2.06 (m, 1H), 1.82 (s, 3H), 1.30 (t, J = 7.2 Hz, 3H); 13 C NMR (125 MHz, CDCl₃) 164.5, 145.7, 137.9, 130.4, 125.9, 124.8, 121.9, 110.4, 110.3, 110.2, 110.2, 109.9, 109.8, 109.4, 109.4, 61.2, 51.5, 47.8, 44.3, 42.1, 39.4, 38.5, 38.2, 37.2, 25.4, 23.4, 21.5, 18.8, 14.1; IR (KBr) 2964, 1718, 1647, 1438, 1384, 1257, 1227, 1189, 1032 cm⁻¹. Anal. Calcd for $C_{29}H_{22}N_8O_2$: C, 67.69; H, 4.31; N, 21.78. Found: C, 67.61; H, 4.44; N, 21.71. The coupling product (**15b**) was isolated as a single isomer, but the stereochemistry has not been determined.

16d: purified by silica gel column chromatography (hexane/AcOEt 10:1); red solid; mp 162-163 °C; ${}^{1}H$ NMR (300 MHz, CDCl₃) 8.13-8.11 (m, 1H), 8.05-8.01 (m, 1H), 7.75-7.68 (m, 2H), 6.38 (d, J = 11.5 Hz, 1H), 5.88 (dt, J = 11.5, 3.9 Hz, 1H), 5.26 (s, 1H), 4.94 (s, 1H), 4.02 (q, J = 7.2 Hz, 2H), 3.46-3.09 (m, 4H), 3.05-2.88 (m, 1H), 2.41-2.29 (m, 1H), 1.85 (s, 3H), 1.16 (t, J = 7.2 Hz, 3H); ${}^{13}C$ NMR (125 MHz, CDCl₃) 184.0, 183.1, 166.4, 164.4, 143.5, 142.9, 133.8, 133.7, 132.1, 132.0, 130.7, 128.1, 127.3, 126.7, 126.5, 126.4, 113.4, 59.7, 43.2, 33.3, 31.2, 25.2, 18.7, 14.2; IR (KBr) 3444, 1705, 1659, 1631, 1591, 1331, 1294, 1216, 1175, 1142 cm⁻¹. HR-MS (EI) Calcd for $C_{24}H_{22}O_4$: 374.1518. Found: 374.1518.

References

- (1) Kortmann, I.; Westermann, B. Synthesis. 1995, 931-933.
- (2) Gevorgyan, V.; Takeda, A.; Homma, M.; Sadayori, N.; Radhakrishnan, U.; Yamamoto, Y. *J. Am. Chem. Soc.* **1999**, *121*, 6391-6402.
- (3) Iso, Y.; Grajkowska, E.; Wroblewski, J. T.; Davis, J.; Goeders, N. E.; Johnson, K. M.; Sanker, S.; Roth, B. L.; Tueckmantel, W.; Kozikowski, A. P. *J. Med. Chem.* **2006**, *49*, 1080-1100.
- (4) Schwab, J. M.; Lin, D. C. T. J. Am. Chem. Soc. 1985, 107, 6046-6052.
- (5) Ager, D. J. Synthesis, 1984, 384-398.
- (6) Ren, H.; Krasovskiy, A.; Knochel, P. Org. Lett. 2004, 6, 4215-4217.
- (7) Negishi, E.; Kotora, M.; Xu, C. J. Org. Chem. 1997, 62, 8957-8960.
- (8) Saito, S.; Kawasaki, T.; Tsuboya, N.; Yamamoto, Y. J. Org. Chem. 2001, 66, 796-802.
- (9) Ishihara, T.; Kuroboshi, M.; Okada, Y. Chem. Lett. 1986, 1895-1896.
- (10) Miller, K. M.; Luanphaisarnnant, T.; Molinaro, C.; Jamison, T. F. J. Am. Chem. Soc. 2004, 126, 4130-4131.
- (11) Mikami, K.; Terada, M.; Nakai, T. J. Am. Chem. Soc. 1990, 112, 3949-3954.

Figure S1. A NOESY experiment of 12a.




Figure S2. An ORTEP view of *endo-***14c**. Crystal data for *endo-***14c**: C₃₆H₃₃NO₄; M = 543.63 g mol⁻¹, monoclinic, $P2_1/c$, colorless prism measuring $0.40 \times 0.30 \times 0.30$ mm, T = 150 K, a = 11.818 (1), b = 8.5654(7), c = 28.864(3) Å, $β = 101.646(1)^\circ$, V = 2861.5 (4) Å³, Z = 4, $D_c = 1.262$ Mg m⁻³, μ(CuKα) = 0.082 mm⁻¹, $T_{\text{max}} = 0.9759$, $T_{\text{min}} = 0.9681$, GOF on $F^2 = 1.080$, $R_1 = 0.0469$, w $R_2 = 0.1213$ [I > 2σ(I)], $R_1 = 0.0641$, and w $R_2 = 0.1324$ (all data).






Figure S3. An ORTEP view of **16d**. Crystal data for **16d**: C₂₄H₂₂O₄; M = 374.42 g mol⁻¹, monoclinic, $P2_1/n$, colorless prism measuring $0.30 \times 0.10 \times 0.10$ mm, T = 120 K, a = 8.319(3), b = 15.649(5), c = 14.325(4) Å, $β = 93.669(2)^\circ$, V = 1861(1) Å³, Z = 4, $D_c = 1.336$ Mg m⁻³, μ(CuKα) = 0.090 mm⁻¹, $T_{\text{max}} = 0.9910$, $T_{\text{min}} = 0.9734$, GOF on $F^2 = 0.821$, $R_1 = 0.0419$, w $R_2 = 0.1197$ [I > 2σ (I)], $R_1 = 0.0603$, and w $R_2 = 0.1382$ (all data).

¹H NMR spectrum of 3a


Z: \psi Noma\pmrdata\pmrdata \pmrdata \


DFILE Z: **Koma**Inmrdata**SK14-01-4.3 CCMNT
DATIM
CBNUC 1H
EXMOD
OBFRQ 300.01 MHz
OBFRQ 1.85 KHz
OBFIN 2.7 Hz
PPRQU 6172.8 Hz
SCANS 6172.8 Hz
ACQTM 0.000 Sec
PD
ACQTM 10.0 Sec
PD
ACQTM 10.0 Sec
RM 10.0 US
IRNUC 0.000 C
SILNUT 10.0 US
SILNUT 10.0 US
REREF 1.20 Hz
RGAIN 0








¹³C NMR spectrum of 3a
















Z: \koma\mrdata\kSK11-60-3-13C.als


















ÇO₂Et





















168.861— 675.861—




¹³C NMR spectrum of 6a











E:\\$SK12-60-G1.dx
















¹H NMR spectrum of 8









¹H NMR spectrum of 10

E:\SK13-52-4.dx




¹³C NMR spectrum of 10







¹H NMR spectrum of endo-12a










¹³C NMR spectrum of endo-12a













¹³C NMR spectrum of 12b










¹H NMR spectrum of endo-12c



















¹H NMR spectrum of endo-12d















¹H NMR spectrum of 13b










¹²C NMR spectrum of 13b













E: \SK14-61-3-50.dx























E: \SK14-61-3-90b2.dx











E:\SK14-61-3-13C90.dx







E:\SK14-39-6r-1H.dx





¹²C NMR spectrum of *exo-*14c

















¹H NMR spectrum of 16d

E:\SK13-09-4.dx



¹³C NMR spectrum of 16d

