

RandLib documentation

Aleksandr Samarin

November 12, 2017

Contents

I	General information	2
1	Calculation of sample moments	3
II	Continuous univariate distributions	4
2	Beta distribution	4
2.1	Arcsine distribution	5
2.2	Balding-Nichols distribution	6
2.3	Uniform distribution	6
3	Beta-prime distribution	7
4	Exponentially-modified Gaussian distribution	9
5	F-distribution	10
6	Gamma distribution	10
6.1	Chi-squared distribution	12
6.2	Erlang distribution	12
6.3	Exponential distribution	12
7	Geometric Stable distribution	13
7.1	Asymmetric Laplace distribution	13
7.2	Laplace distribution	13
8	Noncentral Chi-Squared distribution	13
9	Planck distribution	13

10 Stable distribution	13
10.1 Normal distribution	13
10.2 Cauchy distribution	15
10.3 Levy distribution	15
10.4 Holtsmark distribution	15
10.5 Landau distribution	16
11 Pareto distribution	16
12 Weibull	19
 III Discrete univariate distributions	 21
13 Beta-binomial distribution	21
14 Binomial distribution	21
14.1 Bernoulli	21
15 Poisson distribution	22
 IV Bivariate distributions	 26
16 Bivariate Normal distribution	26
17 Normal-Inverse-Gamma distribution	26
18 Trinomial distribution	26
 V Circular distributions	 27
19 von Mises distribution	27
20 Wrapped Exponential distribution	27
 VI Singular distributions	 28
21 Cantor distribution	28

Part I

General information

1 Calculation of sample moments

We use extension of Welford's method from Knuth. For every n -th element x we have


$$\begin{aligned}\delta &= x - m_1, \\ m'_1 &= m_1 + \frac{\delta}{n}, \\ m'_2 &= m_2 + \delta^2 \frac{n-1}{n}, \\ m'_3 &= m_3 + \delta^3 \frac{(n-1)(n-2)}{n^2} - 3\delta \frac{m_2}{n}, \\ m'_4 &= m_4 + \delta^4 \frac{(n-1)(n^2-3n+3)}{n^3} + 6\delta^2 \frac{m_2}{n^2} - 4\delta \frac{m_3}{n}.\end{aligned}$$

Then m'_1 , $\frac{m_2}{n}$, $\text{Skew}(X) = \frac{\sqrt{n}m'_3}{m_2^{3/2}}$ and $\text{Kurt}(X) = \frac{nm'_4}{m_2^2}$ (we return excess kurtosis).

Part II

Continuous univariate distributions

2 Beta distribution



Notation	$X \sim B(\alpha, \beta, a, b)$ $X \sim B(\alpha, \beta)$ with $a = 0, b = 1$
Parameters	$\alpha, \beta > 0, a, b \in \mathbb{R}$
Support	$x \in [a, b]$
$f(x)$	$\frac{y^{\alpha-1}(1-y)^{\beta-1}}{(b-a)B(\alpha, \beta)}$ with $y = \frac{x-a}{b-a}$
$F(x)$	$I_y(\alpha, \beta)$ for $y = \frac{x-a}{b-a}$
$\mathbb{E}[X]$	$a + (b-a)\frac{\alpha}{\alpha+\beta}$
$\text{Var}(X)$	$(b-a)^2 \frac{\alpha\beta}{(\alpha+\beta)^2(\alpha+\beta+1)}$
Median	Searched numerically
Mode	$a + (b-a)\frac{\alpha-\beta}{\alpha+\beta-2}$ for $\alpha, \beta > 1$.
$\phi(t)$	Calculated numerically

Estimation of shapes with known support. Assume that $a = 0, b = 1$ and we have a sample $X = (X_1, \dots, X_n)$. Then a log-likelihood function is

$$\begin{aligned}
 \ln \mathcal{L}(\alpha, \beta | X) &= \sum_{i=1}^n \ln f(X_i; \alpha, \beta) \\
 &= (\alpha - 1) \sum_{i=1}^n \ln X_i + (\beta - 1) \sum_{i=1}^n \ln(1 - X_i) - n \ln B(\alpha, \beta).
 \end{aligned} \tag{1}$$

Differentiating with respect to the shapes, we obtain

$$\begin{aligned}
 \frac{\partial \ln \mathcal{L}(\alpha, \beta | X)}{\partial \alpha} &= \sum_{i=1}^n \ln X_i + n(\psi(\alpha + \beta) - \psi(\alpha)), \\
 \frac{\partial \ln \mathcal{L}(\alpha, \beta | X)}{\partial \beta} &= \sum_{i=1}^n \ln(1 - X_i) + n(\psi(\alpha + \beta) - \psi(\beta)).
 \end{aligned}$$

Differentiating again we get the Hessian matrix:

$$H(\ln \mathcal{L}(\alpha, \beta|X)) = n \cdot \begin{pmatrix} \psi_1(\alpha + \beta) - \psi_1(\alpha) & \psi_1(\alpha + \beta) \\ \psi_1(\alpha + \beta) & \psi_1(\alpha + \beta) - \psi_1(\beta) \end{pmatrix}.$$

Then we can find the estimators numerically, using Newton's procedure. The initial values of estimators are found via method of moments:

$$\hat{\alpha}_0 = \bar{X}_n \left(\frac{\bar{X}_n(1 - \bar{X}_n)}{\hat{s}_n^2} - 1 \right),$$

$$\hat{\beta}_0 = (1 - \bar{X}_n) \left(\frac{\bar{X}_n(1 - \bar{X}_n)}{\hat{s}_n^2} - 1 \right).$$

These values are applicable only if $\hat{s}_n^2 < \bar{X}_n(1 - \bar{X}_n)$. If this condition is not satisfied, we set $\hat{\alpha}_0 = \hat{\beta}_0 = 0.001$.

In the general case, when $a \neq 0$ or $b \neq 1$, we use the following transformation:

$$Y_i = \frac{X_i - a}{b - a}$$

and estimate parameters, using sample Y .

2.1 Arcsine distribution

Relation to Beta distribution:

$$X \sim B(1 - \alpha, \alpha, a, b).$$

Estimation of shape. For Arcsine distribution log-likelihood function (1) turns into

$$\ln \mathcal{L}(\alpha|X) = -\alpha \sum_{i=1}^n \ln X_i + (\alpha - 1) \sum_{i=1}^n \ln(1 - X_i) - n \ln B(1 - \alpha, \alpha).$$

Taking the derivative with respect to α we get

$$\frac{\partial \ln \mathcal{L}(\alpha|X)}{\partial \alpha} = \sum_{i=1}^n \ln \frac{1 - X_i}{X_i} + n\pi \cot(\pi\alpha).$$

Therefore, maximum-likelihood function is

$$\hat{\alpha} = -\frac{1}{\pi} \operatorname{atan} \left(\frac{n\pi}{\sum_{i=1}^n \ln \frac{1 - X_i}{X_i}} \right).$$

If $\hat{\alpha}$ is negative, we add 1, because $\frac{\operatorname{atan}}{\pi} \in (-0.5, 0.5)$, while $\alpha \in (0, 1)$.

2.2 Balding-Nichols distribution

Notation:


$$X \sim \text{Balding} - \text{Nichols}(p, F)$$

with $p, F \in (0, 1)$. Relation to Beta distribution:

$$X \sim B(pF', (1-p)F')$$

with $F' = (1-F)/F$.

2.3 Uniform distribution



Notation	$X \sim \mathcal{U}(a, b)$
Parameters	$a, b \in \mathbb{R}$
Support	$x \in [a, b]$
$f(x)$	$\frac{1}{b-a}$
$F(x)$	$\frac{x-a}{b-a}$
$\mathbb{E}[X]$	$\frac{a+b}{2}$
$\text{Var}(X)$	$\frac{(b-a)^2}{12}$
Median	$\frac{a+b}{2}$
Mode	doesn't exist
$\phi(t)$	$\frac{e^{itb} - e^{ita}}{it(b-a)}$

Relation to Beta distribution:

$$X \sim B(1, 1, a, b).$$

Estimation of support.

Frequentist inference. Likelihood function is

$$\mathcal{L}(a, b|X) = \frac{1}{(b-a)^n} \mathbf{1}_{\{X_i \in [a, b] \ \forall i=1, \dots, n\}}.$$

Therefore, $\mathcal{L}(a, b|X)$ is the largest for $\hat{b} = X_{(n)}$ and $\hat{a} = X_{(1)}$. However, using the fact that $X_{(k)} \sim B(k, n+1-k, a, b)$, these are biased estimators:

$$\mathbb{E}[X_{(1)}] = \frac{an+b}{n+1} \quad \text{and} \quad \mathbb{E}[X_{(n)}] = \frac{a+bn}{n+1}.$$

To get unbiased estimators we make the transformations:

$$\tilde{a} = \frac{nX_{(1)} - X_{(n)}}{n-1} \quad \text{and} \quad \tilde{b} = \frac{nX_{(n)} - X_{(1)}}{n-1}.$$

Then we get

$$\mathbb{E}[\tilde{a}] = \frac{n\mathbb{E}[X_{(1)}] - \mathbb{E}[X_{(n)}]}{n-1} = \frac{n(an+b) - (a+bn)}{n^2-1} = a.$$

Analogously, $\mathbb{E}[\tilde{b}] = b$.

Bayesian inference. Let us say, we try to estimate $\theta = b - a$ with known a . We set the prior distribution $\theta \sim \text{Pareto}(\alpha, \sigma)$:

$$h(\theta|\alpha, \sigma) = \frac{\alpha\sigma^\alpha}{\theta^{\alpha+1}} \mathbf{1}_{\{\theta \geq \sigma\}}.$$

The density of posterior distribution is

$$f(\theta|X) \propto \frac{\alpha\sigma^\alpha}{\theta^{\alpha+n+1}} \mathbf{1}_{\{\theta \geq \max(\sigma, X_{(n)} - a)\}} \sim \text{Pareto}(\alpha + n, \max(\sigma, X_{(n)} - a)).$$

Hence, Bayesian estimator is

$$\mathbb{E}[\theta|X] = \frac{\alpha + n}{\alpha + n - 1} \max(\sigma, X_{(n)} - a)$$

and MAP estimator is

$$\theta_{MAP} = \max(\sigma, X_{(n)} - a).$$

3 Beta-prime distribution


Relation to other distributions:

$$\begin{aligned} \frac{X}{1+X} &\sim B(\alpha, \beta), \\ \frac{\beta}{\alpha} X &\sim F(2\alpha, 2\beta). \end{aligned}$$

Estimation of shapes. Using relationship with Beta distribution we transform the sample:

$$Y_i = \frac{X_i}{1+X_i}, \quad 1 \leq i \leq N,$$

and run BetaRand estimation for Y .



Notation	$X \sim B'(\alpha, \beta)$
Parameters	$\alpha, \beta > 0$
Support	$x \in \mathbb{R}^+$
$f(x)$	$\frac{x^{\alpha-1}(1+x)^{-\alpha-\beta}}{B(\alpha, \beta)}$
$F(x)$	$I_{\frac{x}{1+x}}(\alpha, \beta)$
$\mathbb{E}[X]$	$\frac{\alpha}{\beta-1} \mathbf{1}_{\{\beta>1\}} + \infty \mathbf{1}_{\{\beta \leq 1\}}$
$\text{Var}(X)$	$\frac{\alpha(\alpha+\beta-1)}{(\beta-2)(\beta-1)^2}$, if $\beta > 1$
Median	Searched numerically
Mode	$\max\left(\frac{\alpha-1}{\beta+1}, 0\right)$.
$\phi(t)$	Calculated numerically

4 Exponentially-modified Gaussian distribution

Notation	$X \sim \text{EMG}(\mu, \sigma, \lambda)$
Parameters	$\mu \in \mathbb{R}, \sigma > 0, \lambda > 0$
Support	$x \in \mathbb{R}$
$f(x)$...
$F(x)$...
$\mathbb{E}[X]$	$\mu + 1/\lambda$
$\text{Var}(X)$	$\sigma^2 + 1/\lambda^2$
Median	Searched numerically
Mode	Searched numerically
$\phi(t)$...

5 F-distribution

Notation	$X \sim F(d_1, d_2)$
Parameters	$d_1, d_2 > 0$
Support	$x \in \mathbb{R}^+$
$f(x)$	$\frac{\sqrt{\frac{(d_1 x)^{d_1} d_2^{d_2}}{(d_1 x + d_2)^{d_1 + d_2}}}}{x B\left(\frac{d_1}{2}, \frac{d_2}{2}\right)}$
$F(x)$	$I_{\frac{d_1 x}{d_1 x + d_2}}\left(\frac{d_1}{2}, \frac{d_2}{2}\right)$
$\mathbb{E}[X]$	$\frac{d_2}{d_2 - 2}$ for $d_2 > 2$
$\text{Var}(X)$	$\frac{2d_2^2(d_1 + d_2 - 2)}{d_1(d_2 - 2)^2(d_2 - 4)}$ for $d_2 > 4$
Median	Searched numerically
Mode	$\max\left(\frac{d_2(d_1 - 2)}{d_1(d_1 + 2)}, 0\right)$
$\phi(t)$	Calculated numerically

Relation to other distributions:

$$\frac{d_1 X}{d_2 + d_1 X} \sim B\left(\frac{d_1}{2}, \frac{d_2}{2}\right),$$

$$\frac{d_1}{d_2} X \sim B'\left(\frac{d_1}{2}, \frac{d_2}{2}\right).$$

6 Gamma distribution


Estimation of parameters.

Frequentist inference. Log-likelihood function:

$$\ln \mathcal{L}(\alpha, \beta | X) = n\alpha \ln \beta - n \ln \Gamma(\alpha) + (\alpha - 1) \sum_{i=1}^n \ln X_i - \beta \sum_{i=1}^n X_i.$$

Derivatives:

$$\frac{\partial \ln \mathcal{L}(\alpha, \beta | X)}{\partial \alpha} = n \ln \beta - n\psi(\alpha) + \sum_{i=1}^n \ln X_i,$$



Notation	$X \sim \Gamma(\alpha, \beta)$
Parameters	$\alpha > 0, \beta > 0$
Support	$x \in \mathbb{R}^+$
$f(x)$	$\frac{\beta^\alpha}{\Gamma(\alpha)} x^{\alpha-1} e^{-\beta x}$
$F(x)$	$P(\alpha, \beta x)$
$\mathbb{E}[X]$	$\frac{\alpha}{\beta}$
$\text{Var}(X)$	$\frac{\alpha}{\beta^2}$
Median	Searched numerically
Mode	$\max\left(\frac{\alpha-1}{\beta}, 0\right)$
$\phi(t)$	$\left(1 - \frac{it}{\beta}\right)^{-\alpha}$

$$\frac{\partial \ln \mathcal{L}(\alpha, \beta | X)}{\partial \beta} = \frac{n\alpha}{\beta} - \sum_{i=1}^n X_i.$$

While the solution for the second equation is analytic:

$$\hat{\beta} = \frac{\alpha}{\bar{X}_n},$$

the first equation is solved numerically, using second derivative:

$$\frac{\partial^2 \ln \mathcal{L}(\alpha, \beta | X)}{\partial \alpha^2} = -n\psi_1(\alpha),$$

or if β is unknown:

$$\frac{\partial^2 \ln \mathcal{L}(\alpha, \beta | X)}{\partial \alpha^2} = -n\psi_1(\alpha) + \frac{n}{\alpha},$$

Moreover, the maximum-likelihood estimation of rate β is biased. Unbiased estimator would be

$$\tilde{\beta} = \frac{\alpha}{\bar{X}_n} \left(1 - \frac{1}{n}\right).$$

Bayesian inference. We suppose that prior distribution of rate β is $\Gamma(\kappa, \theta)$:

$$h(\beta) = \frac{\theta^\kappa}{\Gamma(\kappa)} \beta^{\kappa-1} e^{-\theta\beta}.$$

Then

$$f(\beta|X) \propto \beta^{\alpha n} e^{-\beta \sum_{i=1}^n X_i} \cdot \beta^{\kappa-1} e^{-\theta\beta} \sim \Gamma\left(\alpha n + \kappa, \theta + \sum_{i=1}^n X_i\right).$$

Therefore, Bayesian estimator is

$$\mathbb{E}[\beta|X] = \frac{\alpha n + \kappa}{\theta + \sum_{i=1}^n X_i},$$

and MAP estimator is

$$\beta_{MAP} = \frac{\alpha n + \kappa - 1}{\theta + \sum_{i=1}^n X_i}.$$


6.1 Chi-squared distribution

Relation to Gamma distribution:

6.2 Erlang distribution

The only difference between Gamma and Erlang distributions is that a second one takes an integer shape parameter k .


6.3 Exponential distribution



Notation	$X \sim \text{Exp}(\lambda)$
Parameters	$\lambda > 0$
Support	$x \in \mathbb{R}^+$
$f(x)$	$\lambda e^{-\lambda x}$
$F(x)$	$1 - e^{-\lambda x}$
$\mathbb{E}[X]$	$\frac{1}{\lambda}$
$\text{Var}(X)$	$\frac{1}{\lambda^2}$
Median	$\frac{\ln(2)}{\lambda}$
Mode	0
$\phi(t)$	$\frac{\lambda}{\lambda - it}$

Relation to Gamma distribution: $X \sim \Gamma(1, \lambda)$. Hence, estimation of parameter λ is the particular case of estimation of rate β for Gamma distribution.

7 Geometric Stable distribution



Notation	$X \sim \text{GS}_\alpha(\beta, \gamma, \mu)$
Parameters	$\alpha \in (0, 2], \beta \in [-1, 1], \gamma > 0, \mu \in \mathbb{R}$
Support	$x \in \dots$
$f(x)$	Calculated numerically
$F(x)$	Calculated numerically
$\mathbb{E}[X]$	$k + \lambda$
$\text{Var}(X)$	$2(k + 2\lambda)$
Median	Searched numerically
Mode	Searched numerically
$\phi(t)$	\dots

7.1 Asymmetric Laplace distribution

7.2 Laplace distribution

8 Noncentral Chi-Squared distribution


9 Planck distribution

10 Stable distribution


10.1 Normal distribution

Relation to Stable distribution:


$$X \sim S_2(\cdot, \sigma^2/2, \mu)$$



Notation	$X \sim \chi_k'^2(\lambda)$
Parameters	$k > 0, \lambda > 0$
Support	$x \in \mathbb{R}^+$
$f(x)$...
$F(x)$	$P_{\frac{k}{2}}(\dots)$
$\mathbb{E}[X]$	$k + \lambda$
$\text{Var}(X)$	$2(k + 2\lambda)$
Median	Searched numerically
Mode	Searched numerically
$\phi(t)$	$\frac{\exp \frac{it\lambda}{1-2it}}{(1-2it)^{k/2}}$



Notation	$X \sim \text{Planck}(a, b)$
Parameters	$a, b > 0$
Support	$x \in \mathbb{R}^+$
$f(x)$	$\frac{b^{a+1}}{\Gamma(a+1)\zeta(a+1)} \cdot \frac{x^a}{e^{bx}-1}$
$F(x)$	Calculated numerically
$\mathbb{E}[X]$	$\frac{(a+1)\zeta(a+2)}{b\zeta(a+1)}$
$\text{Var}(X)$	$\frac{(a+1)(a+2)\zeta(a+3)}{b^2\zeta(a+1)} - (\mathbb{E}[X])^2$
Median	Searched numerically
Mode	$\frac{W_0(-ae^{-a})+a}{b}$, if $a > 1$, otherwise 0
$\phi(t)$	Calculated numerically



Notation	$X \sim S_{\alpha}(\beta, \gamma, \mu)$
Parameters	$\alpha \in (0, 2], \beta \in [-1, 1],$ $\gamma > 0, \mu \in \mathbb{R}$
Support	$x \in \mathbb{R}$, if $\beta \neq 1$, $x \in [\mu, \infty)$, if $\beta = 1, \alpha < 2$, $x \in (-\infty, \mu]$, if $\beta = -1, \alpha < 2$
$f(x)$	Calculated numerically
$F(x)$	Calculated numerically
$\mathbb{E}[X]$	μ for $\alpha > 1$, otherwise undefined
$\text{Var}(X)$	$2\gamma^2 1_{\{\alpha=2\}} + \infty 1_{\{\alpha<2\}}$
Median	μ for $\beta = 0$, otherwise searched numerically
Mode	μ , if $\beta = 0$ or $\alpha = 2$, $\mu + \frac{\beta\gamma}{3}$, if $ \beta = 1$ and $\alpha = \frac{1}{2}$, otherwise searched numerically
$\phi(t)$...

10.2 Cauchy distribution

Relation to Stable distribution:

$$X \sim S_1(0, \gamma, \mu)$$

10.3 Levy distribution


Relation to Stable distribution:

$$X \sim S_{\frac{1}{2}}(1, \gamma, \mu)$$

10.4 Holtsmark distribution

Relation to Stable distribution:

$$X \sim S_{\frac{3}{2}}(0, \gamma, \mu)$$



Notation	$X \sim \mathcal{N}(\mu, \sigma^2)$
Parameters	$\mu \in \mathbb{R}, \sigma^2 > 0$
Support	$x \in \mathbb{R}$
$f(x)$	$\frac{1}{\sqrt{2\sigma^2\pi}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$
$F(x)$	$\frac{1}{2}\text{erfc}\left(\frac{\mu-x}{\sqrt{2\sigma^2}}\right)$
$\mathbb{E}[X]$	μ
$\text{Var}(X)$	σ^2
Median	μ
Mode	μ
$\phi(t)$	$e^{i\mu t - \frac{1}{2}\sigma^2 t^2}$

10.5 Landau distribution

Relation to Stable distribution:

$$X \sim S_1(1, \gamma, \mu)$$

11 Pareto distribution


Estimation of parameters.

Frequentist inference. Log-likelihood function is

$$\ln \mathcal{L}(\alpha, \sigma | X) = n \ln \alpha + n \alpha \ln \sigma - (\alpha + 1) \sum_{i=1}^n \ln X_i.$$

We assume that $\sigma \leq X_{(1)}$, otherwise sample X couldn't have been generated from such distribution. It is obvious, that $\ln \mathcal{L}(\alpha, \sigma | X)$ is an increasing function in terms of σ , therefore $\hat{\sigma} = X_{(1)}$ is an optimal estimator. Let's take derivative with respect to α :

$$\frac{\partial \ln \mathcal{L}(\alpha, \sigma | X)}{\partial \alpha} = \frac{n}{\alpha} + n \ln \sigma - \sum_{i=1}^n \ln X_i.$$



Notation	$X \sim \text{Cauchy}(\mu, \gamma)$
Parameters	$\mu \in \mathbb{R}, \gamma^2 > 0$
Support	$x \in \mathbb{R}$
$f(x)$	$\frac{1}{\pi\gamma \left[1 + \left(\frac{x-\mu}{\gamma}\right)^2\right]}$
$F(x)$	$\frac{1}{\pi} \text{atan}\left(\frac{x-\mu}{\gamma}\right) + \frac{1}{2}$
$\mathbb{E}[X]$	Undefined
$\text{Var}(X)$	∞
Median	μ
Mode	μ
$\phi(t)$	$e^{i\mu t - \gamma t }$

From this we conclude that the maximum-likelihood estimator of shape is

$$\hat{\alpha} = \frac{1}{\frac{1}{n}(\sum_{i=1}^n \ln X_i) - \ln \hat{\sigma}}.$$

It is known that $\hat{\sigma} \sim \text{Pareto}(n\alpha, \sigma)$ and $\hat{\alpha} \sim \text{Inv-}\Gamma(n-1, n\alpha)$ and they are independent. Then

$$\mathbb{E}[\hat{\sigma}] = \frac{\sigma}{1 - \frac{1}{n\alpha}}$$

and

$$\mathbb{E}[\hat{\alpha}] = \frac{n\alpha}{n-2}.$$


Therefore, in order to get unbiased estimators we need to make the following transformations:

$$\tilde{\alpha} = \frac{n-2}{n} \hat{\alpha} \quad \text{and} \quad \tilde{\sigma} = \hat{\sigma} \left(1 - \frac{1}{(n-1)\hat{\alpha}}\right).$$

Note that if we estimate parameters separately, then $\hat{\alpha} \sim \text{Inv-}\Gamma(n, n\alpha)$ and transformations are different.

Bayesian inference. We now assume that σ is known and prior distribution of α is $\Gamma(\kappa, \beta)$:

$$h(\alpha) = \frac{\beta^\kappa}{\Gamma(\kappa)} \alpha^{\kappa-1} e^{-\beta\alpha}.$$



Notation	$X \sim \text{Pareto}(\alpha, \sigma)$
Parameters	$\alpha, \sigma > 0$
Support	$x \geq \sigma$
$f(x)$	$\frac{\alpha \sigma^\alpha}{x^{\alpha+1}}$
$F(x)$	$1 - \left(\frac{\sigma}{x}\right)^\alpha$
$\mathbb{E}[X]$	$\frac{\alpha \sigma}{\alpha - 1}$ for $\alpha > 1$, ∞ otherwise
$\text{Var}(X)$	$\frac{\sigma^2 \alpha}{(\alpha - 1)^2 (\alpha - 2)}$ for $\alpha > 2$, ∞ otherwise
Median	$\sigma 2^{1/\alpha}$
Mode	σ
$\phi(t)$	Calculated numerically

The density of posterior distribution is

$$f(\alpha|X) \propto \prod_{i=1}^n \frac{\sigma^\alpha}{X_i^{\alpha+1}} \cdot \alpha^{\kappa+n-1} e^{-\beta\alpha} \propto \alpha^{\kappa+n-1} e^{-(\beta + \sum_{i=1}^n \ln(X_i/\sigma))\alpha}.$$

Therefore, $\alpha|X \sim \Gamma(\kappa + n, \beta + \sum_{i=1}^n \ln(X_i/\sigma))$ and Bayesian estimator is

$$\mathbb{E}[\alpha|X] = \frac{\kappa + n}{\beta + \sum_{i=1}^n \ln(X_i/\sigma)}.$$

MAP estimator is

$$\alpha_{MAP} = \frac{\kappa + n - 1}{\beta + \sum_{i=1}^n \ln(X_i/\sigma)}.$$

Note on fitting scale with Bayes: let it be vice versa, α is known while σ is not. Then we say that a priori $\sigma \sim \text{Pareto}(\kappa, \theta)$:

$$h(\sigma) = \frac{\kappa \theta^\kappa}{\sigma^{\kappa+1}}.$$

Then posterior distribution is:

$$f(\sigma|X) \propto \prod_{i=1}^n \frac{1}{X_i^{\alpha+1}} \cdot \sigma^{\alpha n - \kappa - 1} \mathbf{1}_{\{\theta < \sigma < X_{(1)}\}} \sim \text{Bounded-Pareto}(\kappa - \alpha n, \theta, X_{(1)}).$$

This imposes the following additional constraints on the prior hyperparameters: $\kappa > \alpha n$ and $\theta < X_{(1)}$. Bayesian estimator:


$$\mathbb{E}[\sigma|X] = \frac{\theta^{\alpha'}}{1 - \left(\frac{\theta}{X_{(1)}}\right)^{\alpha'}} \cdot \left(\frac{\alpha'}{\alpha' - 1}\right) \cdot \left(\frac{1}{\theta^{\alpha'}} - \frac{1}{X_{(1)}^{\alpha'}}\right)$$

with $\alpha' = \kappa - \alpha n$. MAP estimator is just

$$\sigma_{MAP} = \theta.$$

However, Bounded-Pareto distribution is not yet supported.

12 Weibull



Notation	$X \sim \text{Weibull}(\lambda, k)$
Parameters	$\lambda, k > 0$
Support	$x \in \mathbb{R}^+$
$f(x)$	$\frac{k}{\lambda} \left(\frac{x}{\lambda}\right)^{k-1} \exp(-(x/\lambda)^k)$
$F(x)$	$1 - \exp(-(x/\lambda)^k)$
$\mathbb{E}[X]$	$\lambda \Gamma(1 + 1/k)$
$\text{Var}(X)$	$\lambda^2 \Gamma(1 + 2/k) - (\mathbb{E}[X])^2$
Median	$\lambda (\ln 2)^{\frac{1}{k}}$
Mode	$\lambda \left(1 - \frac{1}{k}\right)^{\frac{1}{k}}$
$\phi(t)$	Calculated numerically

Estimation of scale

Frequentist inference. Log-likelihood function:

$$\ln \mathcal{L}(\lambda, k|X) = n(\ln k - \ln \lambda) + (k - 1) \sum_{i=1}^n (\ln X_i - \ln \lambda) - \frac{1}{\lambda^k} \sum_{i=1}^n X_i^k.$$

The derivative with respect to scale:

$$\frac{\partial \ln \mathcal{L}(\lambda, k|X)}{\partial \lambda} = -\frac{nk}{\lambda} + \frac{k}{\lambda^{k+1}} \sum_{i=1}^n X_i^k = 0.$$

Therefore, maximum-likelihood estimation for λ is

$$\hat{\lambda} = \left(\sum_{i=1}^n X_i^k \right)^{\frac{1}{k}}.$$

Bayesian inference. Assume k is known. Instead of estimating λ we give an estimation for λ^k . Let's say that prior distribution of λ^k is $\text{Inv-}\Gamma(\alpha, \beta)$:

$$h(\lambda^k) = \frac{\beta^\alpha}{\Gamma(\alpha)} \lambda^{-k(\alpha+1)} e^{-\beta/\lambda^k}.$$

Posterior distribution then:

$$f(\lambda^k|X) \propto \lambda^{-k(\alpha+1+n)} e^{-\frac{1}{\lambda^k}(\beta + \sum_{i=1}^n X_i^k)} \sim \text{Inv-}\Gamma(\alpha + n, \beta + \sum_{i=1}^n X_i^k).$$

Bayesian estimator:

$$\mathbb{E}[\lambda^k|X] = \frac{\beta + \sum_{i=1}^n X_i^k}{\alpha + n - 1},$$

MAP estimator:

$$\lambda_{MAP}^k = \frac{\beta + \sum_{i=1}^n X_i^k}{\alpha + n + 1}.$$

Part III

Discrete univariate distributions

13 Beta-binomial distribution

14 Binomial distribution

Notation	$X \sim \text{Bin}(n, p)$
Parameters	$n \in \mathbb{N}, p \in [0, 1]$
Support	$k \in \{0, \dots, n\}$
P.m.f.	$\mathbb{P}(X = k) = \binom{n}{k} p^k (1 - p)^{n-k}$
$F(x)$	$\mathbb{P}(X \leq k) = I_{1-p}(n - k, 1 + k)$
$\mathbb{E}[X]$	np
$\text{Var}(X)$	$np(1 - p)$
Median	$[np]$
Mode	$[(n + 1)p]$
$\phi(t)$	$(1 - p + pe^{it})^n$

14.1 Bernoulli

Notation:

$$X \sim \text{Bernoulli}(p).$$

Relation to Binomial distribution:

$$X \sim \text{Bin}(1, p).$$

15 Poisson distribution

Notation	$X \sim \text{Po}(\lambda)$
Parameters	$\lambda > 0$
Support	$k \in \mathbb{N}_0$
P.m.f.	$\mathbb{P}(X = k) = \frac{\lambda^k e^{-\lambda}}{k!}$
$F(x)$	$\mathbb{P}(X \leq k) = Q(k+1, \lambda)$
$\mathbb{E}[X]$	λ
$\text{Var}(X)$	λ
Median	$\sim \max\left([\lambda + \frac{1}{3} - \frac{0.02}{\lambda}], 0\right)$
Mode	$[\lambda]$
$\phi(t)$	$\exp\{\lambda(e^{it} - 1)\}$

Generator (let $\delta = \mu \in \mathbb{Z}$). (There is a mistake in Lemma 3.8 in first inequality). Recall that

$$q(X) = X \ln(\lambda) - \ln\left(\frac{(\mu + X)!}{\mu!}\right).$$

We denote acceptance probability $\mathbb{P}(W \leq q(X))$ by p .

- $k = \mu$. Probability to be in this setting is $1/c$.

$$\mathbb{P}(X = 0 | W \leq q(X)) = \frac{\mathbb{P}(X = 0, W \leq q(X))}{\mathbb{P}(W \leq q(X))} = \frac{1}{pc}.$$

On the other hand it should be equal to:

$$\frac{1}{pc} = \frac{\lambda^\mu e^{-\lambda}}{\mu!}.$$

- $k = \mu + 1$.

$$\begin{aligned} \mathbb{P}(X = 1 | W \leq q(X)) &= \frac{\mathbb{P}(X = 1, W \leq q(X))}{\mathbb{P}(W \leq q(X))} = \frac{\lambda}{p(\mu + 1)c} \\ &= \frac{\lambda^{\mu+1} e^{-\lambda}}{(\mu + 1)!}. \end{aligned}$$

- $k < \mu$. Here was mistake in the book. We adjust the probabilities. Probability to be in this setting is $\sqrt{\pi\mu/2e}/c$.

$$\begin{aligned}
\mathbb{P}(W \leq q(X), X = k - \mu | U \leq c_1) &= \mathbb{P}\left(-\frac{N^2}{2} + \frac{1}{2} - E < q(\lfloor -|N|\sqrt{\mu} \rfloor), \lceil |N|\sqrt{\mu} \rceil = \mu - k\right) \\
&= \mathbb{P}\left(-\frac{N^2}{2} + \frac{1}{2} - E < \lfloor -|N|\sqrt{\mu} \rfloor \ln(\lambda) - \ln\left(\frac{(\mu + \lfloor -|N|\sqrt{\mu} \rfloor)!}{\mu!}\right), \right. \\
&\quad \left. \frac{\mu - k - 1}{\sqrt{\mu}} \leq |N| < \frac{\mu - k}{\sqrt{\mu}}\right) \\
&= \mathbb{P}\left(U < \exp\left\{\frac{N^2}{2} - \frac{1}{2} + \lfloor -|N|\sqrt{\mu} \rfloor \ln(\lambda) - \ln\left(\frac{(\mu + \lfloor -|N|\sqrt{\mu} \rfloor)!}{\mu!}\right)\right\}, \right. \\
&\quad \left. \frac{\mu - k - 1}{\sqrt{\mu}} \leq |N| < \frac{\mu - k}{\sqrt{\mu}}\right) \\
&= \sqrt{\frac{2}{e\pi}} \int_{\frac{\mu-k-1}{\sqrt{\mu}}}^{\frac{\mu-k}{\sqrt{\mu}}} \exp\left\{\lfloor -|n|\sqrt{\mu} \rfloor \ln(\lambda) - \ln\left(\frac{(\mu + \lfloor -|n|\sqrt{\mu} \rfloor)!}{\mu!}\right)\right\} dn \\
&= \sqrt{\frac{2}{e\pi\mu}} \int_{\mu-k-1}^{\mu-k} \exp\left\{\lfloor -z \rfloor \ln(\lambda) - \ln\left(\frac{(\mu + \lfloor -z \rfloor)!}{\mu!}\right)\right\} dz \\
&= \sqrt{\frac{2}{e\pi\mu}} \exp\left\{(k - \mu) \ln(\lambda) - \ln\left(\frac{k!}{\mu!}\right)\right\} \\
&= \sqrt{\frac{2}{e\pi\mu}} \lambda^{k-\mu} \frac{\mu!}{k!}
\end{aligned}$$

Hence,

$$\begin{aligned}
\mathbb{P}(X = k - \mu | W \leq q(X)) &= \frac{\mathbb{P}(W \leq q(X), X = k - \mu)}{\mathbb{P}(W \leq q(X))} \\
&= \sqrt{\frac{2}{\pi\mu e}} \lambda^{k-\mu} \frac{\mu!}{k!} \cdot \sqrt{\pi\mu e/2} \frac{\lambda^\mu e^{-\lambda}}{\mu!} \\
&= \frac{\lambda^k e^{-\lambda}}{k!}
\end{aligned}$$

- $k \in [\mu + 2, 2\mu]$. Probability to be in this setting is $\sqrt{\frac{3\pi\mu}{4}} e^{\frac{1}{3\mu}}/c$. We also have

$$W = \frac{-Y^2 + 2Y}{3\mu} - E = \frac{1}{3\mu} - \frac{N^2}{2} - E.$$

Then

$$\begin{aligned}
\mathbb{P}(W \leq q(X)|X = k - \mu|U \in \dots) &= \mathbb{P}\left(\frac{1}{3\mu} - \frac{N^2}{2} - E < q(\lceil 1 + |N|\sqrt{3\mu/2} \rceil), \lceil 1 + |N|\sqrt{3\mu/2} \rceil = k - \mu\right) \\
&= \mathbb{P}\left(U < \exp\left\{-\frac{1}{3\mu} + \frac{N^2}{2} + q(\lceil 1 + |N|\sqrt{3\mu/2} \rceil)\right\}, \right. \\
&\quad \left. \frac{k - \mu - 2}{\sqrt{3\mu/2}} < |N| \leq \frac{k - \mu - 1}{\sqrt{3\mu/2}}\right) \\
&= \sqrt{\frac{2}{\pi}} e^{-\frac{1}{3\mu}} \int_{\frac{k-\mu-2}{\sqrt{3\mu/2}}}^{\frac{k-\mu-1}{\sqrt{3\mu/2}}} \exp\left\{q(\lceil 1 + |n|\sqrt{3\mu/2} \rceil)\right\} dn \\
&= \sqrt{\frac{4}{3\pi\mu}} e^{-\frac{1}{3\mu}} \int_{k-\mu-1}^{k-\mu} \exp\left\{\lceil z \rceil \ln(\lambda) - \ln\left(\frac{(\mu + \lceil z \rceil)!}{\mu!}\right)\right\} dz \\
&= \sqrt{\frac{4}{3\pi\mu}} e^{-\frac{1}{3\mu}} \mu! \frac{\lambda^{k-\mu}}{k!}.
\end{aligned}$$

- $k > 2\mu$. Probability to be in this setting is $6e^{-\frac{2+\mu}{6}}/c$.

$$\begin{aligned}
\mathbb{P}(W \leq q(X)|X = k - \mu|U \in \dots) &= \mathbb{P}\left(-\frac{2+\mu}{6} - V - E < q(\lceil \mu + 6V \rceil), \lceil \mu + 6V \rceil = k - \mu\right) \\
&= \mathbb{P}\left(-\frac{2+\mu}{6} - V + \ln(U) < \lceil \mu + 6V \rceil \ln(\lambda) - \ln\left(\frac{(\mu + \lceil \mu + 6V \rceil)!}{\mu!}\right), \right. \\
&\quad \left. \lceil \mu + 6V \rceil = k - \mu\right) \\
&= \mathbb{P}\left(U < \exp\left\{\frac{2+\mu}{6} + V + \lceil \mu + 6V \rceil \ln(\lambda) - \ln\left(\frac{(\mu + \lceil \mu + 6V \rceil)!}{\mu!}\right)\right\}, \right. \\
&\quad \left. \frac{k - 2\mu - 1}{6} < V \leq \frac{k - 2\mu}{6}\right) \\
&= \int_{\frac{k-2\mu-1}{6}}^{\frac{k-2\mu}{6}} \exp\left\{\frac{2+\mu}{6} + \lceil \mu + 6v \rceil \ln(\lambda) - \ln\left(\frac{(\mu + \lceil \mu + 6v \rceil)!}{\mu!}\right)\right\} dv \\
&= \frac{e^{\frac{2+\lambda}{6}}}{6} \int_{k-\mu-1}^{k-\mu} \exp\left\{\lceil z \rceil \ln(\lambda) - \ln\left(\frac{(\mu + \lceil z \rceil)!}{\mu!}\right)\right\} dz \\
&= \frac{e^{\frac{2+\lambda}{6}}}{6} \exp\left\{(k - \mu) \ln(\lambda) - \ln\left(\frac{k!}{\mu!}\right)\right\} \\
&= \frac{e^{\frac{2+\lambda}{6}}}{6} \lambda^{k-\mu} \frac{\mu!}{k!}
\end{aligned}$$

Hence,

$$\begin{aligned}
\mathbb{P}(X = k - \mu | W \leq q(X)) &= \frac{\mathbb{P}(W \leq q(X), X = k - \mu)}{\mathbb{P}(W \leq q(X))} \\
&= \frac{e^{\frac{2+\lambda}{6}}}{6} \lambda^{k-\mu} \frac{\mu!}{k!} \cdot \frac{6e^{-\frac{2+\mu}{6}}}{pc} \\
&= \frac{\lambda^k e^{-\lambda}}{k!}
\end{aligned}$$

Part IV

Bivariate distributions

16 Bivariate Normal distribution

17 Normal-Inverse-Gamma distribution

18 Trinomial distribution

Part V

Circular distributions

19 von Mises distribution

20 Wrapped Exponential distribution

Part VI

Singular distributions

21 Cantor distribution