

RandLib documentation

Aleksandr Samarin

November 11, 2017

Contents

I	General information	2
1	Calculation of sample moments	3
II	Continuous univariate distributions	4
2	Beta distribution	4
2.1	Arcsine distribution	5
2.2	Balding-Nichols distribution	6
2.3	Uniform distribution	6
3	Beta-prime distribution	7
4	Degenerate distribution	8
5	Exponentially-modified Gaussian distribution	9
6	F-distribution	10
7	Gamma distribution	10
7.1	Chi-squared distribution	10
7.2	Erlang distribution	10
7.3	Exponential distribution	11
8	Geometric Stable distribution	12
8.1	Asymmetric Laplace distribution	12
8.2	Laplace distribution	12
9	Noncentral Chi-Squared distribution	12

10 Planck distribution	12
11 Stable distribution	12
11.1 Normal distribution	12
11.2 Cauchy distribution	14
11.3 Levy distribution	14
11.4 Holtsmark distribution	14
11.5 Landau distribution	15
12 Pareto distribution	15
13 Weibull	18
 III Discrete univariate distributions	 20
14 Beta-binomial distribution	20
15 Binomial distribution	20
15.1 Bernoulli	20
16 Poisson distribution	21
 IV Bivariate distributions	 25
17 Bivariate Normal distribution	25
18 Normal-Inverse-Gamma distribution	25
19 Trinomial distribution	25
 V Circular distributions	 26
20 von Mises distribution	26
21 Wrapped Exponential distribution	26
 VI Singular distributions	 27
22 Cantor distribution	27

Part I

General information

1 Calculation of sample moments

We use extension of Welford's method from Knuth. For every n -th element x we have


$$\begin{aligned}\delta &= x - m_1, \\ m'_1 &= m_1 + \frac{\delta}{n}, \\ m'_2 &= m_2 + \delta^2 \frac{n-1}{n}, \\ m'_3 &= m_3 + \delta^3 \frac{(n-1)(n-2)}{n^2} - 3\delta \frac{m_2}{n}, \\ m'_4 &= m_4 + \delta^4 \frac{(n-1)(n^2-3n+3)}{n^3} + 6\delta^2 \frac{m_2}{n^2} - 4\delta \frac{m_3}{n}.\end{aligned}$$

Then $\mathbb{E}[X] = m'_1$, $\text{Var}(X) = \frac{m_2}{n}$, $\text{Skew}(X) = \frac{\sqrt{n}m'_3}{m_2^{3/2}}$ and $\text{Kurt}(X) = \frac{nm'_4}{m_2'^2}$ (we return excess kurtosis).

Part II

Continuous univariate distributions

2 Beta distribution



Notation	$X \sim B(\alpha, \beta, a, b)$ $X \sim B(\alpha, \beta)$ with $a = 0, b = 1$
Parameters	$\alpha, \beta > 0, a, b \in \mathbb{R}$
Support	$x \in [a, b]$
P.d.f.	$f(x) = \frac{y^{\alpha-1}(1-y)^{\beta-1}}{(b-a)B(\alpha, \beta)}$ with $y = \frac{x-a}{b-a}$
C.d.f.	$F(x) = I_y(\alpha, \beta)$ for $y = \frac{x-a}{b-a}$
Expectation	$\mathbb{E}[X] = a + (b-a) \frac{\alpha}{\alpha+\beta}$
Variance	$\text{Var}(X) = (b-a)^2 \frac{\alpha\beta}{(\alpha+\beta)^2(\alpha+\beta+1)}$
Median	Searched numerically
Mode	$a + (b-a) \frac{\alpha-\beta}{\alpha+\beta-2}$ for $\alpha, \beta > 1$.
C.f.	Calculated numerically

Estimation of shapes with known support. Assume that $a = 0, b = 1$ and we have a sample $X = (X_1, \dots, X_n)$. Then a log-likelihood function is

$$\begin{aligned}
 \ln \mathcal{L}(\alpha, \beta | X) &= \sum_{i=1}^n \ln f(X_i; \alpha, \beta) \\
 &= (\alpha - 1) \sum_{i=1}^n \ln X_i + (\beta - 1) \sum_{i=1}^n \ln(1 - X_i) - n \ln B(\alpha, \beta).
 \end{aligned} \tag{1}$$

Differentiating with respect to the shapes, we obtain

$$\begin{aligned}
 \frac{\partial \ln \mathcal{L}(\alpha, \beta | X)}{\partial \alpha} &= \sum_{i=1}^n \ln X_i + n(\psi(\alpha + \beta) - \psi(\alpha)), \\
 \frac{\partial \ln \mathcal{L}(\alpha, \beta | X)}{\partial \beta} &= \sum_{i=1}^n \ln(1 - X_i) + n(\psi(\alpha + \beta) - \psi(\beta)).
 \end{aligned}$$

Differentiating again we get the Hessian matrix:

$$H(\ln \mathcal{L}(\alpha, \beta|X)) = n \cdot \begin{pmatrix} \psi_1(\alpha + \beta) - \psi_1(\alpha) & \psi_1(\alpha + \beta) \\ \psi_1(\alpha + \beta) & \psi_1(\alpha + \beta) - \psi_1(\beta) \end{pmatrix}.$$

Then we can find the estimators numerically, using Newton's procedure. The initial values of estimators are found via method of moments:

$$\hat{\alpha}_0 = \bar{X}_n \left(\frac{\bar{X}_n(1 - \bar{X}_n)}{\hat{s}_n^2} - 1 \right),$$

$$\hat{\beta}_0 = (1 - \bar{X}_n) \left(\frac{\bar{X}_n(1 - \bar{X}_n)}{\hat{s}_n^2} - 1 \right).$$

These values are applicable only if $\hat{s}_n^2 < \bar{X}_n(1 - \bar{X}_n)$. If this condition is not satisfied, we set $\hat{\alpha}_0 = \hat{\beta}_0 = 0.001$.

In the general case, when $a \neq 0$ or $b \neq 1$, we use the following transformation:

$$Y_i = \frac{X_i - a}{b - a}$$

and estimate parameters, using sample Y .

2.1 Arcsine distribution

Relation to Beta distribution:

$$X \sim B(1 - \alpha, \alpha, a, b).$$

Estimation of shape. For Arcsine distribution log-likelihood function (1) turns into

$$\ln \mathcal{L}(\alpha|X) = -\alpha \sum_{i=1}^n \ln X_i + (\alpha - 1) \sum_{i=1}^n \ln(1 - X_i) - n \ln B(1 - \alpha, \alpha).$$

Taking the derivative with respect to α we get

$$\frac{\partial \ln \mathcal{L}(\alpha|X)}{\partial \alpha} = \sum_{i=1}^n \ln \frac{1 - X_i}{X_i} + n\pi \cot(\pi\alpha).$$

Therefore, maximum-likelihood function is

$$\hat{\alpha} = -\frac{1}{\pi} \operatorname{atan} \left(\frac{n\pi}{\sum_{i=1}^n \ln \frac{1 - X_i}{X_i}} \right).$$

If $\hat{\alpha}$ is negative, we add 1, because $\frac{\operatorname{atan}}{\pi} \in (-0.5, 0.5)$, while $\alpha \in (0, 1)$.

2.2 Balding-Nichols distribution

Notation:


$$X \sim \text{Balding} - \text{Nichols}(p, F)$$

with $p, F \in (0, 1)$. Relation to Beta distribution:

$$X \sim B(pF', (1-p)F')$$

with $F' = (1-F)/F$.

2.3 Uniform distribution



Notation	$X \sim \mathcal{U}(a, b)$
Parameters	$a, b \in \mathbb{R}$
Support	$x \in [a, b]$
P.d.f.	$f(x) = \frac{1}{b-a}$
C.d.f.	$F(x) = \frac{x-a}{b-a}$
Expectation	$\mathbb{E}[X] = \frac{a+b}{2}$
Variance	$\text{Var}(X) = \frac{(b-a)^2}{12}$
Median	$\frac{a+b}{2}$
Mode	doesn't exist
C.f.	$\phi(t) = \frac{e^{itb} - e^{ita}}{it(b-a)}$

Relation to Beta distribution:

$$X \sim B(1, 1, a, b).$$

Estimation of support.

Frequentist inference. Likelihood function is

$$\mathcal{L}(a, b|X) = \frac{1}{(b-a)^n} \mathbf{1}_{\{X_i \in [a, b] \ \forall i=1, \dots, n\}}.$$

Therefore, $\mathcal{L}(a, b|X)$ is the largest for $\hat{b} = X_{(n)}$ and $\hat{a} = X_{(1)}$. However, using the fact that $X_{(k)} \sim B(k, n+1-k, a, b)$, these are biased estimators:

$$\mathbb{E}[X_{(1)}] = \frac{an+b}{n+1} \quad \text{and} \quad \mathbb{E}[X_{(n)}] = \frac{a+bn}{n+1}.$$

To get unbiased estimators we make the transformations:

$$\tilde{a} = \frac{nX_{(1)} - X_{(n)}}{n-1} \quad \text{and} \quad \tilde{b} = \frac{nX_{(n)} - X_{(1)}}{n-1}.$$

Then we get

$$\mathbb{E}[\tilde{a}] = \frac{n\mathbb{E}[X_{(1)}] - \mathbb{E}[X_{(n)}]}{n-1} = \frac{n(an+b) - (a+bn)}{n^2-1} = a.$$

Analogously, $\mathbb{E}[\tilde{b}] = b$.

Bayesian inference. Let us say, we try to estimate $\theta = b - a$ with known a . We set the prior distribution $\theta \sim \text{Pareto}(\alpha, \sigma)$:

$$h(\theta|\alpha, \sigma) = \frac{\alpha\sigma^\alpha}{\theta^{\alpha+1}} \mathbf{1}_{\{\theta \geq \sigma\}}.$$

The density of posterior distribution is

$$f(\theta|X) \propto \frac{\alpha\sigma^\alpha}{\theta^{\alpha+n+1}} \mathbf{1}_{\{\theta \geq \max(\sigma, X_{(n)} - a)\}} \sim \text{Pareto}(\alpha+n, \max(\sigma, X_{(n)} - a)).$$

Hence, Bayesian estimator is

$$\mathbb{E}[\theta|X] = \frac{\alpha+n}{\alpha+n-1} \max(\sigma, X_{(n)} - a)$$

and MAP estimator is

$$\theta_{MAP} = \max(\sigma, X_{(n)} - a).$$

3 Beta-prime distribution


Relation to other distributions:

$$\begin{aligned} \frac{X}{1+X} &\sim B(\alpha, \beta), \\ \frac{\beta}{\alpha}X &\sim F(2\alpha, 2\beta). \end{aligned}$$

Estimation of shapes. Using relationship with Beta distribution we transform the sample:

$$Y_i = \frac{X_i}{1+X_i}, \quad 1 \leq i \leq N,$$

and run BetaRand estimation for Y .



Notation	$X \sim B'(\alpha, \beta)$
Parameters	$\alpha, \beta > 0$
Support	$x \in \mathbb{R}^+$
P.d.f.	$f(x) = \frac{x^{\alpha-1}(1+x)^{-\alpha-\beta}}{B(\alpha, \beta)}$
C.d.f.	$F(x) = I_{\frac{x}{1+x}}(\alpha, \beta)$
Expectation	$\mathbb{E}[X] = \frac{\alpha}{\beta-1} \mathbf{1}_{\{\beta>1\}} + \infty \mathbf{1}_{\{\beta \leq 1\}}$
Variance	$\text{Var}(X) = \frac{\alpha(\alpha+\beta-1)}{(\beta-2)(\beta-1)^2}, \text{ if } \beta > 1$
Median	Searched numerically
Mode	$\max\left(\frac{\alpha-1}{\beta+1}, 0\right).$
C.f.	Calculated numerically

4 Degenerate distribution

Notation	$X \sim \delta(a)$
Parameters	$a \in \mathbb{R}$
Support	$x = a$
P.d.f.	$f(x) = \delta(a)$
C.d.f.	$F(x) = 1_{\{x \leq a\}}$
Expectation	$\mathbb{E}[X] = a$
Variance	$\text{Var}(X) = 0$
Median	a
Mode	a
C.f.	$\phi(t) = e^{ita}$

5 Exponentially-modified Gaussian distribution

Notation	$X \sim \text{EMG}(\mu, \sigma, \lambda)$
Parameters	$\mu \in \mathbb{R}, \sigma > 0, \lambda > 0$
Support	$x \in \mathbb{R}$
P.d.f.	$f(x) = \dots$
C.d.f.	$F(x) = \dots$
Expectation	$\mathbb{E}[X] = \mu + 1/\lambda$
Variance	$\text{Var}(X) = \sigma^2 + 1/\lambda^2$
Median	Searched numerically
Mode	Searched numerically
C.f.	$\phi(t) = \dots$

6 F-distribution

Notation	$X \sim F(d_1, d_2)$
Parameters	$d_1, d_2 > 0$
Support	$x \in \mathbb{R}^+$
P.d.f.	$f(x) = \frac{\sqrt{\frac{(d_1 x)^{d_1} d_2^{d_2}}{(d_1 x + d_2)^{d_1 + d_2}}}}{x B\left(\frac{d_1}{2}, \frac{d_2}{2}\right)}$
C.d.f.	$F(x) = I_{\frac{d_1 x}{d_1 x + d_2}}\left(\frac{d_1}{2}, \frac{d_2}{2}\right)$
Expectation	$\mathbb{E}[X] = \frac{d_2}{d_2 - 2}$ for $d_2 > 2$
Variance	$\text{Var}(X) = \frac{2d_2^2(d_1 + d_2 - 2)}{d_1(d_2 - 2)^2(d_2 - 4)}$ for $d_2 > 4$
Median	Searched numerically
Mode	$\max\left(\frac{d_2(d_1 - 2)}{d_1(d_1 + 2)}, 0\right)$
C.f.	Calculated numerically

Relation to other distributions:

$$\frac{d_1 X}{d_2 + d_1 X} \sim B\left(\frac{d_1}{2}, \frac{d_2}{2}\right),$$

$$\frac{d_1}{d_2} X \sim B'\left(\frac{d_1}{2}, \frac{d_2}{2}\right).$$


7 Gamma distribution

7.1 Chi-squared distribution

Relation to Gamma distribution:

7.2 Erlang distribution

The only difference between Gamma and Erlang distributions is that a second one takes an integer shape parameter k .




Notation	$X \sim \Gamma(\alpha, \beta)$
Parameters	$\alpha > 0, \beta > 0$
Support	$x \in \mathbb{R}^+$
P.d.f.	$f(x) = \frac{\beta^\alpha}{\Gamma(\alpha)} x^{\alpha-1} e^{-\beta x}$
C.d.f.	$F(x) = P(\alpha, \beta x)$
Expectation	$\mathbb{E}[X] = \frac{\alpha}{\beta}$
Variance	$\text{Var}(X) = \frac{\alpha}{\beta^2}$
Median	Searched numerically
Mode	$\max\left(\frac{\alpha-1}{\beta}, 0\right)$
C.f.	$\phi(t) = \left(1 - \frac{it}{\beta}\right)^{-\alpha}$

7.3 Exponential distribution

Notation	$X \sim \text{Exp}(\lambda)$
Parameters	$\lambda > 0$
Support	$x \in \mathbb{R}^+$
P.d.f.	$f(x) = \lambda e^{-\lambda x}$
C.d.f.	$F(x) = 1 - e^{-\lambda x}$
Expectation	$\mathbb{E}[X] = \frac{1}{\lambda}$
Variance	$\text{Var}(X) = \frac{1}{\lambda^2}$
Median	$\frac{\ln(2)}{\lambda}$
Mode	0
C.f.	$\phi(t) = \frac{\lambda}{\lambda - it}$

Relation to Gamma distribution: $X \sim \Gamma(1, \beta)$.

8 Geometric Stable distribution



Notation	$X \sim \text{GS}_\alpha(\beta, \gamma, \mu)$
Parameters	$\alpha \in (0, 2], \beta \in [-1, 1], \gamma > 0, \mu \in \mathbb{R}$
Support	$x \in \dots$
P.d.f.	Calculated numerically
C.d.f.	Calculated numerically
Expectation	$\mathbb{E}[X] = k + \lambda$
Variance	$\text{Var}(X) = 2(k + 2\lambda)$
Median	Searched numerically
Mode	Searched numerically
C.f.	$\phi(t) = \dots$

8.1 Asymmetric Laplace distribution

8.2 Laplace distribution

9 Noncentral Chi-Squared distribution


10 Planck distribution

11 Stable distribution


11.1 Normal distribution

Relation to Stable distribution:


$$X \sim S_2(\cdot, \sigma^2/2, \mu)$$



Notation	$X \sim \chi_k'^2(\lambda)$
Parameters	$k > 0, \lambda > 0$
Support	$x \in \mathbb{R}^+$
P.d.f.	$f(x) = \dots$
C.d.f.	$F(x) = P_{\frac{k}{2}}(\dots)$
Expectation	$\mathbb{E}[X] = k + \lambda$
Variance	$\text{Var}(X) = 2(k + 2\lambda)$
Median	Searched numerically
Mode	Searched numerically
C.f.	$\phi(t) = \frac{\exp \frac{it\lambda}{1-2it}}{(1-2it)^{k/2}}$



Notation	$X \sim \text{Planck}(a, b)$
Parameters	$a, b > 0$
Support	$x \in \mathbb{R}^+$
P.d.f.	$f(x) = \frac{b^{a+1}}{\Gamma(a+1)\zeta(a+1)} \cdot \frac{x^a}{e^{bx}-1}$
C.d.f.	Calculated numerically
Expectation	$\mathbb{E}[X] = \frac{(a+1)\zeta(a+2)}{b\zeta(a+1)}$
Variance	$\text{Var}(X) = \frac{(a+1)(a+2)\zeta(a+3)}{b^2\zeta(a+1)} - (\mathbb{E}[X])^2$
Median	Searched numerically
Mode	$\frac{W_0(-ae^{-a})+a}{b}$, if $a > 1$, otherwise 0
C.f.	Calculated numerically



Notation	$X \sim S_{\alpha}(\beta, \gamma, \mu)$
Parameters	$\alpha \in (0, 2], \beta \in [-1, 1],$ $\gamma > 0, \mu \in \mathbb{R}$
Support	$x \in \mathbb{R}$, if $\beta \neq 1$, $x \in [\mu, \infty)$, if $\beta = 1, \alpha < 2$, $x \in (-\infty, \mu]$, if $\beta = -1, \alpha < 2$
P.d.f.	Calculated numerically
C.d.f.	Calculated numerically
Expectation	$\mathbb{E}[X] = \mu$ for $\alpha > 1$, otherwise undefined
Variance	$\text{Var}(X) = 2\gamma^2 1_{\{\alpha=2\}} + \infty 1_{\{\alpha<2\}}$
Median	μ for $\beta = 0$, otherwise searched numerically
Mode	μ , if $\beta = 0$ or $\alpha = 2$, $\mu + \frac{\beta\gamma}{3}$, if $ \beta = 1$ and $\alpha = \frac{1}{2}$, otherwise searched numerically
C.f.	$\phi(t) = \dots$

11.2 Cauchy distribution

Relation to Stable distribution:

$$X \sim S_1(0, \gamma, \mu)$$

11.3 Levy distribution


Relation to Stable distribution:

$$X \sim S_{\frac{1}{2}}(1, \gamma, \mu)$$

11.4 Holtsmark distribution

Relation to Stable distribution:

$$X \sim S_{\frac{3}{2}}(0, \gamma, \mu)$$



Notation	$X \sim \mathcal{N}(\mu, \sigma^2)$
Parameters	$\mu \in \mathbb{R}, \sigma^2 > 0$
Support	$x \in \mathbb{R}$
P.d.f.	$f(x) = \frac{1}{\sqrt{2\sigma^2\pi}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$
C.d.f.	$F(x) = \frac{1}{2} \operatorname{erfc}\left(\frac{\mu-x}{\sqrt{2\sigma^2}}\right)$
Expectation	$\mathbb{E}[X] = \mu$
Variance	$\operatorname{Var}(X) = \sigma^2$
Median	μ
Mode	μ
C.f.	$\phi(t) = e^{i\mu t - \frac{1}{2}\sigma^2 t^2}$

11.5 Landau distribution

Relation to Stable distribution:

$$X \sim S_1(1, \gamma, \mu)$$

12 Pareto distribution


Estimation of parameters.

Frequentist inference. Log-likelihood function is

$$\ln \mathcal{L}(\alpha, \sigma | X) = n \ln \alpha + n \alpha \ln \sigma - (\alpha + 1) \sum_{i=1}^n \ln X_i.$$

We assume that $\sigma \leq X_{(1)}$, otherwise sample X couldn't have been generated from such distribution. It is obvious, that $\ln \mathcal{L}(\alpha, \sigma | X)$ is an increasing function in terms of σ , therefore $\hat{\sigma} = X_{(1)}$ is an optimal estimator. Let's take derivative with respect to α :

$$\frac{\partial \ln \mathcal{L}(\alpha, \sigma | X)}{\partial \alpha} = \frac{n}{\alpha} + n \ln \sigma - \sum_{i=1}^n \ln X_i.$$



Notation	$X \sim \text{Cauchy}(\mu, \gamma)$
Parameters	$\mu \in \mathbb{R}, \gamma^2 > 0$
Support	$x \in \mathbb{R}$
P.d.f.	$f(x) = \frac{1}{\pi\gamma} \frac{1}{1 + \left(\frac{x-\mu}{\gamma}\right)^2}$
C.d.f.	$F(x) = \frac{1}{\pi} \text{atan}\left(\frac{x-\mu}{\gamma}\right) + \frac{1}{2}$
Expectation	Undefined
Variance	$\text{Var}(X) = \infty$
Median	μ
Mode	μ
C.f.	$\phi(t) = e^{i\mu t - \gamma t }$

From this we conclude that the maximum-likelihood estimator of shape is

$$\hat{\alpha} = \frac{1}{\frac{1}{n}(\sum_{i=1}^n \ln X_i) - \ln \hat{\sigma}}.$$

It is known that $\hat{\sigma} \sim \text{Pareto}(n\alpha, \sigma)$ and $\hat{\alpha} \sim \text{Inv-}\Gamma(n-1, n\alpha)$ and they are independent. Then

$$\mathbb{E}[\hat{\sigma}] = \frac{\sigma}{1 - \frac{1}{n\alpha}}$$

and

$$\mathbb{E}[\hat{\alpha}] = \frac{n\alpha}{n-2}.$$


Therefore, in order to get unbiased estimators we need to make the following transformations:

$$\tilde{\alpha} = \frac{n-2}{n} \hat{\alpha} \quad \text{and} \quad \tilde{\sigma} = \hat{\sigma} \left(1 - \frac{1}{(n-1)\hat{\alpha}}\right).$$

Note that if we estimate parameters separately, then $\hat{\alpha} \sim \text{Inv-}\Gamma(n, n\alpha)$ and transformations are different.

Bayesian inference. We now assume that σ is known and prior distribution of α is $\Gamma(\kappa, \beta)$:

$$h(\alpha) = \frac{\beta^\kappa}{\Gamma(\kappa)} \alpha^{\kappa-1} e^{-\beta\alpha}.$$



Notation	$X \sim \text{Pareto}(\alpha, \sigma)$
Parameters	$\alpha, \sigma > 0$
Support	$x \geq \sigma$
P.d.f.	$f(x) = \frac{\alpha \sigma^\alpha}{x^{\alpha+1}}$
C.d.f.	$F(x) = 1 - \left(\frac{\sigma}{x}\right)^\alpha$
Expectation	$\mathbb{E}[X] = \frac{\alpha \sigma}{\alpha - 1}$ for $\alpha > 1$, ∞ otherwise
Variance	$\text{Var}(X) = \frac{\sigma^2 \alpha}{(\alpha - 1)^2 (\alpha - 2)}$ for $\alpha > 2$, ∞ otherwise
Median	$\sigma 2^{1/\alpha}$
Mode	σ
C.f.	Calculated numerically

The density of posterior distribution is

$$f(\alpha|X) \propto \prod_{i=1}^n \frac{\sigma^\alpha}{X_i^{\alpha+1}} \cdot \alpha^{\kappa+n-1} e^{-\beta\alpha} \propto \alpha^{\kappa+n-1} e^{-(\beta + \sum_{i=1}^n \ln(X_i/\sigma))\alpha}.$$

Therefore, $\alpha|X \sim \Gamma(\kappa + n, \beta + \sum_{i=1}^n \ln(X_i/\sigma))$ and Bayesian estimator is

$$\mathbb{E}[\alpha|X] = \frac{\kappa + n}{\beta + \sum_{i=1}^n \ln(X_i/\sigma)}.$$

MAP estimator is

$$\alpha_{MAP} = \frac{\kappa + n - 1}{\beta + \sum_{i=1}^n \ln(X_i/\sigma)}.$$

Now let it be vice versa, α is known while σ is not. Then we say that a priori $\sigma \sim \text{Pareto}(\kappa, \theta)$:

$$h(\sigma) = \frac{\kappa \theta^\kappa}{\sigma^{\kappa+1}}.$$

Then posterior distribution is:

$$f(\sigma|X) \propto \prod_{i=1}^n \frac{1}{X_i^{\alpha+1}} \cdot \sigma^{\alpha n - \kappa - 1} \sim \text{Pareto}(\kappa - \alpha n, \theta).$$


This imposes the following additional constraint on the prior shape hyperparameter:
 $\kappa > \alpha n$. Bayesian estimator:

$$\mathbb{E}[\sigma|X] = \frac{(\kappa - \alpha n)\theta}{\kappa - \alpha n - 1}.$$

MAP estimator:

$$\sigma_{MAP} = \theta.$$

13 Weibull



Notation	$X \sim \text{Weibull}(\lambda, k)$
Parameters	$\lambda, k > 0$
Support	$x \in \mathbb{R}^+$
P.d.f.	$f(x) = \frac{k}{\lambda} \left(\frac{x}{\lambda}\right)^{k-1} \exp(-(x/\lambda)^k)$
C.d.f.	$F(x) = 1 - \exp(-(x/\lambda)^k)$
Expectation	$\mathbb{E}[X] = \lambda \Gamma(1 + 1/k)$
Variance	$\text{Var}(X) = \lambda^2 \Gamma(1 + 2/k) - (\mathbb{E}[X])^2$
Median	$\lambda (\ln 2)^{\frac{1}{k}}$
Mode	$\lambda \left(1 - \frac{1}{k}\right)^{\frac{1}{k}}$
C.f.	Calculated numerically

Estimation of scale

Frequentist inference. Log-likelihood function:

$$\ln \mathcal{L}(\lambda, k|X) = n(\ln k - \ln \lambda) + (k-1) \sum_{i=1}^n (\ln X_i - \ln \lambda) - \frac{1}{\lambda^k} \sum_{i=1}^n X_i^k.$$

The derivative with respect to scale:

$$\frac{\partial \ln \mathcal{L}(\lambda, k|X)}{\partial \lambda} = -\frac{nk}{\lambda} + \frac{k}{\lambda^{k+1}} \sum_{i=1}^n X_i^k = 0.$$

Therefore, maximum-likelihood estimation for λ is

$$\hat{\lambda} = \left(\sum_{i=1}^n X_i^k \right)^{\frac{1}{k}}.$$

Bayesian inference. Assume k is known. Instead of estimating λ we give an estimation for λ^k . Let's say that prior distribution of λ^k is $\text{Inv-}\Gamma(\alpha, \beta)$:

$$h(\lambda^k) = \frac{\beta^\alpha}{\Gamma(\alpha)} \lambda^{-k(\alpha+1)} e^{-\beta/\lambda^k}.$$

Posterior distribution then:

$$f(\lambda^k|X) \propto \lambda^{-k(\alpha+1+n)} e^{-\frac{1}{\lambda^k}(\beta + \sum_{i=1}^n X_i^k)} \sim \text{Inv-}\Gamma(\alpha + n, \beta + \sum_{i=1}^n X_i^k).$$

Bayesian estimator:

$$\mathbb{E}[\lambda^k|X] = \frac{\beta + \sum_{i=1}^n X_i^k}{\alpha + n - 1},$$

MAP estimator:

$$\lambda_{MAP}^k = \frac{\beta + \sum_{i=1}^n X_i^k}{\alpha + n + 1}.$$

Part III

Discrete univariate distributions

14 Beta-binomial distribution

15 Binomial distribution

Notation	$X \sim \text{Bin}(n, p)$
Parameters	$n \in \mathbb{N}, p \in [0, 1]$
Support	$k \in \{0, \dots, n\}$
P.m.f.	$\mathbb{P}(X = k) = \binom{n}{k} p^k (1 - p)^{n-k}$
C.d.f.	$\mathbb{P}(X \leq k) = I_{1-p}(n - k, 1 + k)$
Expectation	$\mathbb{E}[X] = np$
Variance	$\text{Var}(X) = np(1 - p)$
Median	$[np]$
Mode	$[(n + 1)p]$
C.f.	$\phi(t) = (1 - p + pe^{it})^n$

15.1 Bernoulli

Notation:

$$X \sim \text{Bernoulli}(p).$$

Relation to Binomial distribution:

$$X \sim \text{Bin}(1, p).$$

16 Poisson distribution

Notation	$X \sim \text{Po}(\lambda)$
Parameters	$\lambda > 0$
Support	$k \in \mathbb{N}_0$
P.m.f.	$\mathbb{P}(X = k) = \frac{\lambda^k e^{-\lambda}}{k!}$
C.d.f.	$\mathbb{P}(X \leq k) = Q(k + 1, \lambda)$
Expectation	$\mathbb{E}[X] = \lambda$
Variance	$\text{Var}(X) = \lambda$
Median	$\sim \max\left([\lambda + \frac{1}{3} - \frac{0.02}{\lambda}], 0\right)$
Mode	$[\lambda]$
C.f.	$\phi(t) = \exp\{\lambda(e^{it} - 1)\}$

Generator (let $\delta = \mu \in \mathbb{Z}$). (There is a mistake in Lemma 3.8 in first inequality). Recall that

$$q(X) = X \ln(\lambda) - \ln\left(\frac{(\mu + X)!}{\mu!}\right).$$

We denote acceptance probability $\mathbb{P}(W \leq q(X))$ by p .

- $k = \mu$. Probability to be in this setting is $1/c$.

$$\mathbb{P}(X = 0 | W \leq q(X)) = \frac{\mathbb{P}(X = 0, W \leq q(X))}{\mathbb{P}(W \leq q(X))} = \frac{1}{pc}.$$

On the other hand it should be equal to:

$$\frac{1}{pc} = \frac{\lambda^\mu e^{-\lambda}}{\mu!}.$$

- $k = \mu + 1$.

$$\begin{aligned} \mathbb{P}(X = 1 | W \leq q(X)) &= \frac{\mathbb{P}(X = 1, W \leq q(X))}{\mathbb{P}(W \leq q(X))} = \frac{\lambda}{p(\mu + 1)c} \\ &= \frac{\lambda^{\mu+1} e^{-\lambda}}{(\mu + 1)!}. \end{aligned}$$

- $k < \mu$. Here was mistake in the book. We adjust the probabilities. Probability to be in this setting is $\sqrt{\pi\mu/2e}/c$.

$$\begin{aligned}
\mathbb{P}(W \leq q(X), X = k - \mu | U \leq c_1) &= \mathbb{P}\left(-\frac{N^2}{2} + \frac{1}{2} - E < q(\lfloor -|N|\sqrt{\mu} \rfloor), \lceil |N|\sqrt{\mu} \rceil = \mu - k\right) \\
&= \mathbb{P}\left(-\frac{N^2}{2} + \frac{1}{2} - E < \lfloor -|N|\sqrt{\mu} \rfloor \ln(\lambda) - \ln\left(\frac{(\mu + \lfloor -|N|\sqrt{\mu} \rfloor)!}{\mu!}\right), \right. \\
&\quad \left. \frac{\mu - k - 1}{\sqrt{\mu}} \leq |N| < \frac{\mu - k}{\sqrt{\mu}}\right) \\
&= \mathbb{P}\left(U < \exp\left\{\frac{N^2}{2} - \frac{1}{2} + \lfloor -|N|\sqrt{\mu} \rfloor \ln(\lambda) - \ln\left(\frac{(\mu + \lfloor -|N|\sqrt{\mu} \rfloor)!}{\mu!}\right)\right\}, \right. \\
&\quad \left. \frac{\mu - k - 1}{\sqrt{\mu}} \leq |N| < \frac{\mu - k}{\sqrt{\mu}}\right) \\
&= \sqrt{\frac{2}{e\pi}} \int_{\frac{\mu-k-1}{\sqrt{\mu}}}^{\frac{\mu-k}{\sqrt{\mu}}} \exp\left\{\lfloor -|n|\sqrt{\mu} \rfloor \ln(\lambda) - \ln\left(\frac{(\mu + \lfloor -|n|\sqrt{\mu} \rfloor)!}{\mu!}\right)\right\} dn \\
&= \sqrt{\frac{2}{e\pi\mu}} \int_{\mu-k-1}^{\mu-k} \exp\left\{\lfloor -z \rfloor \ln(\lambda) - \ln\left(\frac{(\mu + \lfloor -z \rfloor)!}{\mu!}\right)\right\} dz \\
&= \sqrt{\frac{2}{e\pi\mu}} \exp\left\{(k - \mu) \ln(\lambda) - \ln\left(\frac{k!}{\mu!}\right)\right\} \\
&= \sqrt{\frac{2}{e\pi\mu}} \lambda^{k-\mu} \frac{\mu!}{k!}
\end{aligned}$$

Hence,

$$\begin{aligned}
\mathbb{P}(X = k - \mu | W \leq q(X)) &= \frac{\mathbb{P}(W \leq q(X), X = k - \mu)}{\mathbb{P}(W \leq q(X))} \\
&= \sqrt{\frac{2}{\pi\mu e}} \lambda^{k-\mu} \frac{\mu!}{k!} \cdot \sqrt{\pi\mu e/2} \frac{\lambda^\mu e^{-\lambda}}{\mu!} \\
&= \frac{\lambda^k e^{-\lambda}}{k!}
\end{aligned}$$

- $k \in [\mu + 2, 2\mu]$. Probability to be in this setting is $\sqrt{\frac{3\pi\mu}{4}} e^{\frac{1}{3\mu}}/c$. We also have

$$W = \frac{-Y^2 + 2Y}{3\mu} - E = \frac{1}{3\mu} - \frac{N^2}{2} - E.$$

Then

$$\begin{aligned}
\mathbb{P}(W \leq q(X)|X = k - \mu|U \in \dots) &= \mathbb{P}\left(\frac{1}{3\mu} - \frac{N^2}{2} - E < q(\lceil 1 + |N|\sqrt{3\mu/2} \rceil), \lceil 1 + |N|\sqrt{3\mu/2} \rceil = k - \mu\right) \\
&= \mathbb{P}\left(U < \exp\left\{-\frac{1}{3\mu} + \frac{N^2}{2} + q(\lceil 1 + |N|\sqrt{3\mu/2} \rceil)\right\}, \right. \\
&\quad \left. \frac{k - \mu - 2}{\sqrt{3\mu/2}} < |N| \leq \frac{k - \mu - 1}{\sqrt{3\mu/2}}\right) \\
&= \sqrt{\frac{2}{\pi}} e^{-\frac{1}{3\mu}} \int_{\frac{k-\mu-2}{\sqrt{3\mu/2}}}^{\frac{k-\mu-1}{\sqrt{3\mu/2}}} \exp\left\{q(\lceil 1 + |n|\sqrt{3\mu/2} \rceil)\right\} dn \\
&= \sqrt{\frac{4}{3\pi\mu}} e^{-\frac{1}{3\mu}} \int_{k-\mu-1}^{k-\mu} \exp\left\{\lceil z \rceil \ln(\lambda) - \ln\left(\frac{(\mu + \lceil z \rceil)!}{\mu!}\right)\right\} dz \\
&= \sqrt{\frac{4}{3\pi\mu}} e^{-\frac{1}{3\mu}} \mu! \frac{\lambda^{k-\mu}}{k!}.
\end{aligned}$$

- $k > 2\mu$. Probability to be in this setting is $6e^{-\frac{2+\mu}{6}}/c$.

$$\begin{aligned}
\mathbb{P}(W \leq q(X)|X = k - \mu|U \in \dots) &= \mathbb{P}\left(-\frac{2+\mu}{6} - V - E < q(\lceil \mu + 6V \rceil), \lceil \mu + 6V \rceil = k - \mu\right) \\
&= \mathbb{P}\left(-\frac{2+\mu}{6} - V + \ln(U) < \lceil \mu + 6V \rceil \ln(\lambda) - \ln\left(\frac{(\mu + \lceil \mu + 6V \rceil)!}{\mu!}\right), \right. \\
&\quad \left. \lceil \mu + 6V \rceil = k - \mu\right) \\
&= \mathbb{P}\left(U < \exp\left\{\frac{2+\mu}{6} + V + \lceil \mu + 6V \rceil \ln(\lambda) - \ln\left(\frac{(\mu + \lceil \mu + 6V \rceil)!}{\mu!}\right)\right\}, \right. \\
&\quad \left. \frac{k - 2\mu - 1}{6} < V \leq \frac{k - 2\mu}{6}\right) \\
&= \int_{\frac{k-2\mu-1}{6}}^{\frac{k-2\mu}{6}} \exp\left\{\frac{2+\mu}{6} + \lceil \mu + 6v \rceil \ln(\lambda) - \ln\left(\frac{(\mu + \lceil \mu + 6v \rceil)!}{\mu!}\right)\right\} dv \\
&= \frac{e^{\frac{2+\lambda}{6}}}{6} \int_{k-\mu-1}^{k-\mu} \exp\left\{\lceil z \rceil \ln(\lambda) - \ln\left(\frac{(\mu + \lceil z \rceil)!}{\mu!}\right)\right\} dz \\
&= \frac{e^{\frac{2+\lambda}{6}}}{6} \exp\left\{(k - \mu) \ln(\lambda) - \ln\left(\frac{k!}{\mu!}\right)\right\} \\
&= \frac{e^{\frac{2+\lambda}{6}}}{6} \lambda^{k-\mu} \frac{\mu!}{k!}
\end{aligned}$$

Hence,

$$\begin{aligned}
\mathbb{P}(X = k - \mu | W \leq q(X)) &= \frac{\mathbb{P}(W \leq q(X), X = k - \mu)}{\mathbb{P}(W \leq q(X))} \\
&= \frac{e^{\frac{2+\lambda}{6}}}{6} \lambda^{k-\mu} \frac{\mu!}{k!} \cdot \frac{6e^{-\frac{2+\mu}{6}}}{pc} \\
&= \frac{\lambda^k e^{-\lambda}}{k!}
\end{aligned}$$

Part IV

Bivariate distributions

- 17 Bivariate Normal distribution
- 18 Normal-Inverse-Gamma distribution
- 19 Trinomial distribution

Part V

Circular distributions

20 von Mises distribution

21 Wrapped Exponential distribution

Part VI

Singular distributions

22 Cantor distribution

Implemented distributions (under titles special cases are listed):

- Continuous distributions
 - Generalised extreme value ???????
 - Gumbel
 - Frechet
 - Weibull
 - Geometric-Stable distribution
 - (Asymmetric) Laplace distribution
 - Inverse-Gamma
 - Inverse-Gaussian
 - Irwin-Hall
 - Kolmogorov-Smirnov

Logistic Log-Normal Nakagami Chi Maxwell-Boltzmann Rayleigh Noncentral Chi-Squared Pareto Planck Raised-cosine Raab-Green Sech Stable Cauchy Holtsmark Landau Levy Normal t-distribution Triangular Von-Mises Wigner-Semicircle Discrete: Beta-binomial Binomial Bernoulli Categorical Hypergeometric Logarithmic Negative-binomial (Polya) Pascal Geometric Negative-hypergeometric Poisson Rademacher Skellam Uniform discrete Yule Zeta Zipf Singular: Cantor Bivariate: Bivariate Normal Normal-inverse-Gamma