

# RandLib documentation

Aleksandr Samarin

October 14, 2018

## Contents

<b>I</b>	<b>General information</b>	<b>4</b>
1	Calculation of sample moments	4
<b>II</b>	<b>Continuous univariate distributions</b>	<b>5</b>
<b>2</b>	<b>Beta distribution</b>	<b>5</b>
2.1	Arcsine distribution . . . . .	7
2.2	Balding-Nichols distribution . . . . .	8
2.3	Uniform distribution . . . . .	8
<b>3</b>	<b>Beta-prime distribution</b>	<b>10</b>
<b>4</b>	<b>Exponentially-modified Gaussian distribution</b>	<b>12</b>
<b>5</b>	<b>F-distribution</b>	<b>13</b>
<b>6</b>	<b>Gamma distribution</b>	<b>14</b>
6.1	Chi-squared distribution . . . . .	15
6.2	Erlang distribution . . . . .	15
6.3	Exponential distribution . . . . .	16
<b>7</b>	<b>Geometric Stable distribution</b>	<b>17</b>
7.1	Asymmetric Laplace distribution . . . . .	17
7.2	Laplace distribution . . . . .	17
<b>8</b>	<b>Kolmogorov-Smirnov distribution</b>	<b>18</b>
<b>9</b>	<b>Logistic distribution</b>	<b>19</b>

<b>10 Log-normal distribution</b>	<b>20</b>
<b>11 Marchenko-Pastur distribution</b>	<b>21</b>
<b>12 Nakagami distribution</b>	<b>22</b>
12.1 Chi distribution . . . . .	22
12.2 Maxwell-Boltzmann distribution . . . . .	23
12.3 Rayleigh distribution . . . . .	23
<b>13 Noncentral Chi-Squared distribution</b>	<b>24</b>
<b>14 Pareto distribution</b>	<b>25</b>
<b>15 Planck distribution</b>	<b>28</b>
<b>16 Stable distribution</b>	<b>29</b>
16.1 Cauchy distribution . . . . .	29
16.2 Levy distribution . . . . .	29
16.3 Normal distribution . . . . .	30
16.4 Holtsmark distribution . . . . .	31
16.5 Landau distribution . . . . .	31
<b>17 Weibull</b>	<b>32</b>
 <b>III Discrete univariate distributions</b>	 <b>34</b>
<b>18 Beta-binomial distribution</b>	<b>34</b>
<b>19 Binomial distribution</b>	<b>35</b>
19.1 Bernoulli . . . . .	36
<b>20 Hypergeometric distribution</b>	<b>37</b>
<b>21 Negative-Binomial (Polya) distribution</b>	<b>38</b>
21.1 Geometric distribution . . . . .	38
21.2 Pascal distribution . . . . .	38
<b>22 Poisson distribution</b>	<b>39</b>
<b>23 Skellam distribution</b>	<b>41</b>
<b>24 Yule distribution</b>	<b>42</b>
<b>25 Zeta distribution</b>	<b>43</b>
<b>26 Zipf distribution</b>	<b>44</b>

<b>IV</b>	<b>Bivariate distributions</b>	<b>45</b>
<b>27</b>	<b>Bivariate Normal distribution</b>	<b>45</b>
<b>28</b>	<b>Normal-Inverse-Gamma distribution</b>	<b>45</b>
<b>29</b>	<b>Trinomial distribution</b>	<b>45</b>
<b>V</b>	<b>Circular distributions</b>	<b>46</b>
<b>30</b>	<b>von Mises distribution</b>	<b>46</b>
<b>31</b>	<b>Wrapped Exponential distribution</b>	<b>46</b>
<b>VI</b>	<b>Singular distributions</b>	<b>47</b>
<b>32</b>	<b>Cantor distribution</b>	<b>47</b>

## Part I

# General information

## 1 Calculation of sample moments

We use extension of Welford's method from Knuth. For every  $n$ -th element  $x$  we have


$$\begin{aligned}\delta &= x - m_1, \\ m'_1 &= m_1 + \frac{\delta}{n}, \\ m'_2 &= m_2 + \delta^2 \frac{n-1}{n}, \\ m'_3 &= m_3 + \delta^3 \frac{(n-1)(n-2)}{n^2} - 3\delta \frac{m_2}{n}, \\ m'_4 &= m_4 + \delta^4 \frac{(n-1)(n^2-3n+3)}{n^3} + 6\delta^2 \frac{m_2}{n^2} - 4\delta \frac{m_3}{n}.\end{aligned}$$

Then  $m'_1, \frac{m_2}{n}, \text{Skew}(X) = \frac{\sqrt{n}m'_3}{m_2^{3/2}}$  and  $\text{Kurt}(X) = \frac{nm'_4}{m_2^2}$  (we return excess kurtosis).

## Part II

# Continuous univariate distributions

## 2 Beta distribution



Notation	$X \sim \mathcal{B}(\alpha, \beta, a, b)$ $X \sim \mathcal{B}(\alpha, \beta)$ with $a = 0, b = 1$
Parameters	$\alpha, \beta > 0, a, b \in \mathbb{R}$
Domain	$x \in [a, b]$
$f(x)$	$\frac{y^{\alpha-1}(1-y)^{\beta-1}}{(b-a)B(\alpha, \beta)}$ with $y = \frac{x-a}{b-a}$
$F(x)$	$I_y(\alpha, \beta)$ for $y = \frac{x-a}{b-a}$
$\mathbb{E}[X]$	$a + (b-a)\frac{\alpha}{\alpha+\beta}$
$\text{Var}(X)$	$(b-a)^2 \frac{\alpha\beta}{(\alpha+\beta)^2(\alpha+\beta+1)}$
Median	Searched numerically
Mode	$a + (b-a)\frac{\alpha-1}{\alpha+\beta-2}$ for $\alpha, \beta > 1$ .
$\phi(t)$	Calculated numerically

**Search of the median.** In general, the value of median is unknown and searched numerically with initial value:

$$m \approx a + (b-a) \frac{\alpha - \frac{1}{3}}{\alpha + \beta - \frac{2}{3}}$$

for  $\alpha, \beta \geq 1$ . However, there are analytical solutions for some particular values:

- $m = \frac{a+b}{2}$ , for  $\alpha = \beta$ ,
- $m = a + (b-a)(1 - 2^{-\frac{1}{\beta}})$ , for  $\alpha = 1$ ,
- $m = a + (b-a)2^{-\frac{1}{\alpha}}$ , for  $\beta = 1$ .

**Calculation of characteristic function.** For  $\alpha, \beta \geq 1$  we use numerical integration by definition

$$\phi(t) = \int_a^b \cos(tx) f(x) dx + i \int_a^b \sin(tx) f(x) dx.$$

For shape parameters  $< 1$ ,  $f(x)$  has singularity points at 0 or 1 or both of them, and numerical integration is impossible. Then we use the following technique: firstly, we can show that

$$\phi(t|a, b) = \mathbb{E}[e^{it(a+(b-a)X)}] = e^{ita} \phi(z|0, 1)$$

with  $z = (b - a)t$ . Hence, w.l.o.g. we can consider standard case  $a = 0, b = 1$ . Then

$$\begin{aligned} \Re(\phi(z)) &= \frac{1}{B(\alpha, \beta)} \int_0^1 \cos(zx) x^{\alpha-1} (1-x)^{\beta-1} dx \\ &= \frac{1}{B(\alpha, \beta)} \int_0^1 (\cos(zx) - 1) x^{\alpha-1} (1-x)^{\beta-1} dx + 1 \\ &= \frac{1}{B(\alpha, \beta)} \int_0^1 \frac{(\cos(zx) - 1) x^{\alpha-1} - (\cos(z) - 1)}{(1-x)^{1-\beta}} dx + 1 + \frac{\cos(z) - 1}{bB(\alpha, \beta)}. \end{aligned}$$

The integrand now doesn't have any singularities, neither for  $\alpha < 1$ , nor for  $\beta < 1$ . Analogously we transform the imaginary part:

$$\begin{aligned} \Im(\phi(z)) &= \frac{1}{B(\alpha, \beta)} \int_0^1 \sin(zx) x^{\alpha-1} (1-x)^{\beta-1} dx \\ &= \frac{1}{B(\alpha, \beta)} \int_0^1 \frac{\sin(zx) x^{\alpha-1} - \sin(z)}{(1-x)^{1-\beta}} dx + \frac{\sin(z)}{bB(\alpha, \beta)}. \end{aligned}$$

**Estimation of shapes with known support.** Assume that  $a = 0, b = 1$  and we have a sample  $X = (X_1, \dots, X_n)$ . Then a log-likelihood function is

$$\begin{aligned} \ln \mathcal{L}(\alpha, \beta | X) &= \sum_{i=1}^n \ln f(X_i; \alpha, \beta) \\ &= (\alpha - 1) \sum_{i=1}^n \ln X_i + (\beta - 1) \sum_{i=1}^n \ln(1 - X_i) - n \ln B(\alpha, \beta). \end{aligned} \tag{1}$$

Differentiating with respect to the shapes, we obtain

$$\begin{aligned} \frac{\partial \ln \mathcal{L}(\alpha, \beta | X)}{\partial \alpha} &= \sum_{i=1}^n \ln X_i + n(\psi(\alpha + \beta) - \psi(\alpha)), \\ \frac{\partial \ln \mathcal{L}(\alpha, \beta | X)}{\partial \beta} &= \sum_{i=1}^n \ln(1 - X_i) + n(\psi(\alpha + \beta) - \psi(\beta)). \end{aligned}$$

Differentiating again we get the Hessian matrix:

$$H(\ln \mathcal{L}(\alpha, \beta|X)) = n \cdot \begin{pmatrix} \psi_1(\alpha + \beta) - \psi_1(\alpha) & \psi_1(\alpha + \beta) \\ \psi_1(\alpha + \beta) & \psi_1(\alpha + \beta) - \psi_1(\beta) \end{pmatrix}.$$

Then we can find the estimators numerically, using Newton's procedure. The initial values of estimators are found via method of moments:

$$\hat{\alpha}_0 = \bar{X}_n \left( \frac{\bar{X}_n(1 - \bar{X}_n)}{\hat{s}_n^2} - 1 \right),$$

$$\hat{\beta}_0 = (1 - \bar{X}_n) \left( \frac{\bar{X}_n(1 - \bar{X}_n)}{\hat{s}_n^2} - 1 \right).$$

These values are applicable only if  $\hat{s}_n^2 < \bar{X}_n(1 - \bar{X}_n)$ . If this condition is not satisfied, we set  $\hat{\alpha}_0 = \hat{\beta}_0 = 0.001$ .

In the general case, when  $a \neq 0$  or  $b \neq 1$ , we use the following transformation:

$$Y_i = \frac{X_i - a}{b - a}$$

and estimate parameters, using sample  $Y$ .

## 2.1 Arcsine distribution

Notation:

$$X \sim \text{Arcsine}(\alpha).$$

Relation to Beta distribution:

$$X \sim \mathcal{B}(1 - \alpha, \alpha, a, b).$$

**Estimation of shape.** For Arcsine distribution log-likelihood function (1) turns into

$$\ln \mathcal{L}(\alpha|X) = -\alpha \sum_{i=1}^n \ln X_i + (\alpha - 1) \sum_{i=1}^n \ln(1 - X_i) - n \ln B(1 - \alpha, \alpha).$$

Taking the derivative with respect to  $\alpha$  we get

$$\frac{\partial \ln \mathcal{L}(\alpha|X)}{\partial \alpha} = \sum_{i=1}^n \ln \frac{1 - X_i}{X_i} + n\pi \cot(\pi\alpha).$$

Therefore, maximum-likelihood function is

$$\hat{\alpha} = -\frac{1}{\pi} \operatorname{atan} \left( \frac{n\pi}{\sum_{i=1}^n \ln \frac{1-X_i}{X_i}} \right).$$

If  $\hat{\alpha}$  is negative, we add 1, because  $\frac{\operatorname{atan}}{\pi} \in (-0.5, 0.5)$ , while  $\alpha \in (0, 1)$ .

## 2.2 Balding-Nichols distribution

Notation:


$$X \sim \text{Balding-Nichols}(p, F)$$

with  $p, F \in (0, 1)$ . Relation to Beta distribution:

$$X \sim \mathcal{B}(pF', (1-p)F')$$

with  $F' = (1-F)/F$ .

## 2.3 Uniform distribution



Notation	$X \sim \mathcal{U}(a, b)$
Parameters	$a, b \in \mathbb{R}$
Domain	$x \in [a, b]$
$f(x)$	$\frac{1}{b-a}$
$F(x)$	$\frac{x-a}{b-a}$
$\mathbb{E}[X]$	$\frac{a+b}{2}$
$\text{Var}(X)$	$\frac{(b-a)^2}{12}$
Median	$\frac{a+b}{2}$
Mode	doesn't exist
$\phi(t)$	$\frac{e^{itb} - e^{ita}}{it(b-a)}$

Relation to Beta distribution:

$$X \sim \mathcal{B}(1, 1, a, b).$$

**Estimation of support.**

**Frequentist inference.** Likelihood function is

$$\mathcal{L}(a, b|X) = \frac{1}{(b-a)^n} \mathbf{1}_{\{X_i \in [a, b] \ \forall i=1, \dots, n\}}.$$



Therefore,  $\mathcal{L}(a, b|X)$  is the largest for  $\hat{b} = X_{(n)}$  and  $\hat{a} = X_{(1)}$ . However, using the fact that  $X_{(k)} \sim B(k, n+1-k, a, b)$ , these are biased estimators:

$$\mathbb{E}[X_{(1)}] = \frac{an+b}{n+1} \quad \text{and} \quad \mathbb{E}[X_{(n)}] = \frac{a+bn}{n+1}.$$

To get unbiased estimators we make the transformations:

$$\tilde{a} = \frac{nX_{(1)} - X_{(n)}}{n-1} \quad \text{and} \quad \tilde{b} = \frac{nX_{(n)} - X_{(1)}}{n-1}.$$

Then we get

$$\mathbb{E}[\tilde{a}] = \frac{n\mathbb{E}[X_{(1)}] - \mathbb{E}[X_{(n)}]}{n-1} = \frac{n(an+b) - (a+bn)}{n^2-1} = a.$$

Analogously,  $\mathbb{E}[\tilde{b}] = b$ .

**Bayesian inference.** Let us say, we try to estimate  $\theta = b - a$  with known  $a$ . We set the prior distribution  $\theta \sim \text{Pareto}(\alpha, \sigma)$ :

$$h(\theta|\alpha, \sigma) = \frac{\alpha\sigma^\alpha}{\theta^{\alpha+1}} \mathbf{1}_{\{\theta \geq \sigma\}}.$$

The density of posterior distribution is

$$f(\theta|X) \propto \frac{\alpha\sigma^\alpha}{\theta^{\alpha+n+1}} \mathbf{1}_{\{\theta \geq \max(\sigma, X_{(n)} - a)\}} \sim \text{Pareto}(\alpha + n, \max(\sigma, X_{(n)} - a)).$$


Hence, Bayesian estimator is

$$\mathbb{E}[\theta|X] = \frac{\alpha + n}{\alpha + n - 1} \max(\sigma, X_{(n)} - a)$$

and MAP estimator is

$$\theta_{MAP} = \max(\sigma, X_{(n)} - a).$$

### 3 Beta-prime distribution



Notation	$X \sim \mathcal{B}'(\alpha, \beta)$
Parameters	$\alpha, \beta > 0$
Domain	$x \in \mathbb{R}^+$
$f(x)$	$\frac{x^{\alpha-1}(1+x)^{-\alpha-\beta}}{B(\alpha, \beta)}$
$F(x)$	$I_{\frac{x}{1+x}}(\alpha, \beta)$
$\mathbb{E}[X]$	$\frac{\alpha}{\beta-1} \mathbf{1}_{\{\beta>1\}} + \infty \mathbf{1}_{\{\beta \leq 1\}}$
$\text{Var}(X)$	$\frac{\alpha(\alpha+\beta-1)}{(\beta-2)(\beta-1)^2}$ , if $\beta > 1$
Median	Searched numerically
Mode	$\max\left(\frac{\alpha-1}{\beta+1}, 0\right)$ .
$\phi(t)$	Calculated numerically

Relation to other distributions:

$$\frac{X}{1+X} \sim \mathcal{B}(\alpha, \beta),$$

$$\frac{\beta}{\alpha} X \sim F(2\alpha, 2\beta).$$

**Search of the median.** For  $\alpha = \beta$  we have  $m = 1$ . Otherwise, we use the relation  $m = \frac{m'}{1-m'}$ , where  $m'$  is the median of beta-distribution  $\mathcal{B}(\alpha, \beta)$ .

**Calculation of characteristic function.** For  $\alpha \geq 1$  one can use numerical integration from section For  $\alpha < 1$  we have  $\lim_{x \rightarrow 0} f(x) \rightarrow \infty$  and  $\int_0^\infty \cos(tx)f(x)dx$  is impossible to compute directly. Then we split the integral:

$$\int_0^\infty \cos(tx)f(x)dx = \int_0^\infty (\cos(tx) - 1)f(x)dx + 1.$$


The limit of the integrand for  $x \rightarrow 0$  is 0 now, regardless of the value of the shape  $\alpha$ .

**Estimation of shapes.** Using relationship with Beta distribution we transform the sample:

$$Y_i = \frac{X_i}{1 + X_i}, \quad 1 \leq i \leq N,$$

and run estimation for beta-distributed  $Y$ .

## 4 Exponentially-modified Gaussian distribution



Notation	$X \sim \text{EMG}(\mu, \sigma, \lambda)$
Parameters	$\mu \in \mathbb{R}, \sigma > 0, \lambda > 0$
Domain	$x \in \mathbb{R}$
$f(x)$	$\frac{\lambda}{2} e^{\frac{\lambda}{2}(2\mu + \lambda\sigma^2 - 2x)} \operatorname{erfc}\left(\frac{\mu + \lambda\sigma^2 - x}{\sqrt{2}\sigma^2}\right)$
$F(x)$	$\Phi(u, 0, v) - e^{-u + \frac{v^2}{2} + \log \Phi(u, v^2, v)}$ , where $\Phi(x, \mu, \sigma)$ is Gaussian CDF, $u = \lambda(x - \mu)$ , $v = \lambda\sigma$ .
$\mathbb{E}[X]$	$\mu + 1/\lambda$
$\operatorname{Var}(X)$	$\sigma^2 + 1/\lambda^2$
Median	Searched numerically
Mode	Searched numerically
$\phi(t)$	$(1 - \frac{it}{\lambda})^{-1} \exp(i\mu t - \frac{1}{2}\sigma^2 t^2)$

## 5 F-distribution


Notation	$X \sim F(d_1, d_2)$
Parameters	$d_1, d_2 > 0$
Domain	$x \in \mathbb{R}^+$
$f(x)$	$\frac{\sqrt{\frac{(d_1 x)^{d_1} d_2^{d_2}}{(d_1 x + d_2)^{d_1 + d_2}}}}{x B\left(\frac{d_1}{2}, \frac{d_2}{2}\right)}$
$F(x)$	$I_{\frac{d_1 x}{d_1 x + d_2}}\left(\frac{d_1}{2}, \frac{d_2}{2}\right)$
$\mathbb{E}[X]$	$\frac{d_2}{d_2 - 2}$ for $d_2 > 2$
$\text{Var}(X)$	$\frac{2d_2^2(d_1 + d_2 - 2)}{d_1(d_2 - 2)^2(d_2 - 4)}$ for $d_2 > 4$
Median	Searched numerically
Mode	$\max\left(\frac{d_2(d_1 - 2)}{d_1(d_1 + 2)}, 0\right)$
$\phi(t)$	Calculated numerically

Relation to other distributions:

$$\frac{d_1 X}{d_2 + d_1 X} \sim \mathcal{B}\left(\frac{d_1}{2}, \frac{d_2}{2}\right),$$

$$\frac{d_1}{d_2} X \sim \mathcal{B}'\left(\frac{d_1}{2}, \frac{d_2}{2}\right).$$

## 6 Gamma distribution



Notation	$X \sim \Gamma(\alpha, \beta)$
Parameters	$\alpha > 0, \beta > 0$
Domain	$x \in \mathbb{R}^+$
$f(x)$	$\frac{\beta^\alpha}{\Gamma(\alpha)} x^{\alpha-1} e^{-\beta x}$
$F(x)$	$P(\alpha, \beta x)$
$\mathbb{E}[X]$	$\frac{\alpha}{\beta}$
$\text{Var}(X)$	$\frac{\alpha}{\beta^2}$
Median	Searched numerically
Mode	$\max\left(\frac{\alpha-1}{\beta}, 0\right)$
$\phi(t)$	$\left(1 - \frac{it}{\beta}\right)^{-\alpha}$

**Estimation of parameters.**

**Frequentist inference.** Log-likelihood function:

$$\ln \mathcal{L}(\alpha, \beta | X) = n\alpha \ln \beta - n \ln \Gamma(\alpha) + (\alpha - 1) \sum_{i=1}^n \ln X_i - \beta \sum_{i=1}^n X_i.$$

Derivatives:

$$\frac{\partial \ln \mathcal{L}(\alpha, \beta | X)}{\partial \alpha} = n \ln \beta - n\psi(\alpha) + \sum_{i=1}^n \ln X_i,$$

$$\frac{\partial \ln \mathcal{L}(\alpha, \beta | X)}{\partial \beta} = \frac{n\alpha}{\beta} - \sum_{i=1}^n X_i.$$

While the solution for the second equation is analytic:

$$\hat{\beta} = \frac{\alpha}{\bar{X}_n},$$

the first equation is solved numerically, using second derivative:

$$\frac{\partial^2 \ln \mathcal{L}(\alpha, \beta | X)}{\partial \alpha^2} = -n\psi_1(\alpha),$$

or if  $\beta$  is unknown:

$$\frac{\partial^2 \ln \mathcal{L}(\alpha, \beta | X)}{\partial \alpha^2} = -n\psi_1(\alpha) + \frac{n}{\alpha},$$

Moreover, the maximum-likelihood estimation of rate  $\beta$  is biased. Unbiased estimator would be

$$\tilde{\beta} = \frac{\alpha}{\bar{X}_n} \left(1 - \frac{1}{n}\right).$$

**Bayesian inference.** We suppose that prior distribution of rate  $\beta$  is  $\Gamma(\kappa, \gamma)$ :

$$h(\beta) = \frac{\gamma^\kappa}{\Gamma(\kappa)} \beta^{\kappa-1} e^{-\gamma\beta}.$$

Then

$$f(\beta | X) \propto \beta^{\alpha n} e^{-\beta \sum_{i=1}^n X_i} \cdot \beta^{\kappa-1} e^{-\gamma\beta} \sim \Gamma\left(\alpha n + \kappa, \gamma + \sum_{i=1}^n X_i\right).$$

Therefore, Bayesian estimator is

$$\mathbb{E}[\beta | X] = \frac{\alpha n + \kappa}{\gamma + \sum_{i=1}^n X_i},$$

and MAP estimator is

$$\beta_{MAP} = \frac{\alpha n + \kappa - 1}{\gamma + \sum_{i=1}^n X_i}.$$

## 6.1 Chi-squared distribution

Notation:

$$X \sim \chi_k^2.$$

Relation to Gamma distribution:


$$X \sim \Gamma\left(\frac{k}{2}, \frac{1}{2}\right).$$

## 6.2 Erlang distribution

Notation:

$$X \sim \text{Erlang}(k, \beta).$$

The only difference between Gamma and Erlang distributions is that a second one takes an integer shape parameter  $k$ .



Notation	$X \sim \text{Exp}(\lambda)$
Parameters	$\lambda > 0$
Domain	$x \in \mathbb{R}^+$
$f(x)$	$\lambda e^{-\lambda x}$
$F(x)$	$1 - e^{-\lambda x}$
$\mathbb{E}[X]$	$\frac{1}{\lambda}$
$\text{Var}(X)$	$\frac{1}{\lambda^2}$
Median	$\frac{\ln(2)}{\lambda}$
Mode	0
$\phi(t)$	$\frac{\lambda}{\lambda - it}$

### 6.3 Exponential distribution


Relation to Gamma distribution:

$$X \sim \Gamma(1, \lambda).$$

Hence, estimation of parameter  $\lambda$  is the particular case of estimation of rate  $\beta$  for Gamma distribution.



## 7 Geometric Stable distribution



Notation	$X \sim \text{GS}_\alpha(\beta, \gamma, \mu)$
Parameters	$\alpha \in (0, 2], \beta \in [-1, 1], \gamma > 0, \mu \in \mathbb{R}$
Domain	$x \in \dots$
$f(x)$	Calculated numerically
$F(x)$	Calculated numerically
$\mathbb{E}[X]$	$k + \lambda$
$\text{Var}(X)$	$2(k + 2\lambda)$
Median	Searched numerically
Mode	Searched numerically
$\phi(t)$	...

### 7.1 Asymmetric Laplace distribution

### 7.2 Laplace distribution

## 8 Kolmogorov-Smirnov distribution

## 9 Logistic distribution

## 10 Log-normal distribution

## 11 Marchenko-Pastur distribution

Notation	$X \sim \mathcal{MP}(\lambda, \sigma^2)$
Parameters	$\lambda, \sigma^2 > 0$
Domain	$x \in [\sigma^2 a, \sigma^2 b]$ , if $\lambda < 1$ , $x \in [\sigma^2 a, \sigma^2 b] \cup \{0\}$ , otherwise, where $a = (1 - \sqrt{\lambda})^2$ and $b = (1 + \sqrt{\lambda})^2$
$f(x)$	...
$F(x)$	...
$\mathbb{E}[X]$	$\sigma^2$
$\text{Var}(X)$	$\sigma^4 \lambda$
Median	0 if $\lambda > 2$ , otherwise searched numerically
Mode	$\frac{\sigma^2(\lambda-1)^2}{\lambda+1}$ , if $\lambda < 1$ , 0, otherwise
$\phi(t)$	Calculated numerically

**Calculation of characteristic function.** For  $\lambda > 1$  we use numerical integration by definition

$$\phi(t) = \int_{\sigma^2 a}^{\sigma^2 b} \cos(tx) f(x) dx + i \int_{\sigma^2 a}^{\sigma^2 b} \sin(tx) f(x) dx.$$


For  $\lambda = 1$  we split the integrand for real part by  $(\cos(tx) - 1)f(x)$  and  $f(x)$ :

$$\Re(\phi(t)) = \int_{\sigma^2 a}^{\sigma^2 b} (\cos(tx) - 1)f(x) dx + 1.$$

And for  $\lambda < 1$  we calculate integral at point 0 separately:

$$\begin{aligned} \phi(t) &= \int_{\{0\} \cup [\sigma^2 a, \sigma^2 b]} \cos(tx) f(x) dx + i \int_{\{0\} \cup [\sigma^2 a, \sigma^2 b]} \sin(tx) f(x) dx \\ &= 1 - \frac{1}{\lambda} + \int_{\sigma^2 a}^{\sigma^2 b} \cos(tx) f(x) dx + i \int_{\sigma^2 a}^{\sigma^2 b} \sin(tx) f(x) dx. \end{aligned}$$

## 12 Nakagami distribution



Notation	$X \sim \text{Nakagami}(\mu, \omega)$
Parameters	$\mu, \omega > 0$
Domain	$x \in \mathbb{R}^+$
$f(x)$	$\frac{2\mu^\mu}{\Gamma(\mu)\omega^\mu} x^{2\mu-1} e^{-\frac{\mu}{\omega}x^2}$
$F(x)$	$P(\mu, \mu x^2 / \omega)$
$\mathbb{E}[X]$	$\frac{\Gamma(\mu+1/2)}{\Gamma(\mu)} \sqrt{\frac{\omega}{\mu}}$
$\text{Var}(X)$	$\omega - (\mathbb{E}[X])^2$
Median	Searched numerically
Mode	$\max\left(\sqrt{\frac{(2\mu-1)\omega}{2\mu}}, 0\right)$
$\phi(t)$	Calculated numerically

**Calculation of characteristic function.** For  $\mu < 1$   $\lim_{x \rightarrow 0} f(x) \rightarrow \infty$ . Then we use the following transformation for real part of characteristic function:

$$\begin{aligned} \Re(\phi(t)) &= \int_0^\infty \cos(tx) f(x) dx \\ &= \int_0^\infty (\cos(tx) - 1) f(x) dx + 1 \end{aligned}$$

Relation to other distributions: if  $Y \sim \Gamma(\mu, \mu/\omega)$ , then

$$X \sim \text{Nakagami}(\mu, \omega).$$

### 12.1 Chi distribution

Notation:

$$X \sim \chi_k$$

Relation to Nakagami distribution:

$$X \sim \text{Nakagami}(k/2, k).$$

## 12.2 Maxwell-Boltzmann distribution

Notation:

$$X \sim \text{MB}(\sigma)$$

Relation to Nakagami distribution:

$$X \sim \text{Nakagami}(3/2, \sigma^2).$$

## 12.3 Rayleigh distribution

Notation:


$$X \sim \text{Rayleigh}(\sigma)$$

Relation to Nakagami distribution:

$$X \sim \text{Nakagami}(1, 2\sigma^2).$$

**Estimation of scale.** ...

## 13 Noncentral Chi-Squared distribution



Notation	$X \sim \chi_k'^2(\lambda)$
Parameters	$k > 0, \lambda > 0$
Domain	$x \in \mathbb{R}^+$
$f(x)$	$\frac{1}{2} e^{-\frac{x+\lambda}{2}} \left(\frac{x}{\lambda}\right)^{\frac{k-2}{4}} I_{\frac{k}{2}-1}(\sqrt{\lambda x})$
$F(x)$	$\text{MarcumP}_{\frac{k}{2}}\left(\frac{\lambda}{2}, \frac{x}{2}\right)$
$\mathbb{E}[X]$	$k + \lambda$
$\text{Var}(X)$	$2(k + 2\lambda)$
Median	Searched numerically
Mode	Searched numerically for $k > 2$ , 0, otherwise
$\phi(t)$	$\frac{\exp \frac{-it\lambda}{1-2it}}{(1-2it)^{k/2}}$

Relation to other distributions:


- Let  $X_1, \dots, X_k$  be independent with  $X_i \sim \mathcal{N}(\mu_i, 1)$ ,  $i = 1, \dots, k$ . Then

$$\sum_{i=1}^k X_i^2 \sim \chi_k'^2\left(\sum_{i=1}^k \mu_i^2\right).$$

- If  $\lambda = 0$ , then  $X \sim \chi_k^2$ .
- If  $J \sim \text{Po}(\lambda)$ , then  $\chi_{k+2J}^2 \sim \chi_k'^2(\lambda)$ .



## 14 Pareto distribution



Notation	$X \sim \text{Pareto}(\alpha, \sigma)$
Parameters	$\alpha, \sigma > 0$
Domain	$x \geq \sigma$
$f(x)$	$\frac{\alpha \sigma^\alpha}{x^{\alpha+1}}$
$F(x)$	$1 - \left(\frac{\sigma}{x}\right)^\alpha$
$\mathbb{E}[X]$	$\frac{\alpha \sigma}{\alpha - 1}$ for $\alpha > 1$ , $\infty$ otherwise
$\text{Var}(X)$	$\frac{\sigma^2 \alpha}{(\alpha - 1)^2 (\alpha - 2)}$ for $\alpha > 2$ , $\infty$ otherwise
Median	$\sigma 2^{1/\alpha}$
Mode	$\sigma$
$\phi(t)$	Calculated numerically

### Estimation of parameters.

**Frequentist inference.** Log-likelihood function is

$$\ln \mathcal{L}(\alpha, \sigma | X) = n \ln \alpha + n \alpha \ln \sigma - (\alpha + 1) \sum_{i=1}^n \ln X_i.$$

We assume that  $\sigma \leq X_{(1)}$ , otherwise sample  $X$  couldn't have been generated from such distribution. It is obvious, that  $\ln \mathcal{L}(\alpha, \sigma | X)$  is an increasing function in terms of  $\sigma$ , therefore  $\hat{\sigma} = X_{(1)}$  is an optimal estimator. Let's take derivative with respect to  $\alpha$ :

$$\frac{\partial \ln \mathcal{L}(\alpha, \sigma | X)}{\partial \alpha} = \frac{n}{\alpha} + n \ln \sigma - \sum_{i=1}^n \ln X_i.$$

From this we conclude that the maximum-likelihood estimator of shape is

$$\hat{\alpha} = \frac{1}{\frac{1}{n} \left( \sum_{i=1}^n \ln X_i \right) - \ln \hat{\sigma}}.$$

It is known that  $\hat{\sigma} \sim \text{Pareto}(n\alpha, \sigma)$  and  $\hat{\alpha} \sim \text{Inv-}\Gamma(n-1, n\alpha)$  and they are independent. Then

$$\mathbb{E}[\hat{\sigma}] = \frac{\sigma}{1 - \frac{1}{n\alpha}}$$

and

$$\mathbb{E}[\hat{\alpha}] = \frac{n\alpha}{n-2}.$$

Therefore, in order to get unbiased estimators we need to make the following transformations:

$$\tilde{\alpha} = \frac{n-2}{n}\hat{\alpha} \quad \text{and} \quad \tilde{\sigma} = \hat{\sigma} \left(1 - \frac{1}{(n-1)\hat{\alpha}}\right).$$

Note that if we estimate parameters separately, then  $\hat{\alpha} \sim \text{Inv-}\Gamma(n, n\alpha)$  and transformations are different.

**Bayesian inference.** We now assume that  $\sigma$  is known and prior distribution of  $\alpha$  is  $\Gamma(\kappa, \beta)$ :

$$h(\alpha) = \frac{\beta^\kappa}{\Gamma(\kappa)} \alpha^{\kappa-1} e^{-\beta\alpha}.$$

The density of posterior distribution is

$$f(\alpha|X) \propto \prod_{i=1}^n \frac{\sigma^\alpha}{X_i^{\alpha-1}} \cdot \alpha^{\kappa+n-1} e^{-\beta\alpha} \propto \alpha^{\kappa+n-1} e^{-(\beta + \sum_{i=1}^n \ln(X_i/\sigma))\alpha}.$$

Therefore,  $\alpha|X \sim \Gamma(\kappa+n, \beta + \sum_{i=1}^n \ln(X_i/\sigma))$  and Bayesian estimator is

$$\mathbb{E}[\alpha|X] = \frac{\kappa+n}{\beta + \sum_{i=1}^n \ln(X_i/\sigma)}.$$

MAP estimator is

$$\alpha_{MAP} = \frac{\kappa+n-1}{\beta + \sum_{i=1}^n \ln(X_i/\sigma)}.$$

*Note on fitting scale with Bayes:* let it be vice versa,  $\alpha$  is known while  $\sigma$  is not. Then we say that a priori  $\sigma \sim \text{Pareto}(\kappa, \theta)$ :

$$h(\sigma) = \frac{\kappa\theta^\kappa}{\sigma^{\kappa+1}}.$$

Then posterior distribution is:

$$f(\sigma|X) \propto \prod_{i=1}^n \frac{1}{X_i^{\alpha-1}} \cdot \sigma^{\alpha n - \kappa - 1} \mathbf{1}_{\{\theta < \sigma < X_{(1)}\}} \sim \text{Bounded-Pareto}(\kappa - \alpha n, \theta, X_{(1)}).$$

This imposes the following additional constraints on the prior hyperparameters:  $\kappa > \alpha n$  and  $\theta < X_{(1)}$ . Bayesian estimator:


$$\mathbb{E}[\sigma|X] = \frac{\theta^{\alpha'}}{1 - \left(\frac{\theta}{X_{(1)}}\right)^{\alpha'}} \cdot \left(\frac{\alpha'}{\alpha' - 1}\right) \cdot \left(\frac{1}{\theta^{\alpha'}} - \frac{1}{X_{(1)}^{\alpha'}}\right)$$

with  $\alpha' = \kappa - \alpha n$ . MAP estimator is just

$$\sigma_{MAP} = \theta.$$

However, Bounded-Pareto distribution is not yet supported in RandLib.

## 15 Planck distribution



Notation	$X \sim \text{Planck}(a, b)$
Parameters	$a, b > 0$
Domain	$x \in \mathbb{R}^+$
$f(x)$	$\frac{b^{a+1}}{\Gamma(a+1)\zeta(a+1)} \cdot \frac{x^a}{e^{bx}-1}$
$F(x)$	Calculated numerically
$\mathbb{E}[X]$	$\frac{(a+1)\zeta(a+2)}{b\zeta(a+1)}$
$\text{Var}(X)$	$\frac{(a+1)(a+2)\zeta(a+3)}{b^2\zeta(a+1)} - (\mathbb{E}[X])^2$
Median	Searched numerically
Mode	$\frac{W_0(-ae^{-a})+a}{b}$ , if $a > 1$ , otherwise 0
$\phi(t)$	Calculated numerically

**Calculation of cumulative distribution function.** For  $a \geq 1$   $F(x)$  can be calculated by straightforward numerical integration:

$$F(x) = \frac{b^{a+1}}{\Gamma(a+1)\zeta(a+1)} \int_0^x \frac{t^a}{e^{bt}-1} dt.$$

Note that for  $a < 1$  integrand has a singularity point at  $t = 0$ . In such case we define

$$h(t) = \frac{b^{a+2}t^{a+1}}{\Gamma(a+1)\zeta(a+1)} \cdot \left( \frac{1}{e^{bt}-1} - \frac{1}{bt} \right)$$

and then


$$F(x) = \int_0^x h(t) dt + \frac{(bx)^a}{a\Gamma(a+1)\zeta(a+1)}.$$

**Calculation of characteristic function.** The idea of calculations for  $a < 1$  is near the same. We split the real part of  $\phi(t)$  into 3 different integrals:

$$\Re(\phi(t)) = \int_0^1 \cos(tx)h(x)dx + \int_1^\infty \cos(tx)f(x)dx + \frac{b^a}{a\Gamma(a+1)\zeta(a+1)} \left( \cos(t) + t \int_0^1 \sin(tx)x^a dx \right).$$

All the integrands now have no singularity points.

## 16 Stable distribution



Notation	$X \sim S_\alpha(\beta, \gamma, \mu)$
Parameters	$\alpha \in (0, 2], \beta \in [-1, 1],$ $\gamma > 0, \mu \in \mathbb{R}$
Domain	$x \in \mathbb{R}$ , if $\beta \neq 1$ , $x \in [\mu, \infty)$ , if $\beta = 1, \alpha < 2$ , $x \in (-\infty, \mu]$ , if $\beta = -1, \alpha < 2$
$f(x)$	Calculated numerically
$F(x)$	Calculated numerically
$\mathbb{E}[X]$	$\mu$ for $\alpha > 1$ , otherwise undefined
$\text{Var}(X)$	$2\gamma^2 1_{\{\alpha=2\}} + \infty 1_{\{\alpha<2\}}$
Median	$\mu$ for $\beta = 0$ , otherwise searched numerically
Mode	$\mu$ , if $\beta = 0$ or $\alpha = 2$ , $\mu + \frac{\beta\gamma}{3}$ , if $ \beta  = 1$ and $\alpha = \frac{1}{2}$ , otherwise searched numerically
$\phi(t)$	...

Calculation of p.d.f.

Calculation of c.d.f.

### 16.1 Cauchy distribution


Relation to Stable distribution:

$$X \sim S_1(0, \gamma, \mu)$$

### 16.2 Levy distribution

Relation to Stable distribution:

$$X \sim S_{\frac{1}{2}}(1, \gamma, \mu)$$



Notation	$X \sim \text{Cauchy}(\mu, \gamma)$
Parameters	$\mu \in \mathbb{R}, \gamma^2 > 0$
Domain	$x \in \mathbb{R}$
$f(x)$	$\frac{1}{\pi\gamma \left[ 1 + \left( \frac{x-\mu}{\gamma} \right)^2 \right]}$
$F(x)$	$\frac{1}{\pi} \text{atan} \left( \frac{x-\mu}{\gamma} \right) + \frac{1}{2}$
$\mathbb{E}[X]$	Undefined
$\text{Var}(X)$	$\infty$
Median	$\mu$
Mode	$\mu$
$\phi(t)$	$e^{i\mu t - \gamma t }$

### 16.3 Normal distribution

Relation to Stable distribution:

$$X \sim S_2(\cdot, \sigma^2/2, \mu)$$

#### Estimation of parameters

**Frequentist inference.** Maximum-likelihood estimators for Normal distribution are very well-known:


$$\hat{\mu} = \overline{X}_n \quad \text{and} \quad \hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \mu)^2.$$

However, for unknown  $\mu$  the value of  $\hat{\sigma}^2 \sim \frac{\sigma^2}{n} \chi_{n-1}^2$ . Therefore, unbiased estimator in this case would be

$$\widetilde{\sigma}^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \overline{X}_n)^2.$$

Moreover, if one is interested in estimating scale  $\sigma$  with known  $\mu$ , then maximum likelihood estimator is

$$\hat{\sigma} = \sqrt{\hat{\sigma}^2} \sim \frac{\sigma}{\sqrt{n}} \chi_n$$



Notation	$X \sim \mathcal{N}(\mu, \sigma^2)$
Parameters	$\mu \in \mathbb{R}, \sigma^2 > 0$
Domain	$x \in \mathbb{R}$
$f(x)$	$\frac{1}{\sqrt{2\sigma^2\pi}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$
$F(x)$	$\frac{1}{2} \operatorname{erfc}\left(\frac{\mu-x}{\sqrt{2\sigma^2}}\right)$
$\mathbb{E}[X]$	$\mu$
$\operatorname{Var}(X)$	$\sigma^2$
Median	$\mu$
Mode	$\mu$
$\phi(t)$	$e^{i\mu t - \frac{1}{2}\sigma^2 t^2}$

and

$$\mathbb{E}[\hat{\sigma}] = \frac{\sigma}{\sqrt{n}} \sqrt{2} \frac{\Gamma((n+1)/2)}{\Gamma(n/2)}.$$

Then unbiased estimator is

$$\tilde{\sigma} = \hat{\sigma} \sqrt{\frac{n}{2}} \frac{\Gamma(n/2)}{\Gamma((n+1)/2)}$$

**Bayesian inference.** ...

## 16.4 Holtsmark distribution

Relation to Stable distribution:


$$X \sim S_{\frac{3}{2}}(0, \gamma, \mu)$$

## 16.5 Landau distribution

Relation to Stable distribution:

$$X \sim S_1(1, \gamma, \mu)$$

## 17 Weibull



Notation	$X \sim \text{Weibull}(\lambda, k)$
Parameters	$\lambda, k > 0$
Domain	$x \in \mathbb{R}^+$
$f(x)$	$\frac{k}{\lambda} \left(\frac{x}{\lambda}\right)^{k-1} \exp(-(x/\lambda)^k)$
$F(x)$	$1 - \exp(-(x/\lambda)^k)$
$\mathbb{E}[X]$	$\lambda \Gamma(1 + 1/k)$
$\text{Var}(X)$	$\lambda^2 \Gamma(1 + 2/k) - (\mathbb{E}[X])^2$
Median	$\lambda (\ln 2)^{\frac{1}{k}}$
Mode	$\lambda \left(1 - \frac{1}{k}\right)^{\frac{1}{k}}$
$\phi(t)$	Calculated numerically

### Estimation of scale

**Frequentist inference.** Log-likelihood function:

$$\ln \mathcal{L}(\lambda, k | X) = n(\ln k - \ln \lambda) + (k-1) \sum_{i=1}^n (\ln X_i - \ln \lambda) - \frac{1}{\lambda^k} \sum_{i=1}^n X_i^k.$$

The derivative with respect to scale:

$$\frac{\partial \ln \mathcal{L}(\lambda, k | X)}{\partial \lambda} = -\frac{nk}{\lambda} + \frac{k}{\lambda^{k+1}} \sum_{i=1}^n X_i^k = 0.$$

Therefore, maximum-likelihood estimation for  $\lambda$  is

$$\hat{\lambda} = \left( \frac{1}{n} \sum_{i=1}^n X_i^k \right)^{\frac{1}{k}}.$$



**Bayesian inference.** Assume  $k$  is known. Instead of estimating  $\lambda$  we give an estimation for  $\lambda^k$ . Let's say that prior distribution of  $\lambda^k$  is  $\text{Inv-}\Gamma(\alpha, \beta)$ :

$$h(\lambda^k) = \frac{\beta^\alpha}{\Gamma(\alpha)} \lambda^{-k(\alpha+1)} e^{-\beta/\lambda^k}.$$

Posterior distribution then:

$$f(\lambda^k|X) \propto \lambda^{-k(\alpha+1+n)} e^{-\frac{1}{\lambda^k}(\beta + \sum_{i=1}^n X_i^k)} \sim \text{Inv-}\Gamma\left(\alpha + n, \beta + \sum_{i=1}^n X_i^k\right).$$

Bayesian estimator:

$$\mathbb{E}[\lambda^k|X] = \frac{\beta + \sum_{i=1}^n X_i^k}{\alpha + n - 1},$$


MAP estimator:

$$\lambda_{MAP}^k = \frac{\beta + \sum_{i=1}^n X_i^k}{\alpha + n + 1}.$$

## Part III

# Discrete univariate distributions


## 18 Beta-binomial distribution



Notation	$X \sim \text{BB}(n, \alpha, \beta)$
Parameters	$n \in \mathbb{N}, \alpha, \beta > 0$
Domain	$k \in \{0, \dots, n\}$
$\mathbb{P}(X = k)$	$\binom{n}{k} \frac{B(k+\alpha, n-k+\beta)}{B(\alpha, \beta)}$
$\mathbb{P}(X \leq k)$	Calculated numerically
$\mathbb{E}[X]$	$n \frac{\alpha}{\alpha + \beta}$
$\text{Var}(X)$	$\frac{n\alpha\beta(\alpha+\beta+n)}{(\alpha+\beta)^2(\alpha+\beta+1)}$
Median	Searched numerically
Mode	Searched numerically
$\phi(t)$	Calculated numerically

Relation to other distributions: if  $p \sim \mathcal{B}(\alpha, \beta)$ , then  $\text{Bin}(n, p) \sim \text{BB}(n, \alpha, \beta)$ .

## 19 Binomial distribution



Notation	$X \sim \text{Bin}(n, p)$
Parameters	$n \in \mathbb{N}, p \in [0, 1]$
Domain	$k \in \{0, \dots, n\}$
$\mathbb{P}(X = k)$	$\binom{n}{k} p^k (1 - p)^{n-k}$
$\mathbb{P}(X \leq k)$	$I_{1-p}(n - k, 1 + k)$
$\mathbb{E}[X]$	$np$
$\text{Var}(X)$	$np(1 - p)$
Median	$[np]$
Mode	$[(n + 1)p]$
$\phi(t)$	$(1 - p + pe^{it})^n$

**Estimation of probability  $p$  with known number  $n$ .**

**Frequentist inference.** Log-likelihood function:

$$\ln \mathcal{L}(p|X) \propto \sum_{i=1}^k (X_i \log p + (n - X_i) \log(1 - p))$$

The derivative with respect to  $p$  is:

$$\frac{\partial \ln \mathcal{L}(p|X)}{\partial p} = \frac{\sum_{i=1}^k X_i}{p} - \frac{nk - \sum_{i=1}^k X_i}{1 - p}.$$

Therefore we reach the maximum value of log-likelihood if

$$p = \frac{\bar{X}_k}{n}.$$

**Bayesian inference.** We set prior Beta distribution  $\mathcal{B}(\alpha, \beta)$ :

$$h(p) = \frac{p^{\alpha-1} (1 - p)^{\beta-1}}{B(\alpha, \beta)}.$$

Then posterior is

$$f(p|X) \propto p^{\alpha-1+\sum_{i=1}^k X_i} (1-p)^{\beta-1+\sum_{i=1}^k (n-X_i)} \sim \mathcal{B}\left(\alpha + \sum_{i=1}^k X_i, \beta + nk - \sum_{i=1}^k X_i\right).$$

Thus Bayesian estimator is

$$\mathbb{E}[p|X] = \frac{\alpha + \sum_{i=1}^k X_i}{\alpha + \beta + nk}$$

and MAP estimator is

$$p_{MAP} = \frac{\alpha + \sum_{i=1}^k X_i - 1}{\alpha + \beta + nk - 2}.$$

Also, Minimax estimator is equal to Bayes estimator if  $\alpha = \beta = \frac{1}{2}\sqrt{n}$ .

## 19.1 Bernoulli


Notation:

$$X \sim \text{Bernoulli}(p).$$

Relation to Binomial distribution:

$$X \sim \text{Bin}(1, p).$$

## 20 Hypergeometric distribution



Notation	$X \sim \text{HG}(N, K, n)$
Parameters	$N \in \mathbb{N}, K \in \{1, 2, \dots, N\},$ $n \in \{1, 2, \dots, N\}$
Domain	$\max(0, n + K - N) \leq k \leq \min(n, K)$
$\mathbb{P}(X = k)$	$\frac{\binom{K}{k} \binom{N-K}{n-k}}{\binom{N}{n}}$
$\mathbb{P}(X \leq k)$	Calculated numerically
$\mathbb{E}[X]$	$\frac{nK}{N}$
$\text{Var}(X)$	$\frac{nK(N-K)(N-n)}{N^2(N-1)}$
Median	Searched numerically
Mode	$\left\lfloor \frac{(n+1)(K+1)}{N+2} \right\rfloor$
$\phi(t)$	Calculated numerically

### Estimation of number of target members of population $K$ .

**Bayesian inference.** Let prior distribution of  $K$  be Beta-Binomial distribution  $BB(N, \alpha, \beta)$ :

$$h(K) = \binom{N}{K} \frac{B(K + \alpha, N - K + \beta)}{B(\alpha, \beta)}.$$

Then for one sample  $X$ :


$$K - X \sim BB(N - n, \alpha + X, \beta + nk - X)$$

and therefore

$$\mathbb{E}[K|X] = X + (N - n) \frac{\alpha}{\alpha + \beta}.$$

However, RandLib doesn't support Bayesian fitting for Hypergeometric distribution yet.

## 21 Negative-Binomial (Polya) distribution



Notation	$X \sim \text{NB}(r, p)$
Parameters	$r > 0, p \in (0, 1)$
Domain	$k \in \mathbb{N}_0$
$\mathbb{P}(X = k)$	$\binom{k+r-1}{k} p^r (1-p)^k$
$\mathbb{P}(X \leq k)$	$I_p(r, k+1)$
$\mathbb{E}[X]$	$\frac{1-p}{p} r$
$\text{Var}(X)$	$\frac{1-p}{p^2} r$
Median	Searched numerically
Mode	$\max \left( \left\lfloor \frac{(r-1)(1-p)}{p} \right\rfloor, 0 \right)$
$\phi(t)$	$\left( \frac{p}{1-(1-p)e^{it}} \right)^r$

Relation to other distributions: if  $\lambda \sim \text{Gamma} \left( r, \frac{p}{1-p} \right)$ , then  $\text{Po}(\lambda) \sim \text{NB}(r, p)$ .

### 21.1 Geometric distribution

Notation:

$$X \sim \text{Geometric}(p).$$

Relation to Negative-Binomial distribution:

$$X \sim \text{NB}(1, p).$$


### 21.2 Pascal distribution

Notation:

$$X \sim \text{Pascal}(r, p).$$

The only difference with Negative-Binomial distribution is that for Pascal distribution shape  $r$  is an integer.

## 22 Poisson distribution



Notation	$X \sim \text{Po}(\lambda)$
Parameters	$\lambda > 0$
Domain	$k \in \mathbb{N}_0$
$\mathbb{P}(X = k)$	$\frac{\lambda^k e^{-\lambda}}{k!}$
$\mathbb{P}(X \leq k)$	$Q(k + 1, \lambda)$
$\mathbb{E}[X]$	$\lambda$
$\text{Var}(X)$	$\lambda$
Median	$\sim \max\left(\left[\lambda + \frac{1}{3} - \frac{0.02}{\lambda}\right], 0\right)$
Mode	$[\lambda]$
$\phi(t)$	$\exp\{\lambda(e^{it} - 1)\}$

**Estimation of rate.**

**Frequentist inference.** Log-likelihood function:

$$\ln \mathcal{L}(\lambda|X) \propto -\lambda n + \sum_{i=1}^n X_i \log \lambda.$$

Setting the derivative w.r.t. rate to 0 we get the optimal value:

$$\lambda = \overline{X}_n.$$

**Bayesian inference.** Let set prior distribution of  $\lambda \sim \Gamma(\alpha, \beta)$ :

$$h(\lambda) = \frac{\beta^\alpha}{\Gamma(\alpha)} \lambda^{\alpha-1} e^{-\beta\lambda}.$$

Posterior distribution:

$$f(\lambda|X) \propto e^{-\lambda(\beta+n)} \lambda^{\alpha-1+\sum_{i=1}^n X_i} \sim \Gamma\left(\alpha + \sum_{i=1}^n X_i, \beta + n\right).$$

Therefore, Bayesian estimator:


$$\mathbb{E}[\lambda|X] = \frac{\alpha + \sum_{i=1}^n X_i}{\beta + n}.$$

And MAP estimator:

$$\lambda_{MAP} = \max\left(\frac{\alpha + \sum_{i=1}^n X_i - 1}{\beta + n}, 0\right).$$



## 23 Skellam distribution



Notation	$X \sim \text{Skellam}(\mu_1, \mu_2)$
Parameters	$\mu_1, \mu_2 > 0$
Domain	$k \in \mathbb{Z}$
$\mathbb{P}(X = k)$	$e^{-(\mu_1 + \mu_2)} \left(\frac{\mu_1}{\mu_2}\right)^{\frac{k}{2}} I_k(2\sqrt{\mu_1 \mu_2})$
$\mathbb{P}(X \leq k)$	$\text{MarcumP}_{k+1}(\mu_2, \mu_1), k \geq 0$ $\text{MarcumQ}_{-k}(\mu_1, \mu_2), k < 0$
$\mathbb{E}[X]$	$\mu_1 - \mu_2$
$\text{Var}(X)$	$\mu_1 + \mu_2$
Median	Searched numerically
Mode	$[\mu_1 - \mu_2]$
$\phi(t)$	$\exp\{\mu_1(e^{it} - 1) - \mu_2(e^{it} - 1)\}$

Relation to other distributions: if  $Y \sim \text{Po}(\mu_1)$  and  $Z \sim \text{Po}(\mu_2)$ , then

$$Y - Z \sim \text{Skellam}(\mu_1, \mu_2).$$

## 24 Yule distribution

## 25 Zeta distribution

## 26 Zipf distribution

## Part IV

# Bivariate distributions

- 27 Bivariate Normal distribution
- 28 Normal-Inverse-Gamma distribution
- 29 Trinomial distribution

## Part V

# Circular distributions

30 von Mises distribution

31 Wrapped Exponential distribution

## Part VI

# Singular distributions

## 32 Cantor distribution