

Variation

November 7, 2018

1 The principle of least action

Let the system occupy positions defined by two sets of values of the coordinates, q_1 and q_2 , at the instants t_1 and t_2 . The *action* of the system takes the form of integral of its *Lagrangian* as

$$S = \int_{t_1}^{t_2} L(q, \dot{q}, t) dt \quad (1)$$

The necessary condition for S to have an extremum is that the variation δS should be zero:

$$\begin{aligned} 0 = \delta S &= \delta \int_{t_1}^{t_2} L(q, \dot{q}, t) dt \\ &= \delta \int_{t_1}^{t_2} \left(\frac{\partial L}{\partial q} \delta q + \frac{\partial L}{\partial \dot{q}} \delta \dot{q} \right) dt \end{aligned} \quad (2)$$

Since $\delta \dot{q} = d\delta q / dt$, we obtain, on integrating the second term by parts,

$$\delta S = \left[\frac{\partial L}{\partial \dot{q}} \delta q \right]_{t_1}^{t_2} + \int_{t_1}^{t_2} \left(\frac{\partial L}{\partial q} - \frac{d}{dt} \frac{\partial L}{\partial \dot{q}} \right) \delta q dt = 0 \quad (3)$$

For $t = t_1$ and $t = t_2$, the values of $q(t)$ should be fixed, i.e.

$$\delta q(t_1) = \delta q(t_2) = 0 \quad (4)$$

Thus the first term in (3) vanishes, so we have

$$\frac{d}{dt} \frac{\partial L}{\partial \dot{q}} - \frac{\partial L}{\partial q} = 0 \quad (5)$$

Considering the degrees of the system, we then obtain s equations of the form

$$\frac{d}{dt} \frac{\partial L}{\partial \dot{q}_i} - \frac{\partial L}{\partial q_i} = 0 \quad (i = 1, 2, \dots, s). \quad (6)$$

These are called *Euler-Lagrange's equations* in mechanics.

2 Gauge invariance

Considering two functions $L'(q, \dot{q}, t)$ and $L(q, \dot{q}, t)$, differing by the total derivative with respect to time of some function $f(q, t)$ as

$$L'(q, \dot{q}, t) = L(q, \dot{q}, t) + \frac{d}{dt}f(q, t) \quad (7)$$

The integrals are

$$S' = \int_{t_1}^{t_2} L'(q, \dot{q}, t) dt = \int_{t_1}^{t_2} L(q, \dot{q}, t) dt + \int_{t_1}^{t_2} \frac{df}{dt} dt = S + f(q_2, t_2) - f(q_1, t_1)$$

Where $f(q_2, t_2)$ and $f(q_1, t_1)$ are fixed values, i.e. $\delta f(q_2, t_2) = \delta f(q_1, t_1) = 0$. This leads to

$$\delta S' = \delta S \quad (8)$$

So the conditions $\delta S' = 0$ and $\delta S = 0$ are equivalent.

3 Law of inertia

A frame of reference in which space is homogeneous and isotropic and time is homogeneous is called an *inertial frame*.

Considering a particle moving freely in such a frame. The homogeneity of space and time implies that the Lagrangian cannot contain explicitly either the radius vector \mathbf{r} or the time t , i.e. L must be a function of the velocity \mathbf{v} only; Since space is isotropic, the Lagrangian must also be independent of the direction of \mathbf{v} , and is therefore a function only of its magnitude, i.e. of $\mathbf{v}^2 = v^2$:

$$L = L(v^2) \quad (9)$$

Bringing $\partial L / \partial \mathbf{r} = 0$ into Lagrange's equation (5), we have

$$0 = \frac{d}{dt} \left(\frac{\partial L}{\partial \mathbf{v}} \right) = \frac{d\mathbf{v}}{dt} \frac{d}{d\mathbf{v}} \left(\frac{\partial L}{\partial \mathbf{v}} \right) \quad (10)$$

whence $\partial L / \partial \mathbf{v} = \text{constant}$. Since $\partial L / \partial \mathbf{v}$ is a function of the velocity only, it follows that

$$\frac{d\mathbf{v}}{dt} = 0, \quad \text{or} \quad \mathbf{v} = \text{constant}. \quad (11)$$

Thus we conclude that, in an inertial frame, any free motion takes place with a velocity which is constant in both magnitude and direction. This is the *law of inertia*.