

Pregunta:

¿En pacientes adultos con diabetes mellitus tipo 2 (P), los hábitos alimentarios inadecuados y un estado nutricional alterado (I), en comparación con hábitos y estado nutricional adecuados (C), se relacionan con las alteraciones en los niveles de HbA1c (O)?

Fórmula: 34 resultados

((young adult) OR (diabetes type 2) OR (diabetes mellitus type 2) OR (glucose intolerance)) AND ((feeding behavior) OR (diet) OR (unhealthy diet) OR (nutritional status)) AND ((healthy lifestyle) OR (food security)) AND ((glycated hemoglobin) OR (blood glucose) OR (glycemic control))

https://pubmed.ncbi.nlm.nih.gov/?term=%28%28young+adult%29+OR+%28diabetes+type+2%29+OR+%28diabetes+mellitus+type+2%29+OR+%28glucose+intolerance%29%29+AND+%28%28feeding+behavior%29+OR+%28diet%29+OR+%28unhealthy+diet%29+OR+%28nutritional+status%29%29+AND+%28%28healthy+lifestyle%29+OR+%28food+security%29%29+AND+%28%28glycated+hemoglobin%29+OR+%28blood+glucose%29+OR+%28glycemic+control%29%29&filter=datesearch.y_5&filter=pubt.clinicaltrial&filter=pubt.randomizedcontrolledtrial