

---

# To those physics students who asked why $q$ and $\dot{q}$ are independent in Lagrangian Mechanics

Niklas Zorbach<sup>1</sup> and Marco Knipfer<sup>1, 2</sup>

<sup>1</sup> Institute for Theoretical Physics, Goethe-University Frankfurt, Germany

<sup>2</sup> Institute for Physics and Astronomy, The University of Alabama, USA

---

May 26, 2019

**L**orem ipsum dolor sit amet, consectetur adipiscing elit. Fusce maximus nisi ligula. Morbi laoreet ex ligula, vitae lobortis purus mattis vel. Vestibulum ante ipsum primis in faucibus orci luctus et ultrices posuere cubilia Curae; Donec ac metus ut turpis mollis placerat et nec enim. Duis tristique nibh maximus faucibus facilisis. Praesent in consequat leo. Maecenas condimentum ex rhoncus, elementum diam vel, malesuada ante.

## 1 Manifolds

This chapter is basically taken from (Schuller, 2015) with our remarks to it.

We start with a set  $\mathcal{M}$  which is supposed to be the space where physics happens. The weakest structure we need in order to talk about continuity (of curves or fields) is called a topology.

*Definition 1.1* (Power set  $\mathcal{P}$ )  
The set of all subsets of  $\mathcal{M}$ .

*Definition 1.2* (Topology)  
A Topology  $\mathcal{O}$  is a subset  $\mathcal{O} \subseteq \mathcal{P}(\mathcal{M})$  satisfying:

1.  $\emptyset \in \mathcal{O}, \mathcal{M} \in \mathcal{O}$ ,

2.  $U \in \mathcal{O}, V \in \mathcal{O} \Rightarrow U \cap V \in \mathcal{O}$
3.  $U_\alpha \in \mathcal{O}, \alpha \in A \Rightarrow \left( \bigcup_{\alpha \in A} U_\alpha \right) \in \mathcal{O}$

Every set has the *chaotic topology*

$$\mathcal{O}_{\text{chaotic}} := \{\emptyset, \mathcal{M}\}, \quad (1)$$

and the *discrete topology*

$$\mathcal{O}_{\text{discrete}} := \mathcal{P}(\mathcal{M}), \quad (2)$$

which are both useless.

The special case  $\mathcal{M} = \mathbb{R}^d = \mathbb{R} \times \dots \times \mathbb{R}$  has a standard topology for which we need the definition of a soft ball.

*Definition 1.3* (Soft Ball in  $\mathbb{R}^d$ )

$$B_r(p) := \left\{ (q_1, \dots, q_d) \mid \sum_{i=1}^d (p_i - q_i)^2 < r^2 \right\}, \quad (3)$$

with  $r \in \mathcal{R}^+, p \in \mathbb{R}^d$ . Note: This does not need a norm or vector space structure on  $\mathbb{R}^d$ .

*Definition 1.4* ( $\mathcal{O}_{\text{standard}}$  on  $\mathbb{R}^d$ )

$$U \in \mathcal{O}_{\text{standard}} :\Leftrightarrow \forall p \in U : \exists r \in \mathcal{R}^+ : B_r(p) \subseteq U \quad (4)$$

*To those physics students who asked why  $q$  and  $\dot{q}$  are independent in Lagrangian Mechanics*

## **Bibliography**

Schuller, Frederic (Feb. 2015). *International Winter School on Gravity and Light 2015*.  
[https://www.youtube.com/watch?v=7G4SqIboeig&list=PLFeEvEPtX\\_OS6vxxiiNPrJbLu9aK1UVC\\_](https://www.youtube.com/watch?v=7G4SqIboeig&list=PLFeEvEPtX_OS6vxxiiNPrJbLu9aK1UVC_).