
To those physics students who asked why q and \dot{q} are independent in Lagrangian Mechanics

Niklas Zorbach¹ and Marco Knipfer^{1, 2}

¹ Institute for Theoretical Physics, Goethe-University Frankfurt, Germany

² Institute for Physics and Astronomy, The University of Alabama, USA

May 27, 2019

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Fusce maximus nisi ligula. Morbi laoreet ex ligula, vitae lobortis purus mattis vel. Vestibulum ante ipsum primis in faucibus orci luctus et ultrices posuere cubilia Curae; Donec ac metus ut turpis mollis placerat et nec enim. Duis tristique nibh maximus faucibus facilisis. Praesent in consequat leo. Maecenas condimentum ex rhoncus, elementum diam vel, malesuada ante.

1 Manifolds

1.1 Topology

This chapter is basically taken from (Schuller, 2015) with our remarks to it.

We start with a set M which is supposed to be the space where physics happens. The weakest structure we need in order to talk about continuity (of curves or fields) is called a topology.

Definition 1.1 (Power set \mathcal{P})

The set of all subsets of M .

Definition 1.2 (Topology)

A topology \mathcal{O} is a subset $\mathcal{O} \subseteq \mathcal{P}(M)$ satisfying:

1. $\emptyset \in \mathcal{O}, M \in \mathcal{O}$,
2. $U \in \mathcal{O}, V \in \mathcal{O} \Rightarrow U \cap V \in \mathcal{O}$
3. $U_\alpha \in \mathcal{O}, \alpha \in A \Rightarrow \left(\bigcup_{\alpha \in A} U_\alpha \right) \in \mathcal{O}$

Every set has the *chaotic topology*

$$\mathcal{O}_{\text{chaotic}} := \{\emptyset, M\}, \quad (1)$$

and the *discrete topology*

$$\mathcal{O}_{\text{discrete}} := \mathcal{P}(M), \quad (2)$$

which are both useless.

The special case $M = \mathbb{R}^d = \mathbb{R} \times \dots \times \mathbb{R}$ has a standard topology for which we need the definition of a soft ball.

Definition 1.3 (Soft Ball in \mathbb{R}^d)

$$B_r(p) := \left\{ (q_1, \dots, q_d) \mid \sum_{i=1}^d (p_i - q_i)^2 < r^2 \right\}, \quad (3)$$

with $r \in \mathbb{R}^+, p \in \mathbb{R}^d$. Note: This does not need a norm or vector space structure on \mathbb{R}^d .

Definition 1.4 ($\mathcal{O}_{\text{standard}}$ on \mathbb{R}^d)

$$U \in \mathcal{O}_{\text{standard}} :\Leftrightarrow \forall p \in U : \exists r \in \mathbb{R}^+ : B_r(p) \subseteq U \quad (4)$$




Figure 1: The set U is in the standard topology, V not.

Some terminology: Let M be a set with a topology $\mathcal{O} =$ set of open sets. We call (M, \mathcal{O}) a *topological space* and:

- $U \in \mathcal{O} \Leftrightarrow$: call $U \subseteq M$ an *open set*
- $M \setminus A \in \mathcal{O} \Leftrightarrow$: call $A \subseteq M$ a *closed set*

Note: The empty set is open and closed. If a set is open we cannot directly follow that it is not closed or vice versa.

1.2 Continuous maps

A map

$$f : M \rightarrow N, \quad (5)$$

takes every point from the domain M (a set) to the target N (a set). If one point $p \in N$ is not reached, the map is not *surjective*. If a point is hit twice, the map is not *injective*. A map that is injective and surjective is called *surjective*.

Definition 1.5 (Preimage)

$$f : M \rightarrow N \supseteq V$$

$$\text{preim}_f(V) := \{m \in M \mid f(m) \in V\} \quad (6)$$

Definition 1.6 (Continuity)

(M, \mathcal{O}_M) and (N, \mathcal{O}_N) topological spaces. Then a map $f : M \rightarrow N$ is called *continuous with respect to \mathcal{O}_M and \mathcal{O}_N* if

$$\forall V \in \mathcal{O}_N : \text{preim}_f(V) \in \mathcal{O}_M. \quad (7)$$

“A map is open iff the preimages of all open sets are open sets.”

Note: If a map is not surjective there are sets with preimage \emptyset , thus we need to have \emptyset in \mathcal{O} , otherwise only surjective maps could be continuous.

Bibliography

Schuller, Frederic (Feb. 2015). *International Winter School on Gravity and Light 2015*.
https://www.youtube.com/watch?v=7G4SqIboeig&list=PLFeEvEPtX_0S6vxxiiNPrJbLu9aK1UVC_.