Dosage Form

amp(s)	ampule(s)
cap(s)	capsule(s)
cr	cream
el ; elix	elixir
fl	fluid
liq	liquid
lot	lotion
pulv	powder
sol; soln	solution
spr	spray
supp	suppository
susp	suspension
syr	syrup
tab(s)	tablet(s)
tinc	tincture
una	ointment

System of Measure

cc; cm ³	cubic centimeter (liquid)
fl oz ; ℥	fluid ounce (liquid)
g;gm	gram (solid)
gr	grain (solid)
gtt(s)	drop(s) (liquid)
IU	International Unit (solid)
mcg	microgram (solid)
kg	kilogram (solid)
L	Liter (liquid)
mEq	milliequivalent (solid)
mg	milligram (solid)
mL	milliliter (liquid)
OZ	ounce (solid)
t ; tsp ; 💈	teaspoonful (liquid)
T; tbsp	tablespoonful (liquid)
u	unit (liquid)

Medication

acetaminophen
aspirin
Calcium
Epinephrine
hydrochlorothiazide
ibuprofen
potassium chloride
morphine sulfate
sodium
sodium fluoride
normal saline
nitroglycerine
penicillin
phenobarbital
(generic) Bactrim ³
Tetracycline

Frequency

ac	before meals		
AM	in the morning		
ad lib	as often as necessary		
ASAP	immediately		
atc	around the clock		
bid	twice daily		
d	daily		
h; hr; ⁰	hour, on the hour of		
hs	at bedtime		
noc	at night		
р	after		
рс	after meals		
PM	in the evening		
q	every		
qid	four times daily		
qod	every other day		
q*h	every * hours ²		
stat	immediately		
tid	three times daily		
wk	weekly		

Route of Administration

Э

Miscellaneous

а	apply
aa	affected area(s)
aff	affected
ВС	birth control
С	with
DC	discontinue
DS	double-strength
EC	enteric coated
ES	extra strength

Miscellaneous (cont.)

headache
no known drug allergies
no refills
nausea/vomiting
pain
without
single-strength ¹
shortness of breath
until gone

¹Certain sig codes have multiple meanings

²For example: q<u>8</u>h : "every <u>8</u> hours"

³Sulfamethoxazole/Trimethoprim