

$f: \mathbb{R}^{n+1} \setminus \{0\} \rightarrow \mathbb{R}$ given by $f(x) = |x|^2$ is smooth and is a submersion this means $\forall r \in \mathbb{R}_{>0}$ we have $S^n(r) = f^{-1}(r)$ is a smooth submanifold.

$$T_p S^n(r) = \ker(df_p)$$

$$df_p: \mathbb{R}^{n+1} \rightarrow \mathbb{R} \quad \text{note} \quad df_p = [2p^1 \ \dots \ 2p^{n+1}] \Rightarrow v \in T_p S^n(r) \Leftrightarrow df_p(v) = 0 \Leftrightarrow 2p \cdot v = 0 \Leftrightarrow p \cdot v = 0$$

Now let $A \in O(p, q) \Leftrightarrow A^T A = g$. Note $O(p, q) = f^{-1}(g)$ where $f(A) = A^T A$ we showed that $df_A(X) = A^T X + X^T A$ (took the curve $\gamma(t) = A + tX$ note $\gamma(0) = A$, $\gamma'(0) = X$)

$$\text{This means } T_A O(p, q) = \ker(df_A) = \{X \in \mathbb{R}^{n^2} \cong \text{Mat}(n, \mathbb{R}) \mid A^T X + X^T A = 0\}$$

Now let us compute $T_{\text{Id}} SL(n, \mathbb{R}) = \mathfrak{sl}(n)$. From the exercises note that $D(\det)(\text{Id})(A) = \text{tr}(A)$ so $T_{\text{Id}} SL(n, \mathbb{R}) = \{X \in \mathbb{R}^{n^2} \mid \text{tr}(X) = 0\}$

$L_B: SL(n, \mathbb{R}) \rightarrow SL(n, \mathbb{R})$ is a diffeomorphism this means $T_B SL(n, \mathbb{R}) \cong dL_B(T_{\text{Id}} SL(n, \mathbb{R}))$

$$L_B^{-1} \circ L_B = L_B^{-1} \circ L_B = \text{Id}_{SL(n, \mathbb{R})}$$

Let $\gamma: I \rightarrow SL(n, \mathbb{R})$ be a curve st. $\gamma(0) = \text{Id}$ and $\gamma'(0) = X \Leftrightarrow X = \left. \frac{d}{dt} \right|_{t=0} \gamma(t)$ then consider

$\psi: I \rightarrow SL(n, \mathbb{R})$ given by $\psi(t) = B\gamma(t)$ then $\psi(0) = B$ and $\psi'(0) = \left. \frac{d}{dt} \right|_{t=0} B\gamma(t) = B\gamma'(0) = BX$

So this show that dL_B can be thought of as $L_B \Rightarrow T_B SL(n, \mathbb{R}) = \{BX \mid \text{tr}(X) = 0\}$

Exercise: Show that $T_B SL(n, \mathbb{R}) \cong \{XB \mid \text{tr}(X) = 0\}$.

Let us start with the cotangent bundle. So its fibres at each point will be the $T_p^* M = (T_p M)^*$.

Suppose $f: M \rightarrow \mathbb{R}$ smooth then $df_p: T_p M \rightarrow T_{f(p)} \mathbb{R} \cong \mathbb{R} \Rightarrow df_p \in T_p^* M$

If we have some configuration space M then we care about $T^* M$ since it is the phase space and hence we can talk about Hamiltonian dynamics.

Start with a chart (U, φ) around $p \in M$. Then we have a basis for $T_p M$ given by $\left. \frac{\partial}{\partial x^i} \right|_p = d\varphi_p^{-1} \left(\left. \frac{\partial}{\partial y^i} \right|_{\varphi(p)} \right)$ where $\left. \frac{\partial}{\partial y^i} \right|_{\varphi(p)}$ is the standard basis for $T_{\varphi(p)} \mathbb{R}^m$. Now we define the

dual vectors $dx^i_p \in T_p^* M$ s.t. $dx^i_p \left(\left. \frac{\partial}{\partial x^j} \right|_p \right) = \delta^i_j$, $dx^i_p = d(\pi^i \circ \varphi)_p$. Check this!!

Now let (U, φ) be another chart around $p \in M$. With basis tangent vectors

$$\frac{\partial}{\partial \tilde{x}^i} \Big|_p = d\varphi^{-1}_{\varphi(p)} \left(\frac{\partial}{\partial y^i} \Big|_{\varphi(p)} \right)$$

$$\frac{\partial}{\partial x^i} \Big|_p = d\varphi^{-1}_{\varphi(p)} \left(\frac{\partial}{\partial y^i} \Big|_{\varphi(p)} \right)$$

$$= d(\varphi^{-1} \circ \varphi \circ \varphi^{-1})_{\varphi(p)} \left(\frac{\partial}{\partial y^i} \Big|_{\varphi(p)} \right)$$

$$= d\varphi^{-1}_{\varphi(p)} \left(d(\varphi \circ \varphi^{-1})_{\varphi(p)} \left(\frac{\partial}{\partial y^i} \Big|_{\varphi(p)} \right) \right)$$

$$= \frac{\partial \tilde{x}^j}{\partial x^i}(\varphi(p)) \frac{\partial}{\partial \tilde{x}^j} \Big|_p$$

Exercise: Fill in the steps.

This tells us that $\frac{\partial}{\partial x^i} = \frac{\partial \tilde{x}^j}{\partial x^i} \frac{\partial}{\partial \tilde{x}^j} \Leftrightarrow \frac{\partial}{\partial \tilde{x}^j} = \frac{\partial x^i}{\partial \tilde{x}^j} \frac{\partial}{\partial x^i}$

let $\omega_p = \omega_i dx^i_p = \tilde{\omega}_j d\tilde{x}^j_p$ note $\omega_i = \omega_j dx^j_p \left(\frac{\partial}{\partial x^i} \Big|_p \right) = \omega \left(\frac{\partial}{\partial x^i} \Big|_p \right) = \tilde{\omega}_j d\tilde{x}^j_p \left(\frac{\partial \tilde{x}^k}{\partial x^i} \frac{\partial}{\partial \tilde{x}^k} \Big|_p \right)$

$$= \tilde{\omega}_j \left(\frac{\partial \tilde{x}^k}{\partial x^i} \right) \delta^j_k$$

$$= \tilde{\omega}_j \frac{\partial \tilde{x}^j}{\partial x^i}$$

this means $dx^i_p = \frac{\partial x^i}{\partial \tilde{x}^j} d\tilde{x}^j$

Defining the cotangent bundle:

let $(U_\alpha, \varphi_\alpha)$ be an atlas on M . Then let $V_\alpha = \varphi_\alpha^{-1}(U_\alpha)$ and $T_\alpha = \varphi_\alpha^{-1}(U_\alpha \cap V_\beta)$

then the cocycle defining TM was $g_{\beta\alpha}: U_\alpha \cap U_\beta \rightarrow GL(n, \mathbb{R})$ given by $g_{\beta\alpha}(x) = D\varphi_{\beta\alpha} \circ \varphi_\alpha^{-1}$.

We look for a representation $\rho: GL(n, \mathbb{R}) \rightarrow GL(l, \mathbb{R})$ then the cocycles $\rho \circ g_{\beta\alpha} = t_{\beta\alpha}$

Exercise: Check $t_{\beta\alpha}$ satisfies the cocycle condition.

So for the cotangent bundle $\rho(A) = (A^*)^{-1}$. Then doing a similar construction as TM we define T^*M .

In any chart (U, φ) we can write $\omega \in T^*M$ as $\omega_i dx^i_p$ where $\omega_i \in \mathbb{R}$ and dx^i_p are the duals of $\frac{\partial}{\partial x^i}|_p$.

A vector field: the idea is that to each point associate a tangent vector.

Def: A section σ of a surjection $\pi: E \rightarrow M$ is a right inverse $\pi \circ \sigma = \text{id}_M$

Then vector fields are sections of TM .

A vector field $X: M \rightarrow TM$ is smooth iff $\forall f \in C^\infty(M)$ $Xf: M \rightarrow \mathbb{R}$ is smooth.

$(Xf)(p) = X_p f$. We denote the space of smooth vector fields on M by $\mathfrak{X}(M)$
 \uparrow
 $\{\text{mathfrak{X}}\}$

In general if we have a smooth vector bundle $\pi: E \rightarrow M$ then its space of sections is denoted by $\Gamma(E)$

$$\mathfrak{X}(M) = \Gamma(TM)$$

Smooth sections $\Gamma(T^*M) = \mathcal{S}^1(M)$ are called differential 1-forms

$M \times \mathbb{R} \rightarrow M$ is a vector bundle. Hence the space of smooth sections is

$$\Gamma(M \times \mathbb{R}) = C^\infty(M)$$

$$\parallel$$

$$\mathcal{S}^0(M)$$

Note: $\omega \in \mathcal{S}^1(M) \Leftrightarrow \forall X \in \mathfrak{X}(M)$ we have $\omega(X)$ defined by $\omega(X)(p) = \omega_p(X_p)$ is smooth