



$$L(\gamma) = \int_{t_i}^{t_f} \sqrt{\dot{x}^2 + \dot{y}^2} dt$$

$$= \int_{t_i}^{t_f} \sqrt{\delta_{ij} \dot{x}^i \dot{x}^j} dt$$




$$\|\dot{\varphi}(p)\|^2 = \langle \dot{\varphi}(p), \dot{\varphi}(p) \rangle_p = g_{ij} \dot{\varphi}^i \dot{\varphi}^j$$

$$\Rightarrow \|\varphi(p)\| = \sqrt{g_{ij} \dot{\varphi}^i \dot{\varphi}^j}$$

$$\Rightarrow L(\varphi) := \int_{t_i}^{t_f} \sqrt{g_{ij} \dot{\varphi}^i \dot{\varphi}^j} dt$$

There might not be a curve that minimizes distances like in the above.



Geodesics are curves that attain minimum length.

length function takes smooth curves to a real number.

$$S[\varphi] = \int_{t_i}^{t_f} L(\varphi, \dot{\varphi}) dt$$

$$L(\varphi + \delta\varphi, \dot{\varphi} + \delta\dot{\varphi})$$

$$= L(\varphi, \dot{\varphi}) + (DL)(\varphi, \dot{\varphi})(\delta\varphi, \delta\dot{\varphi}) + \text{other}$$

$$\varphi \rightarrow \varphi + \delta\varphi$$

$$\dot{\varphi} \rightarrow \dot{\varphi} + \delta\dot{\varphi}$$

$$\Leftrightarrow L(\varphi + \delta\varphi, \dot{\varphi} + \delta\dot{\varphi}) - L(\varphi, \dot{\varphi})$$

$$= \frac{\partial L}{\partial \varphi} \delta\varphi + \frac{\partial L}{\partial \dot{\varphi}} \delta\dot{\varphi} + \text{other}$$

$$S[\varphi + \delta\varphi] - S[\varphi] = \int_{t_i}^{t_f} L(\varphi + \delta\varphi, \dot{\varphi} + \delta\dot{\varphi}) - L(\varphi, \dot{\varphi}) dt$$

$$\approx \int_{t_i}^{t_f} \frac{\partial L}{\partial \varphi} \delta\varphi + \frac{\partial L}{\partial \dot{\varphi}} \delta\dot{\varphi} dt$$

$$= \int_{t_i}^{t_f} \frac{\partial L}{\partial \varphi} \delta\varphi + \frac{\partial L}{\partial \dot{\varphi}} \frac{d}{dt} \delta\varphi dt$$

$$\begin{aligned}
&= \int_{t_i}^{t_f} \frac{\partial L}{\partial \varphi} \delta \varphi dt + \frac{\partial L}{\partial \dot{\varphi}} \delta \varphi \Big|_{t_i}^{t_f} - \int_{t_i}^{t_f} \frac{d}{dt} \frac{\partial L}{\partial \dot{\varphi}} \delta \varphi dt \\
&= \int_{t_i}^{t_f} \left(\frac{\partial L}{\partial \varphi} - \frac{d}{dt} \frac{\partial L}{\partial \dot{\varphi}} \right) \delta \varphi dt + \frac{\partial L}{\partial \dot{\varphi}} \delta \varphi \Big|_{t_i}^{t_f} \\
&= 0
\end{aligned}$$

In the case that the end points stay fixed $\delta \varphi(t_i) = 0 = \delta \varphi(t_f)$ hence no boundary terms.

Hence $\int_{t_i}^{t_f} \left(\frac{\partial L}{\partial \varphi} - \frac{d}{dt} \frac{\partial L}{\partial \dot{\varphi}} \right) \delta \varphi dt = 0$ for all variations keeping endpoints fixed.

Exercise: Show that if $\varphi \in C_c^\infty((t_i, t_f))$ and f is integrable and $\int_{t_i}^{t_f} f \varphi = 0$ \forall such φ then $f = 0$.

$$\Rightarrow \begin{cases} \frac{\partial L}{\partial \varphi} - \frac{d}{dt} \frac{\partial L}{\partial \dot{\varphi}} = 0 \\ \varphi(t_i) = a \\ \varphi(t_f) = b \end{cases}$$

This is called the Euler-Lagrange equation.

Going back to the distance function. Note $L(x, y, \dot{x}, \dot{y}) = \sqrt{\dot{x}^2 + \dot{y}^2}$

$$\frac{\partial L}{\partial x} = 0 \quad \frac{\partial L}{\partial \dot{x}} = \dot{x}(\dot{x}^2 + \dot{y}^2)^{-1/2}$$

$$\frac{\partial L}{\partial y} = 0 \quad \frac{\partial L}{\partial \dot{y}} = \dot{y}(\dot{x}^2 + \dot{y}^2)^{-1/2}$$

$$\frac{d}{dt} \frac{\partial L}{\partial \dot{x}} = \frac{\partial L}{\partial x} \Rightarrow \ddot{x}(\dot{x}^2 + \dot{y}^2)^{-1/2} + \dot{x} \frac{d}{dt} ((\dot{x}^2 + \dot{y}^2)^{-1/2}) = 0$$

$$\frac{d}{dt} \frac{\partial L}{\partial \dot{y}} = \frac{\partial L}{\partial y} \Rightarrow \ddot{y}(\dot{x}^2 + \dot{y}^2)^{-1/2} + \dot{y} \frac{d}{dt} ((\dot{x}^2 + \dot{y}^2)^{-1/2}) = 0$$

$$\begin{aligned} x(t) &= tx_1 + (1-t)x_2 \Rightarrow \dot{x} = x_1 - x_2 \Rightarrow \ddot{x} = 0 \\ y(t) &= ty_1 + (1-t)y_2 \Rightarrow \dot{y} = y_1 - y_2 \Rightarrow \ddot{y} = 0 \end{aligned}$$

Hence this satisfies the E-L equations.

Exercise: How do we show this is a minimum?

From a mathematical standpoint $L: TM \rightarrow \mathbb{R}$. we call M the configuration space and L the lagrangian.

In classical physics we usually take $L = K.E - P.E$. which in standard \mathbb{R}^n is: $K.E = \frac{1}{2}mv^2$ and $P.E: TM \rightarrow \mathbb{R}$

If we had a particle of mass m in free space then its configuration space is \mathbb{R}^n with a Riemannian metric $g_{ij} = m\delta_{ij}$

$$K.E = \frac{1}{2}g_{ij}v^i v^j$$

If we have a free particle then $L = K.E = \frac{1}{2}g_{ij}\dot{x}^i \dot{x}^j = \frac{1}{2}m\delta_{ij}\dot{x}^i \dot{x}^j$

$$\frac{\partial L}{\partial x^k} = 0$$

$$\frac{\partial L}{\partial \dot{x}^k} = \frac{1}{2}m\delta_{ij} \frac{\partial}{\partial \dot{x}^k} (\dot{x}^i \dot{x}^j) = \frac{1}{2}m\delta_{ij} (2\delta^i_k \dot{x}^j) = m\delta_{kj} \dot{x}^j$$

hence $m\delta_{kj} \dot{x}^j = 0 \Rightarrow \ddot{x}^k = 0$ hence the particle moves at constant velocity. we get that $m\dot{x}^k$ is conserved along the motion.

Exercise: let $M = S^1(r)$ and $g = r^2 d\theta^2$ what does the coordinates here mean? what does θ mean? What is the kinetic energy? Find Equations of motion.

Hint: $x = r \cos \theta$
 $y = r \sin \theta$

on a general Riemannian manifold we have a nice Lagrangian which in local coordinates is given by $\frac{1}{2} g_{ij} \dot{x}^i \dot{x}^j$

on each tangent space $g_p: T_p M \times T_p M \rightarrow \mathbb{R}$ is non-degenerate

this means the induced map $\hat{g}_p: T_p M \rightarrow T_p^* M$ is an isomorphism.

$$v \mapsto g_p(v, \cdot) = \tau_v g_p$$

$$g = g_{ij} dx^i \otimes dx^j \text{ hence if } v = v^k \partial_k \text{ then } \hat{g}(v, \cdot) = g_{ij} dx^i(v^k \partial_k) dx^j = g_{ij} v^i dx^j$$

$$\text{we get an inverse } \hat{g}_p^{-1}: T_p^* M \rightarrow T_p M \text{ s.t. } \hat{g}_p^{-1} = g^{ij} \frac{\partial}{\partial x^i} \otimes \frac{\partial}{\partial x^j}$$

$$\text{and } g^{ij} g_{jk} = \delta^i_k$$

$$L = \frac{1}{2} g_{ij} \dot{x}^i \dot{x}^j \text{ what are the E-L equations?}$$

$$\frac{\partial L}{\partial x^k} = \frac{1}{2} (\partial_k g_{ij}) \dot{x}^i \dot{x}^j$$

$$\frac{\partial L}{\partial \dot{x}^k} = \frac{1}{2} g_{ij} \partial_k (\dot{x}^i \dot{x}^j) = \frac{1}{2} g_{ij} \delta^i_k \dot{x}^j = g_{kj} \dot{x}^j = \dot{x}_k$$

$$\frac{d}{dt} \frac{\partial L}{\partial \dot{x}^k} = \ddot{x}^j g_{kj} + \dot{x}^j (\partial_i g_{kj}) \dot{x}^i$$

$$\text{Hence } \ddot{x}^j g_{kj} + (\partial_i g_{kj}) \dot{x}^i \dot{x}^j = \frac{1}{2} (\partial_k g_{ij}) \dot{x}^i \dot{x}^j$$

$$\Rightarrow \ddot{x}^j g_{kj} + \frac{1}{2} (\partial_i g_{kj} + \partial_j g_{ki} - \partial_k g_{ij}) \dot{x}^i \dot{x}^j = 0$$

$$\Rightarrow g^{k\alpha} \ddot{x}^j g_{kj} + \frac{1}{2} g^{k\alpha} (\partial_i g_{kj} + \partial_j g_{ki} - \partial_k g_{ij}) \dot{x}^i \dot{x}^j = 0$$

$$\Rightarrow \delta^\alpha_j \ddot{x}^j + \underbrace{\Gamma^\alpha_{ij}}_{\Gamma^\alpha_{ij}} \dot{x}^i \dot{x}^j = 0$$

$$\Rightarrow \ddot{x}^\alpha + \Gamma^\alpha_{ij} \dot{x}^i \dot{x}^j = 0$$

The n^3 functions Γ^α_{ij} are Christoffel symbols.