


October 16, 2025: We have some measure space (X, Σ, μ) we looked at now to integrate functions.

$(\mathbb{R}^n, \lambda^n) \leftarrow$ we mainly worry about this.

$$\int_{[0,1]^2} 1 d^2\lambda = 1 \cdot \lambda^2([0,1]^2) = 1$$

Say we wanted to integrate $f(x,y) = x^2 + y^3$ over $[0,1]^2$



we now look at how to decompose an integral in "higher dimension" into "lower dimensional" integrals.

$$\int_{[0,1]^2} f d^2\lambda = \int_{[0,1]} \left(\int_{[0,1]} f(x,y) d\lambda(y) \right) d\lambda(x) ?$$


\uparrow integrate wrt y . \nwarrow integrate wrt x .

Idea: Start with measurable spaces (X, Σ_X) and (Y, Σ_Y) and try to define a reasonable σ -algebra on $X \times Y$. Then define a reasonable measure on $X \times Y$ if X, Y are measure spaces.


"Reasonable" here means we hope that the theory allows the Lebesgue measure on \mathbb{R}^n to decompose into the Lebesgue measure on \mathbb{R}^k and \mathbb{R}^{n-k} : $\mathbb{R}^n \cong \mathbb{R}^k \times \mathbb{R}^{n-k}$

$\mathbb{R}^n \cong \underbrace{\mathbb{R} \times \mathbb{R} \times \dots \times \mathbb{R}}_{n \text{ times}}$

let us take (X, Σ_X) and (Y, Σ_Y) then note we define the σ -algebra $\Sigma_X \otimes \Sigma_Y$ to be the smallest σ -algebra generated by $\{A \times B \mid A \in \Sigma_X \text{ and } B \in \Sigma_Y\}$



we will call $A \times B$ a rectangle (geometrically this might always be a rectangle)



let $E \subseteq X \times Y$ then the cross-section E over $x \in X$ is defined as $[E]_x = \{(x,y) \in E \mid y \in Y\}$ is defined as $[E]_y = \{(x,y) \in E \mid x \in X\}$

let $f: X \times Y \rightarrow A$ then the cross-section f over $x \in X$ is defined as $[f]_x(y) = f(x,y)$ $y \in Y$ is defined as $[f]_y(x) = f(x,y)$

If $f: X \times Y \rightarrow \mathbb{R}$ is measurable we have $[f]_x$ is measurable in Y
 $[f]_y$ is measurable in X

(σ -finite)
 let (X, Σ_X, μ) and (Y, Σ_Y, ν) be measure spaces. How can we define a measure on $(X \times Y, \Sigma_X \otimes \Sigma_Y)$?

If $f: X \rightarrow \overline{\mathbb{R}}$ then we write $\int_X g \, d\mu = \int_X g(x) \, d\mu(x)$

Now note that if $E \in \Sigma_X$ then $\int_X \chi_E \, d\mu = \mu(E)$. Then suppose $E \in \Sigma_X \otimes \Sigma_Y$

then note first that $\forall x \in X \begin{cases} [E]_x \text{ is measurable} \\ \forall y \in Y [E]_x^y \text{ is measurable} \end{cases} \Leftrightarrow \begin{cases} [\chi_E]_x \text{ is also measurable} \\ \{\chi_E\}_x^y \text{ is also measurable} \end{cases}$

using this define $(\mu \times \nu)(E) = \int_X \left(\int_Y \underbrace{\chi_E(x, y)}_{[\chi_E]_x(y)} \, d\nu(y) \right) d\mu(x)$ another way to look at
 $[\chi_E]_x: Y \rightarrow \overline{\mathbb{R}}$

this is that define $g(x) = \int_Y [\chi_E]_x(y) \, d\nu(y)$ then $(\mu \times \nu)(E) = \int_X g(x) \, d\mu(x)$

So then we can define the iterated integral of $f: X \times Y \rightarrow \overline{\mathbb{R}}$ as $\int_X \int_Y f(x, y) \, d\nu(y) \, d\mu(x)$
 $= \int_X \left(\int_Y [f]_x(y) \, d\nu(y) \right) d\mu(x)$

We could have also defined $(\mu \times \nu)(E) = \int_Y \int_X \chi_E \, d\mu(x) \, d\nu(y)$ but we will see that it will not matter.

$$\text{Example: } \int_{[0,1]} \int_{[0,1]} x^2 + y^3 \, d\lambda(y) \, d\lambda(x) = \int_{[0,1]} \left(\int_{[0,1]} x^2 + y^3 \, d\lambda(y) \right) d\lambda(x) = \int_{[0,1]} x^2 + \frac{1}{4} \, d\lambda(x) = \frac{1}{3} + \frac{1}{4} = \frac{7}{12}$$

$$\int_{[0,1]} \left(\int_{[0,1]} x^2 + y^3 \, d\lambda(x) \right) d\lambda(y) = \int_{[0,1]} \frac{1}{3} + y^3 \, d\lambda(y) = \frac{1}{3} + \frac{1}{4} = \frac{7}{12}$$

$$\text{We would now hope that } \int_{[0,1]^2} f \, d^2\lambda = \frac{7}{12}$$

So three main questions: (i) Is $\mu \times \nu$ even a measure ?? - Yes it is!

(ii) Is the Lebesgue measure on \mathbb{R}^n a product measure of the Lebesgue measure on \mathbb{R}^k and \mathbb{R}^{n-k} ? Yes it is a product of the Lebesgue measures.

(iii) Is the integral over the product measure equal to the iterated integrals?

(σ-finite)

Theorem 1 (Tonelli's Theorem): Suppose (X, Σ_X, μ) and (Y, Σ_Y, ν) are measure spaces. Suppose also that $f: X \times Y \rightarrow \mathbb{R}$ is non-negative and is also $(\mu \times \nu)$ -measurable.

Then consider $g(x) = \int_Y f(x, y) d\nu(y)$ and $h(y) = \int_X f(x, y) d\mu(x)$, g and h are both measurable (in their respective spaces) and

$$\int_{X \times Y} f d(\mu \times \nu) = \int_X \left(\int_Y f(x, y) d\nu(y) \right) d\mu(x) = \int_Y \left(\int_X f(x, y) d\mu(x) \right) d\nu(y)$$

Example: $\int_{[0,1]^2} xy d^2\lambda = \int_{[0,1]} \left(\int_{[0,1]} xy d\lambda(x) \right) d\lambda(y) = \int_{[0,1]} \frac{y}{2} d\lambda(y) = \frac{y^2}{2} \Big|_0^1 = \frac{1}{2}$

(σ-finite)

Theorem 2 (Fubini's Theorem): Suppose (X, Σ_X, μ) and (Y, Σ_Y, ν) are measure spaces. Suppose $f: X \times Y \rightarrow \mathbb{R}$ is $(\mu \times \nu)$ -measurable with $\int_{X \times Y} |f| d(\mu \times \nu) < \infty$ then

- (i) $\int_Y f(x, y) d\nu(y) < \infty$ for almost every $x \in X$
- (ii) $\int_X f(x, y) d\mu(x) < \infty$ for almost every $y \in Y$
- $$\int_{X \times Y} f^+ d(\mu \times \nu) + \int_{X \times Y} f^- d(\mu \times \nu)$$

Then consider $g(x) = \int_Y f(x, y) d\nu(y)$ and $h(y) = \int_X f(x, y) d\mu(x)$, g and h are both measurable (in their respective spaces) and

$$\int_{X \times Y} f d(\mu \times \nu) = \int_X \left(\int_Y f(x, y) d\nu(y) \right) d\mu(x) = \int_Y \left(\int_X f(x, y) d\mu(x) \right) d\nu(y)$$

Example: $f: [0,1] \times [0,1] \rightarrow \mathbb{R}$ by $f(x, y) = \frac{x^2 - y^2}{(x^2 + y^2)^2}$ then $\int_{[0,1]} \int_{[0,1]} f(x, y) d\lambda(y) d\lambda(x) = \frac{\pi}{4}$

$$\int_{[0,1]} \int_{[0,1]} f(x, y) d\lambda(x) d\lambda(y) = -\frac{\pi}{4}$$

$$f^+ = \begin{cases} \frac{x^2 - y^2}{(x^2 + y^2)^2} & ; x > y \\ 0 & ; \text{else} \end{cases} \quad f^- = \begin{cases} \frac{y^2 - x^2}{(x^2 + y^2)^2} & ; y > x \\ 0 & ; \text{else} \end{cases}$$

$$\int_{[0,1]^2} f^+ d\lambda \text{ and } \int_{[0,1]^2} f^- d\lambda \text{ diverges}$$