


Multiple Regression Analysis




This residual plot is a result of having categorical response variable.




Normal QQ Plot of Residuals



The distribution of residuals appear to be bimodal. The QQ plot also suggests the same conclusion, since we can see a very prominent deviation from the diagonal normal line on the left side. Overall, the Normaltiy Assumption seems to be violated. This could be due to the fact that we are dealing with a categorical response variable quality.



We can see that there is one point with a significantly high leverage around 0.4, comparing with other observations. This could be an outlier and might be an influence point.




The standardized residuals show some points with magnitude greater than 2, but overall, in combination with our observation from Cook's Distance and just one data point with high leverage away from other points, we can conclude there isn't any obvious influential points in this model.


kable(tidy(vif(final.model)))

- ## Warning: 'tidy.numeric' is deprecated.
- ## See help("Deprecated")

names	X
residual.sugarCent	1.402748
citric.acid	1.425081
free.sulfur.dioxideCent	3.169724
sulphatesCent	3.922514
densityCent	2.274231
alcoholCent	1.825881
citric.acid:free.sulfur.dioxideCent	3.249458
citric.acid:sulphatesCent	3.832337
free. sulfur. dioxide Cent: alcohol Cent	1.097922
sulphatesCent:alcoholCent	1.187895


Logistic Regression





Most of the binned residual plots show random pattern and raises no major concerns of violations. The binned residual plot of alcoholCent and free.sulfur.dioxideCent seems to show an outlier on the top left, suggesting high average residuals for that bin. There is also one outlier on the binned residual plot of chloridesCent on the far right.

The binned residual plot also reveals that sulphatesCent might need a transformation, since there appears to be some linear trend.




multicollinearity kable(tidy(vif(modellg1.final)))

Warning: 'tidy.numeric' is deprecated.

See help("Deprecated")


names	X
alcoholCent	1.050002
volatile.acidityCent	1.542159
sulphatesCent	1.416972
chloridesCent	1.433665
free.sulfur.dioxideCent	1.018900
citric.acid	1.650265




Area under the curve: 0.8092


From the ROC curve and AUC calculation, we can see the curve is fairly close to the top left corner (area under the curve is close to 1). This shows that the logistic model is able to distinguish between good and not good quality, so this is a pretty good model.

Ordinal Regression Analysis






There is also some linear trends in the binned residual plots for pHCent. We should also tranform this variable.



The binned residual plot shows some potential linear trend and outliers in three categories of quality. We might need to transform sulphates Cent



```
##
## Re-fitting to get Hessian
## z test of coefficients:
##
##
                       Estimate Std. Error z value Pr(>|z|)
## volatile.acidityCent -3.59868 0.37775 -9.5267 < 2.2e-16 ***
                                   0.37520 -1.9717 0.0486457 *
## citric.acid
                       -0.73977
## chloridesCent
                       -5.25929
                                   1.30690 -4.0242 5.716e-05 ***
## sulphatesCent
                        2.73449
                                   0.36207
                                             7.5524 4.274e-14 ***
## pHCent
                       -1.47755
                                   0.41452 -3.5645 0.0003646 ***
## alcoholCent
                        0.94934
                                   0.05799 16.3708 < 2.2e-16 ***
## 3|4
                       -6.08448
                                   0.34309 -17.7344 < 2.2e-16 ***
## 4|5
                                   0.17838 -23.2280 < 2.2e-16 ***
                       -4.14345
## 5|6
                       -0.49521
                                   0.11605 -4.2674 1.978e-05 ***
## 6|7
                        2.31842
                                   0.13664 16.9678 < 2.2e-16 ***
## 7 8
                        5.33090
                                   0.27003 19.7419 < 2.2e-16 ***
## ---
## Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

p-values of all the variables are extremely small, so they are all significant predictors of the log-odds of the wine falling in or below quality j. We should also not extrapolate beyond quality $j=3,4,\ldots,7$.