

# CLASSIFICATION OF FINITE GROUPS

DANIEL LAING

B.SC (HONS) MATHEMATICS

SUPERVISED BY DR. MARTYN QUICK

UNIVERSITY OF ST ANDREWS

2023

*I certify that this project report has been written by me, is a record of work carried out by me, and is essentially different from work undertaken for any other purpose or assessment.*

# Contents

<b>I</b>	<b>Introduction</b>	<b>3</b>
1.1	Abstract . . . . .	3
1.2	Introduction . . . . .	3
1.3	Automorphisms . . . . .	4
1.4	Semidirect Product . . . . .	5
1.5	Group Actions . . . . .	8
<b>II</b>	<b>Prime Power Orders</b>	<b>10</b>
2.1	Prime Order . . . . .	11
2.2	Groups of Order $p^2$ . . . . .	11
2.3	Groups of Order $p^3$ . . . . .	12
<b>III</b>	<b>Composite Orders</b>	<b>15</b>
3.1	Groups of Order $pq$ . . . . .	15
3.1.1	Groups of Order $2p$ . . . . .	16
3.2	Order $4q$ . . . . .	16
3.3	Order $2p^2$ . . . . .	18
<b>IV</b>	<b>Special Cases</b>	<b>19</b>
4.1	Groups of order 12 . . . . .	19
4.2	Groups of Order 24 . . . . .	21
4.3	Groups of Order 30 . . . . .	27
4.4	Groups of Order 16 . . . . .	28
	<b>Bibliography</b>	<b>34</b>
	<b>Appendix</b>	<b>35</b>

# Chapter I

## Introduction

### 1.1 Abstract

This report covers the full classification, including proofs, of all groups of order less than or equal to 31. In some cases, a classification of the general case is given; namely  $p$ -groups up to  $p^3$ , and groups of order  $pq$ ,  $4p$  and  $2p^2$ , where  $p$  and  $q$  are distinct primes. Where possible, the names of specific, and families of groups are given.

A table of the full classification is found in the appendix. Note: not all presentations are included, specifically long ones, and those not especially relevant.

### 1.2 Introduction

The study of groups is an important area of mathematics, and as such, it's quite useful to have quick examples of groups 'in your back pocket' so to speak; even more so, to have an exhaustive list of the possible behaviours of groups. We do this by the notion of an isomorphism class, a kind of equivalence between groups which are essentially the same. However as groups increase in size, this becomes increasingly hard to do, especially for so called ' $p$ -groups': groups which have order the power of a prime. For example, up to isomorphism, there are 51 possible groups of order 32 — wow!

This report will cover the classification, and proof thereof, of groups of order up to, and including, 31. In doing so, we will further our understanding of constructing groups, and where possible, find out the names given to the groups we come across by the wider world of group theory.

To start, let's solidify the notation used in this report. We shall denote groups and sets with capital letters, like  $G$ ,  $H$ , and elements of those groups with lower case letters, like  $g$ ,  $h$ . Greek letters shall denote mappings, generally  $\phi$ ,  $\psi$ , etc. with  $\iota$  reserved for the identity map, and we will write mappings on the right.

To denote the cyclic group of order  $n$  we will use  $C_n$ ,  $D_{2n}$  to denote the dihedral group of order  $2n$ ,  $A_n$  to denote the alternating group over  $n$  elements,  $S_n$  to denote the symmetric group over  $n$  elements, and  $Q_8$  to denote the quaternion group. The trivial group,  $\{1\}$  is denoted by  $\mathbf{1}$ . We will meet other groups as we go on our journey of classification!

## 1.3 Automorphisms

Let's move on to review some facts and theorems which will be valuable later on, as well as introduce some new concepts and prove some new results!

**Definition 1.** If  $G$  and  $H$  are groups with elements  $g_1, g_2 \in G$ , then a map:

$$\phi : G \rightarrow H$$

is a homomorphism if:

$$(g_1 g_2) \phi = (g_1 \phi)(g_2 \phi)$$

If  $\phi$  is bijective, then we call it an isomorphism, with  $G \cong H$  denoting that  $G$  is isomorphic to  $H$ . And if  $\phi$  is an isomorphism from  $G$  to itself, then we call it an automorphism of  $G$ .

**Lemma 1.** *The set of all automorphisms of a group  $G$  form a group under composition. Indeed, this is called the automorphism group of  $G$ , denoted  $\text{Aut } G$ .*

*Proof.* Let  $A = \text{Aut } G = \{ \phi : G \rightarrow G \mid \phi \text{ is an isomorphism} \}$ , and let  $\phi \in A$ . Denote an element of  $G$  by  $g$ .

We know already that the composition of two isomorphisms is an isomorphism, so  $A$  is closed under composition.

The identity map,  $\iota : g \mapsto g$ , is certainly an automorphism of  $G$  and so  $A$  is non-empty.

Indeed,  $\iota : g \mapsto g$  is the identity of  $A$ , since:

$$g\phi\iota = (g\phi)\iota = g\phi \quad \text{and} \quad g\iota\phi = (g\iota)\phi = g\phi$$

And inverses clearly exist, because automorphisms are bijections, and bijections are invertible. Hence  $A = \text{Aut } G$  is a group.  $\square$

**Lemma 2.** *The automorphism group of  $C_n$  is isomorphic to the multiplicative group of integers mod  $n$ .*

*i.e.*  $\text{Aut } C_n \cong (\mathbb{Z}/n\mathbb{Z})^\times$

*Proof.* Let  $C_n = \langle x \rangle$ . Any automorphism,  $\phi$  of  $C_n$  has the property:

$$(x^i)\phi = (x\phi)^i$$

Hence  $\phi$  is determined by its effect on a generator,  $x$ , and preserves element order. In particular,  $\phi$  sends generators to generators. So for  $\phi$  to be an automorphism, it must send  $x$  to another generator, say  $x^k$ . An element  $x^k$  generates  $C_n$  if  $x^k$  has order  $n$ , i.e. when  $k$  and  $n$  are co-prime. Denote the automorphism sending  $x$  to  $x^k$  by  $\phi_k$ .

Let's now investigate how these automorphisms behave. Let  $\phi_k, \phi_l \in \text{Aut } C_n$ , and consider:

$$x\phi_k\phi_l = (x^k)\phi_l = (x^k)^l = x^{(kl)} = x\phi_{kl} \pmod{n}$$

Because multiplication modulo  $n$  is commutative,  $x^{kl} = x^{lk}$ , so  $\text{Aut } C_n$  is abelian.

Now consider  $\theta : \text{Aut } C_n \rightarrow (\mathbb{Z}/n\mathbb{Z})^\times$  defined by  $\phi_k\theta = k$ . We will show  $\theta$  is an isomorphism. Every  $k \in (\mathbb{Z}/n\mathbb{Z})^\times$  is co-prime to  $n$  and so  $x^k$  is a generator of  $C_n$ , hence there is some  $\phi_k \in \text{Aut } C_n$  such that  $\phi_k\theta = k$ . So  $\theta$  is surjective. If  $\phi_k\theta = \phi_l\theta$  then  $k = l$ , so  $\theta$  is also injective. Finally,  $\theta$  is a homomorphism because:

$$(\phi_k\phi_l)\theta = \phi_{kl}\theta = kl = (\phi_k\theta)(\phi_l\theta)$$

So  $\theta : \text{Aut } C_n \rightarrow (\mathbb{Z}/n\mathbb{Z})^\times$  is an isomorphism. □

**Definition 2.** A subgroup  $H$  of a group  $G$  is called characteristic if it is fixed by all automorphisms of  $G$ .

i.e. for an automorphism  $\phi$  of  $G$ ,  $H\phi = H$ .

**Lemma 3.** Let  $G$  be a group with normal subgroup  $H$ , and let  $K$  be characteristic in  $H$ . Then  $K$  is a normal subgroup of  $G$ .

*Proof.* Consider the map  $\phi_g : G \rightarrow G$  defined by  $\phi_g : x \mapsto g^{-1}xg$  for elements  $x, g \in G$ . We will show that this is an automorphism of  $G$ . For  $x, y \in G$ :

$$x\phi_g y\phi_g = (g^{-1}xg)(g^{-1}yg) = g^{-1}(xy)g = (xy)\phi_g$$

Hence  $\phi_g$  is a homomorphism. Moreover,  $\phi_g$  is invertible with inverse  $\phi_{g^{-1}}$ . So  $\phi_g$  is indeed an automorphism of  $G$ .

Because  $H$  is normal,  $H\phi_g = H$ . So  $\phi_g$  is an automorphism of  $H$  too. And so  $\phi_g$  maps  $K$  to itself, because it is characteristic. Hence:

$$\{g^{-1}kg \mid k \in K\} = K$$

So  $K$  is normal in  $G$ . □

## 1.4 Semidirect Product

We already know about the direct product:

**Definition 3** (Direct Product). For groups  $N$  and  $H$ , the direct product,  $G = N \times H$  is a group of ordered pairs of elements  $(n, h)$  where  $n \in N$  and  $h \in H$  with the operation:

$$(n_1, h_1)(n_2, h_2) = (n_1n_2, h_1h_2)$$

Moreover, if  $\bar{N} = N \times \mathbf{1}$  and  $\bar{H} = \mathbf{1} \times H$ , then:

- (i)  $\bar{N} \trianglelefteq G$  and  $\bar{H} \trianglelefteq G$
- (ii)  $\bar{N} \cap \bar{H} = \mathbf{1}$
- (iii)  $\bar{N}\bar{H} = \{nh \mid n \in N, h \in H\} = G$

Now let's seek a slightly more general way to combine groups, by relaxing that  $H$  must be normal. So we have:

$$N \trianglelefteq G, H \leq G, NH = G, \quad \text{and} \quad N \cap H = \mathbf{1}$$

Consider the set, (not the direct product):

$$N \times H = \{ (n, h) \mid n \in N, h \in H \}$$

and a map

$$\phi : N \times H \rightarrow G \quad \text{defined by} \quad (n, h) \mapsto nh$$

We want  $\phi$  to be an isomorphism.

To show  $\phi$  is injective, take  $n_1, n_2 \in N$  and  $h_1, h_2 \in H$ , and assume  $n_1 h_1 = n_2 h_2$ . Then multiplying on the left by  $n_2^{-1}$  and on the right by  $h_1^{-1}$  gives:

$$n_2^{-1} n_1 = h_2 h_1^{-1}$$

On the left we have an element of  $N$  and on the right, an element of  $H$ , so  $n_2^{-1} n_1 = h_2 h_1^{-1} \in N \cap H$ . But  $N \cap H = \mathbf{1}$  so then  $n_2^{-1} n_1 = h_2 h_1^{-1} = 1$ . Hence:

$$n_1 = n_2 \quad \text{and} \quad h_1 = h_2$$

To show  $\phi$  is surjective, consider the image,  $\text{im } \phi = \{ nh \mid n \in N, h \in H \}$ . This is by definition  $NH = G$ , so  $\phi$  is surjective, and hence a bijection.

For  $\phi$  to be a homomorphism, we need:

$$\begin{aligned} [(n_1, h_1)(n_2, h_2)]\phi &= (n_1, h_1)\phi (n_2, h_2)\phi \\ &= n_1 h_1 n_2 h_2 \\ &= n_1 h_1 n_2 h_1^{-1} h_1 h_2 \\ &= (n_1 h_1 n_2 h_1^{-1})(h_1 h_2) \end{aligned}$$

But  $N$  is normal in  $G$  so  $h_1 n_2 h_1^{-1}$  is just another element in  $N$ , say  $n_3$ . So:

$$[(n_1, h_1)(n_2, h_2)]\phi = (n_1 n_3)(h_1 h_2) = (n_1 n_3, h_1 h_2)\phi$$

We know that  $\phi$  is injective, so then:

$$(n_1, h_1)(n_2, h_2) = (n_1 n_3, h_1 h_2)$$

This tells us the multiplication that will make  $NH$  a group. Because  $N \trianglelefteq G$ , the map

$$n_2 \mapsto h_1 n_2 h_1^{-1} = n_3$$

is an automorphism of  $N$ . This gives rise to the definition:

**Definition 4** (Semidirect Product).

- (i) For a group  $G$  with normal subgroup  $N$  and subgroup  $H$  with  $NH = G$  and  $N \cap H = \mathbf{1}$ ,  $G$  is the internal semidirect product of  $N$  by  $H$ , written  $G = N \rtimes H$ .
- (ii) For groups  $N$  and  $H$ , and a homomorphism  $\psi : H \rightarrow \text{Aut } N$ , the external semidirect product of  $N$  by  $H$  via  $\psi$  is the set:

$$N \times H = \{ (n, h) \mid n \in N, h \in H \}$$

with multiplication:

$$(n_1, h_1)(n_2, h_2) = (n_1(n_2^{h_1\psi}), h_1h_2)$$

denoted:

$$N \rtimes_{\psi} H$$

We use the notation  $n_2^{h_1\psi}$  to mean the image of  $n_2$  under the automorphism  $h_1\psi$ , both because it indicates conjugation, and is clearer.

**Lemma 4.** For a group  $G$  with  $N \leq G$  and  $H \leq G$ , with  $N \cap H = \mathbf{1}$  then:

$$|NH| = |\{nh \mid n \in N, h \in H\}| = |N| \cdot |H|$$

*Proof.* We just saw above that for elements  $n \in N$  and  $h \in H$ , the map:

$$\phi : N \times H \rightarrow NH \quad \text{defined by} \quad (n, h) \mapsto nh$$

is a bijection. The result follows immediately from this. □

**Lemma 5.** Let  $N$  and  $H$  be groups, and  $\alpha \in \text{Aut } N$ . Then the semidirect products via the homomorphism  $\phi$ ,  $N \rtimes_{\phi} H$ , and via the homomorphism  $\psi$ ,  $N \rtimes_{\psi} H$ , are isomorphic if for  $\alpha \in \text{Aut } N$  and  $\beta \in \text{Aut } H$ , we have:

$$h^{\beta}\psi = \alpha^{-1}h\phi\alpha \quad (\text{for all } h \in H)$$

That is, we can apply any automorphism to  $H$  and conjugate  $\text{Aut } N$ , and the resulting semidirect product remains in the same isomorphism class.

*Proof.* Let  $G = N \rtimes_{\phi} H$  and  $\bar{G} = N \rtimes_{\psi} H$ , and define:

$$\theta : G \rightarrow \bar{G} \quad \text{by} \quad \theta : (n, h) \mapsto (n^{\alpha}, h^{\beta})$$

We will show that  $\theta$  is an isomorphism.

First,  $\theta^{-1}$  exists because both  $\alpha^{-1}$  and  $\beta^{-1}$  exist, and is given by:

$$\theta^{-1} : (n, h) \mapsto (n^{\alpha^{-1}}, h^{\beta^{-1}})$$

Hence  $\theta$  is a bijection. We also have that:

$$h^\beta \psi = \alpha^{-1} h \phi \alpha$$

which implies:

$$\alpha h^\beta \psi = h \phi \alpha \quad (*)$$

Now for two elements,  $(n_1, h_1), (n_2, h_2) \in G$ , consider:

$$\begin{aligned} (n_1, h_1)\theta (n_2, h_2)\theta &= (n_1^\alpha, h_1^\beta)(n_2^\alpha, h_2^\beta) \\ &= (n_1^\alpha(n_2^\alpha)^{h_1^\beta \psi}, h_1^\beta h_2^\beta) \\ &= (n_1^\alpha(n_2^{h_1 \phi})^\alpha, h_1^\beta h_2^\beta) && \text{(by *)} \\ &= ((n_1(n_2)^{h_1 \phi})^\alpha, (h_1 h_2)^\beta) \\ &= (n_1(n_2)^{h_1 \phi}, h_1 h_2)\theta \\ &= (n_1, h_1)(n_2, h_2)\theta \end{aligned}$$

So  $\theta$  is an isomorphism. □

## 1.5 Group Actions

Another useful piece of group theory technology will be group actions.

**Definition 5.** Let  $G$  be a group, and  $\Omega$  be a set, with elements  $g \in G$  and  $\omega \in \Omega$ . Consider a map  $\mu : \Omega \times G \rightarrow \Omega$ , and write  $\omega^g$  for the image of  $(\omega, g)$  under  $\mu$ . So we have:

$$\mu : \Omega \times G \rightarrow \Omega \quad \text{defined by} \quad (\omega, g) \mapsto \omega^g$$

We say  $G$  acts on  $\Omega$  if for all  $g_1, g_2 \in G$  and all  $\omega \in \Omega$ :

- (i)  $(\omega^{g_1})^{g_2} = \omega^{(g_1 g_2)}$
- (ii)  $\omega^1 = \omega$

We call  $\mu$  the group action of  $G$  on  $\Omega$ .

This might remind you of a homomorphism. Indeed we have a result:

**Lemma 6.** *A group action induces a homomorphism. Specifically, let  $G$  be a group which acts on a set  $\Omega$ , with  $g \in G$  and  $\omega \in \Omega$ , and define:*

$$\rho_g : \Omega \rightarrow \Omega \quad \text{by} \quad \omega \mapsto \omega^g$$

*Then:*

$$\rho : G \rightarrow \text{Sym } \Omega \quad \text{defined by} \quad g \mapsto \rho_g$$

*is a homomorphism.*

*Proof.* Firstly,  $\rho_g$  is indeed a permutation of  $\Omega$  because it is invertible (and therefore a bijection), with:

$$(\rho_g)^{-1} = \rho_{g^{-1}}$$



Consider  $g, h \in G$  and their corresponding maps,  $\rho_g, \rho_h \in \text{Sym } \Omega$ . Then:

$$\omega(g\rho)(h\rho) = \omega\rho_g\rho_h = (\omega^g)^h = \omega^{(gh)} = \omega\rho_{gh} = \omega(gh)\rho$$

Thus  $\rho$  is a homomorphism. □

By this we see that in a semidirect product,  $N \rtimes H$ , the group  $H$  acts on  $N$ !

**Definition 6.** The orbit of an element  $\omega \in \Omega$ , is the set:

$$\omega^G = \{ \omega^x \mid x \in G \}$$

A group acting on the set its cosets will be useful:

**Definition 7.** For a group  $G$  with  $H \leq G$ , let  $\Omega = \{ Hg \mid g \in G \}$ , i.e. the set of cosets of  $H$  in  $G$ . If  $x \in G$ , define a group action:

$$\Omega \times G \rightarrow \Omega \quad \text{by} \quad (Hg, x) \mapsto Hgx$$

**Lemma 7.** *The action above is well defined, meaning the action is independent of our choice of representative  $g$ .*

*Proof.* Omitted. □

# Chapter II

## Prime Power Orders

First, we will prove a few useful lemmas:

**Lemma 8.** *If  $G$  is a  $p$ -group (i.e. a group of prime power order), then every subgroup of index  $p$  is normal.*

*Proof.* Let  $H$  be a subgroup of  $G$ , with index  $p$ . We know kernels are normal subgroups, so we will show that  $H$  is the kernel of some homomorphism. Let  $\Omega$  be the set of all cosets of  $H$ . So by definition,  $|\Omega| = p$ . By Lemma 6, there is a homomorphism:

$$\rho : G \rightarrow S_p$$

Let's investigate the kernel of  $\rho$ . If we have  $x \in \ker \rho$ , then:

$$(H1)x = H1 = H$$

which means  $x \in H$ . So the kernel of  $\rho$  is  $H$ . Hence,  $H \trianglelefteq G$ . □

**Lemma 9.** *If  $G$  is a group of prime power order, the centre of  $G$  is non-trivial. In particular,  $p$  divides the order of the centre.*

*Proof.* Let  $Z$  denote the centre of  $G$ , and consider the action of  $G$  on itself by conjugation. The orbit of an element,  $g \in G$  is:

$$g^G = \{ x^{-1}gx \mid x \in G \}$$

which is the conjugacy class of  $g$ . So the size of each orbit divides some power of  $p$ . In particular, the size of each orbit is divisible by  $p$ . So then the sum of the sizes of all of the conjugacy classes is also divisible by  $p$ . Looking at the class equation:

$$|G| = |Z| + \sum_{i=1}^k |g_i^G|$$

then reducing mod  $p$  gives:

$$|G| \equiv |Z| \pmod{p}$$

Because  $G$  is non-trivial, it follows that  $|Z| \neq 1$ . □

**Lemma 10.** *For a group  $G$  with centre  $Z(G)$ . Then if  $G/Z(G)$  is cyclic,  $G$  is abelian.*

*Proof.* Let  $x \in G$  be the element such that  $xZ(G)$  generates  $G/Z(G)$ . Because  $G$  is the union of cosets of  $Z(G)$ , then indeed  $\langle x, Z(G) \rangle = G$ . The centraliser of  $x$  certainly contains  $x$ , and every element of  $Z(G)$  also commutes with  $x$ . Hence the centre of  $G$  is a subgroup of the centraliser of  $x$ . The result follows by concluding:

$$G = \langle x, Z(G) \rangle = \langle Z(G) \rangle = Z(G)$$

□

## 2.1 Prime Order

Let's start with the easiest case: groups of order 1. Any group  $G$  must have an identity element, and so that's all our possible elements used up! All groups of order 1 are isomorphic to the trivial group,  $\mathbf{1}$ .

What about groups of prime order? Let  $G$  be a group of order  $p$ , where  $p$  is a prime number. Then Lagrange's Theorem tell us all elements must have order 1 or  $p$ . Pick some  $x \in G$  with  $x$  having order  $p$ . Then  $\langle x \rangle = G$  so  $G$  is cyclic.

Any group of order  $p$  is isomorphic to  $C_p$ .

## 2.2 Groups of Order $p^2$

Let  $G$  be a group of order  $p^2$ . By Lagrange's Theorem, the elements of  $G$  have order 1,  $p$  or  $p^2$ . If  $x \in G$  has order  $p^2$ , then  $x$  generates  $G$  so  $G \cong C_{p^2}$ .

If  $G$  does not have an element of order  $p^2$  then all elements, except the identity, have order  $p$ . We know that  $G$  must have a subgroup of order  $p$ ,  $P$ , and because  $p$  is prime,  $P \cong C_p$ . Pick a generator for  $P$ , say  $x$  and an element  $y \in G$  such that  $y \notin P$ . Then  $y \neq x^i$  for any  $i$ .

If  $y^j = x^i$  for some  $i$  and  $j$ , then:

$$(y^j)^{1-j} = (x^i)^{1-j} = y^{j-j+1} = y = x^{i(1-j)} = x^k \quad \text{for some } k, \text{ a contradiction.}$$

So no power of  $y$  is equal to any power of  $x$ . Because  $y$  has order  $p$ , it generates a subgroup of order  $p$ ,  $\bar{P}$ , with  $P \cap \bar{P} = \mathbf{1}$ . Lemma 8 tells us that both  $P$  and  $\bar{P}$  are normal, and by Lemma 4,  $|P\bar{P}| = p^2 = |G|$ , so:

$$G = P \times \bar{P} \cong C_p \times C_p$$

If  $G$  has no elements of order  $p$  or  $p^2$ , then it only has elements of order 1, which is the trivial group.

Any group of order  $p^2$  is isomorphic to one of:

$$C_{p^2} \quad \text{or} \quad C_p \times C_p$$

## 2.3 Groups of Order $p^3$

This classification is based on the one found on the Groupprops subwiki<sup>1</sup>. Let  $G$  be a group of order  $p^3$ , where  $p$  is a prime number. We will first gain a handle on  $G$  by describing its centre, and quotient by it. If  $G$  is abelian, we know by the Fundamental Theorem of Finite Abelian Groups that it is isomorphic to one of:

$$C_{p^3}, \quad C_{p^2} \times C_p \quad \text{or} \quad C_p \times C_p \times C_p$$

So from now on, we will focus on the non-abelian groups.

Denote the centre of  $G$  by  $Z$  and consider its order. Lagrange's Theorem tells us  $Z$  must have order dividing  $p^3$ . It cannot be  $p^3$  because  $G$  is non-abelian, and Lemma 9 tells us that it cannot be 1. If  $|Z| = p^2$ , then  $|G/Z| = p$ , so  $G/Z \cong C_p$ . However Lemma 10 says that then  $G$  must be abelian, so then  $|Z|$  must be  $p$ . By our previous classification,  $G/Z$  is isomorphic to either  $C_{p^2}$  or  $C_p \times C_p$ . Lemma 10 tells us that it must be the latter.

This gives us a handle to start investigating the structure of  $G$ . Another useful tool will be commutators, which we will denote by  $[a, b] = a^{-1}b^{-1}ab$ . The derived subgroup of  $G$ ,  $G' = \langle [x, y] \mid x, y \in G \rangle$ , is the smallest normal subgroup such that  $G/G'$  is abelian. We saw that  $G/Z$  is abelian, so  $G' \leq Z$ , but because  $G'$  is non-trivial, we must have equality.

So far, we know  $G/Z \cong C_p \times C_p$ , and that  $G' = Z$ . Now pick two elements,  $a$  and  $b$  so that  $aZ$  and  $bZ$  generate  $G/Z$ . So then  $G = \langle Z, a, b \rangle$ .

Let  $z = [a, b]$ . If  $z = 1$  then that means  $a$  and  $b$  commute. And by definition,  $a$  commutes with  $Z$ , so  $a \in Z$ , which contradicts our choice of  $a$  as a generator of  $G/Z$ . Hence  $z \neq 1$ , and in particular,  $a$  and  $b$  do not commute. Now we know  $G' = Z$  which has order  $p$ , so  $Z \cong C_p$ . Moreover,  $z \in Z$ , and  $z \neq 1$  so we can conclude that  $\langle z \rangle = Z$ . We can see that although  $a$  and  $b$  are not in  $Z$ ,  $a^p$  and  $b^p$  are, because  $aZ$  and  $bZ$  have order  $p$  in  $G/Z$ . Considering the orders of  $a$  and  $b$  we have 3 cases:

**Case 1:** Both  $a$  and  $b$  have order  $p$ .

The above descriptions give the presentation:

$$G = \langle z, a, b \mid z^p = a^p = b^p = 1, az = za, bz = zb, [a, b] = z \rangle$$

We can write an arbitrary  $g \in G$  as  $a^i b^j z^k$  for integers  $i, j$  and  $k$  taken mod  $p$ . Hence this presentation has order at most  $p^3$ .

Now consider the set:

$$\left\{ \begin{pmatrix} 1 & \alpha & \beta \\ 0 & 1 & \gamma \\ 0 & 0 & 1 \end{pmatrix} \mid \alpha, \beta, \gamma \in \mathbb{F}_p \right\}$$

It can be shown that this is a group under the usual matrix multiplication, and is known as the unitriangular group<sup>2</sup>, denoted  $UT_3(p)$ . Taking:

$$z = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad a = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad \text{and} \quad b = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}$$

1. Groupprops, "Classification of groups of prime-cube order," February 24, 2016, accessed February 23, 2023, [https://groupprops.subwiki.org/wiki/Classification\\_of\\_groups\\_of\\_prime-cube\\_order](https://groupprops.subwiki.org/wiki/Classification_of_groups_of_prime-cube_order).

2. Groupprops, "Unitriangular matrix group:UT(3,p)," August 22, 2014, accessed February 23, 2023, [https://groupprops.subwiki.org/wiki/Unitriangular\\_matrix\\_group:UT\(3,%20p\)](https://groupprops.subwiki.org/wiki/Unitriangular_matrix_group:UT(3,%20p)).

we can see that  $\text{UT}_3(p)$  satisfies this presentation for  $p > 2$ . (Indeed, the above presentation is the standard one defining  $\text{UT}_3(p)$ ). Thus there is a single isomorphism class for this case.

The group behaves differently when  $p = 2$  because we know that a group whose elements all have order either 1 or 2 is abelian. So the elements cannot have order only 1 or 2. In particular:

$$\begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}$$

have order 4. We can check that all other non-identity elements have order 2. Thus  $\text{UT}_2(2) \cong D_8$ .

**Case 2:** One element of each order  $p$  and  $p^2$ .

The roles of  $a$  and  $b$  are interchangeable, so we can take  $a$  to have order  $p^2$  and  $b$  to have order  $p$  without loss of generality. So  $\langle a \rangle \cong C_{p^2}$ , and has index  $p$ , so by Lemma 8 is normal. We noted that  $a^p \neq 1$  is in  $Z$ , and so  $a^p$  is some power of  $z$ , say  $a^p = z^r$ , with  $r$  taken mod  $p$ .

We know that  $0 < r < p$  and I claim that we can take  $r = 1$  without loss of generality. Because  $z^r \neq 1$ , it has order  $p$ . Hence  $z^r$  generates  $Z$ . Because  $p$  is prime,  $Z \cong C_p$  is a finite field. Therefore, there exists some integer  $s$ , with  $0 < s < p$  such that  $sr \equiv 1 \pmod{p}$ . Then say  $\bar{a} = a^s$  so:

$$\bar{a}^p = (a^s)^p = (a^p)^s = (z^r)^s = z \pmod{p}$$

Indeed we can take  $\bar{a}^p$  to be any power of  $z$  we want using a similar substitution. This is actually a sneaky application of Lemma 5, because it will turn out that  $G$  can be written as a semidirect product.

Take  $a^p = z$ . Now because  $G = \langle z, a, b \rangle$ ,  $\langle a \rangle \cap \langle b \rangle = \mathbf{1}$ . So we can conclude that  $G = \langle a \rangle \rtimes \langle b \rangle$ . How does  $b$  conjugate  $a$ ? Consider:

$$a^{-1}b^{-1}ab = [a, b] = z = a^p$$

So:

$$b^{-1}ab = a^{p+1}$$

Thus:

$$G \cong C_{p^2} \rtimes C_p$$

With presentation:

$$\langle a, b \mid a^{p^2} = b^p = 1, b^{-1}ab = a^{p+1} \rangle$$

When  $p = 2$ , this reduces to the dihedral group,  $D_8$ . However, we can see that for  $p > 2$ , this is not isomorphic to the previous case, because  $\text{UT}_3(p)$  has no element of order  $p^2$ .

**Case 3:** Both  $a$  and  $b$  have order  $p^2$ .

We will show that by substitutions, this is equivalent to the above case when  $p > 2$ . In the previous case, we saw that we can take  $a^p$  and  $b^p$  to be arbitrary

powers of  $z$  using substitutions (which is Lemma 5 behind the scenes). So take  $a^p = z$  and  $b^p = z^{-1}$ . Let  $d = ab$ , and consider:

$$d^p = (ab)^p = abab \dots ab$$

We will collect together the  $a$ 's and  $b$ 's, maintaining equality with the commutator,  $[a, b] = z$ :

$$\begin{aligned} d^p &= ababab \dots ab \\ &= zaabbab \dots ab \\ &= z^2aababb \dots ab \\ &= z^3aaabbb \dots ab \\ &\vdots \\ &= z^{\frac{p(p-1)}{2}} a^p b^p \end{aligned} \tag{*}$$

However,  $z^p = 1$  so:

$$d^p = a^p b^p = z z^{-1} = 1$$

So  $d$  has order  $p$ , and we are back in the previous case.

If  $p = 2$ , (\*) does not hold, and the exponent of  $z$  is 1 in that case. So we have the relations:

$$a^4 = b^4 = z^2 = 1, \quad a^2 = z, \quad b^2 = z^{-1} \quad \text{and} \quad [a, b] = z$$

Therefore:

$$b^{-1}ab = az = a^3 = a^{-1}$$

So we obtain the presentation:

$$G = \langle a, b \mid a^4 = b^4 = 1, a^2 = b^2, b^{-1}ab = a^{-1} \rangle$$

Which we recognise as the quaternion group,  $Q_8$ .

Any group of order  $p^3$  is isomorphic to one of:

$$\begin{array}{lll} C_{p^3}, & C_{p^2} \times C_p, & C_p \times C_p \times C_p \\ & D_8 \quad \text{or} \quad Q_8 & (\text{when } p = 2) \\ & \text{UT}_3(p) \quad \text{or} \quad C_{p^2} \rtimes C_p & (\text{when } p \geq 3) \end{array}$$

# Chapter III

## Composite Orders

### 3.1 Groups of Order $pq$

Let  $G$  be a group of order  $pq$  where  $p, q$  are prime numbers with  $p > q$ , and let  $n_p$  and  $n_q$  denote the number of Sylow  $p$ -subgroups and Sylow  $q$ -subgroups of  $G$  respectively. Then by Sylow's Theorems:

$$n_p \equiv 1 \pmod{p} \quad \text{and} \quad n_p \mid q$$

So  $G$  has a unique Sylow  $p$ -subgroup, say  $P \trianglelefteq G$ , and a Sylow  $q$ -subgroup,  $Q \leq G$ . Because  $p$  and  $q$  are prime numbers,  $P \cong C_p$  and  $Q \cong C_q$ . Pick generators for each, say  $\langle x \rangle = P$  and  $\langle y \rangle = Q$ . We have 2 possibilities for  $n_q$ :  $p - 1$  is a multiple of  $q$  or 1.

**Case 1:**  $q \nmid p - 1$ .

If  $p - 1$  is not a multiple of  $q$  then  $n_q = 1$  and  $Q \trianglelefteq G$ , hence:

$$G = P \times Q \cong C_{pq}$$

**Case 2:**  $q \mid p - 1$ .

If  $p - 1$  is a multiple of  $q$  then  $n_q = p$  and so  $Q$  is not normal in  $G$ . By Lagrange's Theorem,  $P \cap Q = 1$  and by Lemma 4,  $|PQ| = pq$ . Hence, as well as the direct product, we have  $G = P \rtimes Q$ , some non-trivial semidirect product.

By Lemma 2,  $\text{Aut } C_p \cong (\mathbb{Z}/p\mathbb{Z})^\times \cong C_{p-1}$ . So if  $\nu \in (\mathbb{Z}/p\mathbb{Z})^\times$ , then  $x \mapsto x^\nu$  is an automorphism. We know also that  $C_{p-1}$  has a unique subgroup of order  $q$ , hence  $G$  has the presentation:

$$G = \langle x, y \mid x^p = y^q = 1, y^{-1}xy = x^a \rangle$$

where  $a$  is a generator for the subgroup of order  $q$  in  $(\mathbb{Z}/p\mathbb{Z})^\times$ .

Notice that picking different generators are equivalent up to isomorphism because the composition of two isomorphisms is an isomorphism.

Any group of order  $pq$  is isomorphic to one of:

$$C_{pq}$$

$$\langle x, y \mid x^p = y^q = 1, y^{-1}xy = x^a \rangle \quad (\text{additionally, if } q \mid p - 1)$$

### 3.1.1 Groups of Order $2p$

To illustrate an example of groups of order  $pq$ , let's take  $q = 2$ . Because every prime greater than 2 is odd,  $p - 1$  is an even number, and so  $2 \mid p - 1$ .

An element  $\alpha \in (\mathbb{Z}/p\mathbb{Z})^\times$  of order 2 satisfies  $\alpha^2 = 1$ , hence  $\alpha = 1$  or  $-1$ . But 1 has order 1, so  $\alpha$  can only be  $-1$ . Side-note: from the proof of Lemma 2, this corresponds to the inverse map.

So, in addition to  $C_{2p}$ , we have:

$$G \cong \langle x, y \mid x^p = y^2 = 1, y^{-1}xy = x^{-1} \rangle$$

Which is the presentation for the dihedral group of order  $2p$ ,  $D_{2p}$ .

Any group of order  $2p$  is isomorphic to one of:

$$C_{2p} \quad \text{or} \quad D_{2p}$$

## 3.2 Order $4q$

Let  $p$  and  $q$  be distinct prime numbers, and  $G$  be a group of order  $p^2q$ . To classify  $G$  in full generality is beyond this report, so we will focus on the cases when  $p = 2$  and when  $q = 2$ .

Let  $G$  be a group of order  $4p$ , and require  $p > 3$ . And let  $n_q$  denote the number of Sylow  $q$ -subgroups. Then  $n_q$  must divide 4, so could be 1, 2 or 4, and must be congruent to 1 mod  $q$ . If  $q = 3$ , then  $G$  could have 4 Sylow  $q$ -subgroups, so we will classify groups of order 12 later. If  $q = 2$ , then we have a group of order  $p^3$ , which we have already classified. This is why we took  $q > 3$ . So  $G$  has a normal Sylow  $q$ -subgroup,  $Q \cong C_q$ . Let  $x$  generate  $Q$ .

Lagrange's Theorem, together with Lemma 4, tell us that a Sylow 2-subgroup,  $T$ , intersects trivially with  $Q$ , and  $|QT| = |G|$ . Hence,  $G = Q \rtimes T$ .

We know by Lemma 2, that  $\text{Aut } Q \cong C_{q-1}$ . So we have two cases:

**Case 1:**  $T \cong V_4$  i.e.  $G \cong C_q \rtimes V_4$ .

We saw in our classification of groups of order  $2p$ , that  $(\mathbb{Z}/q\mathbb{Z})^\times$  has a unique element of order 2, corresponding to the inversion map. So Lemma 5 tells us that there is a single non-trivial homomorphism  $\psi : T \rightarrow \text{Aut } Q$ .

If  $\psi$  is trivial, then we obtain the product:

$$G \cong C_q \times V_4 \cong C_{2q} \times C_2$$

If  $\psi$  is non-trivial, it maps  $T$  to the subgroup generated by the inversion map, isomorphic to  $C_2$ . Therefore the kernel is isomorphic to  $C_2$ , so pick  $z$  such that it generates the kernel. Denote the other generator of  $T$  by  $y$ , then we obtain the following presentation:

$$G = \langle x, y, z \mid x^q = y^2 = z^2 = 1, yz = zy, xz = zx, y^{-1}xy = x^{-1} \rangle$$



Now let  $a = xz$ , and we will show that  $G \cong D_{4p}$ .

Firstly, notice that the order of  $a$  is  $4q$ , and:

$$a^q = x^q z^q = z \quad \text{and} \quad a^{q-1} = x^{q-1} z^{q-1} = x^{q-1}$$

Now consider:

$$y^{-1}ay = y^{-1}xzy = y^{-1}xyz = x^{-1}z = a^{q-1}a^q = a^{2q-1} = a^{-1}$$

Hence:

$$G = \langle a, y \mid a^{2q} = y^2 = 1, y^{-1}ay = a^{-1} \rangle$$

which we recognise as  $D_{4q}$ .

**Case 2:**  $T \cong C_4$  i.e.  $G \cong C_q \rtimes C_4$ .

Let  $t$  generate  $T$ . Assume  $4 \nmid q-1$ , which means  $q \equiv 3 \pmod{4}$ . So then  $\text{Aut } Q$  has no subgroup of order 4, and a homomorphism,  $\psi$  must map  $T$  to either the trivial group, or the group generated by the inverse automorphism.

If  $T\psi$  is trivial, then we recover the direct product,  $C_q \times C_4 \cong C_{4q}$ .

If  $T\psi$  is non-trivial, then  $G$  has the presentation:

$$G = \langle x, t \mid x^q = t^4 = 1, t^{-1}xt = x^{-1} \rangle$$

Let  $a = xt^2$ . Then:

$$a^q = xt^2 \dots xt^2 = x^qt^{2q} = t^{2q}$$

We know  $q \equiv 3 \pmod{4}$ , so for some  $n$ ,  $q = 4n + 3$ . Thus  $2q = 8n + 6 = 4(2n + 1) + 2$ . So then:

$$a^q = t^{4(2n+1)+2} = t^2$$

Additionally:

$$t^{-1}at = t^{-1}xt^2t = (t^{-1}xt)t^2 = x^{-1}t^2 = t^2x^{-1} = a^{-1}$$

Hence:

$$G = \langle a, t \mid a^{2q} = 1, a^q = t^2, t^{-1}at = a^{-1} \rangle$$

This is known as the binary dihedral or dicyclic group<sup>1</sup>, denoted  $\text{Dic}_{4q}$ .

If  $4 \mid q-1$ , i.e.  $q \equiv 1 \pmod{4}$ , then  $\text{Aut } Q$  contains a unique element of order 4, and so has a unique subgroup generated by it. We know by Lemma 2, that  $\text{Aut } Q \cong (\mathbb{Z}/q\mathbb{Z})^\times$ , so say  $\alpha$  is the generator of the subgroup of order 4 in  $(\mathbb{Z}/q\mathbb{Z})^\times$ . Our homomorphism can map  $T$  to this subgroup, and we get a group with the presentation:

$$G = \langle x, t \mid x^q = t^4 = 1, t^{-1}xt = x^\alpha \rangle$$

Any group of order  $4p$  is isomorphic to one of:

$$C_{4q}, \quad C_{2q} \times C_2, \quad D_{4q} \quad \text{or} \quad \text{Dic}_{4q}$$

$$\langle x, t \mid x^q = t^4 = 1, t^{-1}xt = x^\alpha \rangle \quad (\text{additionally, if } 4 \mid q-1)$$

1. Groupprops, "Dicyclic Groups," October 21, 2017, accessed January 19, 2023, [https://groupprops.subwiki.org/wiki/Dicyclic\\_group](https://groupprops.subwiki.org/wiki/Dicyclic_group).

### 3.3 Order $2p^2$

Let  $G$  be a group of order  $2p^2$ , with  $p > 2$ . Denote the number of Sylow  $p$ -subgroups by  $n_p$ . By Sylow's Theorems,  $n_p$  divides 2, and is congruent to 1 mod  $p$ , so must be 1. Hence,  $G$  has a normal Sylow  $p$ -subgroup,  $P$  of order  $p^2$ .

If  $T$  is a Sylow 2-subgroup, then by applying Lagrange's Theorem, and Lemma 4, we can conclude that  $G = P \rtimes T$ . From our classification of groups of order  $p^2$ , we have 2 choices for  $P$ :

**Case 1:**  $P \cong C_{p^2}$  i.e.  $G \cong C_{p^2} \rtimes C_2$ .

From Lemma 2, we know  $|\text{Aut } P| = p^2 - p = p(p-1)$ . Because  $p$  is prime,  $2 \nmid p$ , but  $2 \mid p-1$ , so  $\text{Aut } P$  has a unique element of order 2. Hence, in addition to the direct product,  $G \cong C_{2p^2}$ , we have  $G \cong C_{p^2} \rtimes C_2$ , with  $C_2$  acting by inversion. If  $x$  generates  $P$ , and  $y$  generates  $T$ , we have the presentation:

$$G = \langle x, y \mid x^{p^2} = y^2 = 1, y^{-1}xy = x^{-1} \rangle$$

which we recognise as  $D_{2p^2}$ , the dihedral group of order  $2p^2$ .

**Case 2:**  $P \cong C_p \times C_p$  i.e.  $G \cong C_p \times C_p \rtimes C_2$ .

Consider  $P$  as the product of the subgroups generated by  $a$  and  $b$ , i.e.  $P = \langle a \rangle \times \langle b \rangle$ . Then the action of  $T$  on  $P$  can either be trivial on both subgroups, invert one, or invert both.

If the action is trivial on both subgroups, then we recover the direct product  $G \cong C_p \times C_{2p}$ .

If the action is non-trivial on just one of the subgroups, then we can consider only one case. This is because they are equivalent up to an isomorphism of  $T$ , and Lemma 5 tells us the resulting semidirect products are isomorphic. So we have:

$$G = \langle a \rangle \times (\langle b \rangle \rtimes T) \cong C_p \times D_{2p}$$

Finally, if we choose to invert both subgroups, then we act on all of  $P$  by inversion. So if  $a$  and  $b$  generate  $P$ , then:

$$G = \langle a, b, x \mid a^p = b^p = x^2 = 1, ab = ba, x^{-1}ax = a^{-1}, x^{-1}bx = b^{-1} \rangle$$

Because  $C_p$  has all elements of order  $p$ , excluding 1, and they are all automorphic to each other (meaning that some automorphism maps one to the other),  $x^{-1}gx = g^{-1}$  for all  $g \in P$ . Hence:

$$G = \langle P, x \mid x^2 = 1, x^{-1}gx = g^{-1} \forall g \in P \rangle$$

which is known as the generalised dihedral group<sup>2</sup> for  $P$ , denoted  $\text{Dih}(P)$ .

Any group of order  $2p^2$  is isomorphic to one of:

$$C_{2p^2}, \quad D_{2p^2}, \quad C_p \times C_{2p}, \quad C_p \times D_{2p} \quad \text{or} \quad \text{Dih}(C_p \times C_p)$$

2. Groupprops, "Generalized dihedral group," January 17, 2011, accessed February 15, 2023, [https://groupprops.subwiki.org/wiki/Generalized\\_dihedral\\_group](https://groupprops.subwiki.org/wiki/Generalized_dihedral_group).

# Chapter IV

## Special Cases

### 4.1 Groups of order 12

We have seen that groups of order 12 have slightly different behaviour to groups of order  $4q$  in general, and we will need this classification in order to classify groups of order 24.

Let  $G$  be a group of order  $12 = 2^2 \cdot 3$ , and  $n_3$  and  $n_2$  denote the number of Sylow 3-subgroups and Sylow 2-subgroups of  $G$  respectively. By Sylow's Theorems:

$$n_2 \equiv 1 \pmod{2} \quad \text{and} \quad n_2 \mid 3$$

$$n_3 \equiv 1 \pmod{3} \quad \text{and} \quad n_3 \mid 4$$

Hence:

$$n_2 = 1 \text{ or } 3$$

$$n_3 = 1 \text{ or } 4$$

Let  $H$  be a Sylow 2-subgroup and  $K$  be a Sylow 3-subgroup of  $G$ , generated by  $x$ .

Lagrange's Theorem tells us  $H$  has elements of order 1, 2, and 4, and  $K$  has elements of order 1 and 3. Hence  $H \cap K = \mathbf{1}$ . Lemma 4 tells us:

$$|HK| = |H| \cdot |K| = 12$$

Hence  $G = HK$ ,  $H \trianglelefteq G$ , and  $H \cap K = \mathbf{1}$ .

Since an automorphism,  $\varphi$ , must map generators to generators,  $\text{Aut } C_4 \cong C_2$  because  $C_4$  has two generators. An automorphism of  $V_4$  corresponds to a permutation of the three non-identity elements, hence  $\text{Aut } V_4 \cong S_3$ .

If we consider  $G$  where  $K \trianglelefteq G$ , i.e.  $G = K \rtimes H$ , then we have two cases:

**Case 1:**  $H \cong C_4$  i.e.  $G \cong C_3 \rtimes C_4$ .

Let  $H = \langle y \rangle$ . We know  $\text{Aut } C_3 \cong C_2$  so a homomorphism  $\psi$  maps  $H$  to the trivial group or to  $\langle \beta : x \mapsto x^{-1} \rangle$ .

If  $H\psi = \mathbf{1}$  then  $G = K \times H \cong C_3 \times C_4 \cong C_{12}$ .

If  $H\psi = \langle \beta \rangle$  then we have:

$$G = \langle x, y \mid x^3 = y^4 = 1, y^{-1}xy = x^{-1} \rangle$$

Now let  $a = xy^2$ . And remember,  $y^{-1}xy = x^{-1}$  means  $x$  commutes with  $y^2$ . So now:

$$a^3 = xy^2xy^2xy^2 = x^3y^6 = y^2$$

and:

$$y^{-1}ay = y^{-1}xy^2y = (y^{-1}xy)y^2 = x^{-1}y^2 = y^2x^{-1} = a^{-1}$$

So:

$$G = \langle a, y \mid a^6 = 1, a^3 = y^2, y^{-1}ay = a^{-1} \rangle$$

which we recognise as  $\text{Dic}_{12}$ . This group is also sometimes denoted by  $T$ .

**Case 2:**  $H \cong V_4$  i.e.  $G \cong C_3 \rtimes (C_2 \times C_2)$ .

If  $\psi : H \rightarrow \text{Aut } K$  is trivial then we obtain the direct product again. We know  $\text{Aut } K \cong C_2$ , so there are 3 choices of elements in  $H$  to send to it, but they are all equivalent up to isomorphism, by Lemma 5, taking  $\alpha$  to be the identity map.

We know that  $H/\text{im } \psi \cong \ker \psi$ , so  $\ker \psi$  must be isomorphic to  $C_2$ . Pick  $z$  so that it generates the kernel, and so the remaining generator,  $y$  is not in the kernel. Then:

$$G = \langle x, y, z \mid x^3 = y^2 = z^2 = 1, yz = zy, xz = zx, y^{-1}xy = x \rangle$$

Let  $a = xz$ . So:

$$a^3 = x^3z^3 = z$$

and:

$$y^{-1}ay = y^{-1}xzy = y^{-1}xyz = x^{-1}z = x^2z = a^{-1}$$

Hence:

$$G = \langle y, a \mid a^6 = y^2 = 1, y^{-1}ay = a^{-1} \rangle \cong D_{12}$$

Instead, if  $G$  has 4 Sylow 3-subgroups, then there are 8 elements of order 3 in  $G$ . So the remaining 4 must form the Sylow 2-subgroup, hence it is normal.

**Case 1:**  $H \cong C_4$  i.e.  $G \cong C_4 \rtimes C_3$ .

Let  $H = \langle y \rangle$ . A homomorphism  $\psi : K \rightarrow \text{Aut } H \cong C_2$ , preserves order and together with Lagrange's Theorem means that the only possibility for  $\psi$  is trivial, i.e.  $K\psi = \mathbf{1}$ .

Hence  $G \cong C_4 \times C_3 \cong C_{12}$ .

**Case 2:**  $H \cong V_4$  i.e.  $G \cong (C_2 \times C_2) \rtimes C_3$ .

Let  $H = \langle y, z \rangle$ . A trivial homomorphism  $K\psi = \mathbf{1}$  yields the direct product. What non-trivial homomorphisms are there? The automorphism group,  $\text{Aut } H \cong S_3$  is of order 6, and so has a unique subgroup of order 3, by Sylow's Theorems. We know that a homomorphism  $\psi : K \rightarrow \text{Aut } H$  is determined by where it sends the generator  $x$ , so for  $\psi$  to be non-trivial, it must send  $x$  to an element of order 3 in  $\text{Aut } H$ .

There are 2 such elements. Because  $\text{Aut } H \cong S_3$ , we will think of them as the permutations of order 3 of the set  $\{1, 2, 3\}$ . Denote them  $a = (1\ 2\ 3)$  and  $b = (1\ 3\ 2)$ . Notice that  $b = a^{-1}$ , so we have homomorphisms:

$$\psi_1 : x \mapsto a \quad \text{and} \quad \psi_2 : x \mapsto a^{-1}$$

It appears we have 2 choices, but this is not the case. The inverse map,  $\beta : x \mapsto x^{-1}$ , is an automorphism of  $K$ , and so by Lemma 5, the corresponding semidirect products of  $\psi_1$  and  $\psi_2$  are isomorphic. Hence (up to isomorphism) there is one non-trivial homomorphism  $\psi : K \rightarrow \text{Aut } H$ . So  $x \in K$  acts by permuting the 3 non-identity elements of  $H$ .

We will show that in this case,  $G \cong A_4$ . First, let's check  $A_4$  has the same subgroup structure as  $G$ . There is a subgroup isomorphic to  $C_3$  in  $A_4$ , generated by the 3-cycle  $(1\ 2\ 3)$ :

$$\bar{K} = \langle (1\ 2\ 3) \rangle$$

We can also find a subgroup isomorphic to  $V_4$ :

$$\bar{H} = \{ 1, (1\ 2)(3\ 4), (1\ 3)(2\ 4), (1\ 4)(2\ 3) \}$$

Indeed, we can check that  $\bar{H}$  is normal in  $A_4$ . We can see that  $\bar{H} \cap \bar{K} = 1$  because  $\bar{H}$  contains no 3-cycles, and that  $\bar{H}\bar{K} = A_4$ . So we can conclude that  $A_4 = \bar{H} \rtimes \bar{K}$ .

Let's investigate how conjugation behaves. If we let  $\alpha = (1\ 2)(3\ 4)$ ,  $\beta = (1\ 4)(2\ 3)$  and  $\gamma = (1\ 2\ 3)$ , then we can write an element of  $A_4$  as  $\alpha^i \beta^j \gamma^k$  for some  $i, j$  and  $k$ . Define  $\phi : A_4 \rightarrow G$  by  $\phi : \alpha^i \beta^j \gamma^k \mapsto x^i y^j z^k$ . Then:

$$\beta \phi = (\gamma^{-1} \alpha \gamma) \phi = c^{-1} a c = b$$

So conjugation acts in the same way. Hence we can conclude that  $G \cong A_4$ .

Any group of order 12 is isomorphic to one of:

$$C_{12}, \quad C_2 \times C_6, \quad A_4, \quad D_{12} \quad \text{or} \quad \text{Dic}_{12}$$

## 4.2 Groups of Order 24

Let  $G$  be a group of order 24, and let  $H$  be a Sylow 3-subgroup of  $G$ , so  $H \cong C_3$ , and let  $h$  generate  $H$ . Let  $T$  be a Sylow 2-subgroup of  $G$ , so  $T$  has order 8. By Lagrange's Theorem,  $H \cap T = 1$  and then applying Lemma 4,  $|HT| = 24$ . Now let  $n_3$  denote the number of Sylow 3-subgroups, and by Sylow's Theorems:

$$n_3 \equiv 1 \pmod{3} \quad \text{and} \quad n_3 \mid 8$$

Hence  $n_3$  is either 1 or 4.

If  $n_3 = 1$ , then  $H$  is normal in  $G$ . Thus  $G = H \rtimes T$ . We'll want a homomorphism  $\psi : T \rightarrow \text{Aut } H$ . We know  $\text{Aut } H \cong C_2$ , and from our classification of groups of order 8, we have 5 possibilities. An action of  $T$  on  $H$  will have image isomorphic to  $C_2$ , and a kernel isomorphic to a group of order 4. We can classify the possible actions by considering the kernel.

**Case 1:**  $T \cong C_8$  i.e.  $G \cong C_3 \rtimes C_8$ .

Let  $t$  generate  $T$ , and so its unique subgroup of order 4 is generated by  $t^2$ . Hence  $\langle t^2 \rangle$  is the kernel of  $\psi$ , so  $\psi$  must send  $t$  to the identity or inversion map. Hence a non-trivial action of  $T$  on  $H$  is unique. If the action is trivial, then:

$$G = T \times H \cong C_{24}$$

Otherwise we obtain:

$$G = \langle h, t \mid h^3 = t^8 = 1, h^{-1}th = t^{-1} \rangle \cong C_3 \rtimes C_8$$

**Case 2:**  $T \cong (C_4 \times C_2)$  i.e.  $G \cong C_3 \rtimes (C_4 \times C_2)$ .

In this case,  $T$  has subgroups isomorphic to both  $C_4$  and  $C_2 \times C_2$ , so we have more possibilities for  $\psi$ . Firstly, if  $\psi$  is trivial, then we obtain the direct product:

$$G \cong C_3 \times C_4 \times C_2$$

Let  $T$  be generated by  $x$  and  $y$ , where  $x^4 = y^2 = 1$ , and consider non-trivial  $\psi$ . Say the kernel of  $\psi$  is isomorphic to  $C_2 \times C_2$ . So it must be generated by the elements of order 2 in  $T$ ,  $x^2$  and  $y$ . Then  $\psi$  must map  $x$  to the non-identity element in  $\text{Aut } H$ : inversion. Hence  $\langle x \rangle$  acts by inversion on  $H$ , giving:

$$\begin{aligned} G &= (H \rtimes \langle x \rangle) \times \langle y \rangle \\ &\cong (C_3 \rtimes C_4) \times C_2 \\ &\cong \text{Dic}_{12} \times C_2 \end{aligned}$$

If instead the kernel is isomorphic to  $C_4$ , then it must be generated by an element of order 4 from  $T$ . However, all elements of order 4 are automorphic, and so by Lemma 5, we can pick  $x$  to generate the kernel, without loss of generality. So then  $\psi$  must map  $y$  to inversion. Hence  $\langle x \rangle$  acts trivially on  $H$ , and  $\langle y \rangle$  acts by inversion. Thus:

$$\begin{aligned} G &= (H \rtimes \langle y \rangle) \times \langle x \rangle \\ &\cong (C_3 \rtimes C_2) \times C_4 \\ &\cong S_3 \times C_4 \end{aligned}$$

**Case 3:**  $T \cong (C_2 \times C_2 \times C_2)$  i.e.  $G \cong C_3 \rtimes (C_2 \times C_2 \times C_2)$ .

Let  $\langle a, b, c \rangle = T$ . All elements in  $T$  have order 1 or 2, so cannot have subgroups isomorphic to  $C_4$ . However,  $T$  does have subgroups isomorphic to  $C_2 \times C_2$ , which can be generated by 2 of the 3 generators of  $T$ . This gives us 3 subgroups, but permuting the generators  $a, b$  and  $c$  is an automorphism of  $T$ , so Lemma 5 tells us the resulting semidirect products are isomorphic. So choose  $\psi$  such that  $b$  and  $c$  are in the kernel. Then  $a\psi$  is either the identity map or the inversion map. If  $\psi$  is trivial, then we obtain the direct product:

$$G \cong C_3 \times C_2 \times C_2 \times C_2$$

If  $a\psi$  is inversion, then:

$$G = (C_3 \rtimes \langle a \rangle) \times \langle b \rangle \times \langle c \rangle \cong S_3 \times C_2 \times C_2$$

**Case 4:**  $T \cong D_8$  i.e.  $G \cong C_3 \rtimes D_8$ .

Let  $r$  and  $s$  generate  $T$  with  $r^4 = s^2 = 1$ . A trivial homomorphism will yield the direct product:

$$G \cong C_3 \times D_8$$

So for a non trivial homomorphism, firstly assume  $\ker \psi \cong C_4$ . There is a unique subgroup in  $T$  isomorphic to  $C_4$ , so it's generated by an element of order 4. However the choice of generator is the same up to an isomorphism of  $T$ , so Lemma 5 lets us pick  $r$  to be the generator, without loss of generality. Hence  $s$  cannot be in the kernel, and so  $s\psi$  is the inversion map. We obtain the presentation:

$$G = \langle x, r, s \mid x^3 = r^4 = s^2 = 1, xr = rx, s^{-1}rs = r^{-1}, s^{-1}xs = x^{-1} \rangle$$

Let  $a = xr$ , and consider:

$$s^{-1}as = s^{-1}xrs = s^{-1}xrs^2s^{-1} = (s^{-1}xs)(srs^{-1}) = x^{-1}r^{-1} = r^{-1}x^{-1} = a^{-1}$$

So we have:

$$G = \langle a, s \mid a^{12} = s^2 = 1, s^{-1}as = a^{-1} \rangle$$

Which we recognise as  $D_{24}$ , the dihedral group of order 24.

If instead we consider  $\psi$  with kernel isomorphic to  $C_2 \times C_2$ , then the kernel is generated by two elements of order 2. However,  $T$  only has two elements of order 2,  $r^2$  and  $s$ , so they must generate the kernel. So then  $\psi$  must map  $r$  to inversion. Hence this action is fully specified. So:

$$G \cong C_3 \rtimes_{V_4} D_8$$

We will use the above notation to mean the unique action with kernel isomorphic to  $V_4$ .

**Case 5:**  $T \cong Q_8$  i.e.  $G \cong C_3 \rtimes Q_8$ . Let  $T$  be generated by  $i$  and  $j$ , with the product denoted by  $k$ . That is:

$$T = \langle i, j \mid i^4 = j^4 = 1, i^2 = j^2, j^{-1}ij = i^{-1} \rangle$$

There is a single element of order 2 in  $T$ , hence  $T$  has no subgroup isomorphic to  $C_2 \times C_2$ . The elements  $i, j$  and  $k$  each generate a cyclic subgroup in  $T$ . So  $\psi$  will send one of them to the kernel. We know that permuting these is an automorphism of  $T$ , so Lemma 5 tells us the choice results in isomorphic semidirect products.

So take  $i \in \ker \psi$ . Indeed  $\langle i \rangle = \ker \psi$ . Then for a non-trivial homomorphism, we must have  $j \notin \ker \psi$ . Otherwise:

$$i\psi j\psi = (ij)\psi = k\psi \in \ker \psi$$

making  $\psi$  trivial.

Thus either  $\psi$  is trivial and we obtain:

$$G \cong C_3 \times Q_8$$

or  $\psi$  maps  $j$  to the inversion map and we obtain the presentation:

$$G = \langle x, i, j \mid x^3 = i^4 = j^4 = 1, xi = ix, i^2 = j^2, j^{-1}xj = x^{-1}, j^{-1}ij = i^{-1} \rangle$$

Now let  $a = xi$ . So:

$$a^6 = x^6 i^6 = i^2 = j^2$$

And:

$$j^{-1}aj = j^{-1}xij = j^{-1}xji^{-1} = x^{-1}i^{-1} = i^{-1}x^{-1} = a^{-1}$$

Hence:

$$G = \langle a, j \mid a^{12} = 1, a^6 = j^2, j^{-1}aj = a^{-1} \rangle$$

We recognise this as the dicyclic group of order 24,  $\text{Dic}_{24}$ .

If  $n_3 = 4$  then  $H$  is not normal. We will proceed to show that  $G$  must have a normal Sylow 2-subgroup in a similar way to Borchers<sup>1</sup>.

The normaliser of  $H$ ,  $N_G(H)$  has index 4. Now let  $G$  act on the set of the cosets of  $N_G(H)$  by conjugation. Hence we obtain a homomorphism  $\rho : G \rightarrow S_4$ . The kernel is a subgroup of  $N_G(H)$  so must have order dividing 6 by Lagrange's Theorem.

The kernel cannot be of order 3, because  $G$  has no normal subgroup of order 3 (because kernels of homomorphism are normal in the domain group). Likewise the kernel cannot be of order 6 because every group of order 6 has a unique Sylow 3-subgroup, which is characteristic. So by Lemma 3, it would be normal in  $G$ . Hence the kernel must have order 1 or 2.

If the kernel is of order 1, then  $\rho$  is an isomorphism, so  $G \cong S_4$ .

If the kernel is of order 2, then we know that  $G/\ker \rho \cong \text{im } \rho$ , so then  $\text{im } \rho$  must have order 12. It also cannot have a normal Sylow 3-subgroup, so looking at our classification of groups of order 12, this must be isomorphic to  $A_4$ . We know that  $A_4$  has a normal subgroup of order 4, and so by the Correspondence Theorem,  $G$  must contain a normal subgroup of order 8, say  $T$ . By Lagrange's Theorem and Lemma 4, we can conclude that  $G = T \rtimes H$ . Again, we have 5 cases, but this time we'll exclude the trivial homomorphism, because that will just give us the direct product which we have already seen:

**Case 1:**  $T \cong C_8$  i.e.  $G \cong C_8 \rtimes C_3$ .

An automorphism of  $T$ ,  $\varphi$ , maps generators to generators, so say  $\langle x \rangle = T$ . Then  $x\varphi$  could be  $x, x^3, x^5$  or  $x^7$ . Hence, and Lagrange's Theorem tells us that there are no non-trivial homomorphisms  $\psi : H \rightarrow \text{Aut } T$ . As a bonus: notice that each of these, apart from the identity, has order 2, so  $\text{Aut } C_8 \cong V_4$ .

**Case 2:**  $T \cong (C_4 \times C_2)$  i.e.  $G \cong (C_4 \times C_2) \rtimes C_3$ .

An automorphism of  $T$ ,  $\psi$  preserves element order, so if  $\langle x, y \mid x^4 = y^2 = 1, xy = yx \rangle = T$ , then  $x\psi$  must be of order 4, and  $y\psi$  must be of order 2. Moreover,  $y\psi$  cannot be in  $\langle x\psi \rangle$  because  $\psi$  is injective.

So we are reduced to 2 possible choices for  $y\psi$ , and 4 possible choices for  $x\psi$ . Because an automorphism is determined by its effect on generators, this gives us 8 possible automorphisms. Hence  $|\text{Aut } T| = 8$ , and Lagrange's Theorem tells us that there are no non-trivial homomorphisms  $\psi : H \rightarrow \text{Aut } T$ .

---

1. Richard E. Borchers, "Group theory 21: Groups of order 24," June 30, 2020, accessed February 9, 2023, <https://www.youtube.com/watch?v=6TWuo2NO8vg>.



**Case 3:**  $T \cong (C_2 \times C_2 \times C_2)$  i.e.  $G \cong (C_2 \times C_2 \times C_2) \rtimes C_3$ .

To determine  $\text{Aut } T$  it is helpful to think of  $C_2$  as the finite field with two elements. Then  $T$  is isomorphic a 3 dimensional vector space over two elements. So an automorphism of that vector space is just any linear map, with non-zero determinant. Thus,  $\text{Aut } T \cong \text{GL}_3(2)$ .

We can determine that  $|\text{GL}_3(2)| = 168 = 2^3 \cdot 3 \cdot 7$ , so  $\text{Aut } T$  has a Sylow 3-subgroup of order 3, isomorphic to  $C_3$ . Sylow's Theorems tells us that all subgroups of order 3 are conjugate, so Lemma 5 tells us there is only one unique action (up to isomorphism) of  $H$  on  $T$ . As before, pick a homomorphism,  $\psi$ , which will let us easily classify the resulting semidirect product.

Write  $T = A \times B$  where  $A \cong C_2$  and  $B \cong C_2 \times C_2$ . Then let  $\psi$  map  $H$  to the subgroup generated by the automorphism which fixes  $A$  and permutes the non-identity elements of  $B$  in a 3-cycle. This automorphism has order 3 by construction, so we can write:

$$G \cong C_2 \times (V_4 \rtimes C_3)$$

We know already that  $V_4 \rtimes C_3 \cong A_4$ , so  $G \cong C_2 \times A_4$ .

**Case 4:**  $T \cong D_8$  i.e.  $G \cong D_8 \rtimes C_3$ .

Let  $\langle s, r \mid s^2 = r^4 = 1, s^{-1}rs = r^{-1} \rangle = T$ . An automorphism,  $\psi$ , of  $T$  preserves element order, so for  $r\psi$  we have two choices,  $r$  or  $r^{-1}$ . We can send  $s\psi$  to any element of order 2 which is not in  $\langle r\psi \rangle$ . This leaves only reflections, of which there are 4:  $s, rs, r^2s$  and  $r^3s$ . Hence there are 8 possible automorphisms of  $D_8$ , so  $|\text{Aut } D_8| = 8$ . Lagrange's Theorem tells us that there are no non-trivial homomorphisms  $\psi : H \rightarrow \text{Aut } T$ .

**Case 5:**  $T \cong Q_8$  i.e.  $G \cong Q_8 \rtimes C_3$ .

Firstly, because of the multiplication structure of the quaternions, the image of  $k$  is determined by the images of  $i$  and  $j$ ; it is forced. This reduces the possibilities for an automorphism. Additionally,  $\pm 1$  are fixed by an automorphism, because they are the only elements of their order. So an automorphism could send  $i$  to any of the 6 elements of order 4. The image of  $j$  cannot be in the subgroup generated by the image of  $i$ , otherwise we wouldn't have an automorphism. Thus there are 4 choices for the image of  $j$ , giving us 24 possible automorphisms altogether.

So  $\text{Aut } T$  will have a Sylow subgroup of order 3. Lemma 5 tells us that if a homomorphism maps  $H$  to a given subgroup in  $\text{Aut } T$ , then mapping  $H$  to a conjugate subgroup produces an isomorphic semidirect product. Further, we can map  $H$  into the subgroup any way we want, because we can apply the automorphism  $\beta$ . Hence,  $Q_8 \rtimes C_3$  defines a single isomorphism class.

We will show that  $\text{SL}_2(3)$  is in that isomorphism class. Firstly, let  $e_1$  and  $e_2$  be the basis vectors for the vector space. We can find an element of order 3 in  $\text{SL}_2(3)$ :

$$\bar{h} = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$$

And so  $\bar{h}$  generates a cyclic subgroup of order 3 inside  $\text{SL}_2(3)$ , say  $\bar{H}$ . The centre of  $\text{SL}_2(3)$  (denote by  $\bar{Z}$ ) is just:

$$\bar{Z} = \left\{ \begin{pmatrix} \lambda & 0 \\ 0 & \lambda \end{pmatrix} \mid \lambda^2 = 1 \right\} = \{\pm I\}$$

Now let's find 3 matrices which correspond to  $i, j$  and  $k$  from  $Q_8$ . We need matrices,  $A, B$  and  $C$  such that  $A^2 = -1$ . So the minimal polynomial of  $A$  is  $x^2 + 1$ . Without too much trouble, we can find:

$$A = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$$

Conjugating by  $\bar{h}$  will give us another:

$$B = \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & -1 \\ -1 & -1 \end{pmatrix}$$

So then their product will give us  $C$ :

$$C = AB = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} 1 & -1 \\ -1 & -1 \end{pmatrix} = \begin{pmatrix} -1 & -1 \\ -1 & 1 \end{pmatrix}$$

With  $A, B$  and  $C$  in hand, let's quotient  $\text{SL}_2(3)$  by its centre and find what group we get. It will turn out that  $\text{SL}_2(3)/\bar{Z}$  is isomorphic to  $A_4$ , and we will use this to find a normal copy of  $Q_8$  inside  $\text{SL}_2(3)$ .

In  $\text{SL}_2(3)/\bar{Z}$ , we have 4 1-dimensional subspaces given by:

$$U_1 = \text{Span}(e_1)$$

$$U_2 = \text{Span}(e_2)$$

$$U_3 = \text{Span}(e_1 + e_2)$$

$$U_4 = \text{Span}(e_1 - e_2)$$

Let's investigate how the quotient acts on the set of these subspaces. The element  $A$  sets  $e_1$  to  $e_2$  and  $e_2$  to  $-e_1$ , so swaps  $U_1$  and  $U_2$ . And so  $e_1 + e_2$  becomes  $e_1 - e_2$ , and  $e_1 - e_2$  becomes  $-e_1 - e_2 = -(e_1 + e_2)$ . Thus  $A$  swaps  $U_3$  and  $U_4$  and so it corresponds to the permutation  $(1\ 2)(3\ 4)$ .

By a similar argument, we can determine that  $B$  corresponds to  $(1\ 4)(2\ 3)$ , and  $C$  corresponds to  $(1\ 3)(2\ 4)$ . And so we can conclude that:

$$\text{SL}_2(3)/\bar{Z} \cong A_4$$

We know that  $A_4$  has a normal subgroup isomorphic to  $C_2 \times C_2$ , and so by applying the Correspondence Theorem,  $\text{SL}_2(3)$  has a normal subgroup isomorphic to  $Q_8$ , which contains the centre, and elements  $A, B$  and  $C$ . Hence,  $Q_8 \rtimes C_3 \cong \text{SL}_2(3)$ .

Any group of order 24 is isomorphic to one of:

$C_{24}$	$C_3 \rtimes C_8$	$C_{12} \times C_2$
$\text{Dic}_{12} \times C_2$	$S_3 \times C_4$	$C_3 \times C_2 \times C_2 \times C_2$
$S_3 \times C_2 \times C_2$	$C_3 \times D_8$	$D_{24}$
$C_3 \rtimes_{V_4} D_8$	$C_3 \times Q_8$	$\text{Dic}_{24}$
$S_4$	$C_2 \times A_4$	$\text{SL}_2(3)$

### 4.3 Groups of Order 30

This classification is based on the one given in the cited Stack Exchange post<sup>2</sup>. Let  $G$  be a group of order  $30 = 2 \cdot 3 \cdot 5$ . So then  $G$  has a Sylow 3-subgroup,  $T$ , and a Sylow 5-subgroup,  $F$ . Let  $H = TF$  and by Lagrange's Theorem,  $T \cap F = \mathbf{1}$ , hence  $|H| = 15$  by Lemma 4. We know from our classification of groups of order  $pq$  that  $H \cong C_{15}$ . Because  $|H| = 15 = \frac{30}{2}$ , the index of  $H$  in  $G$  is 2, and we know a subgroup of index 2 is normal, so  $H \trianglelefteq G$ .

A Sylow 2-subgroup  $K \leq G$  has order 2, so  $K \cong C_2$ . Let  $\langle k \rangle = K$  and  $\langle h \rangle = H$ . By the same argument as above,  $H \cap K = \mathbf{1}$  and  $|HK| = 30$ . Hence  $G = HK$ . Moreover,  $G = H \rtimes K$ .

By Lemma 2:

$$\text{Aut } C_{15} = (\mathbb{Z}/15\mathbb{Z})^\times \cong (\mathbb{Z}/3\mathbb{Z} \times \mathbb{Z}/5\mathbb{Z})^\times \cong (\mathbb{Z}/3\mathbb{Z})^\times \times (\mathbb{Z}/5\mathbb{Z})^\times \cong C_2 \times C_4$$

A homomorphism,  $\psi : C_2 \rightarrow C_2 \times C_4$  preserves element order and we know  $\psi$  is determined by its effect on a generator. So then  $k\psi$  has four possibilities: either the identity, or one of the three elements of order 2.

Additionally,  $\psi$  preserves the Sylow subgroups of  $H$ . So write  $H = \langle h^3 \rangle \times \langle h^5 \rangle$ , the direct product of its Sylow subgroups. So the action of  $K$  on  $H$  is either trivial or by inversion on each of the Sylow subgroups of  $H$ , giving us 4 possibilities:

**Case 1:** Trivial action on both Sylow subgroups.

In this case, because the action is trivial on all of  $H$ , we recover the direct product,  $G = H \times K \cong C_{30}$ .

**Case 2:** Inversion on both Sylow subgroups.

Here,  $K$  acts on all of  $H$ , so we obtain:

$$G = \langle h, k \mid h^{15} = k^2 = 1, k^{-1}hk = h^{-1} \rangle$$

which we recognise as  $D_{30}$ .

**Case 3:** Inversion on  $\langle h^5 \rangle$ .

We know already, from our classification of groups of order  $2p$ , that  $C_3 \rtimes C_2 \cong D_6$ . So then because the action on  $\langle h^3 \rangle$  is trivial:

$$G = \langle h^3 \rangle \times (\langle h^5 \rangle \rtimes K) \cong C_5 \times D_6$$

**Case 4:** Inversion on  $\langle h^3 \rangle$ .

Similar to above, we obtain:

$$G = \langle h^5 \rangle \times (\langle h^3 \rangle \rtimes K) \cong C_3 \times D_{10}$$

Any group of order 30 is isomorphic to one of:

$$C_{30}, \quad D_{15}, \quad C_5 \times D_6 \quad \text{or} \quad C_3 \times D_{10}$$

2. Stack Exchange (user azimuth), "Classification of groups of order 30 (duplicate)," December 10, 2020, accessed January 24, 2023, <https://math.stackexchange.com/questions/569226/classification-of-groups-of-order-30>.

## 4.4 Groups of Order 16

And finally, all that remains is to classify groups of order 16. This classification is often not included; *MT4003* classifies groups up to order 15 for example. So to cap of this report, let's tackle one of the hardest cases!

The following classification is based of the one by Wild<sup>3</sup>. For brevity, denote  $C_2 \times C_2 \times C_2 \times C_2$  by  $(C_2)^4$ . We begin by proving a lemma:

**Lemma 11.** *For a group,  $G$ , with order 16, not isomorphic to  $(C_2)^4$ , then  $G$  has a normal subgroup isomorphic to either  $C_8$  or  $C_4 \times C_2$ .*

*Proof.* Firstly, we know that a subgroup with index 2 is normal, so any subgroup of order 8 in  $G$  is normal. So it remains to show  $G$  possesses such subgroups. If some  $x \in G$  has order 8, then  $\langle x \rangle \cong C_8$  and we are done. So assume  $G$  has no element of order 8.

Because  $G \not\cong (C_2)^4$ , there is at least one element of order 4 in  $G$ , say  $y$ . By Lemma 9, there is some element  $z \in Z(G)$  which has order 2. Let  $H = \langle z \rangle$ . Moreover,  $H \trianglelefteq G$  because  $z$  is in the centre.

If  $y^2 \neq z$ , then  $\langle y \rangle \cap H = \mathbf{1}$ , so  $\langle y, z \rangle \cong C_4 \times C_2$ .

If then all elements of order 4 in  $G$  have  $y^2 = z$ , all elements in  $G/H$  have order 2. Thus  $G/H \cong (C_2)^3$ , and in particular, is abelian. So the conjugacy class of  $y$  is a subgroup of  $yH$ . Hence the centraliser of  $y$ ,  $C_G(y)$ , has order 8.

Let  $g \in C_G(y) \setminus \langle y \rangle$ . If  $g$  has order 2, then  $\langle y, g \rangle \cong C_4 \times C_2$ . If  $g$  has order 4, then:

$$g^2 = z \quad \text{and} \quad (yg)^2 = y^2 g^2 = z^2 = 1$$

We can see that  $yg \notin \langle y \rangle$  because then,  $yg = y^2$  which gives the contradiction  $y = g$ . So  $\langle y, yg \rangle \cong C_4 \times C_2$ .  $\square$

Since we know already that  $(C_2)^4$  is a group of order 16 by the Fundamental Theorem of Finite Abelian Groups, we will concentrate on when  $G$  is not isomorphic to  $(C_2)^4$ . If  $N$  is the subgroup of  $G$  specified by Lemma 11, we will classify the groups by extending  $N$  in different ways. Pick some element  $a \in G \setminus N$ . Then the order of  $aN$  is two, (it must be either 1 or 2, and cannot be 1 because that contradicts our choice of  $a$ ) and  $a^2$  is in  $N$ . We know already what the possible automorphisms of  $C_8$  and  $C_4 \times C_2$  are from our classification of groups of order 24. They are summarised in Tables 1 and 2.

So now by considering choices for the order of  $a$  and automorphisms of  $N$ , we will classify  $G$ . Begin with  $\langle x \rangle = N \cong C_8$ .

**Case 1:** Order 2.

If  $a$  has order 2, then  $a^2$  must be 1. So then  $N \cap \langle a \rangle = \mathbf{1}$ , and  $G = N \rtimes \langle a \rangle$ . We have 4 possible semidirect products, because each of the automorphisms of

---

3. Marcel Wild, "The Groups of Order Sixteen Made Easy," *The American Mathematical Monthly* 112, no. 1 (2005): 20–31, ISSN: 00029890, 19300972, accessed February 27, 2023, <http://www.jstor.org/stable/30037381>.

Table 1: Automorphisms of  $C_8$

Aut $C_8$	$x \mapsto$
$\phi_1$	$x$
$\phi_2$	$x^3$
$\phi_3$	$x^5$
$\phi_4$	$x^{-1}$

Table 2: Automorphisms of  $C_4 \times C_2$

Aut $C_8$	$x \mapsto$	$y \mapsto$	Order
$\psi_1$	$x$	$y$	1
$\psi_2$	$x^3y$	$x^2y$	4
$\psi_3$	$x^3$	$y$	2
$\psi_4$	$xy$	$x^2y$	4
$\psi_5$	$xy$	$y$	2
$\psi_6$	$x^3$	$x^2y$	2
$\psi_7$	$x^3y$	$y$	2
$\psi_8$	$x$	$x^2y$	2

$N$  has order 2 (excluding the identity map). Hence we obtain the following 4 groups:

$$\begin{aligned}
 G_1 &= \langle x, a \mid x^8 = a^2 = 1, a^{-1}xa = x \rangle \\
 &= \langle x \rangle \times \langle a \rangle \\
 &\cong C_8 \times C_2 \\
 G_2 &= \langle x, a \mid x^8 = a^2 = 1, a^{-1}xa = x^3 \rangle \\
 &= \langle x \rangle \rtimes_3 \langle a \rangle \\
 &\cong \text{SD}_{16} \\
 G_3 &= \langle x, a \mid x^8 = a^2 = 1, a^{-1}xa = x^5 \rangle \\
 &= \langle x \rangle \rtimes_5 \langle a \rangle \\
 &\cong \text{M}_{16} \\
 G_4 &= \langle x, a \mid x^8 = a^2 = 1, a^{-1}xa = x^{-1} \rangle \\
 &= \langle x \rangle \rtimes_{-1} \langle a \rangle \\
 &\cong D_{16}
 \end{aligned}$$

We are subscripting  $\rtimes$  with the power of  $x$  it gets sent to upon conjugation with  $a$ . The group  $\text{SD}_{16}$  is known as the semidihedral group<sup>4</sup>, of order 16, and  $\text{M}_{16}$  is called the modular group<sup>5</sup> of order 16.

#### Case 2: Order 4.

Assume all elements in  $G \setminus N$  have order at least 4, otherwise we are back in the first case. How does conjugation behave? In particular, what is  $a^{-1}ga$  for an element  $g \in G$ ? I claim the only possibility is the automorphism  $\phi_4 : x \mapsto x^{-1}$ . It cannot be  $\phi_2 : x \mapsto x^3$  because then:

$$(xa)(xa) = x(axa^{-1})a^2 = x(x^3)a^2 = x^4a^2 = 1$$

Likewise, it cannot be  $\phi_1 : x \mapsto x$  or  $\phi_3 : x \mapsto x^5$  because then  $x^2a$  will have order 2:

$$(x^2a)^2 = x^2(ax^2a^{-1})a^2 = x^2(x^2)^5a^2 = x^4a^2 = 1$$

4. Groupprops, “Semidihedral group:SD16,” January 20, 2013, accessed February 27, 2023, [https://groupprops.subwiki.org/wiki/Semidihedral\\_group:SD16](https://groupprops.subwiki.org/wiki/Semidihedral_group:SD16).

5. David Clausen, “Classifying All Groups of Order 16” (2012), accessed February 27, 2023, <http://buzzard.ups.edu/courses/2012spring/projects/clausen-groups-16-ups-434-2012.pdf>.

All of which contradict our assumption. Hence the only possibility for the effect of conjugation by  $a$  is the map  $\phi_4 : x \mapsto x^7 = x^{-1}$ . So we obtain the presentation:

$$G_5 = \langle x, a \mid x^8 = a^4 = 1, a^2 = x^4, a^{-1}xa = x^{-1} \rangle$$

Which we recognise as the dicyclic group,  $\text{Dic}_{24}$ .

**Case 3:** Order 8.

It turns out, we obtain no new groups in this case. We have two choices for  $a$ : either  $a^2 = x^2$  or  $a^2 = x^6$ . Because  $x^2$  and  $x^6$  are automorphic, we only need to consider one case, say  $a^2 = x^2$ , and apply Lemma 5. If conjugation by  $a$  is either  $\phi_2 : x \mapsto x^3$  or  $\phi_4 : x \mapsto x^7$  then we obtain a contradiction:

$$a^2 = a^{-1}a^2a = a^{-1}x^2a = x^6$$

$$a^2 = a^{-1}a^2a = a^{-1}x^2a = x^{14} = x^6$$

Hence the only possibilities for conjugation by  $a$  are the maps  $\phi_1 : x \mapsto x$  and  $\phi_3 : x \mapsto x^5$ . For the first:

$$(x^3a)(x^3a) = x^3(ax^3a^{-1})a^2 = x^3x^3a^2 = x^8 = 1$$

a contradiction. And the second:

$$(xa)(xa) = x(axa^{-1})a^2 = x(x^5)a^2 = x^8 = 1$$

another contradiction. Hence we obtain no new groups.

**Case 4:** Order 16.

The only possibility here is  $G_6 \cong C_{16}$ , generated by  $a$ .

Now let's move on to consider when  $\langle x, y \rangle = N \cong C_4 \times C_2$ . In particular, all elements of  $G$  have order less than 8.

I claim we can disregard automorphisms of order 4, namely  $\psi_2$  and  $\psi_4$ , because they lead to contradictions. Consider an element  $g \in N$ . We can write  $g = x^i y^j$  for some  $0 \leq i \leq 3$  and  $0 \leq j \leq 1$ . Then on the one hand:

$$a^{-2}ga^2 = g$$

Because  $N$  is abelian. On the other:

$$\begin{aligned} a^{-2}ga^2 &= a^{-1}(a^{-1}x^i y^j a)a \\ &= a^{-1}(x^i \psi_2 y^j \psi_2)a \\ &= a^{-1}(x^{-i} y^i x^{2j} y^j)a \\ &= a^{-1}(x^{2j-i} y^{i+j})a \\ &= x^{i-2j} y^{2j-i} x^{2i+2j} y^{i+j} \\ &= x^{3i} y^{3j} \\ &= y^j x^{-i} \end{aligned}$$

Which is  $g^{-1}$ . Hence we have  $g = g^{-1}$  which is a contradiction, because  $x \in N$  has order 4. A similar contradiction can be shown for  $\psi_4$ .

**Case 1:** Order 2.

If  $a$  has order 2, then  $a^2 = 1$ , so  $\langle a \rangle \cap N = \mathbf{1}$ . Hence  $G = N \rtimes \langle a \rangle$ . So we can apply Lemma 5, and only consider one representative from each conjugacy class. We have the following presentations:

$$\begin{aligned}
 G_7 &= \langle x, y, a \mid x^4 = y^2 = a^2 = 1, xy = yx, a^{-1}xa = x, a^{-1}ya = y \rangle \\
 &= \langle x \rangle \times \langle y \rangle \times \langle a \rangle \\
 &\cong C_4 \times C_2 \times C_2 \\
 G_8 &= \langle x, y, a \mid x^4 = y^2 = a^2 = 1, xy = yx, a^{-1}xa = x^{-1}, a^{-1}ya = y \rangle \\
 &= (\langle x \rangle \rtimes \langle a \rangle) \times \langle y \rangle \\
 &\cong D_8 \times C_2 \\
 G_9 &= \langle x, y, a \mid x^4 = y^2 = a^2 = 1, xy = yx, a^{-1}xa = xy, a^{-1}ya = y \rangle \\
 &= (\langle x \rangle \times \langle y \rangle) \rtimes \langle a \rangle \\
 &\cong (C_4 \times C_2) \rtimes C_2 \quad (\text{nameless}) \\
 G_{10} &= \langle x, y, a \mid x^4 = y^2 = a^2 = 1, xy = yx, a^{-1}xa = x^{-1}, a^{-1}ya = x^2y \rangle
 \end{aligned}$$

The final group,  $G_{10}$ , is sometimes called the Pauli group<sup>6</sup>, because it consists of the Pauli Matrices from quantum mechanics, and their products with powers of  $i$ . The group  $G_9$  doesn't appear to have any significant name.

**Case 2:** Order 4.

As before, assume all elements in  $G \setminus N$  have order at least 4. We have 3 choices for  $a$  here, and will break them into subcases:

**A:**  $a^2 = x^2$ .

Firstly, let's show what conjugation by  $a$  cannot be. If it's  $\phi_1$ :

$$(xa)(xa) = xa^2(a^{-1}xa) = xa^2x = 1$$

If it's  $\phi_6$ :

$$(xya)(xya) = xya^2(a^{-1}xya) = xya^2(x^{-1}x^2y) = x^4y^2 = 1$$

If it's  $\phi_8$ :

$$(x^2ya)(x^2ya) = x^2ya^2(a^{-1}x^2ya) = x^2ya^2(x^2x^2y) = x^8y^2 = 1$$

All of which are contradictions to our assumption. So the remaining possibilities are:

$$\begin{aligned}
 G_{11} &= \langle x, y, a \mid x^4 = y^2 = a^4 = 1, xy = yx, a^{-1}xa = x^{-1}, a^{-1}ya = y \rangle \\
 &= \langle x, a \rangle \times \langle y \rangle \\
 &\cong Q_8 \times C_2 \\
 G_{12a} &= \langle x, y, a \mid x^4 = y^2 = a^4 = 1, xy = yx, a^{-1}xa = xy, a^{-1}ya = y \rangle \\
 G_{12b} &= \langle x, y, a \mid x^4 = y^2 = a^4 = 1, xy = yx, a^{-1}xa = x^{-1}y, a^{-1}ya = y \rangle
 \end{aligned}$$

---

6. Clausen, "Classifying All Groups of Order 16."

Concentrating on the latter two groups, we will show they are both isomorphic to  $C_4 \rtimes C_4$ , with the inversion action. First,  $G_{12a}$ . Consider the element  $ax$ , and so  $\langle ax \rangle = \{1, ax, y, axy\}$ . By inspection,  $\langle x \rangle \cap \langle ax \rangle = \mathbf{1}$ . So then:

$$x^{-1}(ax)x = x^{-1}xaxy = axy = (ax)^{-1}$$

Hence,  $G_{12a} = \langle x \rangle \rtimes \langle ax \rangle \cong C_4 \rtimes C_4$ .

Likewise for  $G_{12b}$ ,  $\langle ax \rangle = \{1, ax, a^2y, x^{-1}a^{-1}\}$ , and again  $\langle x \rangle \cap \langle ax \rangle = \mathbf{1}$ . Additionally:

$$x^{-1}(ax)x = ax^3y = ax^{-1}y = xa$$

Multiplying by  $x^4 = 1$  gives:

$$x^4xa = x^{-1}x^2a = x^{-1}a^2a = x^{-1}a^{-1}$$

Hence  $G_{12b}$  is isomorphic to the same semidirect product,  $C_4 \rtimes C_4$ .

We can check that it is indeed valid to write  $C_4 \rtimes C_4$ . We know  $\text{Aut } C_4 \cong C_2$  and so a homomorphism  $\varphi : C_4 \rightarrow \text{Aut } C_4$  can map the generator to either the identity map or the inverse map. Hence we have only one non-trivial semidirect product.

**B:**  $a^2 = y$ .

Conjugation by  $a$  cannot be  $\psi_6$  or  $\psi_8$  because then:

$$a^2 = a^{-1}a^2a = a^{-1}ya = x^2y$$

a contradiction. If it's  $\psi_5$  then let  $\bar{a} = xa$  and so:

$$(xa)(xa) = x(axa^{-1})a^2 = x(x\psi_5)a^2 = x^2y^2 = x^2$$

and we are in the previous case. Likewise if conjugation by  $a$  is  $\psi_3$  then:

$$(x^2a)(x^2a) = x^2(ax^2a^{-1})a^2 = x^2(x^2\psi_3)a^2 = x^2y^2 = x^2$$

Finally,  $\psi_7$  gives:

$$(xa)(xa) = xa^2(a^{-1}xa) = xa^2x^3y = x^4y^2 = 1$$

a contradiction, leaving only  $\psi_1$  as the final standing possibility, giving:

$$G_{13} = \langle x, y, a \mid x^4 = y^2 = a^4 = 1, xy = yx, ax = xa, ay = ya \rangle$$

Letting  $b = xy$ :

$$G_{13} = \langle b, a \mid b^4 = a^4 = 1, ab = ba \rangle \cong C_4 \times C_4$$

**C:**  $a^2 = x^2y$ .

For any group in this case, if we apply the automorphism  $\psi_8$ , then we have:

$$a^2\psi_8 = (x^2y)\psi_8 = x^2x^2y = y$$

So it will be isomorphic to a group from the previous subcase.



Any group of order 16 is isomorphic to one of:

$(C_2)^4$	$C_8 \times C_2$	$SD_{16}$
$M_{16}$	$D_{16}$	$DiC_{24}$
$C_{16}$	$C_4 \times C_2 \times C_2$	$D_8 \times C_2$
$(C_4 \times C_2) \rtimes C_2$	Pauli Group	$Q_8 \times C_2$
$C_4 \rtimes C_4$	$C_4 \times C_4$	

Thus we finish not only the classification of groups of order 16, but this report as a whole.

# Bibliography

- Borcherds, Richard E. “Group theory 21: Groups of order 24,” June 30, 2020. Accessed February 9, 2023. <https://www.youtube.com/watch?v=6TWuo2NO8vg>.
- Clausen, David. “Classifying All Groups of Order 16.” 2012. Accessed February 27, 2023. <http://buzzard.ups.edu/courses/2012spring/projects/clausen-groups-16-ups-434-2012.pdf>.
- Groupprops. “Classification of groups of prime-cube order,” February 24, 2016. Accessed February 23, 2023. [https://groupprops.subwiki.org/wiki/Classification\\_of\\_groups\\_of\\_prime-cube\\_order](https://groupprops.subwiki.org/wiki/Classification_of_groups_of_prime-cube_order).
- . “Dicyclic Groups,” October 21, 2017. Accessed January 19, 2023. [https://groupprops.subwiki.org/wiki/Dicyclic\\_group](https://groupprops.subwiki.org/wiki/Dicyclic_group).
- . “Generalized dihedral group,” January 17, 2011. Accessed February 15, 2023. [https://groupprops.subwiki.org/wiki/Generalized\\_dihedral\\_group](https://groupprops.subwiki.org/wiki/Generalized_dihedral_group).
- . “Semidihedral group:SD16,” January 20, 2013. Accessed February 27, 2023. [https://groupprops.subwiki.org/wiki/Semidihedral\\_group:SD16](https://groupprops.subwiki.org/wiki/Semidihedral_group:SD16).
- . “Unitriangular matrix group:UT(3,p),” August 22, 2014. Accessed February 23, 2023. [https://groupprops.subwiki.org/wiki/Unitriangular\\_matrix\\_group:UT\(3,%20p\)](https://groupprops.subwiki.org/wiki/Unitriangular_matrix_group:UT(3,%20p)).
- Stack Exchange (user azimuth). “Classification of groups of order 30 (duplicate),” December 10, 2020. Accessed January 24, 2023. <https://math.stackexchange.com/questions/569226/classification-of-groups-of-order-30>.
- Wild, Marcel. “The Groups of Order Sixteen Made Easy.” The American Mathematical Monthly 112, no. 1 (2005): 20–31. ISSN: 00029890, 19300972, accessed February 27, 2023. <http://www.jstor.org/stable/30037381>.

# Appendix

Order	Classification	Presentation
1	<b>1</b>	$\langle 1 \rangle$
2	$C_2$	$\langle x \mid x^2 = 1 \rangle$
3	$C_3$	$\langle x \mid x^3 = 1 \rangle$
4	$C_4$ $C_2 \times C_2$	$\langle x \mid x^4 = 1 \rangle$ $\langle x, y \mid x^2 = y^2 = 1, xy = yx \rangle$
5	$C_5$	$\langle x \mid x^5 = 1 \rangle$
6	$D_6 \cong S_3$ $C_6$	$\langle x, y \mid x^3 = y^2 = 1, y^{-1}xy = x^{-1} \rangle$ $\langle x \mid x^6 = 1 \rangle$
7	$C_7$	$\langle x \mid x^7 = 1 \rangle$
8	$C_8$ $C_4 \times C_2$ $D_8$ $Q_8$ $C_2 \times C_2 \times C_2$	$\langle x \mid x^8 = 1 \rangle$ $\langle x, y \mid x^4 = y^2 = 1, xy = yx \rangle$ $\langle x, y \mid x^4 = y^2 = 1, y^{-1}xy = x^{-1} \rangle$ $\langle x, y \mid x^4 = y^4 = 1, x^2 = y^2, y^{-1}xy = x^{-1} \rangle$ $\langle x, y, z \mid x^2 = y^2 = z^2 = 1, xy = yx, xz = zx, yz = zy \rangle$
9	$C_9$ $C_3 \times C_3$	$\langle x \mid x^9 = 1 \rangle$ $\langle x, y \mid x^3 = y^3 = 1, xy = yx \rangle$
10	$D_{10}$ $C_{10}$	$\langle x, y \mid x^5 = y^2 = 1, y^{-1}xy = x^{-1} \rangle$ $\langle x \mid x^{10} = 1 \rangle$
11	$C_{11}$	$\langle x \mid x^{11} = 1 \rangle$
12	$\text{Dic}_{12}$ $C_{12}$ $A_4$ $D_{12}$ $C_6 \times C_2$	$\langle x, y \mid x^6 = 1, x^3 = y^2, y^{-1}xy = x^{-1} \rangle$ $\langle x \mid x^{12} = 1 \rangle$ — $\langle x, y \mid x^6 = y^2 = 1, y^{-1}xy = x^{-1} \rangle$ $\langle x, y \mid x^6 = y^2 = 1, xy = yx \rangle$
13	$C_{13}$	$\langle x \mid x^{13} = 1 \rangle$
14	$D_{14}$ $C_{14}$	$\langle x, y \mid x^7 = y^2 = 1, y^{-1}xy = x^{-1} \rangle$ $\langle x \mid x^{14} = 1 \rangle$
15	$C_{15}$	$\langle x \mid x^{15} = 1 \rangle$

Order	Classification	Presentation
16	$C_{16}$ $C_4 \times C_4$ $(C_4 \times C_2) \rtimes C_2$ $C_4 \rtimes C_4$ $C_8 \times C_2$ $M_{16}$ $D_{16}$ $SD_{16}$ $Dic_{16}$ $C_4 \times C_2 \times C_2$ $D_8 \times C_2$ $Q_8 \times C_2$ Pauli Group $(C_2)^4$	$\langle x \mid x^{16} = 1 \rangle$ $\langle x, y \mid x^4 = y^4, xy = yx \rangle$ $\langle x, y, z \mid x^4 = y^2 = z^2 = 1, xy = yx, z^{-1}xz = xy, z^{-1}yz = y \rangle$ $\langle x, y \mid x^4 = y^4 = 1, x^{-1}yx = y^{-1} \rangle$ $\langle x, y \mid x^8 = y^2 = 1, xy = yx \rangle$ $\langle x, y \mid x^8 = y^2 = 1, y^{-1}xy = x^5 \rangle$ $\langle x, y \mid x^8 = y^2 = 1, y^{-1}xy = x^{-1} \rangle$ $\langle x, y \mid x^8 = y^2 = 1, y^{-1}xy = x^3 \rangle$ $\langle x, y \mid x^8 = 1, x^4 = y^2, y^{-1}xy = x^{-1} \rangle$ $\langle x, y, z \mid x^4 = y^2 = z^2 = 1, xy = yx, xz = zx, yz = zy \rangle$ $\langle x, y, z \mid x^4 = y^2 = z^2 = 1, xy = yx, yz = zy, z^{-1}xz = x^{-1} \rangle$ $\langle x, y, z \mid x^4 = y^4 = z^2 = 1, xz = zx, yz = zy, y^{-1}xy = x^{-1} \rangle$ $\langle x, y, z \mid x^4 = y^2 = z^2 = 1, xy = yx, z^{-1}xz = x^{-1}, z^{-1}yz = x^2y \rangle$ —
17	$C_{17}$	$\langle x \mid x^{17} = 1 \rangle$
18	$D_{18}$ $C_{18}$ $D_6 \times C_3$ $Dih(C_p \times C_p)$ $C_6 \times C_3$	$\langle x, y \mid x^8 = y^2 = 1, y^{-1}xy = x^{-1} \rangle$ $\langle x \mid x^{18} = 1 \rangle$ $\langle x, y, z \mid x^3 = y^3 = z^2 = 1, xy = yx, yz = zy, z^{-1}xz = x^{-1} \rangle$ $\langle x, y, z \mid x^p = y^p = z^2 = 1, xy = yx, z^{-1}xz = x^{-1}, z^{-1}yz = y^{-1} \rangle$ $\langle x, y \mid x^6 = y^3 = 1, xy = yx \rangle$
19	$C_{19}$	$\langle x \mid x^{19} = 1 \rangle$
20	$Dic_{20}$ $C_{20}$ — $D_{20}$ $C_{10} \times C_2$	$\langle x, y \mid x^{10} = 1, x^5 = y^2, y^{-1}xy = x^{-1} \rangle$ $\langle x \mid x^{20} = 1 \rangle$ $\langle x, y \mid x^5 = y^4 = 1, y^{-1}xy = x^2 \rangle$ $\langle x, y \mid x^{10} = y^2 = 1, y^{-1}xy = x^{-1} \rangle$ $\langle x, y \mid x^{10} = y^2 = 1, xy = yx \rangle$
21	$C_7 \rtimes C_3$ $C_{21}$	$\langle x, y \mid x^7 = y^3 = 1, y^{-1}xy = x^4 \rangle$ $\langle x \mid x^{21} = 1 \rangle$
22	$D_{20}$ $C_{22}$	$\langle x, y \mid x^{11} = y^2 = 1, y^{-1}xy = x^{-1} \rangle$ $\langle x \mid x^{22} = 1 \rangle$
23	$C_{23}$	$\langle x \mid x^{23} = 1 \rangle$

Order	Classification	Presentation
24	$C_3 \rtimes C_8$	$\langle x, y \mid x^3 = y^8 = 1, x^{-1}yx = y^{-1} \rangle$
	$C_{24}$	—
	$\text{SL}_2(3)$	—
	$\text{Dic}_{24}$	$\langle x, y \mid x^{12} = 1, x^6 = y^2, y^{-1}xy = x^{-1} \rangle$
	$S_3 \times C_4$	—
	$D_{24}$	$\langle x, y \mid x^{12} = y^2 = 1, y^{-1}xy = x^{-1} \rangle$
	$\text{Dic}_{12} \times C_2$	$\langle x, y, z \mid x^6 = z^2 = 1, x^3 = y^2, y^{-1}xy = x^{-1}, xz = zx, yz = zy \rangle$
	$C_3 \rtimes_{V_4} D_8$	$\langle x, y, z \mid x^3 = y^4 = z^2 = 1, y^{-1}xy = x^{-1}, z^{-1}xz = x^{-1}, z^{-1}yz = y^{-1} \rangle$
	$C_3 \times C_4 \times C_2$	—
	$D_8 \times C_3$	—
	$Q_8 \times C_3$	—
	$S_4$	—
	$A_4 \times C_2$	—
	$S_3 \times C_2 \times C_2$	—
	$C_3 \times (C_2)^3$	—
25	$C_{25}$	—
	$C_5 \times C_5$	—
26	$D_{26}$	—
	$C_{26}$	—
27	$C_{27}$	—
	$C_9 \times C_3$	—
	$\text{UT}_3(3)$	—
	$C_9 \rtimes C_3$	—
	$(C_3)^3$	—
28	$\text{Dic}_{28}$	$\langle x, y \mid x^{14} = 1, x^7 = y^2, y^{-1}xy = x^{-1} \rangle$
	$C_{28}$	—
	$D_{28}$	—
	$C_{14} \times C_2$	—
29	$C_{29}$	—
30	$C_5 \times D_6$	—
	$C_3 \times D_{10}$	—
	$D_{30}$	—
	$C_{30}$	—
31	$C_{31}$	—