

# Interim Report

Daniel Laing

January 20, 2023

## Contents

<b>1</b>	<b>Theorems and Lemmas</b>	<b>1</b>
1.1	Sylow Theorems . . . . .	1
<b>2</b>	<b>Groups of order 6</b>	<b>2</b>
2.1	$C_3 \rtimes_{\text{id}} C_2$ . . . . .	2
2.2	$C_3 \rtimes_{\varphi} C_2$ . . . . .	2
<b>3</b>	<b>Groups of Order 6 (Attempt 2)</b>	<b>3</b>
<b>4</b>	<b>Generalisation to Groups of Order <math>2p</math></b>	<b>3</b>
<b>5</b>	<b>Groups of order 4</b>	<b>4</b>
<b>6</b>	<b>Groups of order 9 (Might skip)</b>	<b>4</b>
<b>7</b>	<b>Generalisation to Groups of Order <math>p^2</math></b>	<b>4</b>
<b>8</b>	<b>Groups of order 12</b>	<b>4</b>

## 1 Theorems and Lemmas

### 1.1 Sylow Theorems

Let  $G$  be a group of order  $p^n m$  where  $p$  is a prime and  $p \nmid m$ .

**Theorem 1.1** (1<sup>st</sup> Sylow Theorem).  *$G$  has a Sylow  $p$ -subgroup, i.e. a subgroup of order  $p^n$ .*

**Theorem 1.2** (2<sup>nd</sup> Sylow Theorem). *All Sylow  $p$ -subgroups of  $G$  are conjugate to each other.*

**Corollary 1.2.1.** *If  $n_p = 1$  then the Sylow  $p$ -subgroup is normal in  $G$ .*

**Theorem 1.3** (3<sup>rd</sup> Sylow Theorem). *Let  $n_p$  denote the number of Sylow  $p$ -subgroups of  $G$ . Then:*

i)  $n_p \mid m$

ii)  $n_p \equiv 1 \pmod{p}$

**Lemma 1.4.** *For a group  $G$  with  $N \leq G$  and  $H \leq G$ , then*

$$|NH| = |\{nh \mid n \in N, h \in H\}| = \frac{|N| \cdot |H|}{|N \cap H|}$$

**Lemma 1.5.** *The automorphism group of  $C_n$  is isomorphic to the multiplicative group of integers mod  $n$ .*

*i.e.*  $\text{Aut } C_n \cong \mathbb{Z}/n\mathbb{Z}^*$

*Proof.* Any automorphism,  $\varphi$  of  $C_n$  has the property:

$$(x^i)\varphi = (x\varphi)^i$$

Hence  $\varphi$  is determined by its effect on the generator,  $x$ , and preserves element order. In particular,  $\varphi$  sends generators to generators.

So for a generator,  $x$ ,  $x\varphi = x^k$  is surjective if  $x^k$  generates  $C_n$ .  $x^k$  generates  $C_n$  if  $o(x^k) = n$  which is when  $\gcd(n, k) = 1$ .

Denote  $\varphi_k : x \mapsto x^k$ .

Consider:

$$x\varphi_k\varphi_l = (x^k)\varphi_l = (x^k)^l = x^{kl} = x^{lk} = (x^l)^k = x\varphi_l\varphi_k$$

So we see that  $\text{Aut } C_n$  is abelian. Moreover,  $x\varphi_k\varphi_l = x\varphi_{kl}$ .

Now consider  $\theta : \text{Aut } C_n \rightarrow \mathbb{Z}/n\mathbb{Z}^*$  defined by  $\varphi_k\theta \mapsto k$ . We will show  $\theta$  is an isomorphism.

$\theta$  is surjective because every  $k \in \mathbb{Z}/n\mathbb{Z}^*$  is coprime to  $n$  and so  $x^k$  is a generator of  $C_n$ , hence  $\exists \varphi_k \in \text{Aut } C_n$  such that  $\varphi_k\theta = k$ .

$\theta$  is also injective because if  $\varphi_k, \varphi_l \in \text{Aut } C_n$  such that  $\varphi_k\theta = \varphi_l\theta$  then  $k = l$ .

Finally,  $\theta$  is a homomorphism because  $(\varphi_k\varphi_l)\theta = \varphi_{kl}\theta = kl = (\varphi_k\theta)(\varphi_l\theta)$ . So  $\theta : \text{Aut } C_n \rightarrow \mathbb{Z}/n\mathbb{Z}^*$  is an isomorphism. □

## 2 Groups of order 6

The prime factorisation of  $6 = 2 \cdot 3$ , so we can construct groups with products of  $C_2$  and  $C_3$ . The automorphism groups of  $C_2$ ,  $\text{Aut } C_2 = \{\text{id}\}$ , containing just the identity map. So any meaningful products will look like  $C_3 \rtimes C_2$ .

$\text{Aut } C_3 = \{\text{id}, \psi\}$  where  $x\psi = x^{-1}$ . So we have two possible products:  $C_3 \rtimes_{\text{id}} C_2$  and  $C_3 \rtimes_{\varphi} C_2$  where  $\varphi$  is the homomorphism  $\varphi : C_2 \rightarrow \text{Aut } C_3$  mapping  $1 \mapsto \text{id}$  and  $x \mapsto \psi$ .

### 2.1 $C_3 \rtimes_{\text{id}} C_2$

By the Fundamental Theorem of Finite Abelian Groups, we know  $C_3 \rtimes_{\text{id}} C_2 \cong C_3 \times C_2 \cong C_6$ .

### 2.2 $C_3 \rtimes_{\varphi} C_2$

So the group operation is

$$(a_1, b_1)(a_2, b_2) = (a_1 \cdot a_2\varphi_{b_1}, b_1 \cdot b_2)$$

Investigating elements:

$$(1, x)(1, x) = (1 \cdot 1\varphi_x, x \cdot x) = (1 \cdot 1^{-1}, x^2) = (1, 1)$$

So  $(1, x)$  is of order 2.

$$(x, 1)(x, 1) = (x \cdot x\varphi_1, 1 \cdot 1) = (x \cdot x, 1) = (x^2, 1)$$

$$(x^2, 1)(x, 1) = (x^2 \cdot x\varphi_1, 1 \cdot 1) = (x^3, 1) = (1, 1)$$

So  $(x, 1)$  is of order 3.

$$(x, 1)(x, x) = (x \cdot x\varphi_1, 1 \cdot x) = (x, x)$$

$$(x, x)(x, 1) = (x \cdot x\varphi_x, x \cdot 1) = (xx^{-1}, x) = (1, x)$$

Hence,  $C_3 \rtimes_{\varphi} C_2$  is non-abelian.

### 3 Groups of Order 6 (Attempt 2)

Let  $G$  be a group of order 6, and  $n_3$  denote the number of Sylow 3-subgroups of  $G$ . Then by Theorem 1.3:

$$n_3 \equiv 1 \pmod{3} \text{ and } n_3 \mid 2 \implies n_3 = 1$$

So  $G$  has one Sylow 3-subgroup, and because 3 is prime, it is isomorphic to  $C_3$ , i.e.

$$C_3 \trianglelefteq G$$

Any Sylow 2-subgroup of  $G$  will have order 2, and so  $C_2 \leq G$ .

Lagrange's Theorem tells us that  $C_3$  has elements of orders 1 and 3, and  $C_2$  has elements of order 1 and 2 hence:

$$C_3 \cap C_2 = \mathbf{1}$$

By Lemma 1.4:

$$|C_3 C_2| = \frac{|C_3| \cdot |C_2|}{|C_3 \cap C_2|} = \frac{3 \cdot 2}{1} = 6$$

So  $G = C_3 C_2$ ,  $C_3 \trianglelefteq G$  and  $C_3 \cap C_2 = \mathbf{1} \implies G = C_3 \rtimes C_2$

Now we need to determine  $\text{Aut } C_3$ .  $C_3 = \{1, x, x^2 = x^{-1}\}$  and so  $\text{Aut } C_3 = \{\text{id}, \psi : x \mapsto x^{-1}\} \cong C_2$ . So if  $C_3 = \langle x \rangle$  and  $C_2 = \langle y \rangle$ , then we have two possibilities for  $G$ :

**Case 1:**

$$\begin{aligned} G &= \langle x, y \mid x^3 = y^2 = 1, y^{-1}xy = x \rangle \\ &= \langle x, y \mid x^3 = y^2 = 1, xy = yx \rangle \\ &= C_3 \times C_2 \cong C_6 \end{aligned}$$

**Case 2:**

$$\begin{aligned} G &= \langle x, y \mid x^3 = y^2 = 1, y^{-1}xy = x^{-1} \rangle \\ &\cong D_6 \end{aligned}$$

Hence  $G$  is isomorphic to either  $C_6$  or  $D_6$ .

### 4 Generalisation to Groups of Order $2p$

Let  $G$  be a group of order  $2p$  where  $p$  is a prime number, and  $n_p$  denote the number of Sylow  $p$ -subgroups of  $G$ . Then by Theorem 1.3:

$$n_p \equiv 1 \pmod{p} \text{ and } n_p \mid 2 \implies n_p = 1$$

So  $G$  has one Sylow  $p$ -subgroup, it is isomorphic to  $C_p = \langle x \rangle$  hence:

$$C_p \trianglelefteq G$$

A Sylow 2-subgroup of  $G$  will have order 2 so  $C_2 = \langle y \rangle \leq G$ .

Lagrange's Theorem tells us that  $C_p$  has elements of orders 1 and  $p$ , and  $C_2$  has elements of order 1 and 2 hence:

$$C_p \cap C_2 = \mathbf{1}$$

By Lemma 1.4:

$$|C_p C_2| = \frac{|C_p| \cdot |C_2|}{|C_p \cap C_2|} = \frac{p \cdot 2}{1} = 2p$$

So  $G = C_p C_2$ ,  $C_p \trianglelefteq G$  and  $C_p \cap C_2 = \mathbf{1} \implies G = C_p \rtimes C_2$

We want a homomorphism  $\varphi : \text{Aut } C_p \rightarrow C_2$ . By Lemma 1.5,  $\text{Aut } C_p \cong \mathbb{Z}/p\mathbb{Z}^*$ , so now we need to find elements of order 2 in  $\mathbb{Z}/p\mathbb{Z}^*$ .

An element  $x \in \mathbb{Z}/p\mathbb{Z}^*$  of order 2 satisfies:

$$x^2 = 1 \implies x^2 - 1 = 0 \implies (x - 1)(x + 1) = 0$$

Hence  $x = 1$  or  $-1$ . But 1 has order 1 so  $x$  can only be  $-1$ .

So  $C_2\varphi$  could be  $\mathbf{1}$  or  $\langle \beta : x \mapsto x-1 \rangle$ . This gives us two possibilities:

$$y\varphi = x \mapsto x \quad \text{or} \quad y\varphi = x \mapsto x^{-1}$$

**Case 1:**

$$\begin{aligned} G &= \langle x, y \mid x^p = y^2 = 1, y^{-1}xy = x \rangle \\ &= \langle x, y \mid x^p = y^2 = 1, xy = yx \rangle \\ &= C_p \times C_2 \cong C_{2p} \end{aligned}$$

**Case 2:**

$$\begin{aligned} G &= \langle x, y \mid x^p = y^2 = 1, y^{-1}xy = x^{-1} \rangle \\ &\cong D_{2p} \end{aligned}$$

Hence a group of order  $2p$  is isomorphic to  $C_{2p}$  or  $D_{2p}$ .

## 5 Groups of order 4

From the Fundamental Theorem of Finite Abelian Groups, we know that a group  $G$  of order 4 could be isomorphic to one of  $C_4$  and  $C_2 \times C_2$ . Now to show that these are the only possibilities, i.e. a group of order 4 must be abelian.

The Sylow theorems are not so helpful here, because  $4 = 2^2$  so any Sylow 2-subgroup will be of order 4, which is just  $G$ .

## 6 Groups of order 9 (Might skip)

## 7 Generalisation to Groups of Order $p^2$

## 8 Groups of order 12

Let  $G$  be a group of order  $12 = 2^2 \cdot 3$ , and  $n_3$  and  $n_2$  denote the number of Sylow 3-subgroups and Sylow 2-subgroups of  $G$  respectively. By Theorem 1.3:

$$n_2 \equiv 1 \pmod{2} \text{ and } n_2 \mid 3 \implies n_2 = 1$$

$$n_3 \equiv 1 \pmod{3} \text{ and } n_3 \mid 4 \implies n_3 = 1 \text{ or } 4$$

$G$  has a unique Sylow 2-subgroup of order  $2^2 = 4$ , say  $H \trianglelefteq G$ , and we have already classified groups of order 4, so either  $C_4$  or  $V_4 \trianglelefteq G$ . A Sylow 3-subgroup of  $G$  will have order 3, so  $C_3 \leq G$ .

Lagrange's Theorem tells us  $H$  has elements of order 1, 2, and 4, and  $C_3$  has elements of order 1 and 3. Hence  $H \cap C_3 = \mathbf{1}$ .

Lemma 1.4 tells us:

$$|HC_3| = \frac{|H| \cdot |C_3|}{|H \cap C_3|} = 12$$

Hence  $G = HC_3$ ,  $C_3 \leq G$ ,  $H \trianglelefteq G$ , and  $H \cap C_3 = \mathbf{1} \implies G = H \rtimes C_3$ .

Since an automorphism,  $\varphi$ , must map generators to generators,  $\text{Aut } C_4 \cong C_2$  because the generators of  $C_4$  are  $x$  and  $x^{-1}$ . An automorphism of  $V_4$  corresponds to a permutation of the three non-identity elements, hence  $\text{Aut } V_4 \cong S_3$ .

**Case 1:**  $H = C_4$ .

A homomorphism  $\psi : C_3 \rightarrow \text{Aut } C_4 \cong C_2$ , preserves order and together with Lagrange's Theorem means that the only possibility for  $\psi$  is trivial, i.e.  $C_3\psi = \mathbf{1}$ .

Hence  $G \cong C_4 \times C_3 \cong C_{12}$ .

**Case 2:**  $H = V_4$ .

$\text{Aut } V_4 \cong S_3$  has 2 elements of order 3, giving us 3 possibilities for a homomorphism  $\psi : C_3 \rightarrow \text{Aut } V_4$ :

$$C_3\psi = \mathbf{1}, \quad C_3\psi = (a, b, c) \quad \text{and} \quad C_3\psi = (a, c, b)$$

If  $C_3\psi = \mathbf{1}$ , then  $G \cong C_2 \times C_2 \times C_3 \cong C_2 \times C_6$ .

I don't how to do the other two cases. Something like:

$$G = \langle a, b, x \mid x^3 = a^2 = b^2 = (ab)^2 = 1, \ x^{-1}ax = b, \ x^{-1}bx = c, \ x^{-1}cx = a \rangle$$

$$G = \langle a, b, x \mid x^3 = a^2 = b^2 = (ab)^2 = 1, \ x^{-1}ax = c, \ x^{-1}bx = a, \ x^{-1}cx = b \rangle$$