

# Quasicategories talk

Bogdan Simeonov

# Contents

<b>1</b>	<b>Simplicial sets basics</b>	<b>3</b>
1.1	Definition . . . . .	3
1.2	Yoneda and the standard simplices . . . . .	4
1.3	Nerves, total singular complex and the geometric realization . . . . .	5
1.4	Horn fillings and quasicategories . . . . .	6
<b>2</b>	<b>Towards the infinity category of spaces</b>	<b>8</b>
2.1	Functor categories . . . . .	8
2.2	Mapping spaces . . . . .	13
2.3	The homotopy coherent nerve . . . . .	14
2.4	Infinity Yoneda . . . . .	16
<b>3</b>	<b>Limits and colimits</b>	<b>17</b>

We think of an infinity category as a category whose morphism spaces are actually spaces, such that there are coherent higher composition laws. One possible way of modelling such things is using the notion of a simplicial set.

Resources: Emily Riehl's notes on simplicial sets: <https://math.jhu.edu/~eriehl/ssets.pdf>

# 1 Simplicial sets basics

## 1.1 Definition

We first define  $\Delta$  as the category which has objects  $[n] = \{0, 1, \dots, n\}$  and morphisms which are order-preserving set maps.

**Definition 1.1 (Simplicial set):** A simplicial set is a presheaf on  $\Delta$ , in other words a functor

$$X : \Delta^{op} \rightarrow \mathbf{Set}$$

We denote  $X([n]) = X_n$ .

We notice that the category  $\Delta$  comes with certain natural maps which generate it: namely, the maps

$$d^i : [n-1] \rightarrow [n], \text{ 'misses } i \text{'}$$

$$s^i : [n+1] \rightarrow [n], \text{ 'squishes down } i, i+1 \text{'}$$

These operations satisfy the properties

$$d^j d^i = d^i d^{j-1}, i < j$$

$$s^j s^i = s^i s^{j+1}, i \leq j$$

$$s^j d^i = \begin{cases} 1, i = j, j+1 \\ d^i s^{j-1}, i < j \\ d^{i-1} s^j, i > j+1 \end{cases}$$

Every map can be factorized using the (co)face and (co)degeneracy maps, and hence we can think of a simplicial set  $X$  as a module over the algebra generated by these relations. They define face and degeneracy maps on  $X$  as follows:

$$d_i = X d^i : X_n \rightarrow X_{n-1}, s_i = X s^i : X_n \rightarrow X_{n+1}$$

Sometimes, a simplicial set is drawn as follows:

$$X_0 \rightleftarrows X_1 \rightleftarrows X_2 \dots$$

We see that a map of simplicial sets  $X \rightarrow Y$ , which is a natural transformation, can also be seen as a collection of maps  $X_n \rightarrow Y_n$  which commutes with the face and degeneracy maps.

We say that a simplex  $x \in X_n$  is *degenerate* if it comes from one of the degeneracy maps.

## 1.2 Yoneda and the standard simplices

Recall that the Yoneda embedding

$$\mathcal{Y} : \mathcal{C} \rightarrow \mathbf{Set}^{\mathcal{C}^{op}}, c \mapsto \mathcal{C}(-, c)$$

is a full and faithful embedding such that

$$\mathrm{Hom}(\mathcal{Y}c, F) \simeq F(c)$$

We denote the image under Yoneda of  $[n]$  by  $\Delta^n$ . These are precisely the representable simplicial sets, and every other one can be built from them by what is called the *density theorem*. We see that

$$\mathrm{Hom}_{\mathbf{sSet}}(\Delta^n, X) = X_n$$

More explicitly, the  $m$  simplices of  $\Delta^n$  are (by the Yoneda embedding)

$$\Delta_m^n = \mathrm{Hom}([m], [n])$$

Importantly, the action of the face and degeneracy maps on a morphism  $[k] \rightarrow [n]$  is composition on the left:

$$[k] \rightarrow [n] \mapsto [k-1] \rightarrow [k] \rightarrow [n]$$

or

$$[k] \rightarrow [n] \mapsto [k+1] \rightarrow [k] \rightarrow [n]$$

The nondegenerate simplices are essentially the injective maps  $[m] \rightarrow [n]$ . The unique nondegenerate  $n$ -simplex comes from the identity map. If we identify  $x \in X_n$  as a map  $x : \Delta^n \rightarrow X$ , then

$$d_i(x) \in X_{n-1} \leftrightarrow \Delta^{n-1} \xrightarrow{d^i} \Delta^n \xrightarrow{x} X$$

Analogously to how CW complexes are built out of standard simplices, we have the following:

**Theorem 1.2 (Density theorem):** *Every simplicial set is a colimit of the standard  $n$ -simplices:*

$$\varinjlim_{x \in X_n} \Delta^n \simeq X$$

Essentially, every  $x \in X_n$  contributes a  $x : \Delta^n \rightarrow X$ , but these are glued along in some combinatorial way. The index over which we take the colimit is the category of ends, which has objects  $x \in X_n$  for some  $n$  and morphisms  $f : x \rightarrow y$  precisely when this is induced by some map  $[n] \rightarrow [m]$ , taking  $x$  to  $y$ .

### 1.3 Nerves, total singular complex and the geometric realization

*Example 1.3 (Nerve of a category):* Given a category  $\mathcal{C}$ , we define a simplicial set  $\mathcal{NC}$  as

- $\mathcal{NC}_0 = \text{ob } \mathcal{C}$
- $\mathcal{NC}_1 = \text{mor } \mathcal{C}$
- $\mathcal{NC}_2 = \text{pairs of composable arrows}$
- ...

The degeneracy maps send a string of  $n$  composable arrows to  $n + 1$  composable arrows by inserting the identity map somewhere. The face maps compose two consecutive maps. This is a simplicial set, and in fact a quasicategory (we will see what this means in a moment)

Now we consider the geometric realization of a simplicial set.

We first define

$$\Delta : \Delta \rightarrow \mathbf{Top}$$

by  $\Delta([n]) = \Delta_n$ , the standard  $n$ -simplex. We would like to extend this, via some sort of Kan extension, to a functor on  $\mathbf{sSet}$ :

$$\begin{array}{ccc} & \mathbf{sSet} & \\ \nearrow \mathcal{Y} & & \searrow \mathcal{A} \\ \Delta & \xrightarrow{\Delta} & \mathbf{Top} \end{array}$$

The density theorem will allow us to do this:

*Example 1.4 (The geometric realization):* Given a simplicial set  $X$  we define

$$|X| := \frac{\left( \coprod X_n \times |\Delta^n| \right)}{\sim}$$

where  $|\Delta^n| = \Delta_n$  and the equivalence relation identifies the  $i$ th face of  $\{x\} \times \Delta_n$  with  $d_i(x) \times \Delta_{n-1}$  and also collapses  $\{s_i(x)\} \times \Delta_n$  to  $\{x\} \times \Delta_{n-1}$ . Using the description of Kan extensions and coends, one also realizes this as

$$|X| = \int^n X_n \times \Delta_n = \text{colim} \left( \coprod_f X_m \cdot \Delta_n \xrightarrow[\mathcal{F}]{\mathcal{F}_*} \coprod_{[n]} X_n \cdot \Delta_n \right)$$

Interestingly, the Kan complexes are determined by their geometric realization, up to homotopy!

This is the left adjoint to the functor  $S$  taking a topological space  $Y$  to its total singular complex  $SY$ : given any topological space  $Y$ , we can define

$$SY_n = \mathbf{Top}(\Delta_n, Y)$$

as the space of maps from the  $n$ -simplex to  $Y$ . Pre-composition with the maps which insert a zero in some coordinate, or add the  $i$ -th and  $i+1$ -th coordinates, produce maps which satisfy the simplicial set relations. We get the adjunctions

$$\mathbf{sSet} \begin{array}{c} \xrightarrow{|-|} \\ \xleftarrow{s} \end{array} \mathbf{Top}$$

These in fact form what is called a *Quillen equivalence of model categories*: the homotopy categories of both sides are the same, and the Kan complexes correspond to topological spaces.

*Remark 1.5 (Adjoint of Kan extensions):* This fits into a more general picture: whenever we left Kan extend  $F : \Delta \rightarrow \mathcal{E}$  via Yoneda, we get a right adjoint

$$R : \mathcal{E} \rightarrow \mathbf{sSet}, Re_n = \mathcal{E}(F[n], e)$$

These two functors are extremely useful. For example, the classifying space of a group  $G$  can be constructed by taking the nerve of the category defined by  $G$  and then its geometric realization. Similarly, if one takes the geometric realization of  $SY$ , then it forms a CW complex whose cellular homology is the same as the singular homology of  $Y$ :

$$C_{\bullet}^{\text{cell}}(|SY|) \simeq C_{\bullet}^{\text{sing}}(Y)$$

*Example 1.6 (Internal homs):*

We have a functor

$$\Delta \rightarrow \mathbf{sSet}$$

given by taking the product with a fixed  $Y$ ,  $F[n] = \Delta^n \times Y$ . The Kan extension along the Yoneda embedding is naturally the functor  $L = - \times Y$ . This has as a right adjoint the functor defined by

$$RZ_n = \mathbf{sSet}(\Delta^n \times Y, Z) := [Y, Z]_n$$

The face and degeneracy maps are the ones induced by postcomposition from the ones on  $\Delta^n$ .

## 1.4 Horn fillings and quasicategories

We study some simplicial subsets of  $\Delta^n$ . Recall that we have an element

$$d^i \in \Delta_{n-1}^n = \text{Hom}([n-1], [n])$$

which omits  $i$ . We think of this as an element in the  $n-1$  part of the simplicial set  $\Delta^n$ . This then generates a subsimplicial set. Note that by Yoneda, the map  $d^i$  corresponds to an embedding

$$\Delta^{n-1} \rightarrow \Delta^n$$

and we call the image  $\partial_i \Delta^n$ .

The union of all the boundaries we call  $\partial\Delta^n$  and it can be represented as the subsimplicial set defined as

$$(\partial\Delta^n)_m = \{\alpha \in \text{Hom}([m], [n]) \mid \alpha \text{ not surjective}\}$$

Why? Well, note that what the  $d^i$  generate is precisely all the morphisms one can get by composing a bunch of morphisms  $d^j, s^j$ , one of which is  $d^i : [n-1] \rightarrow [n]$ . But any non-surjective morphism must miss some  $i$ , so factorizes through one of these  $d^i$ . The converse is obviously also true: any composition involving  $d^i$  is nonsurjective.

Similarly, we can define the  $i$ -th horn

$$(\Lambda_i^n)[m] = \{\alpha \mid [n] \not\subseteq \alpha([m]) \cup \{i\}\}$$

or equivalently as the subsimplicial set defined by all faces except the  $i$ -th one, since the morphisms generated by composing with anything but the  $d^i$  will always hit  $i$  and miss something else.

For example, a filling in of  $\Lambda_1^2$  would constitute a 2-simplex  $\sigma$  such that  $d_2\sigma = f, d_0\sigma = g$ .

**Definition 1.7 (Kan complexes and quasicategories):** A Kan complex is a simplicial set for which every horn admits a filling. An infinity category is a simplicial set  $X$  such that any inner horn  $0 < i < n$  can be filled.

**Proposition 1.8 (Singular chains are Kan complexes):** For a topological space  $Y$ , its singular chain  $SY$  is a Kan complex.

*Proof.* By adjunction, we turn the left diagram into the right one, but since  $|\Lambda_k^n|$  is a homotopy retract of  $|\Delta^n|$ , we have a lift.

$$\begin{array}{ccc}
 & \text{adjunction} & \\
 & \text{~~~~~} & \\
 \Lambda_k^n & \xrightarrow{\quad} & SY \\
 \downarrow & \nearrow \text{dashed} & \\
 \Delta^n & & 
 \end{array}
 \qquad
 \begin{array}{ccc}
 |\Lambda_k^n| & \xrightarrow{\quad} & Y \\
 \downarrow & \nearrow \text{dashed} & \\
 |\Delta^n| & & 
 \end{array}$$

□

**Proposition 1.9 (Nerves are quasicategories):** The nerve of a category is an infinity category, in which every inner horn has a unique lift.

*Proof.* Essentially, every sequence of composable arrows has unique composition. Or, we can use the fact that the functor  $\tau_1$  (the homotopy category) is adjoint to the nerve functor. □

We see then that all usual categories sit inside the quasicategories, but have unique compositions. An infinity category can be thought of as a normal category, but where composition laws are not uniquely defined: there are higher relations among them! We will see, though, that even though compositions are not uniquely defined, they still form a contractible set, so in homotopical terms is still a point.

**Definition 1.10 (Homotopy equivalence):** Suppose  $f, g \in X_1$  are morphisms from  $a$  to  $b$ . We have different relations of equivalence  $f \sim_a g, f \sim_b g, g \sim_a f, g \sim_b f$  when there is a triangle filling in  $f, g$  and an identity morphisms. These are all equivalent when  $X$  is a quasicategory.

## 2 Towards the infinity category of spaces

The aim of this chapter is two-fold. First, we introduce functor categories, inner fibrations and mapping spaces. Then, we use the homotopy coherent nerve construction and apply it to the simplicially enriched category of Kan complexes to produce the infinity category of spaces.

### 2.1 Functor categories

Recall that we have an internal hom

$$\underline{\mathrm{Hom}}(X, Y)_n = \mathrm{Hom}_{\mathbf{sSet}}(X \times \Delta^n, Y)$$

So for an infinity category  $C$  and a simplicial set  $K$ , we define

$$\mathrm{Fun}(K, C) := \underline{\mathrm{Hom}}(K, C)$$

Some examples of functor categories:

- $\mathrm{Fun}(\Delta^0, C) \simeq C$
- $\mathrm{Fun}(\Delta^1, C)$  is the arrow category
- $\mathrm{Fun}(\Delta^1 \times \Delta^1, C)$  are the commutative squares in  $C$ , as follows: we have 2-simplices such that  $d_1(\sigma) = d_1(\tau)$ .

To clarify the last point: the product  $\Delta^1 \times \Delta^1$  has 4 0-simplices  $(0,0), (0,1), (1,0), (1,1)$ . The 1 simplices are pairs of 1 simplices of  $\Delta^1$ , which are enumerated by  $\{id, 0, 1\}$ , the identity being the nondegenerate one. However, whilst  $0 \times 0, 0 \times 1, 1 \times 0, 1 \times 1$  are degenerate (come from the zero simplices by precomposing with  $\Delta^1 \rightarrow \Delta^0$ ), the ones where we multiply with  $id$  are not and there are five of them:  $id \times id$  being the diagonal, the rest being the sides of a square. The only nondegenerate 2-simplices come from the different degeneracy maps  $[2] \rightarrow [1]$ , which give the two triangles. These 2-simplices witness the fact that the top and bottom 'compositions' are the same!






Figure 1: Nondegenerate 2-simplex

**Definition 2.1 (Equivalence of infinity categories):** Let  $f : C \rightarrow D$  be a functor between infinity categories. This forms an equivalence if it is an isomorphism in  $\text{Fun}(C, D)$ , i.e.  $\exists g : D \rightarrow C$  so that  $f \circ g \simeq \text{id}$ ,  $g \circ f \simeq \text{id}$ .

This is weaker than a natural isomorphism! For nerves, it means equivalence of 1-cats. For spaces, it means a weak homotopy equivalence. For infinity cats, it means "fully faithful and essentially surjective".

Recall:  $C$  being an infinity category is equivalent to

$$\text{Hom}_{\text{sSet}}(\Delta^n, C) \rightarrow \text{Hom}_{\text{sSet}}(\Lambda_i^n, C)$$

being a surjection of sets of outer hom. But if we think of internal hom instead we actually get the equivalent statement that

$$\underline{\text{Hom}}(\Delta^n, C) \rightarrow \underline{\text{Hom}}(\Lambda_i^n, C)$$

is an epimorphism of simplicial sets! Even more is true: this turns out to be a trivial Kan fibration.

We now turn to the claim that when  $X$  is an infinity category, then so is  $\text{Fun}(K, X)$ . The idea is as follows: want to show inner horns extend to  $n$ -simplex. By definition, the inner hom is adjoint to the product with  $K$ , this is equivalent to a map

$$\Lambda_i^n \rightarrow \underline{\text{Hom}}(K, C) \leftrightarrow \Lambda_i^n \times K \rightarrow C$$

and extending it to  $\Delta^n \times K$ . We add in the structure map to the terminal object:

$$\begin{array}{ccc} \Lambda_i^n \times K & \longrightarrow & C \\ \downarrow & & \downarrow \\ \Delta^n \times K & \longrightarrow & \Delta^0 \end{array}$$

We seek a diagonal lift in this problem.

**Definition 2.2 (Inner fibrations and inner anodyne maps):** An inner fibration is a map  $p : X \rightarrow Y$  with the right lifting property with respect to inner horns, i.e. every diagram like this has a lift:

$$\begin{array}{ccc} \Lambda_i^n & \longrightarrow & X \\ \downarrow & & \downarrow p \\ \Delta^n & \longrightarrow & Y \end{array}$$

In this sense, an infinity category is just an inner fibration  $C \rightarrow \Delta^0$ . An inner anodyne map is a map  $q : A \rightarrow B$  with the left lifting property with respect to inner fibrations.

**Example 2.3 (Functors to nerves):** Any functor  $C \rightarrow ND$  from a quasicategory to a nerve is an inner fibration. First, let  $f : \Lambda_i^n \rightarrow C$  be the top arrow.

$$\begin{array}{ccc} \Lambda_i^n & \xrightarrow{f} & C \\ \downarrow & \nearrow \tilde{f} & \downarrow g \\ \Delta^n & \longrightarrow & ND \end{array}$$

We see that the composition  $g \circ f$  has a unique lift to  $ND$  which is the bottom map, since nerves admit unique lifts. On the other hand, since  $C$  is an infinity category, there is some lift  $\tilde{f} : \Delta^n \rightarrow C$ . The reason this commutes with the bottom arrow is because  $g \circ \tilde{f} \circ \iota = g \circ f$  satisfies the uniquely defining property of the map  $\Delta^n \rightarrow ND$ .

**Definition 2.4 (Trivial Kan fibration):** A map  $p : X \rightarrow Y$  is a trivial Kan fibration if it has the right lifting property with respect to all inclusions of simplicial sets. These are obviously a subset of the inner fibrations. They also admit sections, by taking the inclusion  $\emptyset \rightarrow Y$  and the identity map  $Y \rightarrow Y$ .

The Kan fibrations form the *fibrations in the Quillen model structure*, and the monomorphisms form the cofibrations. Trivial Kan fibrations are stable under pullbacks, by using UMP for fiber products.

So it amounts to showing that  $\text{Fun}(K, -)$  preserves inner fibrations, which is also equivalent to the fact that inner anodyne maps are stable under  $- \times K$ . This is a standard result in the literature. We mention the theorem which allows such arguments to work (Theorem 1.3.37 in Markus Land's book):

**Theorem 2.5 (Stability of inner fibrations):** Suppose we are given maps  $f : X \rightarrow Y, \iota : A \rightarrow B$  of simplicial sets. This induces a diagram

$$\begin{array}{ccc} \text{Fun}(B, X) & \longrightarrow & \text{Fun}(A, X) \\ \downarrow & & \downarrow \\ \text{Fun}(B, Y) & \longrightarrow & \text{Fun}(A, Y) \end{array}$$

and hence a map

$$\langle \iota, f \rangle : \text{Fun}(B, X) \rightarrow \text{Fun}(B, Y) \times_{\text{Fun}(A, Y)} \text{Fun}(A, X)$$

If  $f$  is an inner fibration and  $\iota$  a monomorphism, then  $\langle \iota, f \rangle$  is an inner fibration. If  $\iota$  is anodyne, then  $\langle \iota, f \rangle$  is trivial.

**Corollary 2.6 (Corollary):** When  $A = \emptyset, B = K, X = X, Y = \Delta^0$  we see that  $X$  being an infinity category implies that  $\text{Fun}(K, X)$  is also an infinity category. Moreover, if  $X$  is a Kan complex, then so is  $\text{Fun}(K, X)$ .

**Lemma 2.7 (Equivalent condition for being a quasicategory):** Let  $X$  be a simplicial set. Then TFAE:

- $X$  is an  $\infty$ -category
- For every inner anodyne map  $i : A \rightarrow B$ , the induced map  $i^* : \text{Fun}(B, X) \rightarrow \text{Fun}(A, X)$  is a trivial Kan fibration.
- $\text{Fun}(\Delta^2, X) \rightarrow \text{Fun}(\Lambda_1^2, X)$  is a trivial Kan fibration.

*Proof.* For (1)  $\implies$  (2), we use the theorem 2.5. Namely, we put  $Y = \Delta^0$  and we then see that

$$\text{Fun}(A, X) \times_{\text{Fun}(A, \Delta^0)} \text{Fun}(B, \Delta^0) \simeq \text{Fun}(A, X)$$

: the fiber product informally consists of pairs  $f : A \rightarrow X, g : B \rightarrow \Delta^0$  such that the following commutes:

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ \downarrow & & \downarrow \\ B & \xrightarrow{g} & \Delta^0 \end{array}$$

But  $f$  can be lifted to  $\tilde{f}$  since  $A \rightarrow B$  is inner anodyne which determines  $g$ . So the theorem says that

$$\text{Fun}(B, X) \rightarrow \text{Fun}(A, X)$$

is a trivial Kan fibration. This immediately implies (3).

Finally, to show (3)  $\implies$  (1),  $X$  to be an infinity category, we must verify that  $X \rightarrow \Delta^0$  satisfies the extension property for inner anodynes. We need to appeal to the fact that the inner anodyne maps are generated by all maps of the form  $[K \hookrightarrow L] \boxtimes [\Lambda_1^2 \rightarrow \Delta^2]$ , where  $\boxtimes$  refers to the induced map

from the pushout as in the diagram:

$$\begin{array}{ccc}
 K \times \Lambda_1^2 & \longrightarrow & K \times \Delta^2 \\
 \downarrow & & \downarrow \\
 L \times \Delta^2 & \longrightarrow & \text{pushout} \\
 & \searrow & \nearrow \\
 & L \times \Delta^2 &
 \end{array}$$

In this case, we can think of the pushout as the intersection, which is just  $K \times \Lambda_1^2$  when  $K \hookrightarrow L$  is a monomorphism. So the required extension property is reduced to the diagram:

$$\begin{array}{ccc}
 & \text{adjunction} & \\
 & \text{~~~~~} & \\
 K \times \Lambda_1^2 & \longrightarrow & X \\
 \downarrow & \nearrow ? & \downarrow \\
 L \times \Delta^2 & \longrightarrow & \Delta^0
 \end{array}
 \qquad
 \begin{array}{ccc}
 K & \longrightarrow & \text{Fun}(\Lambda_1^2, X) \\
 \downarrow & \nearrow ? & \downarrow \\
 L & \longrightarrow & \text{Fun}(\Delta^2, X)
 \end{array}$$

But by assumption, the right vertical is a trivial Kan fibration, so has the lifting property for all monomorphisms.  $\square$

With this lemma, we can show that the space of compositions is contractible, one of our original aims:

**Corollary 2.8 (Space of compositions is contractible):**

Suppose  $(g, \bullet, f) : \Lambda_1^2 \rightarrow C$  is a pair of composable arrows in an infinity category. Then the simplicial set of fillers which are lifts to  $\Delta^2$  is the pullback

$$\begin{array}{ccc}
 \text{Comp}(f, g) & \longrightarrow & \text{Fun}(\Delta^2, C) \\
 \downarrow & & \downarrow \\
 \Delta^0 & \longrightarrow & \text{Fun}(\Lambda_1^2, C)
 \end{array}$$

which is a subsimplicial set of  $\underline{\text{Hom}}(\Delta^2, C)$  and this simplicial set is contractible: it is a Kan complex and is weakly homotopy equivalent to  $\Delta^0$ .

*Proof.* By the previous theorem, the right vertical arrow is a trivial Kan fibration. Hence, its pullback to  $\Delta^0$  is a trivial Kan fibration.  $\square$

Generalization: recall that the spine  $\text{Sp}_n$  is generated by adjacent edges in  $\Delta^n$  and the inclusion is inner anodyne. Hence

$$\underline{\text{Hom}}(\Delta^n, C) \rightarrow \underline{\text{Hom}}(\text{Sp}_n, C)$$

is a trivial Kan fibration. A map from the spine is a sequence of  $n$  composable morphisms. Hence we can apply the same argument to show that the space of  $n$ -compositions is contractible.

## 2.2 Mapping spaces

With the tools from the previous section, we are ready to define the mapping spaces between two objects in an infinity category.

**Definition 2.9 (Mapping space):** *Given an infinity category  $C$ , we have a pullback diagram where we evaluate at the two endpoints.*

$$\begin{array}{ccc} \mathrm{Map}(c, d) & \longrightarrow & \mathrm{Fun}(\Delta^1, C) \\ \downarrow & & \downarrow \\ \Delta^0 & \longrightarrow & C \times C \end{array}$$

*The resulting space  $\mathrm{Map}(c, d)$  is a simplicial set, indeed a Kan complex! This is because we can think of  $C \times C = \mathrm{Fun}(\partial\Delta^1, C)$  and  $\partial\Delta^1 \hookrightarrow \Delta^1$  is inner anodyne, implying by 2.7 that the right vertical map is a trivial Kan fibration.*

Example: the loop space  $\Omega(X) = \mathrm{Map}_X(x, x)$ .

In ordinary category theory, the 'mapping space' is just a set, or a 0-category. In the same way, we move from an  $(\infty, 1)$  category to  $(\infty, 0)$  when we take mapping spaces of quasicategories, i.e. the mapping spaces are  $\infty$ -groupoids.

We can see that there is a morphism between two  $f, g : c \rightarrow d$  if and only if  $f \simeq g$ , since basically we have to fill in a square which restricts to  $f, g$  on the sides and the identity maps on the top and bottom, since  $\Delta_0^1$  has degeneracy maps which have  $s(c) = id_c$  and  $s(d) = id_d$ .

**Definition 2.10 (Fully faithful functors):**  *$F : C \rightarrow D$  is fully faithful if, for every  $c, d \in C$ , the induced map on mapping spaces is a homotopy equivalence:*

$$\mathrm{Map}(c, d) \xrightarrow{f} \mathrm{Map}(Fc, Fd)$$

Note that this is not a property of the homotopy category! Take  $X$  to be a noncontractible, simply-connected space. The canonical map  $X \rightarrow \Delta^0$  induces an equivalence on homotopy categories! So, in a way, the homotopy category only sees the first homotopy group, whereas the full infinity groupoid sees everything.

Similarly, we can define essential surjectivity. For spaces, this should be a surjection on  $\pi_0$ .

In particular, a map of spaces is ess. surj and fully faithful iff it is a homotopy equivalence, just like in ordinary category theory.

**Proposition 2.11 (Proposition):**  *$F$  is ess surj and fully faithful iff it is an equivalence.*

## Cores and subcategories

Suppose we have a subcategory of the homotopy category of an infinity category  $C$ . Form a pullback using the nerve construction:

$$\begin{array}{ccc} C' & \longrightarrow & C \\ \downarrow & & \downarrow \\ N(h(C)') & \longrightarrow & N(h(C)) \end{array}$$

This is the subcategory of  $C$  spanned by  $h(C)'$ . This is an infinity category, which can be checked by using the lifting criterion as well as the fact that an  $n$  simplex in a nerve is determined by its restriction to the spine, which is a subsimplicial set of the inner horns.

**Definition 2.12 (Cores):** *The core of an infinity category is the subcategory spanned by the core of  $h(C)$ , i.e. the subcategory spanned by isomorphisms. This is the maximal infinity groupoid in  $C$ .*

## The parametrized composition space

Suppose we have three objects  $c, d, e \in C$ . Can define a triple mapping space, by instead using  $\Delta^2$ .

$$\begin{array}{ccc} \text{Map}(c, d, e) & \longrightarrow & \text{Fun}(\Delta^2, C) \\ \downarrow & & \downarrow \\ \Delta^0 & \longrightarrow & C \times C \times C \end{array}$$

This has projection maps to the usual mapping spaces. Then

$$\text{Map}(c, d, e) \rightarrow \text{Map}(d, e) \times \text{Map}(c, d)$$

is a trivial Kan fibration. The fiber is the space of compositions, which is contractible. We thus have a homotopy inverse section i.e. a choice of composition. But this is not well-defined, but only unique up to homotopy. One has to use the general fact that the sections of a Kan fibration form a contractible set: the idea is that trivial Kan fibrations are stable under  $\text{Fun}$  and hence we can form a pullback

$$\begin{array}{ccc} \Gamma(Y, X) & \longrightarrow & \text{Fun}(Y, X) \\ \downarrow & & \downarrow p \\ \Delta^0 & \xrightarrow{id} & \text{Fun}(Y, Y) \end{array}$$

## 2.3 The homotopy coherent nerve

In this section, we describe an adjunction between the simplicial sets and simplicially enriched categories in the form of the coherent nerve and the thickening functor. This also forms a Quillen equivalence between Quillen's model structure on  $\mathbf{sSet}$  and Segal's model structure on  $\mathbf{sCat}$ .

The idea is that we have a 'thickening' functor

$$\mathfrak{C} : \Delta \rightarrow \mathbf{sCat}$$

This can be left Kan extended to a functor on simplicial sets, and has an adjoint called the *homotopy coherent nerve*.

### The thickening functor

The construction is involved.

Suppose  $0 \leq i \leq j$ . Denote by  $P_{i,j}$  to be the set of subsets

$$P_{i,j} = \{I \subset \{i, \dots, j\} \mid i, j \in I\}$$

partially ordered by inclusion. If  $i > j$  define this to be empty.

Define

$$\mathfrak{C}[\Delta^n] := \begin{cases} 0, 1, \dots, n \text{ as objects} \\ \underline{\text{Hom}}(i, j) = N(P_{i,j}) \end{cases}$$

This is a thickening of  $\Delta^n$ . It is a simplicial category, i.e. a category enriched over simplicial sets. Composition is defined by taking unions of subsets.

*Example 2.13 (Example):* We have that  $\mathfrak{C}\Delta^2$  has three objects 0, 1, 2. The hom sets are:

- $\mathfrak{C}\Delta^2(0, 0) = \mathfrak{C}\Delta^2(0, 1) = \mathfrak{C}\Delta^2(1, 1) = \mathfrak{C}\Delta^2(1, 2) = \mathfrak{C}\Delta^2(2, 2) = \Delta^0$
- $\mathfrak{C}\Delta^2(0, 2) = \Delta^1$  is the nerve of the poset of subsets containing both 0 and 2 i.e.  $N(\{0, 2\} \subset \{0, 1, 2\})$ .

We can extend this functor to  $\mathbf{sSet}$  by using the density theorem and this will be a Kan extension along Yoneda. This defines a functor  $\mathfrak{C}$ , which has an adjoint  $\mathcal{N}$  as in 1.5. We get a picture

$$\begin{array}{ccc} & \mathbf{sSet} & \\ \mathcal{Y} \nearrow & & \searrow \mathfrak{C} \\ \Delta & \xrightarrow{\mathfrak{C}} & \mathbf{sCat} \\ & \nwarrow \mathcal{N} & \end{array}$$

As we saw in 1.5, this forces the value of  $\mathcal{N}$  to be

$$\mathcal{N}C_n = \text{Hom}_{\mathbf{sCat}}(\mathfrak{C}\Delta^n, C)$$

For example, the objects of  $\mathcal{N}C_0$  are the objects of  $C$ , the morphisms are also the same, but  $(\mathcal{N}C)_2$  consists of the data of three objects  $x, y, z$ , three morphisms  $f \in C(x, y)_0, g \in C(y, z)_0, h \in C(x, z)_0$  and a 1 simplex in  $C(x, z)_1 = \text{Fun}(\Delta^1, C(x, z))$  restricting to  $h$  and  $g \circ f$  on the ends. Informally, a 2-simplex in this simplicial set consists of 2 composable morphisms and a homotopy between their composition and a third morphism.

We apply this to the simplicially enriched category of Kan complexes, which has objects Kan complexes and morphism spaces  $\text{Fun}(K, L)$ , which are also Kan complexes!

**Definition 2.14 (Spc):** We define a simplicial set  $\mathcal{N}\text{Kan} := \text{Spc}$  whose  $n$ -simplices are

$$\text{Spc}_n := \text{Hom}_{\text{sCat}}(\mathbb{C}\Delta^n, \text{Kan})$$

**Theorem 2.15 (Theorem):** If a given simplicial category  $\mathcal{C}$  is locally Kan (i.e. all morphism spaces are Kan complexes) then  $\mathcal{N}\mathcal{C}$  is an infinity category. In particular,  $\text{Spc}$  is an infinity category, the infinity category of spaces.

*Proof.* By adjunction, the lifting diagram corresponds to one involving  $\mathbb{C}[\Lambda_i^n]$ ,  $\mathbb{C}[\Delta^n]$  and  $\mathcal{C}$ . These are basically the same except how to extend it to  $\text{Hom}(0, n)$ . But since  $\mathcal{C}$  is locally Kan, this is possible.

$$\begin{array}{ccc}
 & \text{adjunction} & \\
 & \text{~~~~~} & \\
 \Lambda_i^n & \longrightarrow & \mathcal{N}\mathcal{C} \\
 \downarrow & \nearrow & \\
 \Delta^n & & 
 \end{array}
 \qquad
 \begin{array}{ccc}
 \mathbb{C}\Lambda_i^n & \longrightarrow & \mathcal{C} \\
 \downarrow & \nearrow & \\
 \mathbb{C}\Delta^n & & 
 \end{array}$$

Essentially,  $\mathbb{C}\Lambda_i^n$  has the same objects as  $\mathbb{C}\Delta^n$ , but one smaller morphism space:

- $\text{Hom}_{\mathbb{C}\Lambda_i^n}(i, k) = \text{Hom}_{\mathbb{C}\Delta^n}(i, k), (i, k) \neq (0, n)$
- $\text{Hom}_{\mathbb{C}\Lambda_i^n}(0, n) \subset \text{Hom}_{\mathbb{C}\Delta^n}(0, n) = NP_{0,n} = (\Delta^1)^{n-1}$  is given by deleting the interior and bottom  $j$  face. This is an anodyne map, so it lifts if  $\mathcal{C}$  is locally Kan.

□

In fact more is true, even though I am not sure how to prove it: the mapping space in the infinity category of spaces recovers the inner hom

$$\underline{\text{Hom}}(X, Y) = \text{Fun}(X, Y) \simeq \text{Map}_{\text{Spc}}(X, Y)$$

## 2.4 Infinity Yoneda

We have a non-unique but contractible choice of compositions, which makes it hard to create a functorial assignment

$$\text{Map}(c, d) \rightarrow \text{Map}(c', d)$$



**Definition 2.16 (Opposite of an infinity category):** The opposite of an infinity category  $C : \Delta^{op} \rightarrow \text{Set}$  is given by precomposing with the opposite map on  $\Delta$ . This is still an infinity category, since  $\Lambda_i^n$  just becomes  $\Lambda_{n-i}^n$ .

Presheaves on infinity categories

$$\mathcal{P}(C) := \text{Fun}(C^{op}, \text{Spc})$$

### 3 Limits and colimits

**Definition 3.1 (Limits and colimits):** A cone over  $F : I \rightarrow C$  is a natural transformation  $\eta : \underline{l} \rightarrow F$ . A limit cone is one such that

$$\text{Map}(c, l) \rightarrow \text{Map}_C F$$

is a homotopy equivalence. We need to compose with  $\eta$ , and while this is not unique, it is up to homotopy.

Similarly, a cone under  $F$  is a natural transformation  $\epsilon : F \rightarrow \underline{l}$  and a colimit cone is one such that

$$\text{Map}(l, c) \simeq \text{Map}(F, \underline{c})$$

An object is initial iff  $\text{Map}(l, c)$  is contractible for all  $c$  and terminal if  $\text{Map}(c, l)$  is contractible for all  $c$ . An infinity category is called pointed if it has a zero object, i.e. an object that is both initial and final.

#### Pullbacks

A pullback amounts to the data

$$\begin{array}{ccc} c & \xrightarrow{h} & r \\ i \downarrow & & \downarrow f \\ s & \xrightarrow{g} & t \end{array}$$

such that  $g \circ i \simeq f \circ h$ .

In the infinity category of spaces, this is a homotopy pullback!

**Proposition 3.2 (Proposition):** Both  $\text{Spc}$  and  $\text{Cat}_\infty$  admit small limits and colimits. The inclusion of the former into the latter preserves all limits and colimits.

Just as limits  $l$  can be compared to the Yoneda sets in ordinary category theory, so limits in infinity categories can be understood via their mapping spaces. For example, a pushout has the formula

$$\text{Map}(y, \coprod_x z, c) \simeq \text{Map}(y, c) \times_{\text{Map}(x, c)} \text{Map}(z, c)$$

where the pullbacks are happening in the infinity category of spaces, which we just said admits small limits and colimits.

### Alternative definitions

There are alternative definitions of limits and colimits via over/undercategories and joins.

Let

$$\begin{aligned}\star : \Delta \times \Delta &\rightarrow \mathbf{sSet} \\ [n] \star [m] &= N([m] \star [n])\end{aligned}$$

and extend it to simplicial sets by density. For example,  $(\Lambda_0^2)^\Delta \simeq \Delta^1 \times \Delta^1$ .

This allows for the definition of the slice over  $F : I \rightarrow C$  as the adjoint to the star operation:

$$\mathrm{Hom}_{\mathbf{sSet}}(K, C_{/F}) = \mathrm{Hom}_F(K \star I, C)$$

where on the right we consider functors that extend  $F$ . This is an infinity category whenever  $C$  is.

When we take  $F = d : \Delta^0 \rightarrow C$ , we get the overcategory  $C_{/d}$  which has objects morphisms  $c \rightarrow d$ . There is a thickened version of this:

$$\begin{array}{ccccc}\mathrm{Map}(c, d) & \longrightarrow & C^{/d} & \longrightarrow & \mathrm{Fun}(\Delta^1, C) \\ \downarrow & & \downarrow & & \downarrow \\ \Delta^0 & \longrightarrow & C & \longrightarrow & C \times C\end{array}$$

In fact, there is a pullback diagram in  $\mathbf{sSet}$

$$\begin{array}{ccc}\mathrm{Map}(L, F) & \longrightarrow & C^{/F} \\ \downarrow & & \downarrow \\ \Delta^0 & \longrightarrow & C\end{array}$$

**Proposition 3.3 (Equivalent definition of limit cone):** *TFAE:*

- $l$  is a limit cone over  $F$
- Restriction induces a homotopy equivalence

$$\mathrm{Map}(\underline{c}, \bar{F}) \simeq \mathrm{Map}(\underline{c}, F)$$

where  $\bar{F}$  is the extension via  $l$  of  $F$  to  $I^\Delta$ .

- $\bar{\eta} \in C_{/F}$  is final.

**Proposition 3.4 (Proposition):** *The infinity category of limits  $\mathrm{Limits}(F)$  is either empty or contractible.*