

PAPER • OPEN ACCESS

Methods of Generating Entangled Photon Pairs

To cite this article: Yixuan Li 2020 *J. Phys.: Conf. Ser.* **1634** 012172

View the [article online](#) for updates and enhancements.

You may also like

- [Traps contribution on detection time of single electron photodetector \(Photo-SET\)](#)
S Chatbouri, M Troudi, N Sghaier et al.
- [Pulsed photoacoustic system calibration for highly excited molecules: II. Influence of the laser beam profile and the excitation energy decay](#)
M D Rabasovi, J D Nikoli and D D Markushev
- [Machine learning for condensed matter physics](#)
Edwin Bedolla, Luis Carlos Padierna and Ramón Castañeda-Priego



244th Electrochemical Society Meeting

October 8 – 12, 2023 • Gothenburg, Sweden

50 symposia in electrochemistry & solid state science



Deadline Extended!
Last chance to submit!

New deadline:
April 21
submit your abstract!

Methods of Generating Entangled Photon Pairs

Yixuan Li

UC Santa Barbara, Santa Barbara, California 93106

729871390a@gmail.com

Abstract. We examined the regular methods of creating knotted photon sets. Our company initially centered upon the nuclear resource, the very first discovered knotted photon resource that creates the photon sets automatically. Our company at that point researched the nonlinear optics procedure: parametric down-conversion. And various other resources like four-wave mixing were actually gone over. The document is actually separated and it also delivers a thorough study of the natural science as well as appropriate procedures about the complicated photon production.

1. Introduction

Quantum Information and Communication Technology (QICT) attracts many researchers in the areas of applied physics, mathematics, information and so on. Entanglement of the nonlocal quantum correlation between two or more quantum-mechanical objects is one of the key things in QICT. Therefore, the improvement of techniques to generate and characterize entanglement is important in the field.

The mystery of complication was actually accelerated as a disagreement that quantum technicians could certainly not be a real common idea, but must be nutritional supplemented through added variables. That started to be modified in 1956. Hanbury-brown, as well as Twiss, suggested and gauged the strong dependency of light-toned [1]. In their practice, they noted much higher purchase obstruction results when the lighting was actually arranged right into half a tiny light beam as well as sensed through a set of photomultiplier cylinders in a device right now called an HBT interferometer. By doing this, they established the slant of Sirius. In 1964, Alarm displayed that quantum technical nonlocality could be experimentally confirmed along with a set of knotted photons. The transgression of Alarm's disparity was actually experimentally validated, making use of a set of photons given off in a waterfall of photons coming from a solitary atom [2, 3]. In 1988, connection sizes of the combined sign as well as idler photons generated in the method of parametric down-conversion have actually been conducted as functionality of a pair of straight polarizer environments. It was discovered that the Alarm disparity for the pair of split up bits was gone against by around 6 typical variances, and also a timeless possibility for lightweight surges was gone against considerably. Lately, the possibility of utilizing the semiconductor component to create knotted photon discharge diodes has actually drawn in people's interest [4-8]. Reliable creation of knotted photons is important to elevate quantum-related information as well as interaction guidelines, which need a multitude of knotted things as their primary sources. As a result, the production as well as the depiction of such a great deal of knotted photons is additionally a very hot subject matter in the business of speculative and academic quantum infotech.

Today, protons [9-11], atoms [12], and caught ions [13, 14], have actually been utilized to verify quantum complication. Nevertheless, in functional requests, it is still complicated to sustain complexity because of its own susceptibility in a common setting. Photons are still the absolutely most encouraging



as well as efficient channel for the production of complication, due to the fact that the quantum specifies encrypted through photons (consisting of knotted conditions) are actually reasonably insusceptible to the setting.

We are going to assess and review procedures for creating knotted photon sets, like nuclear resources, parametric down-conversion, and semiconductor resources.

2. Three ways to generate entangled photon pairs

2.1 Atomic source

The two crucial executions of photon qubits are polarization qubits and time-slot qubits. The qubits are utilized in significant practices to research a variety of knotted conditions, like the exam of Alarm's disparity. Opportunity-worthy qubits are primarily utilized in quantum interaction methods including quantum cryptography, as a result of them being compatible along with fiber optics transmission. The initially reputable is




Figure. 1. Representational design of the knotted photon set discharge via atomic cascade in calcium

$$|\psi\rangle = \frac{1}{\sqrt{2}} * (|L\rangle_A |L\rangle_B + |R\rangle_A |R\rangle_B) \\ \equiv \frac{1}{\sqrt{2}} * (LL\rangle + |RR\rangle), \quad (1)$$

where L, as well as R, signifies the polarization condition of each photon – L: Left behind Cycle, R: Right Cycle. The polarized knotted condition 1 could be revised in a straight polarization manner as

$$|\psi\rangle = \frac{1}{\sqrt{2}} * (|HH\rangle - |VV\rangle), \quad (2)$$

where H, as well as V, embodies the parallel and upright direct polarization specifically. Although this procedure of creating knotted photons is introducing, it is challenging to gather the sent out sets efficiently since the photons are arbitrarily discharged at a full three-dimensional slant. Moreover, handling private atoms is certainly not valuable for overall speculative requests. For that reason, although nuclear waterfall is the initial resource of knotted photons, it has certainly not been largely made use of in quantum info interaction practices.

2.2 Parametric down-conversion

The primary method to produce knotted photons is actually via spontaneous parametric down-conversion.

A popular procedure for creating knotted photon qubits is using the nonlinearity visual procedure of Spontaneous Parametric Down-Conversion (SPDC). In SPDC, the photon is a tough lightweight area dispersing and utilizing a visual tool along with the second-order nonlinearity. It is feasible to be divided right into a set of photons whose complete electricity as well as energy amount to electricity and also the energy of the pushed photon.

Making use of criteria to forecast solitary photon down-conversion because of one sign photon as well as one abandoned photon created concurrently in SPDC, the diagnosis of abandoned photons "anticipates" the visibility of indicator photons; the light, referred to as "anticipated singular photon resource HSPS", serves for the Quantum Key Distribution (QKD), 21 photon quantum info and also

assessment. Below, our experts present the arrangement of a rhythm higher electrical power scope QKD along with a few of higher photons and an extremely identical high-sensitivity healthy protein for photon quantum data processing.

Parametric Down-Conversion (PDC) is the method of turning the pump reprimand pair of visual indicators as well as a nonlinear optics illumination into a crystal along with nonlinearity [15]. Casual parametric down-conversion, or even parametric fluorescence, has been noted because the 1960s [16, 17]. Coming from a quantum visual viewpoint, the PDC is a procedure through which the paired photons are produced all at once as a mom and dad photon. The communication Hamiltonian of the procedure is shown.




Figure. 2. Picture of the parametric down-conversion in $\chi(2)$ nonlinear crystal.

$$\omega_0 = \omega_1 + \omega_2, \quad (3)$$

$$k_0 = k_1 + k_2, \quad (4)$$

These disorders are actually looked at to become the preservation of electricity, and also the energy-related photons. Because of this, the sign as well as Idler photons possess an organic relationship in power and drive. On top of that, the discharge of indicators as well as abandoned photons must be actually concurrent to the real ailments. Connection in formula (3) and (4) have actually been noticed to validate, featuring that, as a result of these features, PDC is typically utilized as a set of photon resources for several kinds of two-photon interferometers [16, 17, 18].

$$\begin{aligned} |\psi\rangle &= \frac{1}{\sqrt{2}} * \left(|H\rangle_A + i|H\rangle_B \otimes \frac{1}{\sqrt{2}} * (-i|V\rangle_A + |V\rangle_B) \right) \\ &= \frac{1}{\sqrt{2}} * (|H\rangle_A |V\rangle_B + |V\rangle_A |H\rangle_B - i|H\rangle_A |V\rangle_A + i|V\rangle_B |H\rangle_B), \end{aligned} \quad (5)$$

In the development of polarized knotted photons, it is actually quite vital to take into consideration the polarization condition of the Photon, which is actually created through PDC. There are actually 2 kinds of period PDC matching problems, depending upon the polarization condition. An indicator created through a kind I stage matching or even the PDC style possesses an analog, polarization along with the idler photon. One more thus referred to as the Style II unreactive photon possesses upright polarization. Introducing job PDC makes use of kind I period polarization complication, matching, and also a mix of a non-polarized beam of lights and a ray of light splitters. In these practices, the polarization of one photon is actually turned to get a pair of orthogonally polarized photons, which at that point combine with the light beam splitter.

$$|\psi\rangle = \frac{1}{\sqrt{2}} * (|HV\rangle + |VH\rangle) \quad (6)$$

Here, A and B are actually the beam of light splitters. This condition is the item condition and as a result is certainly not knotted. Having said that, if our company picks 2 of these activities, the photon is actually broken down into a pair of outcome slots An and B, as well as the leading condition includes the initial 2 phrases in the formula.

This is the max complexity. The method includes a supposed fallback technique, through which the condition of the preferred condition is picked from the created entire. The assortment treatment is probabilistic (results, 1/2) and also for that reason performs certainly do not put on a bug-free examination for Alarm's discrimination [17]. However, this is the initial practice to illustrate the quantity

of polarized knotted photons that PDC creates. Eventually, comparable methods are related to kind II PDC.

In 1995, the PDC of polarization complexity in photons generated through Style II crystals is actually disclosed without making use of post-selection. For a provided Style II period fit, the indicator, the unreactive Photon, is actually sent out along a pair of cone-like areas: one for the indicator photon along with the parallel polarization, and also the various others for the ineffective photon along with upright polarization. Two cone-shaped areas cross another pair of aspects, in the path A and B, where the sign and the unoccupied photon are actually discharged, to make sure that when one is sent out to the various others, it is actually released to B and also the other way around.

$$|\psi\rangle = \frac{1}{\sqrt{2}} * (|HV\rangle + e^{i\theta}|VH\rangle) \quad (7)$$

Thereby, the polarized condition of a photon discharged in both instructions simultaneously needs to be actually ended! Between the 2 conditions in the formula (7), it may be changed through putting an extra birefringent stage shifter among the sunny pathways responsible for the crystal. This procedure is actually the initial speculative demo of Quantum teleportation [19].

Two exactly same crystals along with kind I period matching are actually utilized; their optical axes are straightened in an upright aircraft. As received body 2, when a pair of crystals are actually brightened through a 45-polarized pump beam of light, the sign and the idler photons are released from all of them, including crystal # 1 or even # 2. Crystal # 1 releases an indicator as well as an additional photon, which is actually flat polarized, while crystal 2 discharges an up and down polarized photon.

Along both conoids, the photons given off through each crystal are actually coaxial as well as, if the crystal is actually, lean sufficient. Thereby, the released polarized condition, the Photon, is actually a superposition of a pair of situations, particularly Cam! In

$$|\psi\rangle = \frac{1}{\sqrt{2}} * (|HH\rangle + e^{i\theta}|VV\rangle), \quad (8)$$

pairs of conditions in the formula (8) could be readjusted through managing the loved one stage, in the straight as well as upright elements of the pump beam of light. This strategy is actually far better than utilizing the kind II stage matching explained previously, given that the photons produced to a much larger sound slant are going to be actually knotted. In birefringent products, the velocity of lightweight rhythms relies on their polarization, and consequently, the age of photon surge packages apiece various others relies on their polarization. If split up, the variety of photon surge packages is actually above their opportunity and size, which is actually a purchase of size of the inverted spooky distance (in common speculative ailments, the moment size is actually 100fs), triggering a decline in the level of complexity. Thus, the fullness of the nonlinear crystals ought to be tiny sufficient to make sure that the splitting up of the photon surge packages is actually a little sufficiently relative to their opportunity sizes. It works to place extra birefringent crystals to make up for photon surge package splitting up [20, 21, 22]. Yet another means to utilize PDC to produce polarized knotted photons is actually to position a nonlinear crystal in an interferometer [21-27]. In an interferometer, indicators and abandoned photons are actually released coming from any sort of PDC crystal, forty-five level polarized pumped beam of lights split through a polarized light beam splitter, which may demonstrate and also transfer, up and down and flat polarized photons. The pump beam of light after that brightens 2 nonlinear crystals whose optical axes are actually orthogonal, upright, yet straight. Each crystal gives off a set of photons, including a parallel polarization as well as an upright polarization, and contacted photons. The 2 upper arms of the interferometer may be changed through transforming the pathway size variation in all of them. This obstruction system provides the concept of the age of complexity, i.e. the superposition of 2 conditions of the shared obstruction. Having said that, the security of the interferometer feels to have ecological obstruction, which confines the functionality of the complexity. In this particular instance, the SAGNAC interferometer has actually been utilized to secure dependable knotted photons. Quasi-phase-matching's use of regular poling's nonlinear crystals is actually a strong and scalable method for nonlinear insight sale. Making use of regularly poled lithium as well as blood potassium regular poling, Periodically Poled KTP (PPKTP) [23, 27, 28] has actually been utilized to produce knotted photons by means of the PDC. Particularly, PPLN waveguides are actually utilized for the dependable age group of telecommunication

insight areas (1.3 and 1.5), and are actually made for make use in quantum cryptography requests. The phrase, parameterized SPDC, implies the method where the down-conversion tool continues to be steady. Consequently, this automatically implies that a collection of preservation rules needs to be actually fulfilled through pushing, signalling, and also idling photons.

2.3 Other sources

As described in section 2.2, the most popular method for obtaining entangled photon pairs is PDC, a second-order nonlinear process in which a high-energy photon is converted to a twin-daughter photon. Higher Order Nonlinear Processes, such as χ_3 are also suitable for obtaining entangled photon pairs; however, higher order generation efficiency is very low unless enhanced in some way. The polarized dispersion surface represents the state which can satisfy the phase matching condition. Furthermore, limiting the light intensity spatiotemporal to optical micro-cavity and waveguide structures can enhance the χ_3 optical nonlinear process. Dispersion-shifted fibres are used to generate correlated photon pairs. In this technique, the pump wavelength is set to near zero dispersion wavelength, avoiding vertical separation of DSF, pumping, signal and idler wheel photon pulse. Thus, the χ_3 process can be a significantly strong interaction by the spatiotemporal limitation of the photon pulses in the DSF. Recently, polarization and spatiotemporal entanglement have been demonstrated using the resulting photon pairs. In addition, photonic crystal fibres have also been used. To address one of the remaining problems with correlated photons, one of which is the presence of background noise, it should mainly from spontaneous Raman scattering. By cooling the fibre, the Raman noise can be suppressed and the performance can be improved. Recently, correlated photons have been produced in nanoscale silicon waveguides.

3 Conclusion

In review, we have talked about the creation, review, and also portrayal of knotted photons. Photon set polarization is the easiest yet the most practical body on which 2-qubit complication is actually encrypted. Although our experts carry out certainly but do not point out the multi-part complexity in between, greater than the pair of qubits in this particular post, this may happen from 2 or even more knotted photon sets. As much as 6 photons have been experimentally knotted thus far. Additionally, refining complexity strategies including purification, swap, filtration, and also attention is vital to the sensible use of QICT. Polarized entangled photons have been made use of in each of these general practices. Because of the initial understanding of knotted photons making use of nuclear resources, numerous research studies have actually been carried out on the production and also the review of knotted photons. One of the most prosperous and well-liked methods today is actually to utilize PDC. Just recently, QPM innovation has been made use of to boost the possibility as well as the versatility of PDCS in the age group of knotted photons. Lately shown semiconductor resources have broken the ice for knotted photon-emitting diodes. One means to create polarization knotted multiphoton conditions is actually via the spontaneous parameter down-conversion (SPDC) of ultrafine lightweight pushing rhythms. Our best target is actually to obtain premium quality polarization knotted multi-photon resources based upon the very same SPDC. The resource may be utilized for simple screening of quantum physics in numerous quantum interaction processes as well as high-dimensional Hilbert Spaces.

References

- [1] R. H. Brown and R. Q. Twiss, (1956), *Nature* 177, 27.
- [2] J. F. Clauser, M. A. Horne, A. Shimony, and R. A. Holt, (1969), *Phys. Rev. Lett.* 23, 880.
- [3] A. Aspect, J. Dalibard, and G. Roger, (1982), *Phys. Rev. Lett.* 49, 1804.
- [4] K. Edamatsu, G. Oohata, R. Shimizu, and T. Itoh, (2004): *Nature* 431 167.
- [5] R. M. Stevenson, R. J. Young, P. Atkinson, K. Cooper, D. A. Ritchie, and A. J. Shields, (2006): *Nature* 439 179.
- [6] R. J. Young, R. M. Stevenson, P. Atkinson, K. Cooper, D. A. Ritchie, and A. J. Shields (2006): *New J. Phys.* 829.

- [7] N. Akopian, N. H. Lindner, E. Poem, Y. Berlatzky, J. Avron, D. Gershoni, B. D. Gerardot, and P. M. Petroff(2006): *Phys. Rev. Lett.* 96 130501.
- [8] G. Oohata, R. Shimizu, and K. Edamatsu(2007): *Phys. Rev. Lett.* 98 140503.
- [9] M. Lamehi-Rachti and W. Mittig (1976): *Phys. Rev. D* 14 2543.
- [10] C. Polachic, C. Rangacharyulu, A. van den Berg, S. Hamieh, M. Harakeh, M. Hunyadi, M. de Huu, H. Wo'rtche, J. Heyse, C. Ba'umer, D. Frekers, S. Rakers, J. A. Brooke, and P. Busch (2004): *Phys. Lett. A* 323176.
- [11] H. Sakai, T. Saito, T. Ikeda, K. Itoh, T. Kawabata, H. Kuboki, Y. Maeda, N. Matsui, C. Rangacharyulu, M. Sasano, Y. Satou, K. Sekiguchi, K. Suda, A. Tamii, T. Uesaka, and K. Yako(2006): *Phys. Rev. Lett.* 97 150405.
- [12] E. Hagley, X. Maître, G. Nogues, C. Wunderlich, M. Brune, J. M. Raimond, and S. Haroche (1997): *Phys. Rev. Lett.* 79 1.
- [13] Q. A. Turchette, C. S. Wood, B. E. King, C. J. Myatt, D. Leibfried, W. M. Itano, C. Monroe, and D. J. Wineland(1998): *Phys. Rev. Lett.* 81 3631.
- [14] C. A. Sackett, D. Kielpinski, B. E. King, C. Langer, V. Meyer, C. J. Myatt, M. Rowe, Q. A. Turchette, W. M. Itano, D. J. Wineland, and C. Monroe (2000): *Nature* 404 256.
- [15] L. Mandel and E. Wolf: *Optical Coherence and Quantum Optics* 1995 (Cambridge University Press, Cambridge, U.K.,) Sect. 22.4.
- [16] D. C. Burnham and D. L. Weinberg(1970): *Phys. Rev. Lett.* 25 84.
- [17] L. Mandel (1999): *Rev. Mod. Phys.* 71 S274.
- [18] C. K. Hong, Z. Y. Ou, and L. Mandel(1987): *Phys. Rev. Lett.* 59 2044.
- [19] D. Bouwmeester, J. Pan, K. Mattle, M. Eibl, H. Weinfurter, and A. Zeilinger(1997): *Nature* 390 575.
- [20] Y. Nambu, K. Usami, Y. Tsuda, K. Matsumoto, and K. Nakamura (2002): *Phys. Rev. A* 66 033816.
- [21] D. Branning, W. Grice, R. Erdmann, and I. A. Walmsley(2000): *Phys. Rev. A* 62 013814.
- [22] Y.-H. Kim, M. V. Chekhova, S. P. Kulik, M. H. Rubin, and Y. Shih(2001): *Phys. Rev. A* 63 062301.
- [23] M. Fiorentino, G. Messin, C. E. Kuklewicz, F. N. C. Wong, and J. H. Shapiro(2004): *Phys. Rev. A* 69 041801.
- [24] A. Yoshizawa and H. Tsuchida (2003): *Electron. Lett.* 39 621.
- [25] A. Yoshizawa and H. Tsuchida(2005): *Jpn. J. Appl. Phys.* 44 L375.
- [26] B.-S. Shi and A. Tomita(2004): *Phys. Rev. A* 69 013803.
- [27] T. Kim, M. Fiorentino, and F. N. C. Wong(2006): *Phys. Rev. A* 73 012316.
- [28] M. Fiorentino, C. E. Kuklewicz, and F. N. C. Wong(2006): *Opt. Express* 13 (2005) 127. 58) B.-S. Shi and A. Tomita: *New J. Phys.* 8 38.