

Generating and teleporting entanglement for quantum networks

Adrian Udovičić
Assoc. prof. dr. Rainer Kaltenbaek

University of Ljubljana, Faculty of Mathematics and Physics

23.05.2024, Ljubljana, Slovenia

Contents and introduction

1. Motivation

2. Theory

2.1 SPDC

Phase Matching

2.2 Phase Matching Temperature

2.3 Entanglement swapping

3. Present state

3.1 Parameters

3.2 Phase Matching Temperature

3.3 Building a Sagnac Interferometer

4. Outlook

Motivation

- ▶ Bright source of entanglement

Motivation

- ▶ Bright source of entanglement
- ▶ Training in quantum technologies in Slovenia

Motivation

- ▶ Bright source of entanglement
- ▶ Training in quantum technologies in Slovenia
- ▶ Quantum Network for Slovenia

Motivation

- ▶ Bright source of entanglement
- ▶ Training in quantum technologies in Slovenia
- ▶ Quantum Network for Slovenia
- ▶ Testbed for industrialized version

Theory

1. SPDC
2. Entanglement swapping

- Spontaneous Parametric Downconversion

- Spontaneous Parametric Downconversion

$$\omega_p = \omega_s + \omega_i$$




Illustration of SPDC

Theory

SPDC

- #### ► Spontaneous Parametric Downconversion

$$\omega_p = \omega_s + \omega_i$$




Illustration of SPDC

- Degenerate $\omega_i = \omega_s$

Theory

SPDC

- #### ► Spontaneous Parametric Downconversion

$$\omega_p = \omega_s + \omega_i$$




Illustration of SPDC

- ▶ Degenerate $\omega_i = \omega_s$
 - ▶ Non-degenerate $\omega_i \neq \omega_s$

Theory

State of the Art

Comparison of different sources


| Who When | [1] 2022 | [2] 2010 | [3] 2007 | [4] 2006 | [5] 2012 |
|---|-------------------|--------------------|-------------------|-----------------|-------------------|
| Type | 0 | II | II | II | 0 |
| Brightness/ $\frac{\text{pairs}}{\text{sm} \text{W} \text{nm}}$ | $2,5 \times 10^6$ | $87,5 \times 10^3$ | 273×10^3 | 5×10^3 | 278×10^3 |
| Bandwidth/nm | 106 | 0,3 | 0,3 | 1 | 2,3 |

SPDC

Phase Matching, Quasi Phase Matching, Bandwidth

- #### ► Birefringent Phase Matching, Quasi Phase Matching

Illustration of a) Birefringent Phase Matching $k_p = k_i + k_s$ and




SPDC


Phase Matching, Quasi Phase Matching, Bandwidth

- #### ► Birefringent Phase Matching, Quasi Phase Matching

Illustration of a) Birifringent Phase Matching $k_p = k_i + k_s$ and







b) Quasi Phase Matching $k_p - k_i - k_s - \Delta k = 0$.



Theory

Phase Matching Temperature


Phase Matching Temperature calculations for a) Type-2 crystal of $9.12\text{ }\mu\text{m}$ polling period, b) Type-0, $19.25\text{ }\mu\text{m}$, c) Type-0, $19.45\text{ }\mu\text{m}$, d), Type-0 $19.65\text{ }\mu\text{m}$



Type-II vs Type-0

Bandwidth


Wavelength bandwidth of a) Type-2 crystal with a polling period of 9,12 μm
b) Type-0 crystals with polling periods of 19,25 μm



Type-II vs Type-0

Bandwidth


Wavelength bandwidth of a) Type-2 crystal with a polling period of 9,12 μm
 b) Type-0 crystals with polling periods of 19,25 μm



Type-II vs Type-0

Bandwidth

Wavelength bandwidth of a) Type-2 crystal with a polling period of 9,12 μm
 b) Type-0 crystals with polling periods of 19,25 μm



$$|\Psi_p\rangle = \frac{1}{\sqrt{2}} (\sin(\alpha) a_H^\dagger + \cos(\alpha) a_V^\dagger) |0\rangle$$

$$|\Psi_{\text{Type-2}}\rangle = \frac{1}{\sqrt{2}} (\sin(\alpha) a_H^\dagger(\omega_s) a_V^\dagger(\omega_i) +$$

$$\cos(\alpha) a_V^\dagger(\omega_i) a_H^\dagger(\omega_s)) |0\rangle$$

(1)

$$|\Psi_{\text{Type-0}}\rangle = \frac{1}{\sqrt{2}} (\sin(\alpha) a_H^\dagger(\omega_s) a_H^\dagger(\omega_i) +$$

$$\cos(\alpha) a_V^\dagger(\omega_i) a_V^\dagger(\omega_s)) |0\rangle$$

SPDC

Type-2 vs Type-0


Brightness comparison

$Hz/mW/nm$


| FMF | | IJS |
|-------------------|-------------------|--------------------|
| Type-II | Type-0 | Type-II |
| $7,8 \times 10^6$ | $2,6 \times 10^7$ | $0,05 \times 10^6$ |
| Bandwidth/nm | | |
| 0,68 | 0,68 | 0,68 |

Different Designs


Different design ideas from other groups. a) [6], b) [7], c) [8], d) [9]




(a)



(b)



(c)



(d)

Entanglement swapping


Introduction

- ▶ No specific form required - arbitrary states can be teleported
 1. Bell State Measurements
 2. Will try to use Quantum Memory from IJS group
- ▶ FMF/IJS

Entanglement Swapping

Teleportation and Swapping


Illustration of Entanglement Teleportation.



Entanglement Swapping

Teleportation and Swapping

Illustration of Entanglement Swapping.



Present state

Parameters

- ▶ Focusing parameters [10]

$$\xi = \frac{L}{kw^2} \quad (2)$$

- ▶ Heraldng

Present state


Phase Matching Temperature

Temperature scans of Type-0 crystals with different polling periods, a) misaligned 19,25 μm , b) 19,25 μm , c) 19,45 μm , d) 19,65 μm

Present state

Phase Matching Temperature

Temperature scans of Type-0 crystals with different polling periods, a) misaligned 19,25 μm , b) 19,25 μm , c) 19,45 μm , d) 19,65 μm




(a)


Present state

Phase Matching Temperature


Temperature scans of Type-0 crystals with different polling periods, a) misaligned 19,25 μm , b) 19,25 μm , c) 19,45 μm , d) 19,65 μm




(a)



(b)




(c)



(d)


Present state

Building a Sagnac Interferometer



Present state

Building a Sagnac Interferometer



Specifications from the crystal manufacturer.

Outlook

- ▶ SiQUID
- ▶ Entanglement swapping between FMF and IJS
- ▶ Building quantum internet
- ▶ Free space link to reactor

Thank you

References

-  S. P. Neumann, M. Selimovic, M. Bohmann, and R. Ursin, "Experimental entanglement generation for quantum key distribution beyond 1 Gbit/s," *Quantum*, vol. 6, p. 822, Sep. 2022.
-  H. D. F. A. e. a. Hübel, H., "Direct generation of photon triplets using cascaded photon-pair sources," *Nature* 466, 601–603 (2010).
-  A. P. T. J. A. Z. A. Fedrizzi, T. Herbst, "A wavelength-tunable fiber-coupled source of narrowband entangled photons," *Opt. Express* 15, 15377-15386 (2007).
-  M. F. T. Kim and F. N. C. Wong, "Phase-stable source of polarization-entangled photons using a polarization sagnac interferometer," *Optica Publishing Group*, (2006), paper JTUh5.
-  M. J. H. W. D. P. T. J. R. U. J. R. M. W. M. J. P. T. H. W. V. P. F. Steinlechner, P. Trojek, "A high-brightness source of polarization-entangled photons optimized for applications in free space," *Opt. Express* 20, 9640-9649 (2012).
-  M. C. Sang Min Lee, Heonoh Kim and H. S. Moon, *Opt. Express* 24, 2941-2953 (2016).
-  G. K. S. M. V. Jabir, "Robust, high brightness, degenerate entangled photon source at room temperature," *Sci Rep* 7, 12613 (2017).
-  Y. Chen, S. Ecker, S. Wengerowsky, L. Bulla, S. K. Joshi, F. Steinlechner, and R. Ursin, "Polarization entanglement by time-reversed hong-ou-mandel interference," *Physical Review Letters*, vol. 121, no. 20, Nov. 2018.
-  E. Bocquillon, C. Couteau, M. Razavi, R. Laflamme, and G. Weihs, "Coherence measures for heralded single-photon sources," *Physical Review A*, vol. 79, no. 3, Mar. 2009.
-  R. S. Bennink, "Optimal collinear gaussian beams for spontaneous parametric down-conversion," *Phys. rev. A* 81, 053805 (2010).