

Lecture 9

Linear independent, basis and dimension

Linear independent

Linear independent

Recall: A set of vector $\vec{v}_1, \dots, \vec{v}_k$ are said to be linearly independent if we don't have non-trivial expression of $\vec{0}$.

Linear independent

Recall: A set of vector $\vec{v}_1, \dots, \vec{v}_k$ are said to be linearly independent if we don't have non-trivial expression of $\vec{0}$.

i.e. if $c_1\vec{v}_1 + \dots + c_k\vec{v}_k = 0$

then $c_1 = 0, \dots, c_k = 0$.

Linear independent

Recall: A set of vector $\vec{v}_1, \dots, \vec{v}_k$ are said to be linearly independent if we don't have non-trivial expression of $\vec{0}$.

i.e. if $c_1\vec{v}_1 + \dots + c_k\vec{v}_k = 0$

then $c_1 = 0, \dots, c_k = 0$.

- Any vector \vec{v} in $\text{Span}\{\vec{v}_1, \dots, \vec{v}_k\}$ can be expressed **uniquely** as linear combination of $\vec{v}_1, \dots, \vec{v}_k$

Linear independent

Recall: A set of vector $\vec{v}_1, \dots, \vec{v}_k$ are said to be linearly independent if we don't have non-trivial expression of $\vec{0}$.

i.e. if $c_1\vec{v}_1 + \dots + c_k\vec{v}_k = 0$

then $c_1 = 0, \dots, c_k = 0$.

- Any vector \vec{v} in $\text{Span}\{\vec{v}_1, \dots, \vec{v}_k\}$ can be expressed **uniquely** as linear combination of $\vec{v}_1, \dots, \vec{v}_k$

Reason: If we have

$$c_1\vec{v}_1 + \dots + c_k\vec{v}_k = \vec{v} = d_1\vec{v}_1 + \dots + d_k\vec{v}_k$$

Linear independent

Recall: A set of vector $\vec{v}_1, \dots, \vec{v}_k$ are said to be linearly independent if we don't have non-trivial expression of $\vec{0}$.

i.e. if $c_1\vec{v}_1 + \dots + c_k\vec{v}_k = 0$

then $c_1 = 0, \dots, c_k = 0$.

- Any vector \vec{v} in $\text{Span}\{\vec{v}_1, \dots, \vec{v}_k\}$ can be expressed **uniquely** as linear combination of $\vec{v}_1, \dots, \vec{v}_k$

Reason: If we have

$$c_1\vec{v}_1 + \dots + c_k\vec{v}_k = \vec{v} = d_1\vec{v}_1 + \dots + d_k\vec{v}_k$$

$$\Rightarrow (c_1 - d_1)\vec{v}_1 + \dots + (c_k - d_k)\vec{v}_k = 0$$

Linear independent

Recall: A set of vector $\vec{v}_1, \dots, \vec{v}_k$ are said to be linearly independent if we don't have non-trivial expression of $\vec{0}$.

i.e. if $c_1\vec{v}_1 + \dots + c_k\vec{v}_k = 0$

then $c_1 = 0, \dots, c_k = 0$.


- Any vector \vec{v} in $\text{Span}\{\vec{v}_1, \dots, \vec{v}_k\}$ can be expressed **uniquely** as linear combination of $\vec{v}_1, \dots, \vec{v}_k$

Reason: If we have

$$c_1\vec{v}_1 + \dots + c_k\vec{v}_k = \vec{v} = d_1\vec{v}_1 + \dots + d_k\vec{v}_k$$

$$\Rightarrow (c_1 - d_1)\vec{v}_1 + \dots + (c_k - d_k)\vec{v}_k = 0 \Rightarrow c_i = d_i$$


Linear independent



Linear independent

- Recall:

Let's consider \mathbb{R}^3 .




Linear independent

- Recall: Let's consider \mathbb{R}^3 .

L

Any vector \vec{v}_i is l.i.



Linear independent

- Recall: Let's consider \mathbb{R}^3 .


1.

Any vector \vec{v}_1 is l.i.

2.

Two vector \vec{v}_1, \vec{v}_2 are l.i.

\iff they are not parallel.



Linear independent

- Recall: Let's consider \mathbb{R}^3 .

1


Any vector \vec{v}_1 is l.i.

2

Two vectors \vec{v}_1, \vec{v}_2 are l.i.

3

\iff they are not parallel.



3

Three vectors $\vec{v}_1, \vec{v}_2, \vec{v}_3$ are l.i. \iff

not one is lying in the plane spanned by other two.

Linear independent

- Recall: Let's consider \mathbb{R}^3 .

1

Any vector \vec{v}_1 is l.i.

2

Two vectors \vec{v}_1, \vec{v}_2 are l.i.


3

Three vectors $\vec{v}_1, \vec{v}_2, \vec{v}_3$ are l.i. \iff

not one is lying in the plane spanned by other two.

4

Four vectors cannot be l.i.



Linear independent

Linear independent

fact

Suppose $\vec{v}_1, \dots, \vec{v}_k$ are l.i.

Linear independent

fact:

Suppose $\vec{v}_1, \dots, \vec{v}_k$ are l.i.

adding \vec{v}_{k+1} are still l.i $\Leftrightarrow \vec{v}_{k+1}$ not in $\text{Span}\{\vec{v}_1, \dots, \vec{v}_k\}$

Linear independent

fact:

Suppose $\vec{v}_1, \dots, \vec{v}_k$ are l.i.

adding \vec{v}_{k+1} are still l.i $\Leftrightarrow \vec{v}_{k+1}$ not in $\text{Span}\{\vec{v}_1, \dots, \vec{v}_k\}$

Pf: \Leftarrow) • If $\vec{v}_1, \dots, \vec{v}_{k+1}$ l.d.

Linear independent

fact:

Suppose $\vec{v}_1, \dots, \vec{v}_k$ are l.i.

adding \vec{v}_{k+1} are still l.i $\Leftrightarrow \vec{v}_{k+1}$ not in $\text{Span}\{\vec{v}_1, \dots, \vec{v}_k\}$

Pf: \Leftarrow • If $\vec{v}_1, \dots, \vec{v}_{k+1}$ l.d.

then $c_1 \vec{v}_1 + \dots + c_{k+1} \vec{v}_{k+1} = 0$ for some c_i not all zero.

Linear independent

fact:

Suppose $\vec{v}_1, \dots, \vec{v}_k$ are l.i.

adding \vec{v}_{k+1} are still l.i $\Leftrightarrow \vec{v}_{k+1}$ not in $\text{Span}\{\vec{v}_1, \dots, \vec{v}_k\}$

Pf: \Leftarrow) • If $\vec{v}_1, \dots, \vec{v}_{k+1}$ l.d.

then $c_1 \vec{v}_1 + \dots + c_{k+1} \vec{v}_{k+1} = 0$ for some c_i not all zero.

• $c_{k+1} \neq 0$ because $\vec{v}_1, \dots, \vec{v}_k$ l.i.

Linear independent

fact:

Suppose $\vec{v}_1, \dots, \vec{v}_k$ are l.i.

adding \vec{v}_{k+1} are still l.i $\Leftrightarrow \vec{v}_{k+1}$ not in $\text{Span}\{\vec{v}_1, \dots, \vec{v}_k\}$

Pf: \Leftarrow) • If $\vec{v}_1, \dots, \vec{v}_{k+1}$ l.d.

then $c_1 \vec{v}_1 + \dots + c_{k+1} \vec{v}_{k+1} = 0$ for some c_i not all zero.

• $c_{k+1} \neq 0$ because $\vec{v}_1, \dots, \vec{v}_k$ l.i.

• then $\vec{v}_{k+1} = -\frac{c_1}{c_{k+1}} \vec{v}_1 - \dots - \frac{c_k}{c_{k+1}} \vec{v}_k$

Linear independent

fact:

Suppose $\vec{v}_1, \dots, \vec{v}_k$ are l.i.

adding \vec{v}_{k+1} are still l.i $\Leftrightarrow \vec{v}_{k+1}$ not in $\text{Span}\{\vec{v}_1, \dots, \vec{v}_k\}$

Pf: \Leftarrow) • If $\vec{v}_1, \dots, \vec{v}_{k+1}$ l.d.

then $c_1 \vec{v}_1 + \dots + c_{k+1} \vec{v}_{k+1} = 0$ for some c_i not all zero.

• $c_{k+1} \neq 0$ because $\vec{v}_1, \dots, \vec{v}_k$ l.i.

• then $\vec{v}_{k+1} = -\frac{c_1}{c_{k+1}} \vec{v}_1 - \dots - \frac{c_k}{c_{k+1}} \vec{v}_k$

in $\text{Span}\{\vec{v}_1, \dots, \vec{v}_k\}$.

Linear independent

fact:

Suppose $\vec{v}_1, \dots, \vec{v}_k$ are l.i.

adding \vec{v}_{k+1} are still l.i $\Leftrightarrow \vec{v}_{k+1}$ not in $\text{Span}\{\vec{v}_1, \dots, \vec{v}_k\}$

Pf: \Leftarrow) • If $\vec{v}_1, \dots, \vec{v}_{k+1}$ l.d.

then $c_1 \vec{v}_1 + \dots + c_{k+1} \vec{v}_{k+1} = 0$ for some c_i not all zero.

• $c_{k+1} \neq 0$ because $\vec{v}_1, \dots, \vec{v}_k$ l.i.

• then $\vec{v}_{k+1} = -\frac{c_1}{c_{k+1}} \vec{v}_1 - \dots - \frac{c_k}{c_{k+1}} \vec{v}_k$

in $\text{Span}\{\vec{v}_1, \dots, \vec{v}_k\}$.

\Rightarrow) similarly.

Basis

Basis

Def:

$\vec{v}_1, \dots, \vec{v}_n$ is called a basis for V if

Basis

Def:

$\vec{v}_1, \dots, \vec{v}_n$ is called a basis for V if

1. $\vec{v}_1, \dots, \vec{v}_n$ linearly independent

Basis

Def:

$\vec{v}_1, \dots, \vec{v}_n$ is called a basis for V if

1. $\vec{v}_1, \dots, \vec{v}_n$ linearly independent
2. $\text{Span}\{\vec{v}_1, \dots, \vec{v}_n\} = V.$

Basis

Def:

$\vec{v}_1, \dots, \vec{v}_n$ is called a basis for V if

1. $\vec{v}_1, \dots, \vec{v}_n$ linearly independent
2. $\text{Span}\{\vec{v}_1, \dots, \vec{v}_n\} = V.$

e.g. $e_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, e_2 = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, e_3 = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$ basis for \mathbb{R}^3 .

Basis

Def:

$\vec{v}_1, \dots, \vec{v}_n$ is called a basis for V if

1. $\vec{v}_1, \dots, \vec{v}_n$ linearly independent
2. $\text{Span}\{\vec{v}_1, \dots, \vec{v}_n\} = V.$

e.g. $e_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, e_2 = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, e_3 = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$ basis for \mathbb{R}^3 .

e.g. $\vec{v}_1 = \begin{bmatrix} 1 \\ 2 \end{bmatrix}, \vec{v}_2 = \begin{bmatrix} 1 \\ 3 \end{bmatrix}$ basis for \mathbb{R}^2 .

Basis

Def:

$\vec{v}_1, \dots, \vec{v}_n$ is called a basis for V if

1. $\vec{v}_1, \dots, \vec{v}_n$ linearly independent
2. $\text{Span}\{\vec{v}_1, \dots, \vec{v}_n\} = V.$

e.g. $e_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, e_2 = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, e_3 = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$ basis for \mathbb{R}^3 .

e.g. $\vec{v}_1 = \begin{bmatrix} 1 \\ 2 \end{bmatrix}, \vec{v}_2 = \begin{bmatrix} 1 \\ 3 \end{bmatrix}$ basis for \mathbb{R}^2 .

Person: $N(A) = 0, C(A) = \mathbb{R}^2$ with $A = [\vec{v}_1, \vec{v}_2]$.

Basis

Basis

- $\vec{V}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ is Not a basis for \mathbb{R}^2
 $\therefore \text{Span} \left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix} \right\} \neq \mathbb{R}^2$

Basis

- $\vec{V}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ is Not a basis for \mathbb{R}^2
 $\therefore \text{Span} \left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix} \right\} \neq \mathbb{R}^2$
- $\vec{V}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \vec{V}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \vec{V}_3 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ is Not a basis \mathbb{R}^2 .
 $\therefore \vec{V}_1, \vec{V}_2, \vec{V}_3$ are l.d.

Basis

- $\vec{V}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ is Not a basis for \mathbb{R}^2
 $\therefore \text{Span} \left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix} \right\} \neq \mathbb{R}^2$
- $\vec{V}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \vec{V}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \vec{V}_3 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ is Not a basis \mathbb{R}^2 .
 $\therefore \vec{V}_1, \vec{V}_2, \vec{V}_3$ are l.d.
- A Basis is an optimal spanning set of V .

Basis

- $\vec{v}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ is Not a basis for \mathbb{R}^2
 $\therefore \text{Span}\left\{\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right\} \neq \mathbb{R}^2$
- $\vec{v}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \vec{v}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \vec{v}_3 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ is Not a basis \mathbb{R}^2 .
 $\therefore \vec{v}_1, \vec{v}_2, \vec{v}_3$ are l.d.
- A Basis is an optimal spanning set of V .
- Basis is NOT unique: $\vec{e}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \vec{e}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$ is a basis for \mathbb{R}^2 .

Basis

- $\vec{v}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ is Not a basis for \mathbb{R}^2
 $\therefore \text{Span}\left\{\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right\} \neq \mathbb{R}^2$
- $\vec{v}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \vec{v}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \vec{v}_3 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ is Not a basis \mathbb{R}^2 .
 $\therefore \vec{v}_1, \vec{v}_2, \vec{v}_3$ are l.d.
- A Basis is an optimal spanning set of V .
- Basis is NOT unique: $\vec{e}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \vec{e}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$ is a basis for \mathbb{R}^2 .
 $\vec{v}_1 = \begin{bmatrix} 1 \\ 2 \end{bmatrix}, \vec{v}_2 = \begin{bmatrix} 1 \\ 3 \end{bmatrix}$ is also.

Basis

Basis

e.g.

Consider $\mathcal{U} = \begin{bmatrix} 1 & 3 & 3 & 2 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 0 \end{bmatrix}$ and $C(\mathcal{U})$.

Basis

e.g.

Consider

$$U = \begin{bmatrix} 1 & 3 & 3 & 2 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

↑
pivot ↑
column.

and $C(U)$.

Basis

e.g.

Consider $\mathcal{U} = \begin{bmatrix} 1 & 3 & 3 & 2 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 0 \end{bmatrix}$ and $C(\mathcal{U})$.

↑
pivot ↑
column.

- $C(\mathcal{U}) = \text{Span} \left\{ \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 3 \\ 1 \\ 0 \end{pmatrix} \right\}$

Basis

e.g.

Consider $\mathcal{U} = \begin{bmatrix} 1 & 3 & 3 & 2 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 0 \end{bmatrix}$ and $C(\mathcal{U})$.

↑
pivot ↑
column.

- $C(\mathcal{U}) = \text{Span} \left\{ \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 3 \\ 1 \\ 0 \end{pmatrix} \right\}$
- $\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 3 \\ 1 \\ 0 \end{pmatrix}$ are l.i. as $N\left(\begin{pmatrix} 1 & 3 \\ 0 & 1 \\ 0 & 0 \end{pmatrix}\right) = \{0\}$.

Basis

e.g.

Consider $\mathcal{U} = \begin{bmatrix} 1 & 3 & 3 & 2 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 0 \end{bmatrix}$ and $C(\mathcal{U})$.

\uparrow \uparrow
pivot column.

- $C(\mathcal{U}) = \text{Span} \left\{ \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 3 \\ 1 \\ 0 \end{pmatrix} \right\}$
- $\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 3 \\ 1 \\ 0 \end{pmatrix}$ are l.i. as $N\left(\begin{pmatrix} 1 & 3 \\ 0 & 1 \\ 0 & 0 \end{pmatrix}\right) = \{0\}$.
- Pivot columns is a set of basis for $C(\mathcal{U})$ in the case that \mathcal{U} is in Row Echelon form.

Basis

Basis

- Assuming we have A be general $m \times n$ matrix.

Basis

- Assuming we have A be general $m \times n$ matrix.
- $EA = U$

Basis

- Assuming we have A be general $m \times n$ matrix.
- $EA = U$
Product of elementary matrices

Basis


- Assuming we have A be general $m \times n$ matrix.
- $EA = U$ Row Echelon form.
↑
Product of elementary matrices

Basis

- Assuming we have A be general $m \times n$ matrix.

- $EA = U$ Row Echelon form.

↑
Product of elementary matrices




Basis

- Assuming we have A be general $m \times n$ matrix.

- $EA = U$ Row Echelon form.

↑
Product of elementary matrices



Basis

- Assuming we have A be general $m \times n$ matrix.
- $EA = U$ Row Echelon form.
↑
Product of elementary matrices
- The columns in A correspond to the pivot column from a basis of (CA) .


A diagram of a 4x4 matrix in Row Echelon form. The matrix is enclosed in large parentheses. It has non-zero entries (marked with *) at positions (1,1), (1,2), (2,3), (3,4), (2,4), and (3,3). Red arrows point from below to the first four columns, labeled "pivot columns".

Basis

- Assuming we have A be general $m \times n$ matrix.

- $EA = U$ Row Echelon form.

↑
Product of elementary matrices



- The columns in A correspond to the pivot column form a basis of $C(A)$.


- $$C(A) = \{ E^{-1} \vec{u} \mid \vec{u} \in C(U) \}$$

Basis

- Assuming we have A be general $m \times n$ matrix.

- $EA = U$ Row Echelon form.

Product of elementary matrices



- The columns in A correspond to the pivot column form a basis of $C(A)$.

- $C(A) = \{E^{-1}\vec{u} \mid \vec{u} \in C(U)\}$

fact: $\vec{v}_1, \dots, \vec{v}_k$ basis for $W \iff E\vec{v}_1, \dots, E\vec{v}_k$ basis for $E(W)$ for invertible E .

Basis

Basis

e.g. 1 Let $V = \{ a_0 + a_1x + a_2x^2 + a_3x^3 \mid a_i \in \mathbb{R} \}$.

then $1, x, x^2, x^3$ for a basis for V .

Basis

e.g. 1 Let $V = \{ a_0 + a_1x + a_2x^2 + a_3x^3 \mid a_i \in \mathbb{R} \}$.

then $1, x, x^2, x^3$ for a basis for V .

2. take $M_{2 \times 2}(\mathbb{R}) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mid a, b, c, d \in \mathbb{R} \right\}$

then $\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$

for a basis for $M_{2 \times 2}(\mathbb{R})$.

Basis

e.g. 1 Let $V = \{ a_0 + a_1x + a_2x^2 + a_3x^3 \mid a_i \in \mathbb{R} \}$.

then $1, x, x^2, x^3$ for a basis for V .

2. take $M_{2 \times 2}(\mathbb{R}) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mid a, b, c, d \in \mathbb{R} \right\}$

then $\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$

for a basis for $M_{2 \times 2}(\mathbb{R})$.

3. let $S_{2 \times 2}(\mathbb{R}) = \{ A \mid A \in M_{2 \times 2}(\mathbb{R}), A^T = A \}$

then $\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ for a basis

Basis

Basis

- A basis $\vec{v}_1, \dots, \vec{v}_k$ for V is an optimal spanning set of V :

Basis

- A basis $\vec{v}_1, \dots, \vec{v}_k$ for V is an optimal spanning set of V :
 1. take any \vec{v}_i away will not span V

Basis

- A basis $\vec{v}_1, \dots, \vec{v}_k$ for V is an optimal spanning set of V :
 1. take any \vec{v}_i away will not span V
 \therefore say \vec{v}_i : if $\text{Span}\{\vec{v}_1, \dots, \vec{v}_k\} = V$.

Basis

- A basis $\vec{v}_1, \dots, \vec{v}_k$ for V is an optimal spanning set of V :

1. take any \vec{v}_i away will not span V

\therefore say \vec{v}_i : if $\text{Span}\{\vec{v}_1, \dots, \vec{v}_k\} = V$.

then $\vec{v}_i \in \text{Span}\{\vec{v}_1, \dots, \vec{v}_k\}$

$\Rightarrow \vec{v}_1, \dots, \vec{v}_k$ are l.d.

Basis

- A basis $\vec{v}_1, \dots, \vec{v}_k$ for V is an optimal spanning set of V :

1. take any \vec{v}_i away will not span V

\therefore say \vec{v}_i : if $\text{Span}\{\vec{v}_1, \dots, \vec{v}_k\} = V$.

then $\vec{v}_i \in \text{Span}\{\vec{v}_1, \dots, \vec{v}_k\}$

$\Rightarrow \vec{v}_1, \dots, \vec{v}_k$ are l.d.

2. Adding any vector \vec{v}_{k+1} cannot be l.i.

Basis

- A basis $\vec{v}_1, \dots, \vec{v}_k$ for V is an optimal spanning set of V :

1. take any \vec{v}_i away will not span V

\therefore say \vec{v}_i : if $\text{Span}\{\vec{v}_1, \dots, \vec{v}_k\} = V$.

then $\vec{v}_i \in \text{Span}\{\vec{v}_1, \dots, \vec{v}_k\}$

$\Rightarrow \vec{v}_1, \dots, \vec{v}_k$ are l.d.

2. Adding any vector \vec{v}_{k+1} cannot be l.i.

$\vec{v}_{k+1} \in V = \text{Span}\{\vec{v}_1, \dots, \vec{v}_k\}$.

Basis

Basis

E.g.: 4. Let $sl_2(\mathbb{R}) := \left\{ A \in M_{2 \times 2}(\mathbb{R}) \mid \text{tr}(A) = 0 \right\}$.

then we have $\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$

being a basis

Basis

E.g.: 4. Let $sl_2(\mathbb{R}) := \left\{ A \in M_{2 \times 2}(\mathbb{R}) \mid \text{tr}(A) = 0 \right\}$.

then we have $\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$

being a basis

5. Let \mathbb{C} be the complex number consisting of
 $\{a+bi \mid a, b \in \mathbb{R}\}$, treated as \mathbb{R} -vector space.

then $1, i$ is a basis for \mathbb{C} .

Basis

E.g.: 4. Let $sl_2(\mathbb{R}) := \{ A \in M_{2 \times 2}(\mathbb{R}) \mid \text{tr}(A) = 0 \}$.

then we have $\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$

being a basis

5. Let \mathbb{C} be the complex number consisting of
 $\{ a+bi \mid a, b \in \mathbb{R} \}$, treated as \mathbb{R} -vector space.

then $1, i$ is a basis for \mathbb{C} .

6. Let $so(3) = \{ A \mid A \in M_{3 \times 3}(\mathbb{R}), A^T = -A \}$

$\begin{pmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & -1 & 0 \end{pmatrix}$ is a basis.

Basis

Basis

Prop: If V is finite dimensional, then V has a basis

Basis

Prop: If V is finite dimensional, then V has a basis

Pf: • Let $\{\vec{v}_1, \dots, \vec{v}_k\}$ be a spanning set.

i.e. $\text{Span}\{\vec{v}_1, \dots, \vec{v}_k\} = V$

Basis

Prop: If V is finite dimensional, then V has a basis

Pf: • Let $\{\vec{v}_1, \dots, \vec{v}_k\}$ be a spanning set.

i.e. $\text{Span}\{\vec{v}_1, \dots, \vec{v}_k\} = V$ (finite dimensional)

Basis

Prop: If V is finite dimensional, then V has a basis

Pf: • Let $\{\vec{v}_1, \dots, \vec{v}_k\}$ be a spanning set.
i.e. $\text{Span}\{\vec{v}_1, \dots, \vec{v}_k\} = V$ (finite dimensional)

• Let say there is a non-zero element, say \vec{v}_i :

Basis

Prop: If V is finite dimensional, then V has a basis

Pf: • Let $\{\vec{v}_1, \dots, \vec{v}_k\}$ be a spanning set.
i.e. $\text{Span}\{\vec{v}_1, \dots, \vec{v}_k\} = V$ (finite dimensional)

• Let say there is a non-zero element, say \vec{v}_1 :
then let $S_1 = \text{Span}\{\vec{v}_1\}$, two cases:

Basis

Prop: If V is finite dimensional, then V has a basis

Pf: • Let $\{\vec{v}_1, \dots, \vec{v}_k\}$ be a spanning set.
i.e. $\text{Span}\{\vec{v}_1, \dots, \vec{v}_k\} = V$ (finite dimensional)

• Let say there is a non-zero element, say \vec{v}_1 :
then let $S_1 = \text{Span}\{\vec{v}_1\}$, two cases:
1. $\text{Span}\{\vec{v}_1\} = V$ done!

Basis

Prop: If V is finite dimensional, then V has a basis

Pf: • Let $\{\vec{v}_1, \dots, \vec{v}_k\}$ be a spanning set.
i.e. $\text{Span}\{\vec{v}_1, \dots, \vec{v}_k\} = V$ (finite dimensional)

- Let say there is a non-zero element, say \vec{v}_1 :
then let $S_1 = \text{Span}\{\vec{v}_1\}$, two cases:
 1. $\text{Span}\{\vec{v}_1\} = V$ done!
 2. $\text{Span}\{\vec{v}_1\} \neq V$. continue.

Basis

Basis

- If $\text{Span}\{\vec{v}_i\} \neq V$, then there must be a \vec{v}_i s.t.
 $\vec{v}_i \notin \text{Span}\{\vec{v}_i\}$, say it is \vec{v}_2

Basis

- If $\text{Span}\{\vec{v}_i\} \neq V$, then there must be a \vec{v}_i s.t.
 $\vec{v}_i \notin \text{Span}\{\vec{v}_j\}$, say it is \vec{v}_2 (after rearranging
the index).

Basis

- If $\text{Span}\{\vec{v}_i\} \neq V$, then there must be a \vec{v}_i s.t.
 $\vec{v}_i \notin \text{Span}\{\vec{v}_j\}$, say it is \vec{v}_2 (after rearranging
the index).
- \vec{v}_1, \vec{v}_2 l.i., let $S_2 = \text{Span}\{\vec{v}_1, \vec{v}_2\}$.

Basis

- If $\text{Span}\{\vec{v}_1\} \neq V$, then there must be a \vec{v}_i s.t.
 $\vec{v}_i \notin \text{Span}\{\vec{v}_1\}$, say it is \vec{v}_2 (after rearranging
the index).
- \vec{v}_1, \vec{v}_2 l.i., let $S_2 = \text{Span}\{\vec{v}_1, \vec{v}_2\}$.
two cases: 1. $S_2 = V$ done!

Basis

- If $\text{Span}\{\vec{v}_1\} \neq V$, then there must be a \vec{v}_i s.t.
 $\vec{v}_i \notin \text{Span}\{\vec{v}_1\}$, say it is \vec{v}_2 (after rearranging
the index).
- \vec{v}_1, \vec{v}_2 l.i., let $S_2 = \text{Span}\{\vec{v}_1, \vec{v}_2\}$.
two cases: 1. $S_2 = V$ done!
2. $S_2 \neq V$ continue!

Basis

- If $\text{Span}\{\vec{v}_1\} \neq V$, then there must be a \vec{v}_i s.t.
 $\vec{v}_i \notin \text{Span}\{\vec{v}_1\}$, say it is \vec{v}_2 (after rearranging the index).
- \vec{v}_1, \vec{v}_2 l.i., let $S_2 = \text{Span}\{\vec{v}_1, \vec{v}_2\}$.
two cases: 1. $S_2 = V$ done!
2. $S_2 \neq V$ continue!
- ... we must arrive at a situation that
 $S_l = V$ for some $l \leq k$.

Basis

- If $\text{Span}\{\vec{v}_1\} \neq V$, then there must be a \vec{v}_i s.t.
 $\vec{v}_i \notin \text{Span}\{\vec{v}_1\}$, say it is \vec{v}_2 (after rearranging the index).
- \vec{v}_1, \vec{v}_2 l.i., let $S_2 = \text{Span}\{\vec{v}_1, \vec{v}_2\}$.
two cases: 1. $S_2 = V$ done!
2. $S_2 \neq V$ continue!
- ... we must arrive at a situation that
 $S_l = V$ for some $l \leq k$.
 $\text{Span}\{\vec{v}_1, \dots, \overset{||}{\vec{v}_l}\}$

Basis

- If $\text{Span}\{\vec{v}_1\} \neq V$, then there must be a \vec{v}_i s.t.
 $\vec{v}_i \notin \text{Span}\{\vec{v}_1\}$, say it is \vec{v}_2 (after rearranging the index).
- \vec{v}_1, \vec{v}_2 l.i., let $S_2 = \text{Span}\{\vec{v}_1, \vec{v}_2\}$.
two cases: 1. $S_2 = V$ done!
2. $S_2 \neq V$ continue!
- ... we must arrive at a situation that
 $S_l = V$ for some $l \leq k$.
 $\text{Span}\{\vec{v}_1, \dots, \overset{||}{\vec{v}_l}\}$ then $\vec{v}_1, \dots, \vec{v}_l$ is a basis for V

Basis

Basis

- Indeed we prove that any spanning set $\{\vec{v}_1, \dots, \vec{v}_k\}$ of V can be reduced to a basis.

Basis

- Indeed we prove that any spanning set $\{\vec{v}_1, \dots, \vec{v}_k\}$ of V can be reduced to a basis.

Prop: Suppose V is finite dimensional, then any l.i. subset $\vec{v}_1, \dots, \vec{v}_k$ can be extended to a basis.

Basis

- Indeed we prove that any spanning set $\{\vec{v}_1, \dots, \vec{v}_k\}$ of V can be reduced to a basis.

Prop: Suppose V is finite dimensional, then any l.i. subset $\vec{v}_1, \dots, \vec{v}_k$ can be extended to a basis.

Pf: • Let $\vec{w}_1, \dots, \vec{w}_n$ be a spanning set for V

Basis

- Indeed we prove that any spanning set $\{\vec{v}_1, \dots, \vec{v}_k\}$ of V can be reduced to a basis.

Prop: Suppose V is finite dimensional, then any l.i. subset $\vec{v}_1, \dots, \vec{v}_k$ can be extended to a basis.

Pf: • Let $\vec{w}_1, \dots, \vec{w}_n$ be a spanning set for V

1. If $\text{Span}\{\vec{v}_1, \dots, \vec{v}_k\} = V$ done!

Basis

- Indeed we prove that any spanning set $\{\vec{v}_1, \dots, \vec{v}_k\}$ of V can be reduced to a basis.

Prop: Suppose V is finite dimensional, then any l.i. subset $\vec{v}_1, \dots, \vec{v}_k$ can be extended to a basis.

Pf:

- Let $\vec{w}_1, \dots, \vec{w}_n$ be a spanning set for V
 - If $\text{Span}\{\vec{v}_1, \dots, \vec{v}_k\} = V$ done!
 - If not, then exists $\vec{w}_i \notin \text{Span}\{\vec{v}_1, \dots, \vec{v}_k\}$
Let $\vec{v}_{k+1} = \vec{w}_i$ and add it to $\vec{v}_1, \dots, \vec{v}_k$.

Basis

- Indeed we prove that any spanning set $\{\vec{v}_1, \dots, \vec{v}_k\}$ of V can be reduced to a basis.

Prop: Suppose V is finite dimensional, then any l.i. subset $\vec{v}_1, \dots, \vec{v}_k$ can be extended to a basis.

- Pf:
- Let $\vec{w}_1, \dots, \vec{w}_n$ be a spanning set for V
 - If $\text{Span}\{\vec{v}_1, \dots, \vec{v}_k\} = V$ done!
 - If not, then exists $\vec{w}_i \notin \text{Span}\{\vec{v}_1, \dots, \vec{v}_k\}$
Let $\vec{v}_{k+1} = \vec{w}_i$ and add it to $\vec{v}_1, \dots, \vec{v}_k$.
 - Repeat 1. until it stop.

Basis

Basis

Side story:

Recall $\mathbb{R}^{\infty} = \{ (a_1, a_2, a_3, \dots) \mid a_i \in \mathbb{R} \}$.

Basis

Side story: Recall $\mathbb{R}^{\infty} = \{(a_1, a_2, a_3, \dots) \mid a_i \in \mathbb{R}\}$.

it is infinite dimensional, does it have a basis?

Basis

Side story: Recall $\mathbb{R}^{\infty} = \{ (a_1, a_2, a_3, \dots) \mid a_i \in \mathbb{R} \}$.

it is infinite dimensional, does it have a basis?

- We know that $e_i = (0 \dots, \underset{i\text{-th place}}{\overset{\uparrow}{1}}, 0 \dots)$

i-th place

is **NOT** a spanning set.

Basis

Side story: Recall $\mathbb{R}^{\infty} = \{ (a_1, a_2, a_3, \dots) \mid a_i \in \mathbb{R} \}$.

it is infinite dimensional, does it have a basis?

- We know that $e_i = (0 \dots, \underset{i\text{-th place}}{\overset{\uparrow}{1}}, 0 \dots)$

i-th place

is **NOT** a spanning set.

In fact: One can prove that any vector space has a basis, but require the use of Zorn's lemma / axiom of choice.

Dimension

Dimension

Def.

If V is a vector space with basis $\vec{v}_1, \dots, \vec{v}_n$

we say that the dimension of V is n .

or V is n -dimensional.

Dimension

Def.

If V is a vector space with basis $\vec{v}_1, \dots, \vec{v}_n$

we say that the dimension of V is n .

or V is n -dimensional.

e.g.: dimension of \mathbb{R}^n is n .

Dimension

Def.

If V is a vector space with basis $\vec{v}_1, \dots, \vec{v}_n$

we say that the dimension of V is n .

or V is n -dimensional.

e.g.: 1. dimension of \mathbb{R}^n is n .

2. dimension of $sl_2(\mathbb{R})$ is 3.

Dimension

Def.

If V is a vector space with basis $\vec{v}_1, \dots, \vec{v}_n$

we say that the dimension of V is n .

or V is n -dimensional.

e.g.: 1. dimension of \mathbb{R}^n is n .

2. dimension of $sl_2(\mathbb{R})$ is 3.

3. dimension of $\underline{so}(3)$ is 3.

Dimension

Dimension

Prop: Dimension of V is well-defined.

Dimension

Prop: Dimension of V is well-defined.

Pf:

Let v_1, \dots, v_m be a basis

w_1, \dots, w_n be another basis.

Dimension

Prop: Dimension of V is well-defined.

Pf: Let v_1, \dots, v_m be a basis
 w_1, \dots, w_n be another basis.

Assume $n > m$: Say $n=3, m=2$:

Dimension

Prop: Dimension of V is well-defined.

Pf: Let v_1, \dots, v_m be a basis
 w_1, \dots, w_n be another basis.

Assume $n > m$: Say $n=3, m=2$:

Write: $w_1 = a_{11}v_1 + a_{21}v_2$

$$w_2 = a_{12}v_1 + a_{22}v_2$$

$$w_3 = a_{13}v_1 + a_{23}v_2.$$

Dimension

Dimension

- Let $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \end{bmatrix}$

Dimension

- Let $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \end{bmatrix}$
- $Ax=0$ has a non-zero solution $\begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix}$.

Dimension

- Let $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \end{bmatrix}$
- $Ax=0$ has a non-zero solution $\begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix}$.

Formally we write:

$$[W_1, W_2, W_3] = [V_1, V_2] \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \end{bmatrix}$$

Dimension

- Let $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \end{bmatrix}$
- $Ax=0$ has a non-zero solution $\begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix}$.

Formally we write:

$$[W_1, W_2, W_3] = [V_1, V_2] \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \end{bmatrix}$$

then we see that:

$$[W_1, W_2, W_3] \begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix} = 0$$

Dimension

- Let $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \end{bmatrix}$
- $Ax=0$ has a non-zero solution $\begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix}$.

Formally we write:

$$[W_1, W_2, W_3] = [V_1, V_2] \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \end{bmatrix}$$

then we see that:

$$[W_1, W_2, W_3] \begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix} = 0 \Rightarrow c_1 W_1 + c_2 W_2 + c_3 W_3 = 0$$

$\overrightarrow{\overleftarrow{}}$

Dimension

Dimension

- Recall that we define the rank of a matrix A to be the # of pivot columns in its Row Echelon form \mathcal{U}

Dimension

- Recall that we define the rank of a matrix A to be the # of pivot columns in its Row Echelon form \mathcal{U} .
- Intrinsic definition: $\text{rk}(A) = \dim(C(A))$.

Dimension

- Recall that we define the rank of a matrix A to be the # of pivot columns in its Row Echelon form U .
- Intrinsic definition: $\text{rk}(A) = \dim(C(A))$.

Reason: 1. For U : we see that

$$\dim(C(U)) = \text{rk}(U) = \# \text{ of pivots columns.}$$

Dimension

- Recall that we define the rank of a matrix A to be the # of pivot columns in its Row Echelon form U .
- Intrinsic definition: $\text{rk}(A) = \dim(C(A))$.

Reason: 1. For U : we see that

$$\dim(C(U)) = \text{rk}(U) = \# \text{ of pivots columns.}$$

2. If $A = EU$ for invertible E .

$$\Rightarrow C(A) = \left\{ E\vec{u} \mid \vec{u} \in C(\vec{u}) \right\}.$$

Dimension

- Recall that we define the rank of a matrix A to be the # of pivot columns in its Row Echelon form U .
- Intrinsic definition: $\text{rk}(A) = \dim(C(A))$.

Reason: 1. For U : we see that

$$\dim(C(U)) = \text{rk}(U) = \# \text{ of pivots columns.}$$

2. If $A = EU$ for invertible E .

$$\Rightarrow C(A) = \left\{ E\vec{u} \mid \vec{u} \in C(\vec{u}) \right\}.$$

and $E\vec{u}_1, \dots, E\vec{u}_r$ is a basis for $C(A)$ if $\vec{u}_1, \dots, \vec{u}_r$ is for $C(U)$.

Dimension

Dimension

Prop:

$\dim(N(A)) = \# \text{ of non-pivot columns}$
in the Row Echelon form.

Dimension

Prop: $\dim(N(A)) = \# \text{ of non-pivot columns}$
in the Row Echelon form.

Pf: • Let say $U = \begin{pmatrix} 0 & 0 & 0 \\ 1 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$

Dimension

Prop: $\dim(N(A)) = \# \text{ of non-pivot columns}$
in the Row Echelon form.

Pf: • Let say $U = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \end{pmatrix}$

Dimension

Prop: $\dim(N(A)) = \# \text{ of non-pivot columns}$
in the Row Echelon form.

Pf: • Let say $U = \begin{pmatrix} x_1 & x_2 & x_3 & x_4 & x_5 & x_6 \\ \left(\begin{array}{c|c|c|c} 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{array} \right) \end{pmatrix}$

Dimension

Prop: $\dim(N(A)) = \# \text{ of non-pivot columns}$
in the Row Echelon form.

Pf: • Let say $U = \left(\begin{array}{cccccc} x_1 & x_2 & x_3 & x_4 & x_5 & x_6 \\ \boxed{0} & \boxed{1} & \boxed{0} & \boxed{1} & \boxed{0} & \boxed{1} \\ \hline 1 & & & 1 & & 1 \end{array} \right)$

• $U = \text{Span} \left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\}$.

Dimension

Prop: $\dim(N(A)) = \# \text{ of non-pivot columns}$
in the Row Echelon form.

Pf: • Let say $U = \begin{pmatrix} x_1 & x_2 & x_3 & x_4 & x_5 & x_6 \\ 0 & 1 & 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 & 1 & 0 \end{pmatrix}$

• $U = \text{Span} \left\{ \begin{bmatrix} * \\ 1 \\ * \\ 0 \\ * \\ 0 \end{bmatrix}, \begin{bmatrix} * \\ 0 \\ * \\ 1 \\ * \\ 0 \end{bmatrix}, \begin{bmatrix} * \\ 0 \\ * \\ 0 \\ * \\ 1 \end{bmatrix} \right\}$.

Dimension

Prop: $\dim(N(A)) = \# \text{ of non-pivot columns}$
in the Row Echelon form.

Pf: • Let say $U = \begin{pmatrix} x_1 & x_2 & x_3 & x_4 & x_5 & x_6 \\ 0 & 1 & 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 & 1 & 0 \end{pmatrix}$

• $U = \text{Span} \left\{ \begin{bmatrix} * \\ -1 \\ * \\ 0 \\ * \\ 0 \end{bmatrix}, \begin{bmatrix} * \\ 0 \\ * \\ 1 \\ * \\ 0 \end{bmatrix}, \begin{bmatrix} * \\ 0 \\ * \\ 0 \\ * \\ -1 \end{bmatrix} \right\}$.

• These three vectors are l.i. !

Dimension

Dimension

- $\dim(C(A)) + \dim(N(A)) = n$ for $m \times n$ A .

Dimension

- $\dim(C(A)) + \dim(N(A)) = n$ for $m \times n$ A.


Dimension

Prop: If W is a subspace of V , then $\dim(W) \leq \dim(V)$.

Dimension

Prop: If W is a subspace of V , then $\dim(W) \leq \dim(V)$.

Pf: • take basis of W , say w_1, \dots, w_k l.i.

Dimension

Prop: If W is a subspace of V , then $\dim(W) \leq \dim(V)$.

Pf: • take basis of W , say w_1, \dots, w_k l.i.

- It can be extended to a basis of V .

$$w_1, \dots, w_k, w_{k+1}, \dots, w_n$$

Dimension

Prop: If W is a subspace of V , then $\dim(W) \leq \dim(V)$.

Pf: • take basis of W , say w_1, \dots, w_k l.i.

- It can be extended to a basis of V .

$w_1, \dots, w_k, \underbrace{w_{k+1}, \dots, w_n}$

adding more vectors in V.

Dimension

Dimension

e.g. • In \mathbb{R}^4 , we can only have dimension = 0, 1, 2, 3, 4
subspaces.

Dimension

e.g. • In \mathbb{R}^4 , we can only have dimension = 0, 1, 2, 3, 4 subspaces.

Rk: • We must say $\vec{v} = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}$ being a four dimensional vector, meaning it is a vector in \mathbb{R}^4

Dimension

e.g. • In \mathbb{R}^4 , we can only have dimension = 0, 1, 2, 3, 4 subspaces.

Rk: • We must say $\vec{v} = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}$ being a four dimensional vector, meaning it is a vector in \mathbb{R}^4 .

• But $\text{Span}\{\vec{v}\}$ is 1-dimensional subspace of \mathbb{R}^4 .

Dimension

e.g.: In \mathbb{R}^4 , we can only have dimension = 0, 1, 2, 3, 4 subspaces.

- Rk:
- We must say $\vec{v} = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}$ being a four dimensional vector, meaning it is a vector in \mathbb{R}^4 .
 - But $\text{Span}\{\vec{v}\}$ is 1-dimensional subspace of \mathbb{R}^4 .
 - So the word "four dimensional vector" here refer to its ambient space