

Lecture 4

Inverse and transpose matrices

Partitioned matrices

Partitioned matrices

- Recall: A : a $m \times k$ matrix
 B : a $k \times n$ matrix

We treat $B = [\vec{b}_1, \vec{b}_2, \dots, \vec{b}_n]$

Partitioned matrices

- Recall: A : a $m \times k$ matrix
 B : a $k \times n$ matrix

We treat $B = [\vec{b}_1, \vec{b}_2, \dots, \vec{b}_n]$

- The multiplication is given by

$$AB = [A\vec{b}_1, \dots, A\vec{b}_n]$$

i.e. column by column

Partitioned matrices

Partitioned matrices

- There is also a picture given by rows:

$$\begin{bmatrix} -a_1 - \\ \vdots \\ -a_i - \\ \vdots \\ -a_m - \end{bmatrix} B = \begin{bmatrix} (-a_1 -)B \\ \vdots \\ (-a_i -)B \\ \vdots \\ (-a_n -)B \end{bmatrix}$$

Partitioned matrices

- There is also a picture given by rows:

$$\begin{bmatrix} -a_1 - \\ \vdots \\ -a_i - \\ \vdots \\ -a_m - \end{bmatrix} B = \begin{bmatrix} (-a_1 -)B \\ \vdots \\ (-a_i -)B \\ \vdots \\ (-a_n -)B \end{bmatrix}$$

- These two are examples of more general block multiplication ('分塊乘法')

Partitioned matrices

Partitioned matrices

- In general we may partition a matrix into smaller block:

Partitioned matrices

- In general we may partition a matrix into smaller block:

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} & a_{15} \\ a_{21} & a_{22} & a_{23} & a_{24} & a_{25} \\ a_{31} & a_{32} & a_{33} & a_{34} & a_{35} \\ a_{41} & a_{42} & a_{43} & a_{44} & a_{45} \end{bmatrix}$$

Partitioned matrices

- In general we may partition a matrix into smaller block:

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} & a_{15} \\ a_{21} & a_{22} & a_{23} & a_{24} & a_{25} \\ a_{31} & a_{32} & a_{33} & a_{34} & a_{35} \\ a_{41} & a_{42} & a_{43} & a_{44} & a_{45} \end{bmatrix} = \begin{bmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{bmatrix}$$

Partitioned matrices

- In general we may partition a matrix into smaller block:

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} & a_{15} \\ a_{21} & a_{22} & a_{23} & a_{24} & a_{25} \\ a_{31} & a_{32} & a_{33} & a_{34} & a_{35} \\ a_{41} & a_{42} & a_{43} & a_{44} & a_{45} \end{bmatrix} = \begin{bmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{bmatrix}$$

A blue curved arrow points from the top-left 3x3 block $[a_{11}, a_{12}, a_{13}; a_{21}, a_{22}, a_{23}; a_{31}, a_{32}, a_{33}]$ to the first column of the matrix on the right. Another blue curved arrow points from the bottom-right 2x2 block $[a_{44}, a_{45}; a_{34}, a_{35}]$ to the second column of the matrix on the right.

Partitioned matrices

- In general we may partition a matrix into smaller block:

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} & a_{15} \\ a_{21} & a_{22} & a_{23} & a_{24} & a_{25} \\ a_{31} & a_{32} & a_{33} & a_{34} & a_{35} \\ a_{41} & a_{42} & a_{43} & a_{44} & a_{45} \end{bmatrix} = \begin{bmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{bmatrix}$$

The matrix on the left is partitioned into four blocks: a_{11}, a_{12}, a_{13} (top-left, blue), a_{14}, a_{15} (top-right, green), a_{21}, a_{22}, a_{23} (bottom-left, blue), and a_{24}, a_{25} (bottom-right, green). A blue curved arrow points from the top-left block to the C_{11} block.

Partitioned matrices

- In general we may partition a matrix into smaller block:

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} & a_{15} \\ a_{21} & a_{22} & a_{23} & a_{24} & a_{25} \\ a_{31} & a_{32} & a_{33} & a_{34} & a_{35} \\ a_{41} & a_{42} & a_{43} & a_{44} & a_{45} \end{bmatrix} = \begin{bmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{bmatrix}$$

The matrix on the left is partitioned into four blocks: a_{11}, a_{12}, a_{13} (blue), a_{14}, a_{15} (green), a_{31}, a_{32}, a_{33} (red), and a_{41}, a_{42}, a_{43} (pink). A blue curved arrow points from the top-right block to the right matrix, indicating the mapping of the partitioned matrix to its block-diagonal form.

Partitioned matrices

- In general we may partition a matrix into smaller block:

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \\ a_{41} & a_{42} & a_{43} \end{bmatrix} = \begin{bmatrix} a_{14} & a_{15} \\ a_{24} & a_{25} \\ a_{34} & a_{35} \\ a_{44} & a_{45} \end{bmatrix} = \begin{bmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{bmatrix}$$

The diagram illustrates the partitioning of a 4x5 matrix into four 2x3 submatrices. The top-left 2x3 block is highlighted in blue. The top-right 2x2 block is highlighted in green. The bottom-left 2x3 block is highlighted in red. The bottom-right 1x2 block is highlighted in purple. A blue curved arrow points from the top-left block to the bottom-right block, indicating a transformation or relationship between them.

Partitioned matrices

Partitioned matrices

- We can write the multiplication by a block partition:

Case 1:

Partitioned matrices

- We can write the multiplication by a block partition:

Case 1): $A [B_1, B_2] = [AB_1, AB_2]$.

Partitioned matrices

- We can write the multiplication by a block partition:

Case 1: $A [B_1, B_2] = [AB_1, AB_2]$.

$\begin{matrix} \uparrow & \uparrow & \uparrow \\ m \times k & k \times n_1 & k \times n_2 \end{matrix}$

Partitioned matrices

- We can write the multiplication by a block partition:

Case 1): $A [B_1, B_2] = [AB_1, AB_2]$.

$\begin{matrix} \uparrow & \uparrow & \uparrow \\ m \times k & k \times n_1 & k \times n_2 \end{matrix}$

i.e.

$$\begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} \begin{bmatrix} b_{11} & b_{12} & b_{13} \\ b_{21} & b_{22} & b_{23} \end{bmatrix} = \begin{bmatrix} (a_{11} a_{12}) \begin{pmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{pmatrix}, (a_{11} a_{12}) \begin{pmatrix} b_{13} \\ b_{23} \end{pmatrix} \\ (a_{21} a_{22}) \begin{pmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{pmatrix}, (a_{21} a_{22}) \begin{pmatrix} b_{13} \\ b_{23} \end{pmatrix} \end{bmatrix}$$

Case 2):

Partitioned matrices

- We can write the multiplication by a block partition:

Case 1): $A [B_1, B_2] = [AB_1, AB_2]$.

$\begin{matrix} \uparrow \\ m \times k \end{matrix}$ $\begin{matrix} \uparrow \\ k \times n_1 \end{matrix}$ $\begin{matrix} \uparrow \\ k \times n_2 \end{matrix}$ e.g.: B_1 B_2

i.e.

$$\begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} \begin{bmatrix} b_{11} & b_{12} & b_{13} \\ b_{21} & b_{22} & b_{23} \end{bmatrix} = \begin{bmatrix} (a_{11} a_{12}) \begin{pmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{pmatrix}, (a_{11} a_{12}) \begin{pmatrix} b_{13} \\ b_{23} \end{pmatrix} \\ (a_{21} a_{22}) \begin{pmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{pmatrix}, (a_{21} a_{22}) \begin{pmatrix} b_{13} \\ b_{23} \end{pmatrix} \end{bmatrix}$$

Case 2):

Partitioned matrices

- We can write the multiplication by a block partition:

Case 1: $A [B_1, B_2] = [AB_1, AB_2]$.

$\begin{matrix} \uparrow \\ m \times k \end{matrix}$ $\begin{matrix} \uparrow \\ k \times n_1 \end{matrix}$ $\begin{matrix} \uparrow \\ k \times n_2 \end{matrix}$ e.g.: B_1 B_2

i.e.

$$\begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} \begin{bmatrix} b_{11} & b_{12} & b_{13} \\ b_{21} & b_{22} & b_{23} \end{bmatrix} = \begin{bmatrix} (a_{11} a_{12}) \begin{pmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{pmatrix}, (a_{11} a_{12}) \begin{pmatrix} b_{13} \\ b_{23} \end{pmatrix} \\ (a_{21} a_{22}) \begin{pmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{pmatrix}, (a_{21} a_{22}) \begin{pmatrix} b_{13} \\ b_{23} \end{pmatrix} \end{bmatrix}$$

Case 2: $\begin{bmatrix} A_1 \\ A_2 \end{bmatrix} B = \begin{bmatrix} A_1 B \\ A_2 B \end{bmatrix}$

Partitioned matrices

- We can write the multiplication by a block partition:

Case 1): $A [B_1, B_2] = [AB_1, AB_2]$.

$\begin{matrix} \uparrow \\ m \times k \end{matrix}$ $\begin{matrix} \uparrow \\ k \times n_1 \end{matrix}$ $\begin{matrix} \uparrow \\ k \times n_2 \end{matrix}$ e.g.: B_1 B_2

i.e.

$$\begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} \begin{bmatrix} b_{11} & b_{12} & b_{13} \\ b_{21} & b_{22} & b_{23} \end{bmatrix} = \begin{bmatrix} (a_{11} a_{12}) (b_{11} & b_{12}) \\ (a_{21} a_{22}) (b_{21} & b_{22}) \end{bmatrix}, \begin{bmatrix} (a_{11} a_{12}) (b_{13}) \\ (a_{21} a_{22}) (b_{23}) \end{bmatrix}$$

Case 2): $\begin{bmatrix} A_1 \\ A_2 \end{bmatrix} B = \begin{bmatrix} A_1 B \\ A_2 B \end{bmatrix}$

$m_1 \times k$ $m_2 \times k$ $k \times n$

Partitioned matrices

Partitioned matrices

- Case 3) :

$$\begin{bmatrix} A_1 & A_2 \end{bmatrix} \begin{bmatrix} B_1 \\ B_2 \end{bmatrix} = A_1 B_1 + A_2 B_2$$

Partitioned matrices

- Case 3) :

$$\begin{bmatrix} A_1 & A_2 \end{bmatrix} \begin{bmatrix} B_1 \\ B_2 \end{bmatrix} = A_1 B_1 + A_2 B_2$$

$m \times k_1$ $m \times k_2$ $k_1 \times n$ $k_2 \times n$

Partitioned matrices

- Case 3) :

$$\begin{bmatrix} A_1 & A_2 \end{bmatrix} \begin{bmatrix} B_1 \\ B_2 \end{bmatrix} = \underbrace{A_1 B_1 + A_2 B_2}_{m \times n}.$$

Annotations:

- Dimensions: $m \times k_1$ for A_1 , $m \times k_2$ for A_2 .
- Dimensions: $k_1 \times n$ for B_1 , $k_2 \times n$ for B_2 .
- The resulting product is $m \times n$.

Partitioned matrices

• Case 3) :

$$\begin{bmatrix} A_1 & A_2 \end{bmatrix} \begin{bmatrix} B_1 \\ B_2 \end{bmatrix} = \underbrace{A_1 B_1 + A_2 B_2}_{m \times n}$$

Annotations:

- Matrix A is $m \times k_1$ and A_1 is $m \times n$.
- Matrix B is $k_1 \times n$ and B_1 is $k_1 \times n$.
- Matrix A is $m \times k_2$ and A_2 is $m \times n$.
- Matrix B is $k_2 \times n$ and B_2 is $k_2 \times n$.

e.g.

$$\begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} \begin{bmatrix} a_{13} \\ a_{23} \end{bmatrix} = b_1 \begin{bmatrix} a_{11} \\ a_{21} \end{bmatrix} + b_2 \begin{bmatrix} a_{12} \\ a_{22} \end{bmatrix} + b_3 \begin{bmatrix} a_{13} \\ a_{23} \end{bmatrix}$$

Partitioned matrices

• Case 3) :

$$\begin{bmatrix} A_1 & A_2 \end{bmatrix} \begin{bmatrix} B_1 \\ B_2 \end{bmatrix} = \underbrace{A_1 B_1}_{m \times n} + \underbrace{A_2 B_2}_{m \times n}$$

Annotations:

- Matrix A is partitioned into A_1 (blue) and A_2 (green), both $m \times k$.
- Matrix B is partitioned into B_1 (blue) and B_2 (green), both $k \times n$.
- The resulting matrix is $m \times n$.

e.g.

$$\begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix} = b_1 \begin{bmatrix} a_{11} \\ a_{21} \end{bmatrix} + b_2 \begin{bmatrix} a_{12} \\ a_{22} \end{bmatrix} + b_3 \begin{bmatrix} a_{13} \\ a_{23} \end{bmatrix}$$

$$= \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} \begin{bmatrix} b_1 \\ b_2 \end{bmatrix} + \begin{bmatrix} a_{13} \\ a_{23} \end{bmatrix} \begin{bmatrix} b_3 \end{bmatrix}$$

Partitioned matrices

• Case 3) :

$$\begin{bmatrix} A_1 & A_2 \end{bmatrix} \begin{bmatrix} B_1 \\ B_2 \end{bmatrix} = \underbrace{A_1 B_1}_{m \times n} + \underbrace{A_2 B_2}_{m \times n}$$

$m \times k_1$ $m \times k_2$ $k_1 \times n$ $k_2 \times n$

e.g.

$$\begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} \begin{bmatrix} a_{13} \\ a_{23} \end{bmatrix} = b_1 \begin{bmatrix} a_{11} \\ a_{21} \end{bmatrix} + b_2 \begin{bmatrix} a_{12} \\ a_{22} \end{bmatrix} + b_3 \begin{bmatrix} a_{13} \\ a_{23} \end{bmatrix}$$
$$= \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} \begin{bmatrix} b_1 \\ b_2 \end{bmatrix} + \begin{bmatrix} a_{13} \\ a_{23} \end{bmatrix} \begin{bmatrix} b_3 \end{bmatrix}$$

Partitioned matrices

Partitioned matrices

- Case 4:

$$\begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix} \begin{bmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{bmatrix} = \begin{bmatrix} A_{11}B_{11} + A_{12}B_{21} & A_{11}B_{12} + A_{12}B_{22} \\ A_{21}B_{11} + A_{22}B_{21} & A_{21}B_{12} + A_{22}B_{22} \end{bmatrix}$$

Partitioned matrices

- Case 4:

$$\begin{matrix} m_1 \\ m_2 \end{matrix} \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix} \begin{bmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{bmatrix} = \begin{bmatrix} A_{11}B_{11} + A_{12}B_{21} & A_{11}B_{12} + A_{12}B_{22} \\ A_{21}B_{11} + A_{22}B_{21} & A_{21}B_{12} + A_{22}B_{22} \end{bmatrix}$$

Partitioned matrices

• Case 4:

$$\begin{matrix} m_1 \\ m_2 \end{matrix} \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix} \begin{bmatrix} n_1 & n_2 \\ B_{11} & B_{12} \\ B_{21} & B_{22} \end{bmatrix} = \begin{bmatrix} A_{11}B_{11} + A_{12}B_{21} & A_{11}B_{12} + A_{12}B_{22} \\ A_{21}B_{11} + A_{22}B_{21} & A_{21}B_{12} + A_{22}B_{22} \end{bmatrix}$$

Partitioned matrices

• Case 4:

$$\begin{matrix} m_1 \\ m_2 \end{matrix} \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix} \begin{bmatrix} n_1 & n_2 \\ k_1 & k_2 \end{bmatrix} \begin{bmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{bmatrix} = \begin{bmatrix} n_1 & n_2 \\ m_1 & m_2 \end{bmatrix} \begin{bmatrix} A_{11}B_{11} + A_{12}B_{21} & A_{11}B_{12} + A_{12}B_{22} \\ A_{21}B_{11} + A_{22}B_{21} & A_{21}B_{12} + A_{22}B_{22} \end{bmatrix}$$

Partitioned matrices

- Case 4:

$$\begin{matrix} m_1 \\ m_2 \end{matrix} \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix} \begin{bmatrix} n_1 & n_2 \\ k_1 & k_2 \end{bmatrix} \begin{bmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{bmatrix} = \begin{bmatrix} n_1 & n_2 \\ m_1 & m_2 \end{bmatrix} \begin{bmatrix} A_{11}B_{11} + A_{12}B_{21} & A_{11}B_{12} + A_{12}B_{22} \\ A_{21}B_{11} + A_{22}B_{21} & A_{21}B_{12} + A_{22}B_{22} \end{bmatrix}$$

- The sizes of the matrices has to match.


Partitioned matrices

- Case 4:

$$\begin{matrix} m_1 \\ m_2 \end{matrix} \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix} \begin{bmatrix} n_1 & n_2 \\ n_1 & n_2 \end{bmatrix} = \begin{bmatrix} n_1 & n_2 \\ n_1 & n_2 \end{bmatrix} \begin{bmatrix} A_{11}B_{11} + A_{12}B_{21} & A_{11}B_{12} + A_{12}B_{22} \\ A_{21}B_{11} + A_{22}B_{21} & A_{21}B_{12} + A_{22}B_{22} \end{bmatrix}$$

- The sizes of the matrices has to match.
- In general: whenever the partition into block has the right sizes
block multiplication works!

Partitioned matrices




Partitioned matrices

• e.g.:

$$A = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 2 & 1 & 1 \\ 3 & 2 & 2 & 2 \end{bmatrix}$$


$$B = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 2 & 1 & 1 \\ 3 & 1 & 1 & 1 \\ 3 & 2 & 1 & 2 \end{bmatrix}$$



Partitioned matrices

• e.g.:

$$A = \left[\begin{array}{cc|cc} 1 & 1 & 1 & 1 \\ 2 & 2 & 1 & 1 \\ \hline 3 & 2 & 2 & 2 \end{array} \right] \quad B = \left[\begin{array}{cccc} 1 & 1 & 1 & 1 \\ 1 & 2 & 1 & 1 \\ 3 & 1 & 1 & 1 \\ 3 & 2 & 1 & 2 \end{array} \right]$$



Partitioned matrices

• e.g.:

$$A = \begin{bmatrix} 1 & 1 & & \\ 2 & 2 & & \\ \hline 3 & 2 & 2 & 2 \end{bmatrix} \quad A_{11} \quad B = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 2 & 1 & 1 \\ 3 & 1 & 1 & 1 \\ 3 & 2 & 1 & 2 \end{bmatrix}$$

Partitioned matrices

• e.g.:

$$A = \begin{bmatrix} 1 & 1 & A_{11} & A_{12} \\ 2 & 2 & 1 & 1 \\ 3 & 2 & 2 & 2 \end{bmatrix}$$
$$B = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 2 & 1 & 1 \\ 3 & 1 & 1 & 1 \\ 3 & 2 & 1 & 2 \end{bmatrix}$$

Partitioned matrices

• e.g.:

$$A = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix} \quad B = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 2 & 1 & 1 \\ 3 & 1 & 1 & 1 \\ 3 & 2 & 1 & 2 \end{bmatrix}$$

The matrix A is partitioned into four submatrices: A_{11} (top-left 2x2 blue box), A_{12} (top-right 2x2 green box), A_{21} (bottom-left 2x2 yellow box), and A_{22} (bottom-right 2x2 black box). A red cross is drawn through the matrix B .

Partitioned matrices

• e.g.:

$$A = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix} \quad B = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 2 & 1 & 1 \\ 3 & 1 & 1 & 1 \\ 3 & 2 & 1 & 2 \end{bmatrix}$$

The matrix A is partitioned into four submatrices: A_{11} (top-left, blue), A_{12} (top-right, green), A_{21} (bottom-left, yellow), and A_{22} (bottom-right, purple). A red cross is drawn through the matrix B .

Partitioned matrices

• e.g.:

$$A = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix}$$

$A_{11} = \begin{bmatrix} 1 & 1 \\ 2 & 2 \end{bmatrix}$ $A_{12} = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$

$A_{21} = \begin{bmatrix} 3 & 2 \end{bmatrix}$ $A_{22} = \begin{bmatrix} 2 & 2 \end{bmatrix}$

$m_1 = 2$

$m_2 = 1$

$$B = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 2 & 1 & 1 \\ 3 & 1 & 1 & 1 \\ 3 & 2 & 1 & 2 \end{bmatrix}$$

Partitioned matrices

• e.g.: $k_1=2 \quad k_2=2$

$m_1=2$

$m_2=1$

$$A = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix}$$
$$B = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 2 & 1 & 1 \\ 3 & 1 & 1 & 1 \\ 3 & 2 & 1 & 2 \end{bmatrix}$$

The diagram illustrates the partitioning of matrix A into four submatrices: A_{11} , A_{12} , A_{21} , and A_{22} . The submatrix A_{11} is highlighted in blue, A_{12} in green, A_{21} in yellow, and A_{22} in purple. Matrix B is shown as a 4x4 matrix with the same row and column indices as the submatrices of A .

Partitioned matrices

• e.g.: $k_1=2 \quad k_2=2$

$m_1=2$

$m_2=1$

$$A = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix}$$
$$B = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 2 & 1 & 1 \\ 3 & 1 & 1 & 1 \\ 3 & 2 & 1 & 2 \end{bmatrix}$$

Partitioned matrices

• e.g.: $k_1=2 \quad k_2=2$

$$A = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix}$$

$m_1=2$

$m_2=1$

$$B = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 2 & 1 & 1 \\ 3 & 1 & 1 & 1 \\ 3 & 2 & 1 & 2 \end{bmatrix}$$

Partitioned matrices

• e.g.: $k_1 = 2 \quad k_2 = 2$

$m_1 = 2$

$m_2 = 1$

$$A = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix}$$

$n_1 = 3 \quad n_2 = 1$.

$$B = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 2 & 1 \\ 3 & 1 & 1 \\ 3 & 2 & 1 \end{bmatrix}$$

Partitioned matrices

• e.g.: $k_1 = 2 \quad k_2 = 2$

$$A = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix}$$

$m_1 = 2$

$m_2 = 1$

$B = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 2 & 1 & 1 \\ 3 & 1 & 1 & 1 \\ 3 & 2 & 1 & 2 \end{bmatrix}$

$n_1 = 3 \quad n_2 = 1$.

or

$$A = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 2 & 1 & 1 \\ 3 & 2 & 2 & 2 \end{bmatrix}$$

$$B = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 2 & 1 & 1 \\ 3 & 1 & 1 & 1 \\ 3 & 2 & 1 & 2 \end{bmatrix}$$

Partitioned matrices

- e.g.: $k_1 = 2 \quad k_2 = 2$

$$A = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix}$$

$m_1 = 2$

$m_2 = 1$

Handwritten annotations:

- A_{11} is highlighted with a blue box.
- A_{12} is highlighted with a green box.
- A_{21} is highlighted with a yellow box.
- A_{22} is highlighted with a purple box.

$$B = \begin{bmatrix} n_1 = 3 & n_2 = 1 \end{bmatrix}$$

k_1

k_2

Handwritten annotations:

- The matrix is partitioned into two vertical blocks: one of width 3 and one of width 1.
- The first block has three rows labeled 1, 2, 3.
- The second block has one row labeled 1.

or

$$A = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 2 & 1 & 1 \\ 3 & 2 & 2 & 2 \end{bmatrix}$$

$$B = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 2 & 1 & 1 \\ 3 & 1 & 1 & 1 \\ 3 & 2 & 1 & 2 \end{bmatrix}$$

Partitioned matrices

- e.g.: $k_1=2 \quad k_2=2$

$A = \begin{bmatrix} 1 & 1 \\ 2 & 2 \\ 3 & 2 \\ 2 & 2 \end{bmatrix}$

$m_1=2$

$m_2=1$

$A = \begin{array}{c|cc|c} A_{11} & A_{12} & & \\ \hline & & & \\ A_{21} & A_{22} & & \end{array}$

$B = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 2 & 1 & 1 \\ 3 & 1 & 1 & 1 \\ 3 & 2 & 1 & 2 \end{bmatrix}$

$n_1=3 \quad n_2=1.$

or

$k_1=3 \quad k_2=1$

$A = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 2 & 1 & 1 \\ 3 & 2 & 2 & 2 \end{bmatrix}$

$B = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 2 & 1 & 1 \\ 3 & 1 & 1 & 1 \\ 3 & 2 & 1 & 2 \end{bmatrix}$

Partitioned matrices

• e.g.: $k_1 = 2 \quad k_2 = 2$

$$A = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix}$$

$m_1 = 2$

$m_2 = 1$

$$B = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 2 & 1 & 1 \\ 3 & 1 & 1 & 1 \\ 3 & 2 & 1 & 2 \end{bmatrix}$$

$n_1 = 3 \quad n_2 = 1$.

or

$k_1 = 3 \quad k_2 = 1$

$$A = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 2 & 1 & 1 \\ 3 & 2 & 2 & 2 \end{bmatrix}$$

$$B = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 2 & 1 & 1 \\ 3 & 1 & 1 & 1 \\ 3 & 2 & 1 & 2 \end{bmatrix}$$

Partitioned matrices

• e.g.: $k_1=2 \quad k_2=2$

$m_1=2$

$$A = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix}$$

$m_2=1$

$n_1=3 \quad n_2=1$.

$$B = \begin{bmatrix} 1 & 1 & 1 & | & 1 \\ 1 & 2 & 1 & | & 1 \\ 3 & 1 & 1 & | & 1 \\ 3 & 2 & 1 & | & 2 \end{bmatrix}$$

or

$k_1=3 \quad k_2=1$

$$A = \begin{bmatrix} 1 & 1 & 1 & | & 1 \\ 2 & 2 & 1 & | & 1 \\ 3 & 2 & 2 & | & 2 \end{bmatrix}$$

$n_1=2 \quad n_2=2$

k_1

$$B = \begin{bmatrix} 1 & 1 & | & 1 & 1 \\ 1 & 2 & | & 1 & 1 \\ 3 & 1 & | & 1 & 1 \\ 3 & 2 & | & 1 & 2 \end{bmatrix}$$

k_2

Partitioned matrices

Partitioned matrices

e.g.

$$\begin{bmatrix} A_{11} & 0 \\ 0 & A_{22} \end{bmatrix} \begin{bmatrix} B_{11} & 0 \\ 0 & B_{22} \end{bmatrix} = \begin{bmatrix} A_{11}B_{11} & 0 \\ 0 & A_{22}B_{22} \end{bmatrix}$$

Partitioned matrices

e.g.

$$\begin{matrix} k_1 & \begin{bmatrix} A_{11} & 0 \\ 0 & A_{22} \end{bmatrix} & \begin{bmatrix} B_{11} & 0 \\ 0 & B_{22} \end{bmatrix} & = & \begin{bmatrix} A_{11}B_{11} & 0 \\ 0 & A_{22}B_{22} \end{bmatrix} \\ k_2 & & & & \end{matrix}$$

Partitioned matrices

~~e.g.~~

$$\begin{bmatrix} k_1 & \\ & k_2 \end{bmatrix} \begin{bmatrix} A_{11} & 0 \\ 0 & A_{22} \end{bmatrix} \begin{bmatrix} B_{11} & 0 \\ 0 & B_{22} \end{bmatrix} = \begin{bmatrix} A_{11}B_{11} & 0 \\ 0 & A_{22}B_{22} \end{bmatrix}$$

Partitioned matrices

e.g.

$$\begin{bmatrix} k_1 & A_{11} \\ k_2 & 0 \end{bmatrix} \begin{bmatrix} 0 & B_{11} \\ A_{22} & 0 \end{bmatrix} = \begin{bmatrix} A_{11}B_{11} & 0 \\ 0 & A_{22}B_{22} \end{bmatrix}$$

Partitioned matrices

e.g.

$$\begin{bmatrix} k_1 & A_{11} \\ k_2 & 0 \end{bmatrix} \begin{bmatrix} k_2 & 0 \\ 0 & A_{22} \end{bmatrix} = \begin{bmatrix} k_1 & B_{11} \\ k_2 & 0 \end{bmatrix} \begin{bmatrix} 0 & B_{22} \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} A_{11}B_{11} & 0 \\ 0 & A_{22}B_{22} \end{bmatrix}$$

Partitioned matrices

e.g.

$$\begin{bmatrix} k_1 & A_{11} \\ k_2 & 0 \end{bmatrix} \begin{bmatrix} k_1 & B_{11} \\ k_2 & 0 \end{bmatrix} = \begin{bmatrix} A_{11}B_{11} & 0 \\ 0 & A_{22}B_{22} \end{bmatrix}$$
$$\begin{bmatrix} 0 & A_{22} \\ 0 & 0 \end{bmatrix}$$
$$\begin{bmatrix} 0 & B_{21} \\ 0 & 0 \end{bmatrix}$$

- with A_{ii}, B_{ii} are square matrices.

Inverse matrices

Inverse matrices

- Recall that when $E = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ l & 0 & 1 \end{pmatrix}$, $E^{-1} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -l & 0 & 1 \end{pmatrix}$ satisfying $EE^{-1} = E^{-1}E = I$.

Inverse matrices

- Recall that when $E = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ l & 0 & 1 \end{pmatrix}$, $E^{-1} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -l & 0 & 1 \end{pmatrix}$ satisfying $EE^{-1} = E^{-1}E = I$.
- For $L = E_k \dots E_1$, $L^{-1} = E_1^{-1} \dots E_k^{-1}$.

Inverse matrices

- Recall that when $E = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ l & 0 & 1 \end{pmatrix}$, $E^{-1} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -l & 0 & 1 \end{pmatrix}$ satisfying $EE^{-1} = E^{-1}E = I$.
- For $L = E_k \dots E_1$, $L^{-1} = E_1^{-1} \dots E_k^{-1}$.
- Def: For a $n \times n$ matrix A , the inverse (逆) is a $n \times n$ matrix B s.t. $AB = BA = I_{n \times n}$.

Inverse matrices

- Recall that when $E = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ l & 0 & 1 \end{pmatrix}$, $E^{-1} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -l & 0 & 1 \end{pmatrix}$ satisfying $EE^{-1} = E^{-1}E = I$.
 - For $L = E_k \dots E_1$, $L^{-1} = E_1^{-1} \dots E_k^{-1}$.
- Def: • For a $n \times n$ matrix A , the inverse is a $n \times n$ matrix B s.t. $AB = BA = I_{n \times n}$.
- If inverse of A exists, we say it is invertible

Inverse matrices

- Recall that when $E = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ l & 0 & 1 \end{pmatrix}$, $E^{-1} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -l & 0 & 1 \end{pmatrix}$ satisfying $EE^{-1} = E^{-1}E = I$.
 - For $L = E_k \dots E_1$, $L^{-1} = E_1^{-1} \dots E_k^{-1}$.
- Def: • For a $n \times n$ matrix A , the **inverse** (逆) is a $n \times n$ matrix B s.t. $AB = BA = I_{n \times n}$.
- If inverse of A exists, we say it is **invertible** (可逆).
- Rk: Inverse, when exists, is unique.

Inverse matrices

- Recall that when $E = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ l & 0 & 1 \end{pmatrix}$, $E^{-1} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -l & 0 & 1 \end{pmatrix}$ satisfying $EE^{-1} = E^{-1}E = I$.
- For $L = E_k \dots E_1$, $L^{-1} = E_1^{-1} \dots E_k^{-1}$.

Def: • For a $n \times n$ matrix A , the inverse is a $n \times n$ matrix B s.t. $AB = BA = I_{n \times n}$.

• If inverse of A exists, we say it is invertible.

Rk: Inverse, when exists, is unique.

Pf: if B, \tilde{B} are inverse, then

Inverse matrices

- Recall that when $E = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ l & 0 & 1 \end{pmatrix}$, $E^{-1} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -l & 0 & 1 \end{pmatrix}$ satisfying $EE^{-1} = E^{-1}E = I$.
- For $L = E_k \dots E_1$, $L^{-1} = E_1^{-1} \dots E_k^{-1}$.

Def: • For a $n \times n$ matrix A , the inverse is a $n \times n$ matrix B s.t. $AB = BA = I_{n \times n}$.

• If inverse of A exists, we say it is invertible.

Rk: Inverse, when exists, is unique.

Pf: if B, \tilde{B} are inverse, then

$$B = BI = B(A\tilde{B})$$

Inverse matrices

- Recall that when $E = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ l & 0 & 1 \end{pmatrix}$, $E^{-1} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -l & 0 & 1 \end{pmatrix}$ satisfying $EE^{-1} = E^{-1}E = I$.
- For $L = E_k \dots E_1$, $L^{-1} = E_1^{-1} \dots E_k^{-1}$.

Def: • For a $n \times n$ matrix A , the inverse (逆) is a $n \times n$ matrix B s.t. $AB = BA = I_{n \times n}$.

• If inverse of A exists, we say it is invertible (可逆).

Rk: Inverse, when exists, is unique.

Pf: if B, \tilde{B} are inverse, then

$$B = BI = B(A\tilde{B}) = (BA)\tilde{B} = \tilde{B}.$$

Inverse matrices

Inverse matrices

- We write A^{-1} for the unique inverse of A . (if exists)

Inverse matrices

- We write A^{-1} for the unique inverse of A (if exists)

e.g.: $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$, then $A^{-1} = \frac{1}{ad-bc} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}$

exists when $ad-bc \neq 0$.

Inverse matrices

- We write A^{-1} for the unique inverse of A . (if exists)

e.g.: $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$, then $A^{-1} = \frac{1}{ad-bc} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}$

exists when

$$ad-bc \neq 0$$

$\nwarrow \det(A)$.

Inverse matrices

- We write A^{-1} for the unique inverse of A . (if exists)

e.g.: $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$, then $A^{-1} = \frac{1}{ad-bc} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}$

exists when $ad-bc \neq 0$.
 $\nwarrow \det(A)$.

e.g.:

$$A = \begin{bmatrix} d_1 & 0 & 0 \\ 0 & d_2 & 0 \\ 0 & 0 & d_3 \end{bmatrix}, \quad A^{-1} = \begin{bmatrix} d_1^{-1} & 0 & 0 \\ 0 & d_2^{-1} & 0 \\ 0 & 0 & d_3^{-1} \end{bmatrix}$$

exists when $d_1 d_2 d_3 \neq 0$.

Inverse matrices

- We write A^{-1} for the unique inverse of A . (if exists)

e.g.: $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$, then $A^{-1} = \frac{1}{ad-bc} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}$

exists when $ad-bc \neq 0$.
 $\nwarrow \det(A)$.

e.g.: $A = \begin{bmatrix} d_1 & 0 & 0 \\ 0 & d_2 & 0 \\ 0 & 0 & d_3 \end{bmatrix}$, $A^{-1} = \begin{bmatrix} d_1^{-1} & 0 & 0 \\ 0 & d_2^{-1} & 0 \\ 0 & 0 & d_3^{-1} \end{bmatrix}$

exists when $d_1 d_2 d_3 \neq 0$.

e.g.: $P = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$, $P^2 = I \Rightarrow P^{-1} = P$.

Inverse matrices

Inverse matrices

Prop:

if A, B are invertible (\exists^{-1}), then so is AB .
Furthermore $(AB)^{-1} = B^{-1}A^{-1}$.

Inverse matrices

Prop:

if A, B are invertible (\exists^{-1}), then so is AB .
Furthermore $(AB)^{-1} = B^{-1}A^{-1}$.

Pf:

$$(AB) \cdot (B^{-1}A^{-1}) =$$

Inverse matrices

Prop:

if A, B are invertible (\exists^{-1}), then so is AB .
Furthermore $(AB)^{-1} = B^{-1}A^{-1}$.

Pf:

$$(AB) \cdot (B^{-1}A^{-1}) = A(BB^{-1})A^{-1}$$

Inverse matrices

Prop:

if A, B are invertible (\exists^{-1}), then so is AB .
Furthermore $(AB)^{-1} = B^{-1}A^{-1}$.

Pf:

$$(AB) \cdot (B^{-1}A^{-1}) = A(BB^{-1})A^{-1} = A(I)A^{-1}$$

Inverse matrices

Prop:

if A, B are invertible (\exists^{-1}), then so is AB .
Furthermore $(AB)^{-1} = B^{-1}A^{-1}$.

Pf:

$$\begin{aligned}(AB) \cdot (B^{-1}A^{-1}) &= A(BB^{-1})A^{-1} = A(I)A^{-1} \\ &= AA^{-1} = I.\end{aligned}$$

Inverse matrices

Prop:

if A, B are invertible (\exists^{-1}), then so is AB .
Furthermore $(AB)^{-1} = B^{-1}A^{-1}$.

Pf:
$$(AB) \cdot (B^{-1}A^{-1}) = A(BB^{-1})A^{-1} = A(I)A^{-1}$$
$$= AA^{-1} = I.$$

- All elementary matrices are invertible.
 \Rightarrow their products are invertible.

Inverse matrices

Prop:

if A, B are invertible ($\exists A^{-1}, B^{-1}$), then so is AB .
Furthermore $(AB)^{-1} = B^{-1}A^{-1}$.

Pf:

$$\begin{aligned}(AB) \cdot (B^{-1}A^{-1}) &= A(BB^{-1})A^{-1} = A(I)A^{-1} \\ &= AA^{-1} = I.\end{aligned}$$

- All elementary matrices are invertible.
 \Rightarrow their products are invertible.

Prop:

A $n \times n$ matrix A is **invertible** if and only if
the elimination process gives n pivots.
(i.e.) **non-singular**.

Inverse matrices

Inverse matrices

Pf: • Recall that we can always have $PA = LU$

Inverse matrices

Pf: • Recall that we can always have $PA = LU$

permutation
matrices' product

Inverse matrices

Pf:

- Recall that we can always have $PA = LU$
permutation
matrices' product elementary
matrices product.

Inverse matrices

Pf:

- Recall that we can always have $PA = LU$
Permutation
matrices' product elementary
matrices product.
- P, L are invertible.

Inverse matrices

Pf:

- Recall that we can always have $PA = LU$
Permutation
matrices' product elementary
matrices product.
- P, L are invertible.
So A is invertible $\iff U$ is invertible.

Inverse matrices

Pf:

- Recall that we can always have $PA = LU$
Permutation
matrices' product elementary
matrices product.
- P, L are invertible.
So A is invertible $\iff U$ is invertible.
- U is upper triangular, and it is invertible
 \iff its diagonal entries are non-zero.

Inverse matrices

Pf:

- Recall that we can always have $PA = LU$
Permutation
matrices' product elementary
matrices product.
- P, L are invertible.
So A is invertible $\iff U$ is invertible.
- U is upper triangular, and it is invertible
 \iff its diagonal entries are non-zero.

Reason \Rightarrow :

Assume U is invertible

but with diagonal entry zero:

Inverse matrices

Pf:

- Recall that we can always have $PA = LU$
Permutation
matrices' product elementary
matrices product.
- P, L are invertible.
So A is invertible $\iff U$ is invertible.
- U is upper triangular, and it is invertible
 \iff its diagonal entries are non-zero.

Reason \Rightarrow :

Assume U is invertible

but with diagonal entry zero:

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix} \underbrace{\begin{pmatrix} * & * & * \\ 0 & * & * \\ 0 & 0 & 0 \end{pmatrix}}_{\tilde{U}} = U$$

E

Inverse matrices

Pf:

- Recall that we can always have $PA = LU$
Permutation
matrices' product elementary
matrices product.
- P, L are invertible.
So A is invertible $\Leftrightarrow U$ is invertible.
- U is upper triangular, and it is invertible
 \Leftrightarrow its diagonal entries are non-zero.

Reason \Rightarrow :

Assume U is invertible

$$\text{but with diagonal entry zero : } \underbrace{\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}}_E \underbrace{\begin{pmatrix} * & * & * \\ 0 & * & * \\ 0 & 0 & 0 \end{pmatrix}}_U = U$$

$$EU = U \Rightarrow E = I \quad \text{contradiction}$$

Inverse matrices

Inverse matrices

Reason \Leftarrow :) if $U = \begin{pmatrix} d_1 & * & * \\ 0 & d_2 & * \\ 0 & 0 & d_3 \end{pmatrix}$, with $d_1 d_2 d_3 \neq 0$.

Inverse matrices

Reason $\leftarrow:$) if $U = \begin{pmatrix} d_1 & * & * \\ 0 & d_2 & * \\ 0 & 0 & d_3 \end{pmatrix}$, with $d_1 d_2 d_3 \neq 0$.

$$U = \begin{pmatrix} d_1 & 0 & 0 \\ 0 & d_2 & 0 \\ 0 & 0 & d_3 \end{pmatrix} \begin{pmatrix} 1 & U_{12} & U_{13} \\ 0 & 1 & U_{23} \\ 0 & 0 & 1 \end{pmatrix}$$

Inverse matrices

Reason \Leftarrow :) if $U = \begin{pmatrix} d_1 & * & * \\ 0 & d_2 & * \\ 0 & 0 & d_3 \end{pmatrix}$, with $d_1 d_2 d_3 \neq 0$.

$$U = \begin{pmatrix} d_1 & 0 & 0 \\ 0 & d_2 & 0 \\ 0 & 0 & d_3 \end{pmatrix} \begin{pmatrix} 1 & U_{12} & U_{13} \\ 0 & 1 & U_{23} \\ 0 & 0 & 1 \end{pmatrix}$$

We can easily write

$$\begin{pmatrix} 1 & U_{12} & U_{13} \\ 0 & 1 & U_{23} \\ 0 & 0 & 1 \end{pmatrix}^{-1} = \begin{pmatrix} 1 & -U_{12} & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & -U_{13} \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & -U_{23} \\ 0 & 0 & 1 \end{pmatrix}$$

Inverse matrices

Reason \Leftarrow :) if $U = \begin{pmatrix} d_1 & * & * \\ 0 & d_2 & * \\ 0 & 0 & d_3 \end{pmatrix}$, with $d_1, d_2, d_3 \neq 0$.

$$U = \begin{pmatrix} d_1 & 0 & 0 \\ 0 & d_2 & 0 \\ 0 & 0 & d_3 \end{pmatrix} \begin{pmatrix} 1 & U_{12} & U_{13} \\ 0 & 1 & U_{23} \\ 0 & 0 & 1 \end{pmatrix}$$

We can easily write

$$\begin{pmatrix} 1 & U_{12} & U_{13} \\ 0 & 1 & U_{23} \\ 0 & 0 & 1 \end{pmatrix}^{-1} = \begin{pmatrix} 1 & -U_{12} & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & -U_{13} \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & -U_{23} \\ 0 & 0 & 1 \end{pmatrix}$$

$$= \begin{pmatrix} 1 & -U_{12} & -U_{13} + U_{12}U_{23} \\ 0 & 1 & -U_{23} \\ 0 & 0 & 1 \end{pmatrix}$$

Inverse matrices

Inverse matrices

- Relation to equation: If A is $n \times n$ matrix, consider

$$Ax = b.$$

Inverse matrices

- Relation to equation: If A is $n \times n$ matrix, consider

$$Ax = b.$$

- if A is **invertible**, then we multiple A^{-1} to both sides

$$A^{-1}Ax = A^{-1}b \Rightarrow x = A^{-1}b.$$

Inverse matrices

- Relation to equation: If A is $n \times n$ matrix, consider

$$Ax = b.$$

- if A is **invertible**, then we multiple A^{-1} to both sides
$$A^{-1}Ax = A^{-1}b \Rightarrow x = A^{-1}b.$$
- In this case, the solution is unique.

Inverse matrices

- Relation to equation: If A is $n \times n$ matrix, consider

$$Ax = b.$$

- if A is **invertible**, then we multiple A^{-1} to both sides
- $$A^{-1}Ax = A^{-1}b \Rightarrow x = A^{-1}b.$$
- In this case, the solution is unique.

eg.

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} e_1 \\ e_2 \end{pmatrix}$$

Inverse matrices

- Relation to equation: If A is $n \times n$ matrix, consider

$$Ax = b.$$

- if A is **invertible**, then we multiple A^{-1} to both sides

$$A^{-1}Ax = A^{-1}b \Rightarrow x = A^{-1}b.$$

- In this case, the solution is unique.

eg. $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} e_1 \\ e_2 \end{pmatrix} \Rightarrow \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \frac{1}{ad-bc} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix} \begin{pmatrix} e_1 \\ e_2 \end{pmatrix}$

Inverse matrices

- Relation to equation: If A is $n \times n$ matrix, consider

$$Ax = b.$$

- if A is **invertible**, then we multiple A^{-1} to both sides


$$A^{-1}Ax = A^{-1}b \Rightarrow x = A^{-1}b.$$

- In this case, the solution is unique.

eg. $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} e_1 \\ e_2 \end{pmatrix} \Rightarrow \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \frac{1}{ad-bc} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix} \begin{pmatrix} e_1 \\ e_2 \end{pmatrix}$

can be written down directly if $ad-bc \neq 0$

The Gauss-Jordan Method



The Gauss-Jordan Method

Q: How to find the inverse A^{-1} if given A?

The Gauss-Jordan Method

Q: How to find the inverse A^{-1} if given A ?

• Recall:

$$A \begin{bmatrix} | & | & | \\ b_1, & b_2, & \dots, & b_n \\ | & | & | \end{bmatrix} = \begin{bmatrix} | \\ Ab_1, & \dots, & Ab_n \\ | \end{bmatrix}$$

i.e. multiplication can be done column by column.

The Gauss-Jordan Method

Q: How to find the inverse A^{-1} if given A ?

- Recall:

$$A \begin{bmatrix} | & | & | \\ b_1, & b_2, & \dots, & b_n \\ | & | & | \end{bmatrix} = \begin{bmatrix} | \\ Ab_1, & \dots, & Ab_n \\ | \end{bmatrix}$$

i.e. multiplication can be done column by column.

- In particular: $A [B, C] = [AB, AC]$

The Gauss-Jordan Method

Q: How to find the inverse A^{-1} if given A ?

- Recall:

$$A \begin{bmatrix} | & | & | \\ b_1, & b_2, & \dots, & b_n \\ | & | & | \end{bmatrix} = \begin{bmatrix} | \\ Ab_1, & \dots, & Ab_n \\ | \end{bmatrix}$$

i.e. multiplication can be done column by column.

- In particular:

$$A [B, C] = [AB, AC]$$

$\uparrow \uparrow$
matrices with same no. of rows

The Gauss-Jordan Method

Q: How to find the inverse A^{-1} if given A ?

- Recall:

$$A \begin{bmatrix} | & | & | \\ b_1, & b_2, & \dots, & b_n \\ | & | & | \end{bmatrix} = \begin{bmatrix} | \\ Ab_1, & \dots, & Ab_n \\ | \end{bmatrix}$$

i.e. multiplication can be done column by column.

- In particular:

$$A [B, C] = [AB, AC]$$

$\uparrow \uparrow$
matrices with same no. of rows

Say 3×3 case:

$$A \begin{bmatrix} | & | & | \\ x_1, & x_2, & x_3 \\ | & | & | \end{bmatrix} = \begin{bmatrix} | & 0 & 0 \\ 0 & | & 0 \\ 0 & 0 & | \end{bmatrix}$$

The Gauss-Jordan Method

Q: How to find the inverse A^{-1} if given A ?

- Recall:

$$A \begin{bmatrix} | & | & | \\ b_1, & b_2, & \dots, & b_n \\ | & | & | \end{bmatrix} = \begin{bmatrix} | \\ Ab_1, & \dots, & | \\ | & | & | \end{bmatrix}$$

i.e. multiplication can be done column by column.

- In particular:

$$A \begin{bmatrix} | & | \\ B, & C \\ | & | \end{bmatrix} = \begin{bmatrix} | \\ AB, & AC \\ | & | \end{bmatrix}$$

matrices with same no. of rows

Say 3×3 case:

// A^{-1} .

$$A \begin{bmatrix} | & | & | \\ x_1, & x_2, & x_3 \\ | & | & | \end{bmatrix} = \begin{bmatrix} | & 0 & 0 \\ 0 & | & 0 \\ 0 & 0 & | \end{bmatrix}$$

The Gauss-Jordan Method

The Gauss-Jordan Method

- $A \begin{bmatrix} | & | & | \\ x_1 & x_2 & x_3 \\ | & | & | \end{bmatrix} = \begin{bmatrix} | & 0 & 0 \\ 0 & | & 0 \\ 0 & 0 & | \end{bmatrix}$

The Gauss-Jordan Method

$$\bullet \quad A \begin{bmatrix} | & | & | \\ x_1 & x_2 & x_3 \\ | & | & | \end{bmatrix} = \begin{bmatrix} | & 0 & 0 \\ 0 & | & 0 \\ 0 & 0 & | \end{bmatrix}$$

$$\Leftrightarrow [A\vec{x}_1, A\vec{x}_2, A\vec{x}_3] = \begin{bmatrix} | & 0 & 0 \\ 0 & | & 0 \\ 0 & 0 & | \end{bmatrix}$$

The Gauss-Jordan Method

$$\bullet \quad A \begin{bmatrix} | & | & | \\ x_1 & x_2 & x_3 \\ | & | & | \end{bmatrix} = \begin{bmatrix} | & 0 & 0 \\ 0 & | & 0 \\ 0 & 0 & | \end{bmatrix}$$

$$\Leftrightarrow [A\vec{x}_1, A\vec{x}_2, A\vec{x}_3] = \begin{bmatrix} | & 0 & 0 \\ 0 & | & 0 \\ 0 & 0 & | \end{bmatrix}$$

$$\Leftrightarrow A\vec{x}_1 = \begin{bmatrix} | \\ 0 \\ 0 \end{bmatrix}, A\vec{x}_2 = \begin{bmatrix} 0 \\ | \\ 0 \end{bmatrix}, A\vec{x}_3 = \begin{bmatrix} 0 \\ 0 \\ | \end{bmatrix}$$

The Gauss-Jordan Method

$$\bullet \quad A \begin{bmatrix} | & | & | \\ x_1 & x_2 & x_3 \\ | & | & | \end{bmatrix} = \begin{bmatrix} | & 0 & 0 \\ 0 & | & 0 \\ 0 & 0 & | \end{bmatrix}$$

$$\Leftrightarrow [A\vec{x}_1, A\vec{x}_2, A\vec{x}_3] = \begin{bmatrix} | & 0 & 0 \\ 0 & | & 0 \\ 0 & 0 & | \end{bmatrix}$$

$$\Leftrightarrow A\vec{x}_1 = \begin{bmatrix} | \\ 0 \\ 0 \end{bmatrix} = \vec{e}_1, A\vec{x}_2 = \begin{bmatrix} 0 \\ | \\ 0 \end{bmatrix} = \vec{e}_2, A\vec{x}_3 = \begin{bmatrix} 0 \\ 0 \\ | \end{bmatrix} = \vec{e}_3$$

The Gauss-Jordan Method

- $A \begin{bmatrix} | & | & | \\ x_1 & x_2 & x_3 \\ | & | & | \end{bmatrix} = \begin{bmatrix} | & 0 & 0 \\ 0 & | & 0 \\ 0 & 0 & | \end{bmatrix}$

$$\Leftrightarrow [A\vec{x}_1, A\vec{x}_2, A\vec{x}_3] = \begin{bmatrix} | & 0 & 0 \\ 0 & | & 0 \\ 0 & 0 & | \end{bmatrix}$$

$$\Leftrightarrow A\vec{x}_1 = \begin{bmatrix} | \\ 0 \\ 0 \end{bmatrix} = \vec{e}_1, A\vec{x}_2 = \begin{bmatrix} 0 \\ | \\ 0 \end{bmatrix} = \vec{e}_2, A\vec{x}_3 = \begin{bmatrix} 0 \\ 0 \\ | \end{bmatrix} = \vec{e}_3$$

- We need to do Gaussian elimination for $[A | \vec{e}_i]$:

The Gauss-Jordan Method

$$\bullet A \begin{bmatrix} | & | & | \\ x_1 & x_2 & x_3 \\ | & | & | \end{bmatrix} = \begin{bmatrix} | & 0 & 0 \\ 0 & | & 0 \\ 0 & 0 & | \end{bmatrix}$$

$$\Leftrightarrow [A\vec{x}_1, A\vec{x}_2, A\vec{x}_3] = \begin{bmatrix} | & 0 & 0 \\ 0 & | & 0 \\ 0 & 0 & | \end{bmatrix}$$

$$\Leftrightarrow A\vec{x}_1 = \begin{bmatrix} | \\ 0 \\ 0 \end{bmatrix} = \vec{e}_1, A\vec{x}_2 = \begin{bmatrix} 0 \\ | \\ 0 \end{bmatrix} = \vec{e}_2, A\vec{x}_3 = \begin{bmatrix} 0 \\ 0 \\ | \end{bmatrix} = \vec{e}_3$$

- We need to do Gaussian elimination for $[A | \vec{e}_i]$:
- Two cases: • if A is singular , elimination fails
no inverse

The Gauss-Jordan Method

$$\bullet A \begin{bmatrix} | & | & | \\ x_1 & x_2 & x_3 \\ | & | & | \end{bmatrix} = \begin{bmatrix} | & 0 & 0 \\ 0 & | & 0 \\ 0 & 0 & | \end{bmatrix}$$

$$\Leftrightarrow [A\vec{x}_1, A\vec{x}_2, A\vec{x}_3] = \begin{bmatrix} | & 0 & 0 \\ 0 & | & 0 \\ 0 & 0 & | \end{bmatrix}$$

$$\Leftrightarrow A\vec{x}_1 = \begin{bmatrix} | \\ 0 \\ 0 \end{bmatrix} = \vec{e}_1, A\vec{x}_2 = \begin{bmatrix} 0 \\ | \\ 0 \end{bmatrix} = \vec{e}_2, A\vec{x}_3 = \begin{bmatrix} 0 \\ 0 \\ | \end{bmatrix} = \vec{e}_3$$

- We need to do Gaussian elimination for $[A | \vec{e}_i]$:
- Two cases:
 - if A is singular, elimination fails
no inverse
 - if A non-singular, find unique \vec{x}_i 's

The Gauss-Jordan Method

The Gauss-Jordan Method

- We do the elimination simultaneously

The Gauss-Jordan Method

- We do the elimination simultaneously

e.g. $[A | \dots] = \left[\begin{array}{ccc|c} 2 & 1 & 1 & \\ 4 & -6 & 0 & \\ -2 & 7 & 2 & \end{array} \right]$

The Gauss-Jordan Method

- We do the elimination simultaneously

e.g. $[A | \vec{e}_1, \dots] = \left[\begin{array}{ccc|c} 2 & 1 & 1 & 1 \\ 4 & -6 & 0 & 0 \\ -2 & 7 & 2 & 0 \end{array} \right]$

The Gauss-Jordan Method

- We do the elimination simultaneously

e.g. $[A | \vec{e}_1, \vec{e}_2] = \left[\begin{array}{ccc|cc} 2 & 1 & 1 & 1 & 0 \\ 4 & -6 & 0 & 0 & 1 \\ -2 & 7 & 2 & 0 & 0 \end{array} \right]$

The Gauss-Jordan Method

- We do the elimination simultaneously

e.g. $[A \mid \vec{e}_1, \vec{e}_2, \vec{e}_3] = \left[\begin{array}{ccc|ccc} 2 & 1 & 1 & 1 & 0 & 0 \\ 4 & -6 & 0 & 0 & 1 & 0 \\ -2 & 7 & 2 & 0 & 0 & 1 \end{array} \right]$

The Gauss-Jordan Method

- We do the elimination simultaneously

e.g. $[A | \vec{e}_1, \vec{e}_2, \vec{e}_3] = \left[\begin{array}{ccc|ccc} 2 & 1 & 1 & 1 & 0 & 0 \\ 4 & -6 & 0 & 0 & 1 & 0 \\ -2 & 7 & 2 & 0 & 0 & 1 \end{array} \right]$


$$\left[\begin{array}{ccc} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 1 & 0 & 1 \end{array} \right] \xrightarrow{\quad} \left[\begin{array}{ccc|ccc} 2 & 1 & 1 & 1 & 0 & 0 \\ 0 & -8 & -2 & -2 & 1 & 0 \\ 0 & 8 & 3 & 1 & 0 & 1 \end{array} \right]$$

The Gauss-Jordan Method

- We do the elimination simultaneously

e.g. $[A | \vec{e}_1, \vec{e}_2, \vec{e}_3] = \left[\begin{array}{ccc|ccc} 2 & 1 & 1 & 1 & 0 & 0 \\ 4 & -6 & 0 & 0 & 1 & 0 \\ -2 & 7 & 2 & 0 & 0 & 1 \end{array} \right]$

$$\left[\begin{array}{ccc} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 1 & 0 & 1 \end{array} \right]$$



$$\left[\begin{array}{ccc|ccc} 2 & 1 & 1 & 1 & 0 & 0 \\ 0 & -8 & -2 & -2 & 1 & 0 \\ 0 & 8 & 3 & 1 & 0 & 1 \end{array} \right]$$

$$\left[\begin{array}{ccc} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{array} \right]$$



$$\left[\begin{array}{ccc|ccc} 2 & 1 & 1 & 1 & 0 & 0 \\ 0 & -8 & -2 & -2 & 1 & 0 \\ 0 & 0 & 1 & -1 & 1 & 1 \end{array} \right]$$

The Gauss-Jordan Method

$$\left[\begin{array}{ccc|ccc} 2 & 1 & 1 & 1 & 0 & 0 \\ 0 & -8 & -2 & -2 & 1 & 0 \\ 0 & 0 & 1 & -1 & 1 & 1 \end{array} \right]$$

The Gauss-Jordan Method

$$\left[\begin{array}{ccc|ccc} 2 & 1 & 1 & 1 & 0 & 0 \\ 0 & -8 & -2 & -2 & 1 & 0 \\ 0 & 0 & 1 & -1 & 1 & 1 \end{array} \right] \xrightarrow{\left[\begin{array}{ccc} 1 & 0 & -1 \\ 0 & 1 & 2 \\ 0 & 0 & 1 \end{array} \right]} \left[\begin{array}{ccc|ccc} 2 & 1 & 0 & 2 & -1 & -1 \\ 0 & -8 & 0 & -4 & 3 & 2 \\ 0 & 0 & 1 & -1 & 1 & 1 \end{array} \right]$$

The Gauss-Jordan Method

$$\left[\begin{array}{ccc|ccc} 2 & 1 & 1 & 1 & 0 & 0 \\ 0 & -8 & -2 & -2 & 1 & 0 \\ 0 & 0 & 1 & -1 & 1 & 1 \end{array} \right] \xrightarrow{\left[\begin{array}{ccc} 1 & 0 & -1 \\ 0 & 1 & 2 \\ 0 & 0 & 1 \end{array} \right]} \left[\begin{array}{ccc|ccc} 2 & 1 & 0 & 2 & -1 & -1 \\ 0 & -8 & 0 & -4 & 3 & 2 \\ 0 & 0 & 1 & -1 & 1 & 1 \end{array} \right]$$

$$\left[\begin{array}{ccc} 1 & \frac{-8}{8} & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{array} \right] \xrightarrow{} \left[\begin{array}{ccc|ccc} 2 & 0 & 0 & \frac{12}{8} & \frac{-5}{8} & \frac{-6}{8} \\ 0 & 8 & 0 & -4 & 3 & 2 \\ 0 & 0 & 1 & -1 & 1 & 1 \end{array} \right]$$

The Gauss-Jordan Method

$$\left[\begin{array}{ccc|ccc} 2 & 1 & 1 & 1 & 0 & 0 \\ 0 & -8 & -2 & -2 & 1 & 0 \\ 0 & 0 & 1 & -1 & 1 & 1 \end{array} \right] \xrightarrow{\left[\begin{array}{ccc} 1 & 0 & -1 \\ 0 & 1 & 2 \\ 0 & 0 & 1 \end{array} \right]} \left[\begin{array}{ccc|ccc} 2 & 1 & 0 & 2 & -1 & -1 \\ 0 & -8 & 0 & -4 & 3 & 2 \\ 0 & 0 & 1 & -1 & 1 & 1 \end{array} \right]$$

$$\left[\begin{array}{ccc} 1 & \frac{1}{8} & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{array} \right] \rightarrow \left[\begin{array}{ccc|ccc} 2 & 0 & 0 & \frac{12}{8} & \frac{-5}{8} & \frac{-6}{8} \\ 0 & 8 & 0 & -4 & 3 & 2 \\ 0 & 0 & 1 & -1 & 1 & 1 \end{array} \right]$$

$$\left[\begin{array}{ccc} 1 & 0 & 0 \\ 0 & \frac{1}{8} & 0 \\ 0 & 0 & 1 \end{array} \right] \rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & \frac{12}{16} & \frac{-5}{16} & \frac{-6}{16} \\ 0 & 1 & 0 & -\frac{1}{2} & \frac{3}{8} & \frac{1}{4} \\ 0 & 0 & 1 & -1 & 1 & 1 \end{array} \right]$$

The Gauss-Jordan Method

$$\left[\begin{array}{ccc|ccc} 2 & 1 & 1 & 1 & 0 & 0 \\ 0 & -8 & -2 & -2 & 1 & 0 \\ 0 & 0 & 1 & -1 & 1 & 1 \end{array} \right] \xrightarrow{\left[\begin{array}{ccc} 1 & 0 & -1 \\ 0 & 1 & 2 \\ 0 & 0 & 1 \end{array} \right]} \left[\begin{array}{ccc|ccc} 2 & 1 & 0 & 2 & -1 & -1 \\ 0 & -8 & 0 & -4 & 3 & 2 \\ 0 & 0 & 1 & -1 & 1 & 1 \end{array} \right]$$

$$\left[\begin{array}{ccc} 1 & \frac{1}{8} & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{array} \right] \rightarrow \left[\begin{array}{ccc|ccc} 2 & 0 & 0 & \frac{12}{8} & \frac{-5}{8} & \frac{-6}{8} \\ 0 & 8 & 0 & -4 & 3 & 2 \\ 0 & 0 & 1 & -1 & 1 & 1 \end{array} \right]$$

$$\left[\begin{array}{ccc} 0 & 0 & 0 \\ 0 & \frac{1}{8} & 0 \\ 0 & 0 & -1 \end{array} \right] \rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & \frac{12}{16} & \frac{-5}{16} & \frac{-6}{16} \\ 0 & 1 & 0 & -\frac{1}{2} & \frac{3}{8} & \frac{1}{4} \\ 0 & 0 & 1 & -1 & 1 & 1 \end{array} \right]$$

= x_1

The Gauss-Jordan Method

$$\left[\begin{array}{ccc|ccc} 2 & 1 & 1 & 1 & 0 & 0 \\ 0 & -8 & -2 & -2 & 1 & 0 \\ 0 & 0 & 1 & -1 & 1 & 1 \end{array} \right] \xrightarrow{\left[\begin{array}{ccc} 1 & 0 & -1 \\ 0 & 1 & 2 \\ 0 & 0 & 1 \end{array} \right]} \left[\begin{array}{ccc|ccc} 2 & 1 & 0 & 2 & -1 & -1 \\ 0 & -8 & 0 & -4 & 3 & 2 \\ 0 & 0 & 1 & -1 & 1 & 1 \end{array} \right]$$

$$\left[\begin{array}{ccc} 1 & \frac{-8}{8} & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{array} \right] \rightarrow \left[\begin{array}{ccc|ccc} 2 & 0 & 0 & \frac{12}{8} & \frac{-5}{8} & \frac{-6}{8} \\ 0 & 8 & 0 & -4 & 3 & 2 \\ 0 & 0 & 1 & -1 & 1 & 1 \end{array} \right]$$

$$\left[\begin{array}{ccc} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{array} \right] \rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & \frac{12}{16} & \frac{-5}{16} & \frac{-6}{16} \\ 0 & 1 & 0 & -\frac{1}{2} & \frac{3}{8} & \frac{1}{4} \\ 0 & 0 & 1 & -1 & 1 & 1 \end{array} \right]$$

$\bar{x}_1 = \frac{12}{16}$ $\bar{x}_2 = \frac{3}{8}$

The Gauss-Jordan Method

$$\left[\begin{array}{ccc|ccc} 2 & 1 & 1 & 1 & 0 & 0 \\ 0 & -8 & -2 & -2 & 1 & 0 \\ 0 & 0 & 1 & -1 & 1 & 1 \end{array} \right] \xrightarrow{\left[\begin{array}{ccc} 1 & 0 & -1 \\ 0 & 1 & 2 \\ 0 & 0 & 1 \end{array} \right]} \left[\begin{array}{ccc|ccc} 2 & 1 & 0 & 2 & -1 & -1 \\ 0 & -8 & 0 & -4 & 3 & 2 \\ 0 & 0 & 1 & -1 & 1 & 1 \end{array} \right]$$

$$\left[\begin{array}{ccc} 1 & \frac{-8}{8} & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{array} \right] \rightarrow \left[\begin{array}{ccc|ccc} 2 & 0 & 0 & \frac{12}{8} & \frac{-5}{8} & \frac{-6}{8} \\ 0 & 8 & 0 & -4 & 3 & 2 \\ 0 & 0 & 1 & -1 & 1 & 1 \end{array} \right]$$

$$\left[\begin{array}{ccc} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{array} \right] \rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & \frac{12}{16} & \frac{-5}{16} & \frac{-6}{16} \\ 0 & 1 & 0 & -\frac{1}{2} & \frac{3}{8} & \frac{1}{4} \\ 0 & 0 & 1 & -1 & 1 & 1 \end{array} \right]$$

$x_1 = \frac{12}{16}$ $x_2 = \frac{-5}{16}$ $x_3 = \frac{-6}{16}$

The Gauss-Jordan Method

$$\left[\begin{array}{ccc|ccc} 2 & 1 & 1 & 1 & 0 & 0 \\ 0 & -8 & -2 & -2 & 1 & 0 \\ 0 & 0 & 1 & -1 & 1 & 1 \end{array} \right] \xrightarrow{\left[\begin{array}{ccc} 1 & 0 & -1 \\ 0 & 1 & 2 \\ 0 & 0 & 1 \end{array} \right]} \left[\begin{array}{ccc|ccc} 2 & 1 & 0 & 2 & -1 & -1 \\ 0 & -8 & 0 & -4 & 3 & 2 \\ 0 & 0 & 1 & -1 & 1 & 1 \end{array} \right]$$

$$\left[\begin{array}{ccc} 1 & \frac{-8}{8} & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{array} \right] \rightarrow \left[\begin{array}{ccc|ccc} 2 & 0 & 0 & \frac{12}{8} & \frac{-5}{8} & \frac{-6}{8} \\ 0 & 8 & 0 & -4 & 3 & 2 \\ 0 & 0 & 1 & -1 & 1 & 1 \end{array} \right]$$

$$\left[\begin{array}{ccc} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{array} \right] \rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & \frac{12}{16} & \frac{-5}{16} & \frac{-6}{16} \\ 0 & 1 & 0 & -\frac{1}{2} & \frac{3}{8} & \frac{1}{4} \\ 0 & 0 & 1 & -1 & 1 & 1 \end{array} \right]$$

$\underbrace{\hspace{10em}}_{A^{-1}}$

The Gauss-Jordan Method

The Gauss-Jordan Method

- In general: $n \times n$ matrix A , we let $\vec{e}_i = \begin{pmatrix} 0 \\ \vdots \\ 1 \\ \vdots \\ 0 \end{pmatrix}$

The Gauss-Jordan Method

- In general: $n \times n$ matrix A , we let $\vec{e}_i = \begin{pmatrix} 0 \\ \vdots \\ 1 \\ \vdots \\ 0 \end{pmatrix}$ i-th place.

The Gauss-Jordan Method

- In general: $n \times n$ matrix A , we let $\vec{e}_i = \begin{pmatrix} 0 \\ \vdots \\ 1 \\ \vdots \\ 0 \end{pmatrix}$ i-th place.

then we look at matrix : $[A | \vec{e}_1, \dots, \vec{e}_n]$

The Gauss-Jordan Method

- In general: $n \times n$ matrix A , we let $\vec{e}_i = \begin{pmatrix} 0 \\ \vdots \\ 1 \\ \vdots \\ 0 \end{pmatrix}$ i-th place.

then we look at matrix : $[A | \vec{e}_1, \dots, \vec{e}_n]$

- Use row operation to turn it into the form.

$$\left[\begin{matrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 1 \end{matrix} \mid B \right]$$

The Gauss-Jordan Method

- In general: $n \times n$ matrix A , we let $\vec{e}_i = \begin{pmatrix} 0 \\ \vdots \\ 1 \\ \vdots \\ 0 \end{pmatrix}$ i-th place.

then we look at matrix : $[A | \vec{e}_1, \dots, \vec{e}_n]$

- Use row operation to turn it into the form.

$$\left[\begin{matrix} 1 & 0 & 0 & | & B \\ 0 & 1 & 0 & | & \\ 0 & 0 & 1 & | & \\ \vdots & \vdots & \vdots & | & \end{matrix} \right]$$

$n \times n$ identity!

The Gauss-Jordan Method

- In general: $n \times n$ matrix A , we let $\vec{e}_i = \begin{pmatrix} 0 \\ \vdots \\ 1 \\ \vdots \\ 0 \end{pmatrix}$ i-th place.

then we look at matrix : $[A | \vec{e}_1, \dots, \vec{e}_n]$

- Use row operation to turn it into the form.

$$\left[\begin{matrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 1 \end{matrix} \mid B \right] \quad \text{it is } A^{-1}.$$

n \times n identity!

The Gauss-Jordan Method

- In general: $n \times n$ matrix A , we let $\vec{e}_i = \begin{pmatrix} 0 \\ \vdots \\ 1 \\ \vdots \\ 0 \end{pmatrix}$ i-th place.

then we look at matrix : $[A | \vec{e}_1, \dots, \vec{e}_n]$

- Use row operation to turn it into the form.

$$\left[\begin{matrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 1 \end{matrix} \mid B \right] \quad \text{it is } A^{-1}.$$

n \times n identity!

- If elimination fails to bring L.H.S to I, A is not invertible!

The Gauss-Jordan Method

$$\left[\begin{array}{c|c} \overbrace{\begin{array}{c} a_1 \\ \vdots \\ a_i \\ \vdots \\ a_n \end{array}} & \begin{array}{c} 0 \\ \vdots \\ 1 \\ \vdots \\ 0 \end{array} \end{array} \right]$$

The Gauss-Jordan Method

- The cost: $L^{-1}A = U$

$$\left[\begin{array}{c|c} \overbrace{\begin{array}{c} a_1 \\ \vdots \\ a_i \\ \vdots \\ a_n \end{array}} & \begin{array}{c} 0 \\ \vdots \\ 1 \\ \vdots \\ 0 \end{array} \end{array} \right]$$

The Gauss-Jordan Method

- The cost: $L^{-1}A = \mathcal{U}$ ← this take $\sim \frac{1}{3}n^3$ steps.

$$\left[\begin{array}{c|c} \overbrace{\begin{array}{c} a_1 \\ \vdots \\ a_i \\ \vdots \\ a_n \end{array}} & \begin{array}{c} 0 \\ \vdots \\ 1 \\ \vdots \\ 0 \end{array} \end{array} \right]$$

The Gauss-Jordan Method

- The cost: $L^{-1}A = \mathcal{U}$ ← this takes $\sim \frac{1}{3}n^3$ steps.
i.e. the cost of row operation to make it upper triangular.

$$\left[\begin{array}{c|c} \overbrace{\begin{array}{c} a_1 \\ \vdots \\ a_i \\ \vdots \\ a_n \end{array}} & \begin{array}{c} 0 \\ \vdots \\ 1 \\ \vdots \\ 0 \end{array} \end{array} \right]$$

The Gauss-Jordan Method

- The cost: $L^{-1}A = \mathcal{U}$ ← this takes $\sim \frac{1}{3}n^3$ steps.
i.e. the cost of row operation to make it upper triangular.
- We also take into account of the R.H.S. of $[A | \vec{e}_i]$:

$$\left[\begin{array}{c|c} \overline{a_1} & \overline{0} \\ \vdots & \vdots \\ \overline{a_i} & \overline{0} \\ \vdots & \vdots \\ \overline{a_n} & \overline{0} \end{array} \right]$$

The Gauss-Jordan Method

- The cost: $L^{-1}A = \mathcal{U}$ ← this takes $\sim \frac{1}{3}n^3$ steps.
i.e. the cost of row operation to make it upper triangular.
- We also take into account of the R.H.S. of $[A | \vec{e}_i]$:

$$\left[\begin{array}{c|c} \overbrace{\begin{array}{c} a_1 \\ \vdots \\ a_i \\ \vdots \\ a_n \end{array}} & \begin{array}{c} 0 \\ \vdots \\ 0 \\ \vdots \\ 0 \end{array} \end{array} \right] \xrightarrow{i\text{-th place}}$$

The Gauss-Jordan Method

- The cost: $L^{-1}A = \mathcal{U}$ ← this takes $\sim \frac{1}{3}n^3$ steps.
i.e. the cost of row operation to make it upper triangular.
- We also take into account of the R.H.S. of $[A | \vec{e}_i]$:

$$\left[\begin{array}{ccc|c} a & & & 0 \\ - & a_i & & \vdots \\ - & & a_n & 0 \end{array} \right] \quad \text{i-th place.}$$

The Gauss-Jordan Method

- The cost: $L^{-1}A = \mathcal{U}$ ← this takes $\sim \frac{1}{3}n^3$ steps.
i.e. the cost of row operation to make it upper triangular.
- We also take into account of the R.H.S. of $[A | \vec{e}_i]$:

$$\left[\begin{array}{c|c} \overbrace{\begin{matrix} a_1 \\ \vdots \\ a_i \\ \vdots \\ a_n \end{matrix}}^0 & \begin{matrix} 0 \\ \vdots \\ 0 \\ \vdots \\ 0 \end{matrix} \end{array} \right] \quad \text{i-th place.}$$

observe only row operations of adding a_j ($j \geq i$)
to other rows will have effects on R.H.S.

The Gauss-Jordan Method

- The cost: $L^{-1}A = \mathcal{U}$ ← this takes $\sim \frac{1}{3}n^3$ steps.
i.e. the cost of row operation to make it upper triangular.
- We also take into account of the R.H.S. of $[A | \vec{e}_i]$:

$$\left[\begin{array}{c|c} \overbrace{\begin{array}{c} a \\ - \\ a_i \\ - \\ a_n \end{array}}^0 & \begin{array}{c} 0 \\ \vdots \\ 0 \\ \vdots \\ 0 \end{array} \end{array} \right] \quad \text{i-th place.}$$

observe only row operations of adding a_j ($j \geq i$)
to other rows will have effects on R.H.S.

$$\Rightarrow (n-i) + (n-i-1) + \dots + 1 = \frac{(n-i)(n-i+1)}{2} \quad \text{steps taken on the R.H.S.}$$

The Gauss-Jordan Method

The Gauss-Jordan Method

- Summing over $i \sim \sum_{i=1}^n \frac{(n-i)(n-i+1)}{2} \sim \frac{n^3}{6}$.

The Gauss-Jordan Method

- Summing over $i \sim \sum_{i=1}^n \frac{(n-i)(n-i+1)}{2} \sim \frac{n^3}{6}$.
- Then we need to solve $U\vec{x}_i = \vec{c}_i$ by backward substitution.

The Gauss-Jordan Method

- Summing over $i \sim \sum_{i=1}^n \frac{(n-i)(n-i+1)}{2} \sim \frac{n^3}{6}$.
- Then we need to solve $\vec{U}\vec{x}_i = \vec{c}_i$ by backward substitution.
obtain from \vec{e}_i via row operation

The Gauss-Jordan Method

- Summing over $i \sim \sum_{i=1}^n \frac{(n-i)(n-i+1)}{2} \sim \frac{n^3}{6}$.
- Then we need to solve $\vec{U}\vec{x}_i = \vec{c}_i$ by backward substitution.
obtain from \vec{e}_i via row operation

For each i : the cost is $1+2+\dots+n-1 = \frac{n(n-1)}{2}$.

The Gauss-Jordan Method

- Summing over $i \sim \sum_{i=1}^n \frac{(n-i)(n-i+1)}{2} \sim \frac{n^3}{6}$.
 - Then we need to solve $U\vec{x}_i = \vec{c}_i$ by backward substitution.
obtain from \vec{e}_i via row operation
- For each i : the cost is $1+2+\dots+n-1 = \frac{n(n-1)}{2}$.
- Total cost of backward substitution $\sim \frac{n^2(n-1)}{2} \sim \frac{n^3}{2}$.

The Gauss-Jordan Method

- Summing over $i \sim \sum_{i=1}^n \frac{(n-i)(n-i+1)}{2} \sim \frac{n^3}{6}$.
- Then we need to solve $U\vec{x}_i = \vec{c}_i$ by backward substitution.
obtain from \vec{e}_i via row operation

For each i : the cost is $1+2+\dots+n-1 = \frac{n(n-1)}{2}$.

- Total cost of backward substitution $\sim \frac{n^2(n-1)}{2} \sim \frac{n^3}{2}$.
- Total cost $\sim \frac{n^3}{6} + \frac{n^3}{3} + \frac{n^3}{2} = n^3$

The Gauss-Jordan Method

.

The Gauss-Jordan Method

Remarks:

1. Computation of A^{-1} is very effective $\sim n^3$
notice if we compute A^2 , the cost is n^3 .

The Gauss-Jordan Method

Remarks: 1. Computation of A^{-1} is very effective $\sim n^3$

notice if we compute A^2 , the cost is n^3 .

Remark 2: • If we just want to solve for $Ax = b$
For efficiency, do not solve A^{-1} and then
use $x = A^{-1}b$ to compute.

The Gauss-Jordan Method

Remarks: 1. Computation of A^{-1} is very effective $\sim n^3$
notice if we compute A^2 , the cost is n^3 .

Remark 2: • If we just want to solve for $Ax = b$
For efficiency, do not solve A^{-1} and then
use $x = A^{-1}b$ to compute.
• Use Gauss Elimination directly.

Transpose matrix

Transpose matrix

- Consider

$$\begin{cases} a_{11}x_1 + a_{12}x_2 + a_{13}x_3 = b_1 \\ a_{21}x_1 + a_{22}x_2 + a_{23}x_3 = b_2 \\ a_{31}x_1 + a_{32}x_2 + a_{33}x_3 = b_3. \end{cases}$$

which we write it as $Ax = b$.

Transpose matrix

- Consider $\begin{cases} a_{11}x_1 + a_{12}x_2 + a_{13}x_3 = b_1 \\ a_{21}x_1 + a_{22}x_2 + a_{23}x_3 = b_2 \\ a_{31}x_1 + a_{32}x_2 + a_{33}x_3 = b_3 \end{cases}$.
which we write it as $Ax = b$.
- It can also be written a $[x_1, x_2, x_3] \tilde{A} = [b_1, b_2, b_3]$

Transpose matrix

- Consider

$$\begin{cases} a_{11}x_1 + a_{12}x_2 + a_{13}x_3 = b_1 \\ a_{21}x_1 + a_{22}x_2 + a_{23}x_3 = b_2 \\ a_{31}x_1 + a_{32}x_2 + a_{33}x_3 = b_3 \end{cases}$$

which we write it as $Ax = b$.

- It can also be written a $[x_1, x_2, x_3] \tilde{A} = [b_1, b_2, b_3]$

the matrix $\tilde{A} = \begin{pmatrix} a_{11} & a_{21} & a_{31} \\ a_{12} & a_{22} & a_{32} \\ a_{13} & a_{23} & a_{33} \end{pmatrix}$ is called

the **transpose** of A .

Transpose matrix

- Consider

$$\begin{cases} a_{11}x_1 + a_{12}x_2 + a_{13}x_3 = b_1 \\ a_{21}x_1 + a_{22}x_2 + a_{23}x_3 = b_2 \\ a_{31}x_1 + a_{32}x_2 + a_{33}x_3 = b_3 \end{cases}$$

which we write it as $Ax = b$.

- It can also be written a $[x_1, x_2, x_3] \tilde{A} = [b_1, b_2, b_3]$

the matrix $\tilde{A} = \begin{pmatrix} a_{11} & a_{21} & a_{31} \\ a_{12} & a_{22} & a_{32} \\ a_{13} & a_{23} & a_{33} \end{pmatrix}$ is called

the **transpose** of A .

- We write A^T for the transpose matrix

Transpose matrix

- Consider

$$\begin{cases} a_{11}x_1 + a_{12}x_2 + a_{13}x_3 = b_1 \\ a_{21}x_1 + a_{22}x_2 + a_{23}x_3 = b_2 \\ a_{31}x_1 + a_{32}x_2 + a_{33}x_3 = b_3 \end{cases}$$

which we write it as $Ax = b$.

- It can also be written a $[x_1, x_2, x_3] \tilde{A} = [b_1, b_2, b_3]$

the matrix $\tilde{A} = \begin{pmatrix} a_{11} & a_{21} & a_{31} \\ a_{12} & a_{22} & a_{32} \\ a_{13} & a_{23} & a_{33} \end{pmatrix}$ is called

the **transpose** of A .

- We write A^T for the transpose matrix

Def: Let $A = (a_{ij})$ be $m \times n$ matrix, we let an $n \times m$ matrix $A^T = (a_{ji}^T)$ by $a_{ji}^T = a_{ij}$.

Transpose matrix

Transpose matrix

e.g.

$$A = \begin{bmatrix} 2 & 1 & 4 \\ 0 & 0 & 3 \end{bmatrix} \Rightarrow A^T = \begin{bmatrix} 2 & 0 \\ 1 & 0 \\ 4 & 3 \end{bmatrix}$$

Transpose matrix

e.g.

$$A = \begin{bmatrix} 2 & 1 & 4 \\ 0 & 0 & 3 \end{bmatrix} \Rightarrow A^T = \begin{bmatrix} 2 & 0 \\ 1 & 0 \\ 4 & 3 \end{bmatrix}$$

is the “flip” of the matrix along diagonal.

Transpose matrix

e.g.

$$A = \begin{bmatrix} 2 & 1 & 4 \\ 0 & 0 & 3 \end{bmatrix} \Rightarrow A^T = \begin{bmatrix} 2 & 0 \\ 1 & 0 \\ 4 & 3 \end{bmatrix}$$

is the “flip” of the matrix along diagonal.

Prop:

$$1. (A+B)^T = A^T + B^T$$

Transpose matrix

e.g.

$$A = \begin{bmatrix} 2 & 1 & 4 \\ 0 & 0 & 3 \end{bmatrix} \Rightarrow A^T = \begin{bmatrix} 2 & 0 \\ 1 & 0 \\ 4 & 3 \end{bmatrix}$$

is the “flip” of the matrix along diagonal.

Prop:

$$1. (A+B)^T = A^T + B^T$$

$$2. (AB)^T = B^T A^T$$

Transpose matrix

e.g.

$$A = \begin{bmatrix} 2 & 1 & 4 \\ 0 & 0 & 3 \end{bmatrix} \Rightarrow A^T = \begin{bmatrix} 2 & 0 \\ 1 & 0 \\ 4 & 3 \end{bmatrix}$$

is the “flip” of the matrix along diagonal.

Prop:

$$1. (A+B)^T = A^T + B^T$$

$$2. (AB)^T = B^T A^T$$

$$3. (A^{-1})^T = (A^T)^{-1}$$

Transpose matrix

e.g.

$$A = \begin{bmatrix} 2 & 1 & 4 \\ 0 & 0 & 3 \end{bmatrix} \Rightarrow A^T = \begin{bmatrix} 2 & 0 \\ 1 & 0 \\ 4 & 3 \end{bmatrix}$$

is the “flip” of the matrix along diagonal.

Prop:

$$1. (A+B)^T = A^T + B^T$$

$$2. (AB)^T = B^T A^T$$

$$3. (A^{-1})^T = (A^T)^{-1}$$

this means A^T is
invertible if A is.

Transpose matrix

e.g.

$$A = \begin{bmatrix} 2 & 1 & 4 \\ 0 & 0 & 3 \end{bmatrix} \Rightarrow A^T = \begin{bmatrix} 2 & 0 \\ 1 & 0 \\ 4 & 3 \end{bmatrix}$$

is the "flip" of the matrix along diagonal.

Prop:

$$1. (A+B)^T = A^T + B^T$$

$$2. (AB)^T = B^T A^T$$

$$3. (A^{-1})^T = (A^T)^{-1}$$

this means A^T is
invertible if A is.

Pf: 1

exercise.

Transpose matrix

Transpose matrix

Pf: 2. e.g.

Transpose matrix

Pf: 2.

E.g.

$$\begin{array}{c} \text{A} & \text{B} \\ \left[\begin{array}{cc} 1 & 0 \\ 1 & 1 \end{array} \right] \left[\begin{array}{ccc} 3 & 3 & 3 \\ 2 & 2 & 2 \end{array} \right] = \left[\begin{array}{ccc} 3 & 3 & 3 \\ 4 & 4 & 4 \end{array} \right] \end{array}$$

Transpose matrix

Pf: 2.

E.g.

$$\begin{matrix} A & B \\ \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} & \begin{bmatrix} 3 & 3 & 3 \\ 2 & 2 & 2 \end{bmatrix} = \begin{bmatrix} 3 & 3 & 3 \\ 4 & 5 & 5 \end{bmatrix} \end{matrix}$$

$$\begin{matrix} B^T & A^T \\ \begin{bmatrix} 3 & 2 \\ 3 & 2 \\ 3 & 2 \end{bmatrix} & \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 3 & 5 \\ 3 & 5 \end{bmatrix} \end{matrix}$$

Transpose matrix

Pf: 2.

E.g.

$$\begin{matrix} & \textcolor{red}{A} & \textcolor{red}{B} \\ \left[\begin{matrix} 1 & 0 \\ 1 & 1 \end{matrix} \right] & \left[\begin{matrix} 3 & 3 & 3 \\ 2 & 2 & 2 \end{matrix} \right] & = & \left[\begin{matrix} 3 & 3 & 3 \\ 4 & 5 & 5 \end{matrix} \right] \end{matrix}$$

$$\begin{matrix} \textcolor{red}{B^T} & \textcolor{red}{A^T} \\ \left[\begin{matrix} 3 & 2 \\ 3 & 2 \\ 3 & 2 \end{matrix} \right] & \left[\begin{matrix} 1 & 1 \\ 0 & 1 \end{matrix} \right] & = & \left[\begin{matrix} 3 & 5 \\ 3 & 5 \end{matrix} \right] \end{matrix}$$

$$AB = [A\vec{b}_1, \dots, A\vec{b}_n]$$

Transpose matrix

Pf: 2.

e.g.

$$\begin{matrix} A & B \\ \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} & \begin{bmatrix} 3 & 3 & 3 \\ 2 & 2 & 2 \end{bmatrix} \end{matrix} = \begin{bmatrix} 3 & 3 & 3 \\ 4 & 5 & 5 \end{bmatrix}$$

$$\begin{matrix} B^T & A^T \\ \begin{bmatrix} 3 & 2 \\ 3 & 2 \\ 3 & 2 \end{bmatrix} & \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \end{matrix} = \begin{bmatrix} 3 & 5 \\ 3 & 5 \\ 3 & 5 \end{bmatrix}$$

$m \times k$
 $k \times n$ matrices

$$\downarrow \quad \downarrow$$
$$AB = [A\vec{b}_1, \dots, A\vec{b}_n]$$

Transpose matrix

Pf: 2.

e.g.

$$\begin{matrix} A & B \\ \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} & \begin{bmatrix} 3 & 3 & 3 \\ 2 & 2 & 2 \end{bmatrix} \end{matrix} = \begin{bmatrix} 3 & 3 & 3 \\ 4 & 4 & 4 \end{bmatrix}$$

$$\begin{matrix} B^T & A^T \\ \begin{bmatrix} 3 & 2 \\ 3 & 2 \\ 3 & 2 \end{bmatrix} & \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \end{matrix} = \begin{bmatrix} 3 & 5 \\ 3 & 5 \\ 3 & 5 \end{bmatrix}$$

$m \times k$
 $k \times n$ matrices

$$\downarrow$$
$$AB = [A\vec{b}_1, \dots, A\vec{b}_n] \Rightarrow (AB)^T = \begin{bmatrix} -(A\vec{b}_1)^T \\ \vdots \\ -(A\vec{b}_n)^T \end{bmatrix}$$

Transpose matrix

Pf: 2.

E.g.

$$\begin{matrix} A & B \\ \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} & \begin{bmatrix} 3 & 3 & 3 \\ 2 & 2 & 2 \end{bmatrix} \end{matrix} = \begin{bmatrix} 3 & 3 & 3 \\ 4 & 4 & 4 \end{bmatrix}$$

$$\begin{matrix} B^T & A^T \\ \begin{bmatrix} 3 & 2 \\ 3 & 2 \\ 3 & 2 \end{bmatrix} & \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \end{matrix} = \begin{bmatrix} 3 & 5 \\ 3 & 5 \\ 3 & 5 \end{bmatrix}$$

$m \times k$
 $k \times n$ matrices

$$AB = [A\vec{b}_1, \dots, A\vec{b}_n] \Rightarrow (AB)^T = \begin{bmatrix} -(A\vec{b}_1)^T \\ \vdots \\ -(A\vec{b}_n)^T \end{bmatrix}$$

only need to know $(A\vec{b}_i)^T = \vec{b}_i^T \cdot A^T$.

Transpose matrix

Pf: 2.

E.g.

$$\begin{matrix} A & B \\ \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} & \begin{bmatrix} 3 & 3 & 3 \\ 2 & 2 & 2 \end{bmatrix} \end{matrix} = \begin{bmatrix} 3 & 3 & 3 \\ 4 & 4 & 4 \end{bmatrix}$$

$$\begin{matrix} B^T & A^T \\ \begin{bmatrix} 3 & 2 \\ 3 & 2 \\ 3 & 2 \end{bmatrix} & \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \end{matrix} = \begin{bmatrix} 3 & 5 \\ 3 & 5 \\ 3 & 5 \end{bmatrix}$$

$m \times k$
 $k \times n$ matrices

$$AB = [A\vec{b}_1, \dots, A\vec{b}_n] \Rightarrow (AB)^T = \begin{bmatrix} -(A\vec{b}_1)^T \\ \vdots \\ -(A\vec{b}_n)^T \end{bmatrix}$$

only need to know $(A\vec{b}_i)^T = \vec{b}_i^T \cdot A^T$.

$$\Rightarrow (AB)^T = \begin{bmatrix} -\vec{b}_1^T A^T \\ -\vec{b}_2^T A^T \\ \vdots \\ -\vec{b}_n^T A^T \end{bmatrix}$$

Transpose matrix

Pf: 2.

E.g.

$$\begin{matrix} A & B \\ \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} & \begin{bmatrix} 3 & 3 & 3 \\ 2 & 2 & 2 \end{bmatrix} \end{matrix} = \begin{bmatrix} 3 & 3 & 3 \\ 4 & 4 & 4 \end{bmatrix}$$

$$\begin{matrix} B^T & A^T \\ \begin{bmatrix} 3 & 2 \\ 3 & 2 \\ 3 & 2 \end{bmatrix} & \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \end{matrix} = \begin{bmatrix} 3 & 5 \\ 3 & 5 \\ 3 & 5 \end{bmatrix}$$

$m \times k$
 $k \times n$ matrices

$$AB = [A\vec{b}_1, \dots, A\vec{b}_n] \Rightarrow (AB)^T = \begin{bmatrix} -(A\vec{b}_1)^T \\ \vdots \\ -(A\vec{b}_n)^T \end{bmatrix}$$

only need to know $(A\vec{b}_i)^T = \vec{b}_i^T \cdot A^T$.

$$\Rightarrow (AB)^T = \begin{bmatrix} -\vec{b}_1^T A^T \\ -\vec{b}_2^T A^T \\ \vdots \\ -\vec{b}_n^T A^T \end{bmatrix} = \begin{bmatrix} -\vec{b}_1^T \\ \vdots \\ -\vec{b}_n^T \end{bmatrix} A^T$$

Transpose matrix

Pf: 2.

E.g.

$$\begin{matrix} A & B \\ \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} & \begin{bmatrix} 3 & 3 & 3 \\ 2 & 2 & 2 \end{bmatrix} \end{matrix} = \begin{bmatrix} 3 & 3 & 3 \\ 4 & 4 & 4 \end{bmatrix}$$

$$\begin{matrix} B^T & A^T \\ \begin{bmatrix} 3 & 2 \\ 3 & 2 \\ 3 & 2 \end{bmatrix} & \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \end{matrix} = \begin{bmatrix} 3 & 5 \\ 3 & 5 \\ 3 & 5 \end{bmatrix}$$

$m \times k$
 $k \times n$ matrices

$$AB = [A\vec{b}_1, \dots, A\vec{b}_n] \Rightarrow (AB)^T = \begin{bmatrix} -(A\vec{b}_1)^T \\ \vdots \\ -(A\vec{b}_n)^T \end{bmatrix}$$

only need to know $(A\vec{b}_i)^T = \vec{b}_i^T \cdot A^T$.

$$\Rightarrow (AB)^T = \begin{bmatrix} -\vec{b}_1^T A^T \\ -\vec{b}_2^T A^T \\ \vdots \\ -\vec{b}_n^T A^T \end{bmatrix} = \begin{bmatrix} -\vec{b}_1^T \\ \vdots \\ -\vec{b}_n^T \end{bmatrix} A^T = B^T A^T.$$

Transpose matrix

Transpose matrix

Pf: 3, $(A^{-1})^T = (A^T)^{-1}$

Transpose matrix

Pf: 3, $(A^{-1})^T = (A^T)^{-1}$

- $AA^{-1} = I$

Transpose matrix

$$\underline{\text{Pf}}: 3, \quad (A^{-1})^T = (A^T)^{-1}$$

$$\bullet \quad AA^{-1} = I \quad \Rightarrow \quad (AA^{-1})^T = I^T = I.$$

Transpose matrix

Pf: 3, $(A^{-1})^T = (A^T)^{-1}$

• $AA^{-1} = I \Rightarrow (AA^{-1})^T = I^T = I.$

$$(A^{-1})^T A^T = I.$$

Transpose matrix

Pf: 3, $(A^{-1})^T = (A^T)^{-1}$

• $AA^{-1} = I \Rightarrow (AA^{-1})^T = I^T = I.$

$$(A^{-1})^T A^T = I.$$

• $A^{-1}A = I \Rightarrow A^T (A^{-1})^T = I.$

Transpose matrix

Pf: 3, $(A^{-1})^T = (A^T)^{-1}$

• $AA^{-1} = I \Rightarrow (AA^{-1})^T = I^T = I.$

$$(A^{-1})^T A^T = I.$$

• $A^{-1}A = I \Rightarrow A^T(A^{-1})^T = I.$

we conclude A^T is invertible and

$$(A^T)^{-1} = (A^{-1})^T$$

Symmetric matrix

Symmetric matrix

Def:

A is called a **symmetric matrix** if $A^T = A$.

Symmetric matrix

Def:

A is called a **symmetric matrix** if $A^T = A$.

e.g:

$$A = \begin{bmatrix} 1 & 2 \\ 2 & 8 \end{bmatrix}, \quad A^{-1} = \frac{1}{4} \begin{bmatrix} 8 & -2 \\ -2 & 1 \end{bmatrix}.$$

Symmetric matrix

Def:

A is called a **symmetric matrix** if $A^T = A$.

e.g: $A = \begin{bmatrix} 1 & 2 \\ 2 & 8 \end{bmatrix}$, $A^{-1} = \frac{1}{4} \begin{bmatrix} 8 & -2 \\ -2 & 1 \end{bmatrix}$.

- A symmetric matrix must be a square matrix.

Symmetric matrix

Def:

A is called a **symmetric matrix** if $A^T = A$.

e.g: $A = \begin{bmatrix} 1 & 2 \\ 2 & 8 \end{bmatrix}$, $A^{-1} = \frac{1}{4} \begin{bmatrix} 8 & -2 \\ -2 & 1 \end{bmatrix}$.

- A symmetric matrix must be a square matrix.
- If A is symmetric and invertible, then A^{-1} is symmetric.

Symmetric matrix

Def:

A is called a **symmetric matrix** if $A^T = A$.

e.g: $A = \begin{bmatrix} 1 & 2 \\ 2 & 8 \end{bmatrix}$, $A^{-1} = \frac{1}{4} \begin{bmatrix} 8 & -2 \\ -2 & 1 \end{bmatrix}$.

- A symmetric matrix must be a square matrix.
- If A is symmetric and invertible, then A^{-1} is symmetric.

e.g.:

AA^T is a symmetric matrix for any $m \times n$ A.

Symmetric matrix

Def:

A is called a **symmetric matrix** if $A^T = A$.

e.g: $A = \begin{bmatrix} 1 & 2 \\ 2 & 8 \end{bmatrix}$, $A^{-1} = \frac{1}{4} \begin{bmatrix} 8 & -2 \\ -2 & 1 \end{bmatrix}$.

- A symmetric matrix must be a square matrix.
- If A is symmetric and invertible, then A^{-1} is symmetric.

e.g.:

AA^T is a symmetric matrix for any $m \times n$ A .

Thm:

Suppose there is a LU factorization for non-singular symmetric $A = LDL^T$, then $L^T = U$.

Symmetric matrix

Symmetric matrix

- Let's look at the elimination process for symmetric matrix

$$A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 3 & -1 \\ 1 & -1 & 0 \end{bmatrix}$$

Symmetric matrix

- Let's look at the elimination process for symmetric matrix

$$A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 3 & -1 \\ 1 & -1 & 0 \end{bmatrix}$$

$$\underbrace{\begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ -1 & 0 & 1 \end{bmatrix}}_{E.} A = \begin{bmatrix} 1 & 2 & 1 \\ 0 & -1 & -3 \\ 0 & -3 & -1 \end{bmatrix}$$

Symmetric matrix

- Let's look at the elimination process for symmetric matrix

$$A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 3 & -1 \\ 1 & -1 & 0 \end{bmatrix} \quad \underbrace{\begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ -1 & 0 & 1 \end{bmatrix}}_{E.} A = \begin{bmatrix} 1 & 2 & 1 \\ 0 & -1 & -3 \\ 0 & -3 & -1 \end{bmatrix}$$

- $EAE^T = \begin{bmatrix} 1 & 2 & 1 \\ 0 & -1 & -3 \\ 0 & -3 & -1 \end{bmatrix} \begin{bmatrix} 1 & -2 & -1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & -1 & -3 \\ 0 & -3 & -1 \end{bmatrix}$

Symmetric matrix

- Let's look at the elimination process for symmetric matrix

$$A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 3 & -1 \\ 1 & -1 & 0 \end{bmatrix} \quad \underbrace{\begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ -1 & 0 & 1 \end{bmatrix}}_{E.} A = \begin{bmatrix} 1 & 2 & 1 \\ 0 & -1 & -3 \\ 0 & -3 & -1 \end{bmatrix}$$

$$EA E^T = \begin{bmatrix} 1 & 2 & 1 \\ 0 & -1 & -3 \\ 0 & -3 & -1 \end{bmatrix} \begin{bmatrix} 1 & -2 & -1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & -1 & -3 \\ 0 & -3 & -1 \end{bmatrix}$$

Symmetric matrix

- Let's look at the elimination process for symmetric matrix

$$A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 3 & -1 \\ 1 & -1 & 0 \end{bmatrix} \quad \underbrace{\begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ -1 & 0 & 1 \end{bmatrix}}_{E.} A = \begin{bmatrix} 1 & 2 & 1 \\ 0 & -1 & -3 \\ 0 & -3 & -1 \end{bmatrix}$$

$$EA E^T = \begin{bmatrix} 1 & 2 & 1 \\ 0 & -1 & -3 \\ 0 & -3 & -1 \end{bmatrix} \begin{bmatrix} 1 & -2 & -1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & -1 & -3 \\ 0 & -3 & -1 \end{bmatrix}$$

↑ remain symmetric

Symmetric matrix

- Let's look at the elimination process for symmetric matrix

$$A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 3 & -1 \\ 1 & -1 & 0 \end{bmatrix} \quad \underbrace{\begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ -1 & 0 & 1 \end{bmatrix}}_{E.} A = \begin{bmatrix} 1 & 2 & 1 \\ 0 & -1 & -3 \\ 0 & -3 & -1 \end{bmatrix}$$

$$EA E^T = \begin{bmatrix} 1 & 2 & 1 \\ 0 & -1 & -3 \\ 0 & -3 & -1 \end{bmatrix} \begin{bmatrix} 1 & -2 & -1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & -1 & -3 \\ 0 & -3 & -1 \end{bmatrix}$$

↑
remain symmetric

- For each elimination step : we notice the next lower right-handed corner block remains symmetric

Symmetric matrix

- Let's look at the elimination process for symmetric matrix

$$A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 3 & -1 \\ 1 & -1 & 0 \end{bmatrix}; \quad \underbrace{\begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ -1 & 0 & 1 \end{bmatrix}}_{E.} A = \begin{bmatrix} 1 & 2 & 1 \\ 0 & -1 & -3 \\ 0 & -3 & -1 \end{bmatrix}$$

$$EA E^T = \begin{bmatrix} 1 & 2 & 1 \\ 0 & -1 & -3 \\ 0 & -3 & -1 \end{bmatrix} \begin{bmatrix} 1 & -2 & -1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & -1 & -3 \\ 0 & -3 & -1 \end{bmatrix}$$

remain symmetric

- For each elimination step : we notice the next lower right-handed corner block remains **symmetric**
- So for symmetric matrices : elimination cost $\sim \frac{n^3}{6}$.

Inner and outer product

Inner and outer product

- Inner product

For $\vec{x}, \vec{y} \in \mathbb{R}^n$ be column vector

we define: $\langle \vec{x}, \vec{y} \rangle := \vec{x}^T y \in \mathbb{R}$.

Inner and outer product

- Inner product

For $\vec{x}, \vec{y} \in \mathbb{R}^n$ be column vector

we define: $\langle \vec{x}, \vec{y} \rangle := \vec{x}^T y \in \mathbb{R}$.

e.g: $\left\langle \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix} \right\rangle = \begin{bmatrix} 1 & 2 & 3 \end{bmatrix} \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix}$

$$= 1 \cdot 4 + 2 \cdot 5 + 3 \cdot 6$$
$$= 32.$$

Inner and outer product

- Inner product

For $\vec{x}, \vec{y} \in \mathbb{R}^n$ be column vector

we define: $\langle \vec{x}, \vec{y} \rangle := \vec{x}^T y \in \mathbb{R}$.

e.g: $\left\langle \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix} \right\rangle = \begin{bmatrix} 1 & 2 & 3 \end{bmatrix} \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix}$

$$= 1 \cdot 4 + 2 \cdot 5 + 3 \cdot 6$$
$$= 32.$$

- Length/Norm: We let $\|\vec{x}\| = \sqrt{\langle \vec{x}, \vec{x} \rangle}$.

Inner and outer product

Inner and outer product

- For $\vec{x}, \vec{y} \in \mathbb{R}^n$, we define the outer product to be the matrix $\vec{x}\vec{y}^T$

Inner and outer product

- For $\vec{x}, \vec{y} \in \mathbb{R}^n$, we define the outer product to be the matrix $\vec{x}\vec{y}^T$

e.g.

$$\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} [4 \ 5 \ 6] = \begin{bmatrix} 4 & 5 & 6 \\ 8 & 10 & 12 \\ 12 & 15 & 18 \end{bmatrix}$$

Inner and outer product

- For $\vec{x}, \vec{y} \in \mathbb{R}^n$, we define the outer product to be the matrix $\vec{x}\vec{y}^T$

e.g.

$$\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} [4 \ 5 \ 6] = \begin{bmatrix} 4 & 5 & 6 \\ 8 & 10 & 12 \\ 12 & 15 & 18 \end{bmatrix}$$

- the columns are multiple of \vec{x} .
The rows are multiple of \vec{y} .

Inner and outer product

Inner and outer product

e.g.:

compute $\left(\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} [1 \ 2 \ 3] \right)^{2020}$

Inner and outer product

e.g.:

compute $\vec{x}^T = \left(\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} [1 \ 2 \ 3] \right)^{2020}$

Inner and outer product

e.g.:

compute

$$\vec{x}^T = \left(\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} [1 \ 2 \ 3] \right)^{2020}$$

observe:

$$\left(\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} [1 \ 2 \ 3] \right)^2 = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} [1 \ 2 \ 3] \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} [1 \ 2 \ 3]$$

Inner and outer product

e.g.:

compute

$$\vec{x}^T = \left(\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} [1, 2, 3] \right)^{2020}$$

observe:

$$\begin{aligned} \left(\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} [1, 2, 3] \right)^2 &= \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} [1, 2, 3] \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} [1, 2, 3] \\ &= \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \langle \vec{x}, \vec{x} \rangle [1, 2, 3] \end{aligned}$$

Inner and outer product

e.g.:

compute

$$\vec{x}^T = \left(\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} [1, 2, 3] \right)^{2020}$$

observe:

$$\begin{aligned} & \left(\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} [1, 2, 3] \right)^2 = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} [1, 2, 3] \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} [1, 2, 3] \\ &= \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \textcircled{< \vec{x}, \vec{x} >} [1, 2, 3] \end{aligned}$$

scalar!

Inner and outer product

e.g.:

compute $\vec{x}^T = \left(\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} [1, 2, 3] \right)^{2020}$

observe:

$$\begin{aligned} & \left(\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} [1, 2, 3] \right)^2 = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} [1, 2, 3] \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} [1, 2, 3] \\ &= \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \textcircled{<}\vec{x}, \vec{x}> [1, 2, 3] \text{ scalar!} \\ &= <\vec{x}, \vec{x}> \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} [1, 2, 3] \end{aligned}$$

Inner and outer product

e.g.: compute $\vec{x}^T = \left(\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} [1 \ 2 \ 3] \right)^{2020}$

observe: $\left(\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} [1 \ 2 \ 3] \right)^2 = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} [1 \ 2 \ 3] \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} [1 \ 2 \ 3]$

$= \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \underbrace{\langle \vec{x}, \vec{x} \rangle}_{\text{scalar!}} [1, 2, 3]$

$$= \langle \vec{x}, \vec{x} \rangle \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} [1 \ 2 \ 3]$$

$$\left(\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} [1 \ 2 \ 3] \right)^3 = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} [1 \ 2 \ 3] \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} [1 \ 2 \ 3] \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} [1 \ 2 \ 3]$$

Inner and outer product

e.g.:

compute

$$\vec{x}^T = \left(\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} [1 \ 2 \ 3] \right)^{2020}$$

observe:

$$\left(\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} [1 \ 2 \ 3] \right)^2 = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} [1 \ 2 \ 3] \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} [1 \ 2 \ 3]$$

scalar!

$$= \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \circled{< \vec{x}, \vec{x} >} [1, 2, 3]$$

$$\left(\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} [1 \ 2 \ 3] \right)^3 = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \circled{[1 \ 2 \ 3]} \circled{\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}} \circled{[1 \ 2 \ 3]} \circled{\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}}$$

Inner and outer product

e.g.:

compute

$$\vec{x}^T = \left(\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} [1, 2, 3] \right)^{2020}$$

observe:

$$\left(\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} [1, 2, 3] \right)^2 = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} [1, 2, 3] \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} [1, 2, 3]$$

scalar!

$$= \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \circled{< \vec{x}, \vec{x} >} [1, 2, 3]$$

$$\left(\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} [1, 2, 3] \right)^3 = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \circled{[1, 2, 3]} \circled{\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}} [1, 2, 3]$$
$$= < \vec{x}, \vec{x} >^2 \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} [1, 2, 3]$$

Inner and outer product

Inner and outer product

$$\left(\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \begin{bmatrix} 1 & 2 & 3 \end{bmatrix} \right)^{2020}$$

Inner and outer product

$$\begin{aligned} \left(\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \begin{bmatrix} 1 & 2 & 3 \end{bmatrix} \right)^{2020} &= \langle \tilde{x}, \tilde{x} \rangle^{2019} \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \begin{bmatrix} 1 & 2 & 3 \end{bmatrix} \\ &= 14^{2019} \begin{bmatrix} 1 & 2 & 3 \\ 2 & 4 & 6 \\ 3 & 4 & 9 \end{bmatrix} \end{aligned}$$

Inner and outer product

$$\left(\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \begin{bmatrix} 1 & 2 & 3 \end{bmatrix} \right)^{2020} = \langle \tilde{x}, \tilde{x} \rangle^{2019} \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \begin{bmatrix} 1 & 2 & 3 \end{bmatrix}$$
$$= 14^{2019} \begin{bmatrix} 1 & 2 & 3 \\ 2 & 4 & 6 \\ 3 & 4 & 9 \end{bmatrix}$$

e.g.: Compute the outer product expansion of $\mathbf{x}\mathbf{Y}^T$

$$\mathbf{x} = \begin{bmatrix} 3 & 1 \\ 2 & 4 \\ 1 & 2 \end{bmatrix}, \quad \mathbf{Y} = \begin{bmatrix} 1 & 2 \\ 2 & 4 \\ 3 & 1 \end{bmatrix}$$

Inner and outer product

$$\left(\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \begin{bmatrix} 1 & 2 & 3 \end{bmatrix} \right)^{2020} = \langle \tilde{x}, \tilde{x} \rangle^{2019} \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \begin{bmatrix} 1 & 2 & 3 \end{bmatrix}$$
$$= 14^{2019} \begin{bmatrix} 1 & 2 & 3 \\ 2 & 4 & 6 \\ 3 & 4 & 9 \end{bmatrix}$$

e.g.: Compute the outer product expansion of $X Y^T$

$$X = \begin{bmatrix} 3 & 1 \\ 2 & 4 \\ 1 & 2 \end{bmatrix}, \quad Y = \begin{bmatrix} 1 & 2 \\ 2 & 4 \\ 3 & 1 \end{bmatrix}$$

$$XY^T = \begin{bmatrix} 3 & 1 \\ 2 & 4 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} 1 & 2 & 3 \\ 2 & 4 & 1 \end{bmatrix}$$

Inner and outer product

$$\left(\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} [1 \ 2 \ 3] \right)^{2020} = \langle \tilde{x}, \tilde{x} \rangle^{2019} \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} [1 \ 2 \ 3]$$
$$= 14^{2019} \begin{bmatrix} 1 & 2 & 3 \\ 2 & 4 & 6 \\ 3 & 4 & 9 \end{bmatrix}$$

e.g.: Compute the outer product expansion of $X Y^T$

$$X = \begin{bmatrix} 3 & 1 \\ 2 & 4 \\ 1 & 2 \end{bmatrix}, \quad Y = \begin{bmatrix} 1 & 2 \\ 2 & 4 \\ 3 & 1 \end{bmatrix}$$

$$XY^T = \begin{bmatrix} 3 & 1 \\ 2 & 4 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} 1 & 2 & 3 \\ 2 & 4 & 1 \end{bmatrix} = \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix} [1 \ 2 \ 3] + \begin{bmatrix} 1 \\ 4 \\ 2 \end{bmatrix} [2 \ 4 \ 1]$$