

Lecture 12

Matrix and linear transformation

Matrix representation of linear transformation

Matrix representation of linear transformation

- Recall that for a linear transformation $L : V \rightarrow W$

Matrix representation of linear transformation

- Recall that for a linear transformation $L : V \rightarrow W$
Fixing a basis $\vec{v}_1, \dots, \vec{v}_n$ of V , and $\vec{w}_1, \dots, \vec{w}_m$ of W .

Matrix representation of linear transformation

- Recall that for a linear transformation $L : V \rightarrow W$

Fixing a basis $\vec{v}_1, \dots, \vec{v}_n$ of V , and $\vec{w}_1, \dots, \vec{w}_m$ of W .

→ a matrix $[L]$ representing L

Matrix representation of linear transformation

- Recall that for a linear transformation $L : V \rightarrow W$
Fixing a basis $\vec{v}_1, \dots, \vec{v}_n$ of V , and $\vec{w}_1, \dots, \vec{w}_m$ of W .
→ a matrix $[L]$ representing L
- $[L(\vec{v}_1), \dots, L(\vec{v}_n)] = [\vec{w}_1, \dots, \vec{w}_m] [L]$

Matrix representation of linear transformation

- Recall that for a linear transformation $L : V \rightarrow W$

Fixing a basis $\vec{v}_1, \dots, \vec{v}_n$ of V , and $\vec{w}_1, \dots, \vec{w}_m$ of W .

→ a matrix $[L]$ representing L

- $[L(\vec{v}_1), \dots, L(\vec{v}_n)] = [\vec{w}_1, \dots, \vec{w}_m] [L]$
- For any general vector $\vec{r} = c_1\vec{v}_1 + \dots + c_n\vec{v}_n$, we have

Matrix representation of linear transformation

- Recall that for a linear transformation $L : V \rightarrow W$

Fixing a basis $\vec{v}_1, \dots, \vec{v}_n$ of V , and $\vec{w}_1, \dots, \vec{w}_m$ of W .

→ a matrix $[L]$ representing L

$$[L(\vec{v}_1), \dots, L(\vec{v}_n)] = [\vec{w}_1, \dots, \vec{w}_m] [L]$$

- For any general vector $\vec{v} = c_1\vec{v}_1 + \dots + c_n\vec{v}_n$, we have

$$L(\vec{v}) = [\vec{w}_1, \dots, \vec{w}_n] [L] \begin{bmatrix} c_1 \\ \vdots \\ c_n \end{bmatrix}$$

Matrix representation of linear transformation

Matrix representation of linear transformation

e.g.: Let $I: P_3 \rightarrow P_4$ given by $I(f(x)) = \int_0^x f(t) dt$

Matrix representation of linear transformation

e.g.: Let $I: P_3 \rightarrow P_4$ given by $I(f(x)) = \int_0^x f(t) dt$

- take a basis $1, x, x^2, x^3$ for P_3

Matrix representation of linear transformation

e.g.: Let $I: P_3 \rightarrow P_4$ given by $I(f(x)) = \int_0^x f(t) dt$

- take a basis $1, x, x^2, x^3$ for P_3

Matrix representation of linear transformation

e.g.: Let $I: P_3 \rightarrow P_4$ given by $I(f(x)) = \int_0^x f(t) dt$

- take a basis $1, x, x^2, x^3$ for P_3
- take a basis $1, x, x^2, x^3, x^4$ for P_4 .

Matrix representation of linear transformation

e.g.: Let $I: P_3 \rightarrow P_4$ given by $I(f(x)) = \int_0^x f(t) dt$

- take a basis $\vec{v}_1, \vec{v}_2, \vec{v}_3, \vec{v}_4$ for P_3
 $1, x, x^2, x^3$ for P_3
- take a basis $\vec{w}_1, \vec{w}_2, \vec{w}_3, \vec{w}_4, \vec{w}_5$ for P_4 .
 $1, x, x^2, x^3, x^4$ for P_4 .

Matrix representation of linear transformation

e.g.: Let $I: P_3 \rightarrow P_4$ given by $I(f(x)) = \int_0^x f(t) dt$

- take a basis $\vec{v}_1, \vec{v}_2, \vec{v}_3, \vec{v}_4$ for P_3
 $1, x, x^2, x^3$
- take a basis $\vec{w}_1, \vec{w}_2, \vec{w}_3, \vec{w}_4, \vec{w}_5$ for P_4 .
 $1, x, x^2, x^3, x^4$

We can compute:

$$I(1) = 0 \cdot 1 + 1 \cdot x + 0 \cdot x^2 + 0 \cdot x^3 + 0 \cdot x^4$$

Matrix representation of linear transformation

e.g.: Let $I: P_3 \rightarrow P_4$ given by $I(f(x)) = \int_0^x f(t) dt$

- take a basis $\vec{v}_1, \vec{v}_2, \vec{v}_3, \vec{v}_4$ for P_3
 $1, x, x^2, x^3$
- take a basis $\vec{w}_1, \vec{w}_2, \vec{w}_3, \vec{w}_4, \vec{w}_5$ for P_4 .
 $1, x, x^2, x^3, x^4$

We can compute:

$$I(1) = 0 \cdot 1 + 1 \cdot x + 0 \cdot x^2 + 0 \cdot x^3 + 0 \cdot x^4 = [w_1, \dots, w_5] \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

Matrix representation of linear transformation

e.g.: Let $I: P_3 \rightarrow P_4$ given by $I(f(x)) = \int_0^x f(t) dt$

- take a basis $\vec{v}_1, \vec{v}_2, \vec{v}_3, \vec{v}_4$ for P_3

- take a basis $1, x, x^2, x^3, x^4$ for P_4 .

We can compute:

$$I(1) = 0 \cdot 1 + 1 \cdot x + 0 \cdot x^2 + 0 \cdot x^3 + 0 \cdot x^4 = [w_1, \dots, w_5] \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

$$I(x) = \frac{x^2}{2} = [w_1, \dots, w_5] \begin{bmatrix} 0 \\ 0 \\ \frac{1}{2} \\ 0 \\ 0 \end{bmatrix}$$

Matrix representation of linear transformation

e.g.: Let $I: P_3 \rightarrow P_4$ given by $I(f(x)) = \int_0^x f(t) dt$

- take a basis $1, x, x^2, x^3$ for P_3

$$\vec{v}_1 \vec{v}_2 \vec{v}_3 \vec{v}_4$$

- take a basis $1, x, x^2, x^3, x^4$ for P_4 .

$$\vec{w}_1 \vec{w}_2 \vec{w}_3 \vec{w}_4 \vec{w}_5$$

We can compute:

$$I(1) = 0 \cdot 1 + 1 \cdot x + 0 \cdot x^2 + 0 \cdot x^3 + 0 \cdot x^4 = [w_1, \dots, w_5] \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

$$I(x) = \frac{x^2}{2} = [w_1, \dots, w_5] \begin{bmatrix} 0 \\ 0 \\ \frac{1}{2} \\ 0 \\ 0 \end{bmatrix}$$

$$I(x^2) = \frac{x^3}{3} = [w_1, \dots, w_5] \begin{bmatrix} 0 \\ 0 \\ 0 \\ \frac{1}{3} \\ 0 \end{bmatrix}$$

Matrix representation of linear transformation

e.g.: Let $I: P_3 \rightarrow P_4$ given by $I(f(x)) = \int_0^x f(t) dt$

- take a basis $1, x, x^2, x^3$ for P_3

$$\vec{v}_1 \vec{v}_2 \vec{v}_3 \vec{v}_4$$

- take a basis $1, x, x^2, x^3, x^4$ for P_4 .

$$\vec{w}_1 \vec{w}_2 \vec{w}_3 \vec{w}_4 \vec{w}_5$$

We can compute:

$$I(1) = 0 \cdot 1 + 1 \cdot x + 0 \cdot x^2 + 0 \cdot x^3 + 0 \cdot x^4 = [w_1, \dots, w_5] \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

$$I(x) = \frac{x^2}{2} = [w_1, \dots, w_5] \begin{bmatrix} 0 \\ 0 \\ \frac{1}{2} \\ 0 \\ 0 \end{bmatrix}$$

$$I(x^2) = \frac{x^3}{3} = [w_1, \dots, w_5] \begin{bmatrix} 0 \\ 0 \\ 0 \\ \frac{1}{3} \\ 0 \end{bmatrix}$$

$$I(x^3) = \frac{x^4}{4} = [w_1, \dots, w_5] \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ \frac{1}{4} \end{bmatrix}$$

Matrix representation of linear transformation

e.g.: Let $I: P_3 \rightarrow P_4$ given by $I(f(x)) = \int_0^x f(t) dt$

- take a basis $1, x, x^2, x^3$ for P_3

$$\vec{v}_1 \vec{v}_2 \vec{v}_3 \vec{v}_4$$

- take a basis $1, x, x^2, x^3, x^4$ for P_4 .

$$\vec{w}_1 \vec{w}_2 \vec{w}_3 \vec{w}_4 \vec{w}_5$$

We can compute:

$$I(1) = 0 \cdot 1 + 1 \cdot x + 0 \cdot x^2 + 0 \cdot x^3 + 0 \cdot x^4 = [w_1, \dots, w_5] \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

$$I(x) = \frac{x^2}{2} = [w_1, \dots, w_5] \begin{bmatrix} 0 \\ 0 \\ \frac{1}{2} \\ 0 \\ 0 \end{bmatrix}$$

$$I(x^2) = \frac{x^3}{3} = [w_1, \dots, w_5] \begin{bmatrix} 0 \\ 0 \\ 0 \\ \frac{1}{3} \\ 0 \end{bmatrix}$$

$$I(x^3) = \frac{x^4}{4} = [w_1, \dots, w_5] \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ \frac{1}{4} \end{bmatrix}$$

$$[I] = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & \frac{1}{2} & 0 & 0 \\ 0 & 0 & \frac{1}{3} & 0 \\ 0 & 0 & 0 & \frac{1}{4} \end{bmatrix}$$

Matrix representation of linear transformation

Matrix representation of linear transformation

- We can do this for $L: \mathbb{R}^2 \rightarrow \mathbb{R}^3$

$$L\begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} x_2 \\ x_1 + x_2 \\ x_1 - x_2 \end{pmatrix} \quad \text{it is linear!}$$

Matrix representation of linear transformation

- We can do this for $L: \mathbb{R}^2 \rightarrow \mathbb{R}^3$

$$L\begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} x_2 \\ x_1 + x_2 \\ x_1 - x_2 \end{pmatrix} \quad \text{it is linear!}$$

We take basis $\vec{v}_1 = \begin{bmatrix} 1 \\ 2 \end{bmatrix}, \vec{v}_2 = \begin{bmatrix} 3 \\ 1 \end{bmatrix}; \vec{w}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \vec{w}_2 = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \vec{w}_3 = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$

Matrix representation of linear transformation

- We can do this for $L: \mathbb{R}^2 \rightarrow \mathbb{R}^3$

$$L\begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} x_2 \\ x_1 + x_2 \\ x_1 - x_2 \end{pmatrix} \quad \text{it is linear!}$$

We take basis $\vec{v}_1 = \begin{bmatrix} 1 \\ 2 \end{bmatrix}, \vec{v}_2 = \begin{bmatrix} 3 \\ 1 \end{bmatrix}; \vec{w}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \vec{w}_2 = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \vec{w}_3 = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$

- We can find the matrix representation w.r.t these basis.

Matrix representation of linear transformation

- We can do this for $L: \mathbb{R}^2 \rightarrow \mathbb{R}^3$

$$L\begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} x_2 \\ x_1 + x_2 \\ x_1 - x_2 \end{pmatrix} \quad \text{it is linear!}$$

We take basis $\vec{v}_1 = \begin{bmatrix} 1 \\ 2 \end{bmatrix}, \vec{v}_2 = \begin{bmatrix} 3 \\ 1 \end{bmatrix}; \vec{w}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \vec{w}_2 = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \vec{w}_3 = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$

- We can find the matrix representation w.r.t these basis.

$$L(v_1) = \begin{bmatrix} 2 \\ 3 \\ -1 \end{bmatrix} = -\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + 4\begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} - \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

Matrix representation of linear transformation

- We can do this for $L: \mathbb{R}^2 \rightarrow \mathbb{R}^3$

$$L\begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} x_2 \\ x_1+x_2 \\ x_1-x_2 \end{pmatrix} \quad \text{it is linear!}$$

We take basis $\vec{v}_1 = \begin{bmatrix} 1 \\ 2 \end{bmatrix}, \vec{v}_2 = \begin{bmatrix} 3 \\ 1 \end{bmatrix}; \vec{w}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \vec{w}_2 = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \vec{w}_3 = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$

- We can find the matrix representation w.r.t these basis.

$$L(v_1) = \begin{bmatrix} 2 \\ 3 \\ -1 \end{bmatrix} = -\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + 4\begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} - \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = [\vec{w}_1, \vec{w}_2, \vec{w}_3] \begin{bmatrix} -1 \\ 4 \\ -1 \end{bmatrix}$$

Matrix representation of linear transformation

- We can do this for $L: \mathbb{R}^2 \rightarrow \mathbb{R}^3$

$$L\begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} x_2 \\ x_1+x_2 \\ x_1-x_2 \end{pmatrix} \quad \text{it is linear!}$$

We take basis $\vec{v}_1 = \begin{bmatrix} 1 \\ 2 \end{bmatrix}, \vec{v}_2 = \begin{bmatrix} 3 \\ 1 \end{bmatrix}; \vec{w}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \vec{w}_2 = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \vec{w}_3 = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$

- We can find the matrix representation w.r.t these basis.

$$L(v_1) = \begin{bmatrix} 2 \\ 3 \\ -1 \end{bmatrix} = -\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + 4\begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} - \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = [\vec{w}_1, \vec{w}_2, \vec{w}_3] \begin{bmatrix} -1 \\ 4 \\ -1 \end{bmatrix}$$

$$L(v_2) = \begin{bmatrix} 1 \\ 4 \\ 2 \end{bmatrix} = -3\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + 2\begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} + 2\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

Matrix representation of linear transformation

- We can do this for $L: \mathbb{R}^2 \rightarrow \mathbb{R}^3$

$$L\begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} x_2 \\ x_1+x_2 \\ x_1-x_2 \end{pmatrix} \quad \text{it is linear!}$$

We take basis $\vec{v}_1 = \begin{bmatrix} 1 \\ 2 \end{bmatrix}, \vec{v}_2 = \begin{bmatrix} 3 \\ 1 \end{bmatrix}; \vec{w}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \vec{w}_2 = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \vec{w}_3 = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$

- We can find the matrix representation w.r.t these basis.

$$L(v_1) = \begin{bmatrix} 2 \\ 3 \\ -1 \end{bmatrix} = -\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + 4\begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} - \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = [\vec{w}_1, \vec{w}_2, \vec{w}_3] \begin{bmatrix} -1 \\ 4 \\ -1 \end{bmatrix}$$

$$L(v_2) = \begin{bmatrix} 1 \\ 4 \\ 2 \end{bmatrix} = -3\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + 2\begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} + 2\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = [\vec{w}_1, \vec{w}_2, \vec{w}_3] \begin{bmatrix} -3 \\ 2 \\ 2 \end{bmatrix}$$

Matrix representation of linear transformation

Matrix representation of linear transformation

- Therefore, the matrix representation of the L is given by

$$[L] = \begin{bmatrix} -1 & -3 \\ 4 & 2 \\ -1 & 2 \end{bmatrix}$$

Matrix representation of linear transformation

- Therefore, the matrix representation of the L is given by

$$[L] = \begin{bmatrix} -1 & -3 \\ 4 & 2 \\ -1 & 2 \end{bmatrix}$$

- Matrix representation relation:

$$[L(\vec{v}_1), L(\vec{v}_2)] = [\vec{w}_1, \vec{w}_2, \vec{w}_3] [L]$$

Matrix representation of linear transformation

- Therefore, the matrix representation of the L is given by

$$[L] = \begin{bmatrix} -1 & -3 \\ 4 & 2 \\ -1 & 2 \end{bmatrix}$$

- Matrix representation relation:

$$[L(\vec{v}_1), L(\vec{v}_2)] = [\vec{w}_1, \vec{w}_2, \vec{w}_3] [L]$$

$$\begin{bmatrix} 2 & 1 \\ 3 & 4 \\ -1 & 2 \end{bmatrix} \underset{\parallel}{=} \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} -1 & -3 \\ 4 & 2 \\ -1 & 2 \end{bmatrix}$$

Matrix representation of linear transformation

- Therefore, the matrix representation of the L is given by

$$[L] = \begin{bmatrix} -1 & -3 \\ 4 & 2 \\ -1 & 2 \end{bmatrix}$$

- Matrix representation relation:

$$[L(\vec{v}_1), L(\vec{v}_2)] = [\vec{w}_1, \vec{w}_2, \vec{w}_3] [L]$$

$$\begin{bmatrix} 2 & 1 \\ 3 & 4 \\ -1 & 2 \end{bmatrix} \underset{\parallel}{=} \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} -1 & -3 \\ 4 & 2 \\ -1 & 2 \end{bmatrix}$$

- In this case, they are honest identity in matrix.

Matrix representation of linear transformation

- Therefore, the matrix representation of the L is given by

$$[L] = \begin{bmatrix} -1 & -3 \\ 4 & 2 \\ -1 & 2 \end{bmatrix}$$

- Matrix representation relation:

$$[L(\vec{v}_1), L(\vec{v}_2)] = [\vec{w}_1, \vec{w}_2, \vec{w}_3] [L]$$

$$\begin{bmatrix} 2 & 1 \\ 3 & 4 \\ -1 & 2 \end{bmatrix} = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} -1 & -3 \\ 4 & 2 \\ -1 & 2 \end{bmatrix}$$

- In this case, they are honest identity in matrix.

- When we write L_A , we can have $L_A(\vec{x}) = A\vec{x} = [\vec{e}_1, \dots, \vec{e}_m] A \vec{x}$

Matrix representation of linear transformation

- Therefore, the matrix representation of the L is given by

$$[L] = \begin{bmatrix} -1 & -3 \\ 4 & 2 \\ -1 & 2 \end{bmatrix}$$

- Matrix representation relation:

$$[L(\vec{v}_1), L(\vec{v}_2)] = [\vec{w}_1, \vec{w}_2, \vec{w}_3] [L]$$

$$\begin{bmatrix} 2 & 1 \\ 3 & 4 \\ -1 & 2 \end{bmatrix} = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} -1 & -3 \\ 4 & 2 \\ -1 & 2 \end{bmatrix}$$

- In this case, they are honest identity in matrix.

- When we write L_A , we can have $L_A(\vec{x}) = A\vec{x} = [\vec{e}_1, \dots, \vec{e}_m] A \vec{x}$

Matrix representation of Linear transformation

Matrix representation of Linear transformation

- Prop: • Given V and W as above, with $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n$ basis for V . Fix $\vec{l}_1, \dots, \vec{l}_n$ be n vectors in W

Matrix representation of Linear transformation

Prop:

- Given V and W as above, with $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n$ basis for V . Fix $\vec{l}_1, \dots, \vec{l}_n$ be n vectors in W
- There exists unique $L: V \rightarrow W$ linear trans. such that $L(\vec{v}_i) = \vec{l}_i$.

Matrix representation of Linear transformation

Prop:

- Given V and W as above, with $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n$ basis for V . Fix $\vec{l}_1, \dots, \vec{l}_n$ be n vectors in W
- There exists unique $L: V \rightarrow W$ linear trans. such that $L(\vec{v}_i) = \vec{l}_i$.

Pf:

- (say $n=3$): Given $\vec{v} \in V$, we can write

$$\vec{v} = a_1 \vec{v}_1 + a_2 \vec{v}_2 + a_3 \vec{v}_3$$

Matrix representation of Linear transformation

Prop:

- Given V and W as above, with $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n$ basis for V . Fix $\vec{l}_1, \dots, \vec{l}_n$ be n vectors in W
- There exists unique $L: V \rightarrow W$ linear trans. such that $L(\vec{v}_i) = \vec{l}_i$.

Pf:

- (say $n=3$): Given $\vec{v} \in V$, we can write $\vec{v} = a_1\vec{v}_1 + a_2\vec{v}_2 + a_3\vec{v}_3$ uniquely (because of basis).

Matrix representation of Linear transformation

Prop:

- Given V and W as above, with $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n$ basis for V . Fix $\vec{l}_1, \dots, \vec{l}_n$ be n vectors in W
- There exists unique $L: V \rightarrow W$ linear trans. such that $L(\vec{v}_i) = \vec{l}_i$.

Pf:

- (say $n=3$): Given $\vec{v} \in V$, we can write $\vec{v} = a_1\vec{v}_1 + a_2\vec{v}_2 + a_3\vec{v}_3$ uniquely (because of basis).
- Define $L(\vec{v}) = a_1\vec{l}_1 + a_2\vec{l}_2 + a_3\vec{l}_3$

Matrix representation of Linear transformation

Prop:

- Given V and W as above, with $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n$ basis for V . Fix $\vec{l}_1, \dots, \vec{l}_n$ be n vectors in W
- There exists unique $L: V \rightarrow W$ linear trans. such that $L(\vec{v}_i) = \vec{l}_i$.

Pf:

- (say $n=3$): Given $\vec{v} \in V$, we can write $\vec{v} = a_1\vec{v}_1 + a_2\vec{v}_2 + a_3\vec{v}_3$ uniquely (because of basis).
- Define $L(\vec{v}) = a_1\vec{l}_1 + a_2\vec{l}_2 + a_3\vec{l}_3$. (well-defined!)

Matrix representation of Linear transformation

Prop:

- Given V and W as above, with $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n$ basis for V . Fix $\vec{l}_1, \dots, \vec{l}_n$ be n vectors in W
- There exists unique $L: V \rightarrow W$ linear trans. such that $L(\vec{v}_i) = \vec{l}_i$.

Pf:

- (say $n=3$): Given $\vec{v} \in V$, we can write $\vec{v} = a_1\vec{v}_1 + a_2\vec{v}_2 + a_3\vec{v}_3$ uniquely (because of basis).
- Define $L(\vec{v}) = a_1\vec{l}_1 + a_2\vec{l}_2 + a_3\vec{l}_3$. (well-defined!)

We check for linearity.

Matrix representation of linear transformation

Matrix representation of linear transformation

Ex : Let $M_{2 \times 2}(\mathbb{R})$ be the space of 2×2 matrices :

Matrix representation of linear transformation

Ex : Let $M_{2 \times 2}(\mathbb{R})$ be the space of 2×2 matrices :

Let $A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$ and we let $L : M_{2 \times 2}(\mathbb{R}) \rightarrow M_{2 \times 2}(\mathbb{R})$
given by $L(B) = B \cdot A$

Matrix representation of linear transformation

- Let $M_{2 \times 2}(\mathbb{R})$ be the space of 2×2 matrices :
- Let $A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$ and we let $L : M_{2 \times 2}(\mathbb{R}) \rightarrow M_{2 \times 2}(\mathbb{R})$
given by $L(B) = B \cdot A$
- It is a linear transformation.

Matrix representation of linear transformation

- Let $M_{2 \times 2}(\mathbb{R})$ be the space of 2×2 matrices :
- Let $A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$ and we let $L : M_{2 \times 2}(\mathbb{R}) \rightarrow M_{2 \times 2}(\mathbb{R})$
given by $L(B) = B \cdot A$
- It is a linear transformation.

$$L(c_1B_1 + c_2B_2) = (c_1B_1 + c_2B_2) \cdot A$$

Matrix representation of linear transformation

- Let $M_{2 \times 2}(\mathbb{R})$ be the space of 2×2 matrices :
- Let $A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$ and we let $L : M_{2 \times 2}(\mathbb{R}) \rightarrow M_{2 \times 2}(\mathbb{R})$ given by $L(B) = B \cdot A$
- It is a linear transformation.

$$\begin{aligned} L(c_1B_1 + c_2B_2) &= (c_1B_1 + c_2B_2) \cdot A \\ &= c_1B_1 \cdot A + c_2B_2 \cdot A \\ &= c_1(B_1 \cdot A) + c_2(B_2 \cdot A) \end{aligned}$$

Matrix representation of linear transformation

• Let $M_{2 \times 2}(\mathbb{R})$ be the space of 2×2 matrices :

Let $A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$ and we let $L : M_{2 \times 2}(\mathbb{R}) \rightarrow M_{2 \times 2}(\mathbb{R})$ given by $L(B) = B \cdot A$

• It is a linear transformation.

$$\begin{aligned} L(c_1B_1 + c_2B_2) &= (c_1B_1 + c_2B_2) \cdot A \\ &= c_1B_1 \cdot A + c_2B_2 \cdot A \\ &= c_1(L(B_1)) + c_2(L(B_2)). \end{aligned}$$

Matrix representation of linear transformation

Matrix representation of linear transformation

- We take the basis $\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$ for $M_{2 \times 2}(\mathbb{R})$.

Matrix representation of linear transformation

- We take the basis $\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$ for $M_{2 \times 2}(\mathbb{R})$.

v_1

v_2

v_3

v_4

w_1

w_2

w_3

w_4

Matrix representation of linear transformation

- We take the basis $\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$ for $M_{2 \times 2}(\mathbb{R})$.
 v_1 v_2 v_3 v_4
 w_1 w_2 w_3 w_4

$$L(v_1) = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$$

Matrix representation of linear transformation

- We take the basis $\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$ for $M_{2 \times 2}(\mathbb{R})$.

$$\begin{array}{c} v_1 \\ w_1 \end{array}, \quad \begin{array}{c} v_2 \\ w_2 \end{array}, \quad \begin{array}{c} v_3 \\ w_3 \end{array}, \quad \begin{array}{c} v_4 \\ w_4 \end{array}$$

$$L(v_1) = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} = w_1 + w_2 = [w_1, w_2, w_3, w_4] \begin{bmatrix} 1 \\ 1 \\ 0 \\ 0 \end{bmatrix}$$

Matrix representation of linear transformation

- We take the basis $\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$ for $M_{2 \times 2}(\mathbb{R})$.

$$\begin{array}{c} v_1 \\ w_1 \end{array}, \quad \begin{array}{c} v_2 \\ w_2 \end{array}, \quad \begin{array}{c} v_3 \\ w_3 \end{array}, \quad \begin{array}{c} v_4 \\ w_4 \end{array}$$

$$L(v_1) = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} = w_1 + w_2 = [w_1, w_2, w_3, w_4] \begin{bmatrix} 1 \\ 1 \\ 0 \\ 0 \end{bmatrix}$$

$$L(v_2) = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$$

Matrix representation of linear transformation

- We take the basis $\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$ for $M_{2 \times 2}(\mathbb{R})$.
 $v_1 = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$, $v_2 = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$, $v_3 = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}$, $v_4 = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$
 $w_1 = \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$, $w_2 = \begin{bmatrix} 0 & 1 \\ 0 & 1 \end{bmatrix}$, $w_3 = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}$, $w_4 = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$

$$L(v_1) = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} = w_1 + w_2 = [w_1, w_2, w_3, w_4] \begin{bmatrix} 1 \\ 1 \\ 0 \\ 0 \end{bmatrix}$$

$$L(v_2) = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} = w_2 = [w_1, w_2, w_3, w_4] \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix}$$

Matrix representation of linear transformation

- We take the basis $\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$ for $M_{2 \times 2}(\mathbb{R})$.

$$L(v_1) = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} = w_1 + w_2 = [w_1, w_2, w_3, w_4] \begin{bmatrix} 1 \\ 1 \\ 0 \\ 0 \end{bmatrix}$$

$$L(v_2) = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} = w_2 = [w_1, w_2, w_3, w_4] \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix}$$

$$L(v_3) = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix} = w_3 + w_4 = [w_1, w_2, w_3, w_4] \begin{bmatrix} 0 \\ 0 \\ 1 \\ 1 \end{bmatrix}$$

Matrix representation of linear transformation

- We take the basis $\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$ for $M_{2 \times 2}(\mathbb{R})$.
 v_1 v_2 v_3 v_4
 w_1 w_2 w_3 w_4

$$L(v_1) = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} = w_1 + w_2 = [w_1, w_2, w_3, w_4] \begin{bmatrix} 1 \\ 1 \\ 0 \\ 0 \end{bmatrix}$$

$$L(v_2) = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} = w_2 = [w_1, w_2, w_3, w_4] \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix}$$

$$L(v_3) = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix} = w_3 + w_4 = [w_1, w_2, w_3, w_4] \begin{bmatrix} 0 \\ 0 \\ 1 \\ 1 \end{bmatrix}$$

$$L(v_4) = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} = w_4 = [w_1, w_2, w_3, w_4] \begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \end{bmatrix}$$

Matrix representation of linear transformation

- We take the basis $\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$
 for $M_{2 \times 2}(\mathbb{R})$.
 $v_1 \quad v_2 \quad v_3 \quad v_4$
 $w_1 \quad w_2 \quad w_3 \quad w_4$

$$L(v_1) = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} = w_1 + w_2 = [w_1, w_2, w_3, w_4] \begin{bmatrix} 1 \\ 1 \\ 0 \\ 0 \end{bmatrix}$$

$$L(v_2) = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} = w_2 = [w_1, w_2, w_3, w_4] \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix}$$

$$L(v_3) = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix} = w_3 + w_4 = [w_1, w_2, w_3, w_4] \begin{bmatrix} 0 \\ 0 \\ 1 \\ 1 \end{bmatrix}$$

$$L(v_4) = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} = w_4 = [w_1, w_2, w_3, w_4] \begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \end{bmatrix}$$

$$[L] = \begin{bmatrix} 1 & 0 & 0 & 0 \\ -1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \text{ is the matrix representation.}$$

Matrix representation of linear transformation

Matrix representation of linear transformation

- Q: What if we choose a basis in \mathbb{R}^n differ from the standard basis $\vec{e}_1, \dots, \vec{e}_n$?

Matrix representation of linear transformation

• Q: What if we choose a basis in \mathbb{R}^n differ from the standard basis $\vec{e}_1, \dots, \vec{e}_n$?

e.g. We recall the map $L_p: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ with $P = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$

Matrix representation of linear transformation

- Q: What if we choose a basis in \mathbb{R}^n differ from the standard basis $\vec{e}_1, \dots, \vec{e}_n$?

e.g. We recall the map $L_p: \mathbb{R}^2 \rightarrow \mathbb{R}^2$
with $P = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ $\vec{v}_1 = \vec{e}_1, \vec{v}_2 = \vec{e}_2$ $\vec{w}_1 = \vec{e}_2, \vec{w}_2 = \vec{e}_1$

Matrix representation of linear transformation

- Q: What if we choose a basis in \mathbb{R}^n differ from the standard basis $\vec{e}_1, \dots, \vec{e}_n$?

e.g. We recall the map $L_p: \mathbb{R}^2 \rightarrow \mathbb{R}^2$
with $P = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ $\vec{v}_1 = \vec{e}_1, \vec{v}_2 = \vec{e}_2$ $\vec{w}_1 = \vec{e}_2, \vec{w}_2 = \vec{e}_1$

$$L_p(\vec{v}_1) = \vec{e}_2 = \vec{w}_1 = [\vec{w}_1, \vec{w}_2] \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

Matrix representation of linear transformation

- Q: What if we choose a basis in \mathbb{R}^n differ from the standard basis $\vec{e}_1, \dots, \vec{e}_n$?

e.g. We recall the map $L_p: \mathbb{R}^2 \rightarrow \mathbb{R}^2$
with $P = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ $\vec{v}_1 = \vec{e}_1, \vec{v}_2 = \vec{e}_2$ $\vec{w}_1 = \vec{e}_2, \vec{w}_2 = \vec{e}_1$

$$L_p(\vec{v}_1) = \vec{e}_2 = \vec{w}_1 = [\vec{w}_1, \vec{w}_2] \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

$$L_p(\vec{v}_2) = \vec{e}_1 = \vec{w}_2 = [\vec{w}_1, \vec{w}_2] \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

Matrix representation of linear transformation

- Q: What if we choose a basis in \mathbb{R}^n differ from the standard basis $\vec{e}_1, \dots, \vec{e}_n$?

e.g. We recall the map $L_p: \mathbb{R}^2 \rightarrow \mathbb{R}^2$
with $P = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ $\vec{v}_1 = \vec{e}_1, \vec{v}_2 = \vec{e}_2$ $\vec{w}_1 = \vec{e}_2, \vec{w}_2 = \vec{e}_1$

$$L_p(\vec{v}_1) = \vec{e}_2 = \vec{w}_1 = [\vec{w}_1, \vec{w}_2] \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

$$L_p(\vec{v}_2) = \vec{e}_1 = \vec{w}_2 = [\vec{w}_1, \vec{w}_2] \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

Therefore the matrix representation is $[L_p] = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$

Matrix representation of linear transformation

Matrix representation of linear transformation

- What happen here is the change of basis: $w_1 = \vec{e}_2$, $w_2 = \vec{e}_1$

Matrix representation of linear transformation

- What happen here is the change of basis: $w_1 = \vec{e}_2, w_2 = \vec{e}_1$
- Consider $L_I: \mathbb{R}^2 \longrightarrow \mathbb{R}^2$ be the identity map.

Matrix representation of linear transformation

- What happen here is the change of basis: $w_1 = \vec{e}_2, w_2 = \vec{e}_1$
- Consider $L_I : \mathbb{R}^2 \xrightarrow{\quad} \mathbb{R}^2$ be the identity map.
 \vec{w}_1, \vec{w}_2 \vec{e}_1, \vec{e}_2 .

Matrix representation of linear transformation

- What happen here is the change of basis: $w_1 = \vec{e}_2$, $w_2 = \vec{e}_1$
- Consider $L_I : \mathbb{R}^2 \xrightarrow{\quad} \mathbb{R}^2$ be the identity map.
 $\begin{matrix} \vec{w}_1, \vec{w}_2 \\ \vec{e}_1, \vec{e}_2 \end{matrix}$.

$$L_I(w_1) = \vec{e}_2 = [\vec{e}_1, \vec{e}_2] \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \quad L_I(w_2) = \vec{e}_1 = [e_1, e_2] \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

Matrix representation of linear transformation

- What happen here is the change of basis: $w_1 = \vec{e}_2, w_2 = \vec{e}_1$
- Consider $L_I : \mathbb{R}^2 \xrightarrow{\quad} \mathbb{R}^2$ be the identity map.
 $\begin{matrix} \vec{w}_1, \vec{w}_2 \\ \vec{e}_1, \vec{e}_2 \end{matrix}$.

$$L_I(w_1) = \vec{e}_2 = [\vec{e}_1, \vec{e}_2] \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \quad L_I(w_2) = \vec{e}_1 = [e_1, e_2] \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

$$[L_I] = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}.$$

Matrix representation of linear transformation

- What happen here is the change of basis: $w_1 = \vec{e}_2, w_2 = \vec{e}_1$
 - Consider $L_I : \mathbb{R}^2 \xrightarrow{\quad} \mathbb{R}^2$ be the identity map.
 $\begin{matrix} \vec{w}_1, \vec{w}_2 \\ \vec{e}_1, \vec{e}_2 \end{matrix}$.
- $L_I(w_1) = \vec{e}_2 = [\vec{e}_1, \vec{e}_2] \begin{bmatrix} 0 \\ 1 \end{bmatrix}, L_I(w_2) = \vec{e}_1 = [e_1, e_2] \begin{bmatrix} 1 \\ 0 \end{bmatrix}$
- $[L_I] = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}.$
- So we consider the composition:


Matrix representation of linear transformation

- What happen here is the change of basis: $w_1 = \vec{e}_2, w_2 = \vec{e}_1$
- Consider $L_I : \mathbb{R}^2 \xrightarrow{\quad} \mathbb{R}^2$ be the identity map.
 $\begin{matrix} \vec{w}_1, \vec{w}_2 \\ \vec{e}_1, \vec{e}_2 \end{matrix}$.

$$L_I(w_1) = \vec{e}_2 = [\vec{e}_1, \vec{e}_2] \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \quad L_I(w_2) = \vec{e}_1 = [e_1, e_2] \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

$$[L_I] = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}.$$

- So we consider the composition:




Matrix representation of linear transformation

- What happen here is the change of basis: $w_1 = \vec{e}_2, w_2 = \vec{e}_1$
- Consider $L_I : \mathbb{R}^2 \xrightarrow{\quad} \mathbb{R}^2$ be the identity map.
 \vec{w}_1, \vec{w}_2 \vec{e}_1, \vec{e}_2 .

$$L_I(w_1) = \vec{e}_2 = [\vec{e}_1, \vec{e}_2] \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \quad L_I(w_2) = \vec{e}_1 = [e_1, e_2] \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

$$[L_I] = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}.$$

- So we consider the composition:




Matrix representation of linear transformation

- What happen here is the change of basis: $w_1 = \vec{e}_2, w_2 = \vec{e}_1$
- Consider $L_I : \mathbb{R}^2 \xrightarrow{\quad} \mathbb{R}^2$ be the identity map.
 \vec{w}_1, \vec{w}_2 \vec{e}_1, \vec{e}_2 .

$$L_I(w_1) = \vec{e}_2 = [\vec{e}_1, \vec{e}_2] \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \quad L_I(w_2) = \vec{e}_1 = [e_1, e_2] \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

$$[L_I] = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}.$$

- So we consider the composition:




Matrix representation of linear transformation

- What happen here is the change of basis: $w_1 = \vec{e}_2$, $w_2 = \vec{e}_1$
- Consider $L_I : \mathbb{R}^2 \xrightarrow{\quad} \mathbb{R}^2$ be the identity map.
 \vec{w}_1, \vec{w}_2 \vec{e}_1, \vec{e}_2 .

$$L_I(w_1) = \vec{e}_2 = [\vec{e}_1, \vec{e}_2] \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \quad L_I(w_2) = \vec{e}_1 = [e_1, e_2] \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

$$[L_I] = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}.$$

- So we consider the composition:



Matrix representation of linear transformation


- What happen here is the change of basis: $w_1 = \vec{e}_2, w_2 = \vec{e}_1$
- Consider $L_I : \mathbb{R}^2 \xrightarrow{\quad} \mathbb{R}^2$ be the identity map.
 \vec{w}_1, \vec{w}_2 \vec{e}_1, \vec{e}_2 .

$$L_I(w_1) = \vec{e}_2 = [\vec{e}_1, \vec{e}_2] \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \quad L_I(w_2) = \vec{e}_1 = [e_1, e_2] \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

$$[L_I] = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}.$$

- So we consider the composition:

$$L_I \circ L_P = L_P.$$



Matrix representation of linear transformation

- What happen here is the change of basis: $w_1 = \vec{e}_2, w_2 = \vec{e}_1$
- Consider $L_I : \mathbb{R}^2 \xrightarrow{\quad} \mathbb{R}^2$ be the identity map.
 \vec{w}_1, \vec{w}_2 \vec{e}_1, \vec{e}_2 .


$$L_I(w_1) = \vec{e}_2 = [\vec{e}_1, \vec{e}_2] \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \quad L_I(w_2) = \vec{e}_1 = [e_1, e_2] \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

$$[L_I] = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}.$$

- So we consider the composition:

$$L_I \circ L_p = L_p.$$

$$\xrightarrow{\quad} \underbrace{[\vec{e}_i]}_{\vec{w}_i} \underbrace{[L_I]}_{\vec{w}_i} \circ \underbrace{[L_p]}_{\vec{v}_i} \xrightarrow{\quad} \underbrace{[\vec{e}_i]}_{\vec{v}_i} [L_p] \xrightarrow{\quad} \underbrace{\vec{e}_i}_{\vec{v}_i}$$



Matrix representation of linear transformation

Matrix representation of linear transformation

- Let $V \xrightarrow{L} W \xrightarrow{M} Z$ be a composition of linear map.

Matrix representation of linear transformation

- Let $V \xrightarrow{L} W \xrightarrow{M} Z$ be a composition of linear maps.
 $\vec{v}_1, \dots, \vec{v}_n$ $\vec{w}_1, \dots, \vec{w}_m$ $\vec{z}_1, \dots, \vec{z}_l$

Matrix representation of linear transformation

- Let $V \xrightarrow{L} W \xrightarrow{M} Z$ be a composition of linear map.
 $\vec{v}_1, \dots, \vec{v}_n$ $\vec{w}_1, \dots, \vec{w}_m$ $\vec{z}_1, \dots, \vec{z}_l$
- With a corresponding choice of basis. then we have the identity

Matrix representation of linear transformation

- Let $V \xrightarrow{L} W \xrightarrow{M} Z$ be a composition of linear maps.
 $\vec{v}_1, \dots, \vec{v}_n$ $\vec{w}_1, \dots, \vec{w}_m$ $\vec{z}_1, \dots, \vec{z}_l$
- With a corresponding choice of basis. then we have the identity

$$[M] \cdot [L] = [M \circ L]$$

Matrix representation of linear transformation

- Let $V \xrightarrow{L} W \xrightarrow{M} Z$ be a composition of linear maps.
 $\vec{v}_1, \dots, \vec{v}_n$ $\vec{w}_1, \dots, \vec{w}_m$ $\vec{z}_1, \dots, \vec{z}_l$
- With a corresponding choice of basis. then we have the identity

$$\sum_i \vec{z}_i [M]_{\vec{w}_i} \cdot \vec{w}_i [L]_{\vec{v}_i} = \sum_i [\mathbf{M} \circ L]_{\vec{v}_i}$$

Matrix representation of linear transformation

- Let $V \xrightarrow{L} W \xrightarrow{M} Z$ be a composition of linear map.
 $\vec{v}_1, \dots, \vec{v}_n$ $\vec{w}_1, \dots, \vec{w}_m$ $\vec{z}_1, \dots, \vec{z}_l$

- With a corresponding choice of basis. then we have the identity

$$\sum_i \vec{z}_i [M]_{\vec{w}_i} \cdot \vec{w}_i [L]_{\vec{v}_i} = \sum_i [\vec{z}_i [M \circ L]]_{\vec{v}_i}$$

- In wording, after the basis are fixed accordingly the composition of linear map correspond to matrix multiplication.

Matrix representation of linear transformation

Matrix representation of linear transformation

- We may look at the change of basis in this way:

Matrix representation of linear transformation

- We may look at the change of basis in this way:

Say : $L : V \longrightarrow W$.

Matrix representation of linear transformation

- We may look at the change of basis in this way:

Say : $L: V \longrightarrow W.$

$$\begin{matrix} \vec{v}_1, \dots, \vec{v}_n \\ \downarrow \\ \vec{u}_1, \dots, \vec{u}_n \end{matrix} \qquad \begin{matrix} \vec{w}_1, \dots, \vec{w}_n \end{matrix}$$

Matrix representation of linear transformation

- We may look at the change of basis in this way:

Say : $L: V \longrightarrow W$.

$$\begin{array}{ccc} \vec{v}_1, \dots, \vec{v}_n & & \vec{w}_1, \dots, \vec{w}_n \\ \downarrow & & \\ \vec{u}_1, \dots, \vec{u}_n & & \end{array}$$

- Then in particular we have

$$V \xrightarrow{\text{id}} V \xrightarrow{L} W$$

Matrix representation of linear transformation

- We may look at the change of basis in this way:

Say : $L: V \longrightarrow W$.

$$\begin{array}{ccc} \vec{v}_1, \dots, \vec{v}_n & & \vec{w}_1, \dots, \vec{w}_n \\ \downarrow & & \\ \vec{u}_1, \dots, \vec{u}_n & & \end{array}$$

- Then in particular we have

$$\begin{array}{ccccc} V & \xrightarrow{\text{id}} & V & \xrightarrow{L} & W \\ \vec{v}_1, \dots, \vec{v}_n & & \vec{u}_1, \dots, \vec{u}_n & & \vec{w}_1, \dots, \vec{w}_n \end{array}$$

Matrix representation of linear transformation

- We may look at the change of basis in this way:

Say : $L: V \longrightarrow W$.

$$\begin{array}{ccc} \vec{v}_1, \dots, \vec{v}_n & & \vec{w}_1, \dots, \vec{w}_n \\ \downarrow & & \\ \vec{u}_1, \dots, \vec{u}_n & & \end{array}$$

- Then in particular we have

$$\begin{array}{ccccc} V & \xrightarrow{\text{id}} & V & \xrightarrow{L} & W \\ \vec{v}_1, \dots, \vec{v}_n & & \vec{u}_1, \dots, \vec{u}_n & & \vec{w}_1, \dots, \vec{w}_n \end{array}$$

and we get $\vec{w}_i [L]_{\vec{u}_i} \cdot \vec{u}_i [\text{id}]_{\vec{v}_i} = \vec{w}_i [L]_{\vec{v}_i}$.

Matrix representation of linear transformation

- We may look at the change of basis in this way:

Say : $L: V \longrightarrow W$.

$$\begin{array}{ccc} \vec{v}_1, \dots, \vec{v}_n & & \vec{w}_1, \dots, \vec{w}_n \\ \downarrow & & \\ \vec{u}_1, \dots, \vec{u}_n & & \end{array}$$

- Then in particular we have

$$\begin{array}{ccccc} & id & & L & \\ V & \xrightarrow{\quad} & V & \xrightarrow{\quad} & W \\ \vec{v}_1, \dots, \vec{v}_n & & \vec{u}_1, \dots, \vec{u}_n & & \vec{w}_1, \dots, \vec{w}_n \end{array}$$

and we get $\vec{w}_i [L] \vec{u}_i \cdot \vec{u}_i [id] \vec{v}_i = \vec{w}_i [L] \vec{v}_i$.

transition matrix


Matrix representation of linear transformation

Matrix representation of linear transformation

- We may use this relation to write down some linear transformation :

Matrix representation of linear transformation

- We may use this relation to write down some linear transformation :
- Want to find.




the projection
onto the θ -
rotation of e_1 .

Matrix representation of linear transformation

- We may use this relation to write down some linear transformation:

- Want to find.




the projection
onto the θ -
rotation of e_1 .

- take a second basis $\vec{w}_1 = \vec{v}_1 = \begin{bmatrix} \cos \theta \\ \sin \theta \end{bmatrix}$, $\vec{w}_2 = \vec{v}_2 = \begin{bmatrix} -\sin \theta \\ \cos \theta \end{bmatrix}$
and then under this basis the Projection P.

Matrix representation of linear transformation

- We may use this relation to write down some linear transformation:

- Want to find.



the projection
onto the θ -
rotation of e_1 .


- take a second basis $\vec{w}_1 = \vec{v}_1 = \begin{bmatrix} \cos\theta \\ \sin\theta \end{bmatrix}$, $\vec{w}_2 = \vec{v}_2 = \begin{bmatrix} -\sin\theta \\ \cos\theta \end{bmatrix}$
and then under this basis the Projection P.

$$\vec{w}_i [P]_{\vec{v}_i} = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$$

Matrix representation of linear transformation

- We may use this relation to write down some linear transformation:

- Want to find.



- take a second basis $\vec{w}_1 = \vec{v}_1 = \begin{bmatrix} \cos\theta \\ \sin\theta \end{bmatrix}$, $\vec{w}_2 = \vec{v}_2 = \begin{bmatrix} -\sin\theta \\ \cos\theta \end{bmatrix}$
and then under this basis the Projection P.

$$\vec{w}_1 [P]_{\vec{v}_1} = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \Leftrightarrow \begin{array}{l} P \vec{v}_1 = \vec{w}_1 \\ P \vec{v}_2 = 0. \end{array}$$

Matrix representation of linear transformation

Matrix representation of linear transformation

- In our earlier discussion :

$$\begin{array}{ccccc} \mathbb{R}^2 & \xrightarrow{\text{id}} & \mathbb{R}^2 & \xrightarrow{P} & \mathbb{R}^2 \\ \vec{e}_1, \vec{e}_2 & & \vec{v}_1, \vec{v}_2 & & \vec{v}_1, \vec{v}_2 \\ & & & & \\ & & & & \xrightarrow{\text{id}} \mathbb{R}^2 \\ & & & & \vec{e}_1, \vec{e}_2 \end{array}$$

Matrix representation of linear transformation

- In our earlier discussion :

$$\begin{array}{ccccc} \mathbb{R}^2 & \xrightarrow{\text{id}} & \mathbb{R}^2 & \xrightarrow{P} & \mathbb{R}^2 \\ \vec{e}_1, \vec{e}_2 & & \vec{v}_1, \vec{v}_2 & & \vec{v}_1, \vec{v}_2 \\ & & & & \vec{e}_1, \vec{e}_2 \end{array} .$$

- Their matrix representation satisfies.

$$\vec{e}_i [\text{id}] \vec{v}_i \cdot \vec{v}_i [P] \vec{v}_i \cdot \vec{v}_i [\text{id}] \vec{e}_i = \vec{e}_i [P] \vec{e}_i$$

Matrix representation of linear transformation

- In our earlier discussion :

$$\begin{array}{ccccc} \mathbb{R}^2 & \xrightarrow{\text{id}} & \mathbb{R}^2 & \xrightarrow{P} & \mathbb{R}^2 \\ \vec{e}_1, \vec{e}_2 & & \vec{v}_1, \vec{v}_2 & & \vec{v}_1, \vec{v}_2 \\ & & & & \vec{e}_1, \vec{e}_2 \end{array} .$$

- Their matrix representation satisfies.

$$\underbrace{\vec{e}_i [\text{id}] \vec{v}_i}_{\begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}} \cdot \underbrace{\vec{v}_i [P] \vec{v}_i}_{\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}} \cdot \underbrace{\vec{v}_i [\text{id}] \vec{e}_i}_{\vec{v}_i} = \vec{e}_i [P] \vec{e}_i$$

Matrix representation of linear transformation

- In our earlier discussion :

$$\begin{array}{ccccc} \mathbb{R}^2 & \xrightarrow{\text{id}} & \mathbb{R}^2 & \xrightarrow{P} & \mathbb{R}^2 \\ \vec{e}_1, \vec{e}_2 & & \vec{v}_1, \vec{v}_2 & & \vec{v}_1, \vec{v}_2 \\ & & & & \vec{e}_1, \vec{e}_2 \end{array} .$$

- Their matrix representation satisfies.

$$\underbrace{\vec{e}_i [\text{id}] \vec{v}_i}_{\begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}} \cdot \underbrace{\vec{v}_i [P] \vec{v}_i}_{\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}} \cdot \underbrace{\vec{v}_i [\text{id}] \vec{e}_i}_{\vec{v}_i} = \vec{e}_i [P] \vec{e}_i$$

$\stackrel{=}{R}$

Matrix representation of linear transformation

- In our earlier discussion :

$$\begin{array}{ccccc} \mathbb{R}^2 & \xrightarrow{\text{id}} & \mathbb{R}^2 & \xrightarrow{P} & \mathbb{R}^2 \\ \vec{e}_1, \vec{e}_2 & & \vec{v}_1, \vec{v}_2 & & \vec{v}_1, \vec{v}_2 \\ & & & & \vec{e}_1, \vec{e}_2 \end{array} .$$

- Their matrix representation satisfies.

$$\begin{array}{cccc} \vec{e}_i [\text{id}] \vec{v}_i & \cdot \vec{v}_i [P] \vec{v}_i & \cdot \vec{v}_i [\text{id}] \vec{e}_i & = \vec{e}_i [P] \vec{e}_i \\ \underbrace{\quad\quad\quad}_{\begin{bmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{bmatrix}} & \underbrace{\quad\quad\quad}_{\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}} & \underbrace{\quad\quad\quad}_{R^{-1}} & \\ & & R^{-1} = \begin{bmatrix} \cos\theta & \sin\theta \\ -\sin\theta & \cos\theta \end{bmatrix} & \end{array}$$

Matrix representation of linear transformation

- In our earlier discussion :

$$\begin{array}{ccccc} \mathbb{R}^2 & \xrightarrow{\text{id}} & \mathbb{R}^2 & \xrightarrow{P} & \mathbb{R}^2 \\ \vec{e}_1, \vec{e}_2 & & \vec{v}_1, \vec{v}_2 & & \vec{v}_1, \vec{v}_2 \\ & & & & \vec{e}_1, \vec{e}_2 \end{array} .$$

- Their matrix representation satisfies.

$$\begin{array}{c} \vec{e}_i [\text{id}] \vec{v}_i \cdot \vec{v}_i [P] \vec{v}_i \cdot \vec{v}_i [\text{id}] \vec{e}_i = \vec{e}_i [P] \vec{e}_i \\ \underbrace{\quad}_{\begin{bmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{bmatrix}} \quad \underbrace{\quad}_{\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}} \quad \underbrace{\quad}_{R^{-1} = \begin{bmatrix} \cos\theta & \sin\theta \\ -\sin\theta & \cos\theta \end{bmatrix}} \end{array}$$

$$\vec{e}_i [P] \vec{e}_i = \begin{bmatrix} \cos\theta & \sin\theta \\ \sin\theta & \cos\theta \end{bmatrix} \begin{bmatrix} \cos\theta & \sin\theta \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} \cos^2\theta & \cos\theta\sin\theta \\ \cos\theta\sin\theta & \sin^2\theta \end{bmatrix}$$

Matrix representation of linear transformation

Matrix representation of linear transformation

- That is we $R \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} R^{-1} = [P]$ give us the matrix representation of the projection P.

Matrix representation of linear transformation

- That is we $R \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} R^{-1} = [P]$ give us the matrix representation of the projection P.
- $[P]^2 = (R \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} R^{-1})(R \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} R^{-1})$
 $= R \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} R^{-1} = [P]$.

Matrix representation of linear transformation

- That is we $R \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} R^{-1} = [P]$ give us the matrix representation of the projection P .
- $[P]^2 = (R \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} R^{-1})(R \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} R^{-1})$
 $= R \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} R^{-1} = [P]$.
- Indeed we can read it in the other way

$$R^{-1}[R]R = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}.$$

Matrix representation of linear transformation

- That is we $R \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} R^{-1} = [P]$ give us the matrix representation of the projection P .
- $[P]^2 = (R \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} R^{-1})(R \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} R^{-1})$
 $= R \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} R^{-1} = [P]$.
- Indeed we can read it in the other way


$$R^{-1} [R] R = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}.$$

this is a
diagonalisation of
[P], later chapter.

Matrix representation of linear transformation


Matrix representation of linear transformation

- Suppose we want to write down reflection along the line



Matrix representation of linear transformation

- Suppose we want to write down reflection along the line




- We take a basis.

$$\vec{v}_1 = \begin{bmatrix} \cos\theta \\ \sin\theta \end{bmatrix}, \quad \vec{v}_2 = \begin{bmatrix} -\sin\theta \\ \cos\theta \end{bmatrix}$$

Matrix representation of linear transformation

- Suppose we want to write down reflection along the line



- We take a basis.


$$\vec{v}_1 = \begin{bmatrix} \cos\theta \\ \sin\theta \end{bmatrix}, \quad \vec{v}_2 = \begin{bmatrix} -\sin\theta \\ \cos\theta \end{bmatrix}$$

-

$$\vec{e}_i [id]_{\vec{v}_i} \cdot \vec{v}_i [H]_{\vec{v}_i} \cdot [id]_{\vec{e}_i} = \vec{e}_i [H]_{\vec{e}_i}$$

Matrix representation of linear transformation

- Suppose we want to write down reflection along the line



- We take a basis.

$$\vec{v}_1 = \begin{bmatrix} \cos\theta \\ \sin\theta \end{bmatrix}, \vec{v}_2 = \begin{bmatrix} -\sin\theta \\ \cos\theta \end{bmatrix}$$


-

$$\vec{e}_i \xrightarrow{\parallel} [\text{id}] \vec{v}_i \cdot \vec{v}_i^\top [\text{H}] \vec{v}_i \cdot \vec{v}_i^\top [\text{id}] \vec{e}_i \xrightarrow{\parallel} \vec{e}_i \xrightarrow{\parallel} [\text{H}] \vec{e}_i$$

$$\begin{bmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} \cos\theta & \sin\theta \\ -\sin\theta & \cos\theta \end{bmatrix}$$

Matrix representation of linear transformation

- Suppose we want to write down reflection along the line



- We take a basis.

$$\vec{v}_1 = \begin{bmatrix} \cos\theta \\ \sin\theta \end{bmatrix}, \vec{v}_2 = \begin{bmatrix} -\sin\theta \\ \cos\theta \end{bmatrix}$$


- $\vec{e}_i [\text{id}]_{\vec{v}_i} \cdot \vec{v}_i [H]_{\vec{v}_i} \cdot [\text{id}]_{\vec{e}_i} = \vec{e}_i [H]_{\vec{e}_i}$

$$\begin{bmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} \cos\theta & \sin\theta \\ -\sin\theta & \cos\theta \end{bmatrix}$$

- $\vec{e}_i [F]_{\vec{e}_i} = \begin{bmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{bmatrix} \begin{bmatrix} \cos\theta & \sin\theta \\ -\sin\theta & \cos\theta \end{bmatrix}$

Matrix representation of linear transformation

- Suppose we want to write down reflection along the line



- We take a basis.

$$\vec{v}_1 = \begin{bmatrix} \cos\theta \\ \sin\theta \end{bmatrix}, \vec{v}_2 = \begin{bmatrix} -\sin\theta \\ \cos\theta \end{bmatrix}$$

- $\vec{e}_i [\text{id}]_{\vec{v}_i} \cdot \vec{v}_i [H]_{\vec{v}_i} \cdot [\text{id}]_{\vec{e}_i} = \vec{e}_i [H]_{\vec{e}_i}$

$$\begin{matrix} \parallel & \parallel & \parallel \\ \begin{bmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{bmatrix} & \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} & \begin{bmatrix} \cos\theta & \sin\theta \\ -\sin\theta & \cos\theta \end{bmatrix} \end{matrix}$$

- $\vec{e}_i [F]_{\vec{e}_i} = \begin{bmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{bmatrix} \begin{bmatrix} \cos\theta & \sin\theta \\ -\sin\theta & \cos\theta \end{bmatrix} = \begin{bmatrix} \cos^2\theta - \sin^2\theta & 2\cos\theta\sin\theta \\ 2\cos\theta\sin\theta & \sin^2\theta - \cos^2\theta \end{bmatrix}$

Image and Kernel of linear transformation

Image and Kernel of linear transformation

Def:

Given a linear transformation $L: V \rightarrow W$

Image and Kernel of linear transformation

Def:

Given a linear transformation $L: V \rightarrow W$

We define 1. $\text{Ker}(L) = \left\{ \vec{v} \mid \vec{v} \in V, L(\vec{v}) = \vec{0} \right\}$.

Image and Kernel of linear transformation

Def:

Given a linear transformation $L: V \rightarrow W$

We define 1. $\text{Ker}(L) = \left\{ \vec{v} \mid \vec{v} \in V, L(\vec{v}) = \vec{0} \right\}$.

2. $\text{Im}(L) = \left\{ \vec{w} \mid \vec{w} \in W, \text{ there exist } \vec{v} \in V \text{ s.t. } L(\vec{v}) = \vec{w} \right\}$

Image and Kernel of linear transformation

Def:

Given a linear transformation $L: V \rightarrow W$

We define 1. $\text{Ker}(L) = \left\{ \vec{v} \mid \vec{v} \in V, L(\vec{v}) = \vec{0} \right\}$.

2. $\text{Im}(L) = \left\{ \vec{w} \mid \vec{w} \in W, \text{ there exist } \vec{v} \in V \text{ s.t. } L(\vec{v}) = \vec{w} \right\}$

e.g.

$L = L_A : \mathbb{R}^n \longrightarrow \mathbb{R}^m$.

Image and Kernel of linear transformation

Def:

Given a linear transformation $L: V \rightarrow W$

We define 1. $\text{Ker}(L) = \left\{ \vec{v} \mid \vec{v} \in V, L(\vec{v}) = \vec{0} \right\}$.

2. $\text{Im}(L) = \left\{ \vec{w} \mid \vec{w} \in W, \text{ there exist } \vec{v} \in V \right. \\ \left. \text{s.t. } L(\vec{v}) = \vec{w} \right\}$

e.g.

$L = L_A : \mathbb{R}^n \longrightarrow \mathbb{R}^m$.

then $\text{Ker}(L_A) = \left\{ \vec{v} \in \mathbb{R}^n \mid L_A(\vec{v}) = \vec{0} \right\} = N(A)$.

Image and Kernel of linear transformation

Def:

Given a linear transformation $L: V \rightarrow W$

We define 1. $\text{Ker}(L) = \left\{ \vec{v} \mid \vec{v} \in V, L(\vec{v}) = \vec{0} \right\}$.

2. $\text{Im}(L) = \left\{ \vec{w} \mid \vec{w} \in W, \text{ there exist } \vec{v} \in V \text{ s.t. } L(\vec{v}) = \vec{w} \right\}$

e.g.

$$L = L_A : \mathbb{R}^n \longrightarrow \mathbb{R}^m.$$

then $\text{Ker}(L_A) = \left\{ \vec{v} \in \mathbb{R}^n \mid L_A(\vec{v}) = \vec{0} \right\} = N(A)$.

$\text{Im}(L_A) = \left\{ \vec{w} \in \mathbb{R}^m \mid \vec{w} = L_A(\vec{v}) \text{ for some } \vec{v} \in \mathbb{R}^n \right\}$

Image and Kernel of linear transformation

Def:

Given a linear transformation $L: V \rightarrow W$

We define 1. $\text{Ker}(L) = \left\{ \vec{v} \mid \vec{v} \in V, L(\vec{v}) = \vec{0} \right\}$.

2. $\text{Im}(L) = \left\{ \vec{w} \mid \vec{w} \in W, \text{ there exist } \vec{v} \in V \text{ s.t. } L(\vec{v}) = \vec{w} \right\}$

e.g.

$$L = L_A : \mathbb{R}^n \longrightarrow \mathbb{R}^m.$$

then $\text{Ker}(L_A) = \left\{ \vec{v} \in \mathbb{R}^n \mid L_A(\vec{v}) = \vec{0} \right\} = N(A)$.

$\text{Im}(L_A) = \left\{ \vec{w} \in \mathbb{R}^m \mid \vec{w} = L_A(\vec{v}) \text{ for some } \vec{v} \in \mathbb{R}^n \right\}$

$= C(A)$.

Image and Kernel of linear transformation

Image and Kernel of linear transformation

- Now with $L: V \longrightarrow W$ equipped with basis $\vec{v}_1, \dots, \vec{v}_n$ for V , $\vec{w}_1, \dots, \vec{w}_m$ for W .

Image and Kernel of linear transformation

- Now with $L: V \rightarrow W$ equipped with basis $\vec{v}_1, \dots, \vec{v}_n$ for V , $\vec{w}_1, \dots, \vec{w}_m$ for W .
- Matrix representation: (say $m=2, n=3$).
write $\vec{v} = c_1\vec{v}_1 + c_2\vec{v}_2 + c_3\vec{v}_3$ then

Image and Kernel of linear transformation

- Now with $L: V \rightarrow W$ equipped with basis $\vec{v}_1, \dots, \vec{v}_n$ for V , $\vec{w}_1, \dots, \vec{w}_m$ for W .
- Matrix representation: (say $m=2, n=3$).
write $\vec{v} = c_1\vec{v}_1 + c_2\vec{v}_2 + c_3\vec{v}_3$ then

$$L(\vec{v}) = [\vec{w}_1, \vec{w}_2] [L] \begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix}$$

Image and Kernel of linear transformation

- Now with $L: V \rightarrow W$ equipped with basis $\vec{v}_1, \dots, \vec{v}_n$ for V , $\vec{w}_1, \dots, \vec{w}_m$ for W .
- Matrix representation : (say $m=2, n=3$).
write $\vec{v} = c_1\vec{v}_1 + c_2\vec{v}_2 + c_3\vec{v}_3$ then

$$L(\vec{v}) = [\vec{w}_1, \vec{w}_2] [L] \begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix}$$


 2×3 matrix.

Image and Kernel of linear transformation

- Now with $L: V \rightarrow W$ equipped with basis $\vec{v}_1, \dots, \vec{v}_n$ for V , $\vec{w}_1, \dots, \vec{w}_m$ for W .

- Matrix representation : (say $m=2, n=3$).

write $\vec{v} = c_1\vec{v}_1 + c_2\vec{v}_2 + c_3\vec{v}_3$ then

$$L(\vec{v}) = [\vec{w}_1, \vec{w}_2] [L] \begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix}$$



2x3 matrix.

- $N([L]) \leftrightarrow \text{Ker}(L)$ given by the 1-1 correspondence.

$$\begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix} \longrightarrow [\vec{v}_1, \vec{v}_2, \vec{v}_3] \begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix}$$

Image and Kernel of linear transformation

Image and Kernel of linear transformation

- $C([L]) \leftrightarrow \text{Im}(L)$ is a 1-1 correspondence.

$$\begin{bmatrix} a_1 \\ a_2 \end{bmatrix} \longrightarrow [w_1, w_2] \begin{bmatrix} a_1 \\ a_2 \end{bmatrix}.$$

Image and Kernel of linear transformation

- $C([L]) \leftrightarrow \text{Im}(L)$ is a 1-1 correspondence.

$$\begin{bmatrix} a_1 \\ a_2 \end{bmatrix} \rightarrow [w_1, w_2] \begin{bmatrix} a_1 \\ a_2 \end{bmatrix}.$$

- $\text{Ker}(L) \subseteq V$ is a subspace
 $\text{Im}(L) \subseteq W$ is a subspace.

Image and Kernel of linear transformation

- $C([L]) \leftrightarrow \text{Im}(L)$ is a 1-1 correspondence.

$$\begin{bmatrix} a_1 \\ a_2 \end{bmatrix} \rightarrow [w_1, w_2] \begin{bmatrix} a_1 \\ a_2 \end{bmatrix}.$$

- $\text{Ker}(L) \subseteq V$ is a subspace
 $\text{Im}(L) \subseteq W$ is a subspace.
- $\text{rk}([L]) = \dim(C[L]) = \dim(\text{Im}(L)).$

Image and Kernel of linear transformation

- $C([L]) \leftrightarrow \text{Im}(L)$ is a 1-1 correspondence.

$$\begin{bmatrix} a_1 \\ a_2 \end{bmatrix} \rightarrow [w_1, w_2] \begin{bmatrix} a_1 \\ a_2 \end{bmatrix}.$$

- $\text{Ker}(L) \subseteq V$ is a subspace

$\text{Im}(L) \subseteq W$ is a subspace.

- $\text{rk}([L]) = \dim(C[L]) = \dim(\text{Im}(L)).$

- $\dim(\text{Ker}(L)) + \dim(\text{Im}(L)) = \dim(V).$

Isomorphism

Isomorphism

Def: $f: V \rightarrow W$ linear map is an **isomorphism**
if we can find another $g: W \rightarrow V$ be a linear map
s.t. $g \circ f = \text{id}_V$, $f \circ g = \text{id}_W$.

Isomorphism

Def: $f: V \rightarrow W$ linear map is an **isomorphism**
if we can find another $g: W \rightarrow V$ be a linear map
s.t. $g \circ f = \text{id}_V$, $f \circ g = \text{id}_W$.

e.g. $f = L_A: \mathbb{R}^n \rightarrow \mathbb{R}^n$ for invertible A
is isomorphism.

Isomorphism

Def: $f: V \rightarrow W$ linear map is an **isomorphism**
if we can find another $g: W \rightarrow V$ be a linear map
s.t. $g \circ f = \text{id}_V$, $f \circ g = \text{id}_W$.

e.g. $f = L_A: \mathbb{R}^n \rightarrow \mathbb{R}^n$ for invertible A
is isomorphism.

$$g = L_{A^{-1}}: \mathbb{R}^n \rightarrow \mathbb{R}^n$$

Isomorphism

Def: $f: V \rightarrow W$ linear map is an **isomorphism**
if we can find another $g: W \rightarrow V$ be a linear map
s.t. $g \circ f = \text{id}_V$, $f \circ g = \text{id}_W$.

e.g. $f = L_A: \mathbb{R}^n \rightarrow \mathbb{R}^n$ for invertible A
is isomorphism.

$$g = L_{A^{-1}}: \mathbb{R}^n \rightarrow \mathbb{R}^n$$

then we have $g \circ f(B) = g(A \cdot B) = A^{-1}AB = B$

Isomorphism

Def: $f: V \rightarrow W$ linear map is an **isomorphism**
if we can find another $g: W \rightarrow V$ be a linear map
s.t. $g \circ f = \text{id}_V$, $f \circ g = \text{id}_W$.

e.g. $f = L_A: \mathbb{R}^n \rightarrow \mathbb{R}^n$ for invertible A
is isomorphism.

$$g = L_{A^{-1}}: \mathbb{R}^n \rightarrow \mathbb{R}^n$$

then we have $g \circ f(B) = g(A \cdot B) = A^{-1}AB = B$

$$f \circ g(B) = A(A^{-1}B) = B.$$

Isomorphism

Isomorphism

e.g. • Recall that we have the vector space $W = \mathbb{R}_+$
with $x+y := xy$, $c \cdot x = x^c$

Isomorphism

- e.g. • Recall that we have the vector space $W = \mathbb{R}_+$
with $x+y := xy$, $c \cdot x = x^c$
- In fact it is isomorphic to $(\mathbb{R}, +, \cdot)$

Isomorphism

e.g. • Recall that we have the vector space $W = \mathbb{R}_+$

with $x+y := x \cdot y$, $c \cdot x = x^c$

• In fact it is isomorphic to $(\mathbb{R}, +, \cdot)$

We take: $f: \mathbb{R} \rightarrow \mathbb{R}_+$ with $f(a) = e^a$

Isomorphism

- e.g. • Recall that we have the vector space $W = \mathbb{R}_+$
with $x+y := xy$, $c \cdot x = x^c$
- In fact it is isomorphic to $(\mathbb{R}, +, \cdot)$

We take:

$$f: \mathbb{R} \rightarrow \mathbb{R}_+ \quad \text{with } f(a) = e^a$$

$$g: \mathbb{R}_+ \rightarrow \mathbb{R} \quad \text{with } g(x) = \log(x)$$

Isomorphism

e.g. • Recall that we have the vector space $W = \mathbb{R}_+$

with $x+y := x \cdot y$, $c \cdot x = x^c$

• In fact it is isomorphic to $(\mathbb{R}, +, \cdot)$

We take:

$$f: \mathbb{R} \rightarrow \mathbb{R}_+ \quad \text{with } f(a) = e^a$$
$$g: \mathbb{R}_+ \rightarrow \mathbb{R} \quad \text{with } g(x) = \log(x)$$

claim: f, g are linear and $f \circ g = id_{\mathbb{R}_+}$

$$g \circ f = id_{\mathbb{R}}$$

Isomorphism

Isomorphism

- $L : V \rightarrow W$, with basis $\vec{v}_1, \dots, \vec{v}_n$ for V
 $\vec{w}_1, \dots, \vec{w}_m$ for W .

Isomorphism

- $L : V \rightarrow W$, with basis $\vec{v}_1, \dots, \vec{v}_n$ for V
 $\vec{w}_1, \dots, \vec{w}_m$ for W .

Claim: L is an isomorphism if and only if $[L]$ is invertible.

Isomorphism

- $L : V \rightarrow W$, with basis $\vec{v}_1, \dots, \vec{v}_n$ for V
 $\vec{w}_1, \dots, \vec{w}_m$ for W .

Claim: L is an isomorphism if and only if $[L]$ is invertible.

Pf: \Rightarrow • Assume L is an isomorphism, then we have

$$T: W \rightarrow V \text{ s.t. } T \circ L = \text{id}_V, L \circ T = \text{id}_W.$$

Isomorphism

- $L : V \rightarrow W$, with basis $\vec{v}_1, \dots, \vec{v}_n$ for V
 $\vec{w}_1, \dots, \vec{w}_m$ for W .

Claim: L is an isomorphism if and only if $[L]$ is invertible.

Pf: \Rightarrow • Assume L is an isomorphism, then we have

$$T: W \rightarrow V \text{ s.t. } T \circ L = \text{id}_V, L \circ T = \text{id}_W.$$

- then we take their matrix representation:

$$\vec{v}_i [T]_{\vec{w}_i} \cdot \vec{w}_i [L]_{\vec{v}_i} = \vec{v}_i [\text{id}]_{\vec{v}_i} = I_{n \times n}.$$

Isomorphism

- $L : V \rightarrow W$, with basis $\vec{v}_1, \dots, \vec{v}_n$ for V
 $\vec{w}_1, \dots, \vec{w}_m$ for W .

Claim: L is an isomorphism if and only if $[L]$ is invertible.

Pf: \Rightarrow • Assume L is an isomorphism, then we have

$$T: W \rightarrow V \text{ s.t. } T \circ L = \text{id}_V, L \circ T = \text{id}_W.$$

- then we take their matrix representation:

$$\vec{v}_i [T]_{\vec{w}_i} \cdot \vec{w}_i [L]_{\vec{v}_i} = \vec{v}_i [\text{id}]_{\vec{v}_i} = I_{n \times n}.$$

and $[L] \cdot [T] = \vec{w}_i [\text{id}]_{\vec{w}_i} = I_{m \times m}.$

Isomorphism

- $L: V \rightarrow W$, with basis $\vec{v}_1, \dots, \vec{v}_n$ for V
 $\vec{w}_1, \dots, \vec{w}_m$ for W .

Claim: L is an isomorphism if and only if $[L]$ is invertible.

Pf: \Rightarrow • Assume L is an isomorphism, then we have

$$T: W \rightarrow V \text{ s.t. } T \circ L = \text{id}_V, L \circ T = \text{id}_W.$$

- then we take their matrix representation:

$$\vec{v}_i [T]_{\vec{w}_i} \cdot \vec{w}_i [L]_{\vec{v}_i} = \vec{v}_i [\text{id}]_{\vec{v}_i} = I_{n \times n}.$$

and $[L] \cdot [T] = \vec{w}_i [\text{id}]_{\vec{w}_i} = I_{m \times m}.$

$$\Rightarrow m=n \text{ and } [L]^{-1} = [T]$$

Isomorphism

- $L : V \rightarrow W$, with basis $\vec{v}_1, \dots, \vec{v}_n$ for V
 $\vec{w}_1, \dots, \vec{w}_m$ for W .

Claim: L is an isomorphism if and only if $[L]$ is invertible.

Pf: \Rightarrow • Assume L is an isomorphism, then we have

$$T: W \rightarrow V \text{ s.t. } T \circ L = \text{id}_V, L \circ T = \text{id}_W.$$

• then we take their matrix representation:

$$m = \text{rk}(\underline{I}_{m \times m}) \leq \text{rk}([L]) \leq n \quad \vec{v}_i [\vec{T}]_{\vec{w}_i} \cdot \vec{w}_i [\vec{L}]_{\vec{v}_i} = \vec{v}_i [\text{id}]_{\vec{v}_i} = \underline{I}_{n \times n}.$$

$$n = \text{rk}(\underline{I}_{n \times n}) \leq \text{rk}([L]) \leq m.$$

and

$$[\vec{L}] \cdot [\vec{T}] = \vec{w}_i [\text{id}]_{\vec{w}_i} = \underline{I}_{m \times m}.$$

$$\Rightarrow m = n \quad \text{and} \quad [L]^{-1} = [T]$$

Isomorphism

Isomorphism

Pf (cont'd) : \Leftarrow if $[L]$ is invertible, then $m=n$
and we have $[L]^{-1}$

Isomorphism

Pf (cont'd): \Leftarrow if $[L]$ is invertible, then $m=n$
and we have $[L]^{-1}$

- We define $T: W \longrightarrow V$ for
 $\vec{w} = a_1 \vec{w}_1 + a_2 \vec{w}_2 + \dots + a_n \vec{w}_n$ by

Isomorphism

Pf (cont'd): \Leftarrow if $[L]$ is invertible, then $m=n$
and we have $[L]^{-1}$

- We define $T: W \longrightarrow V$ for

$$\vec{w} = a_1 \vec{w}_1 + a_2 \vec{w}_2 + \dots + a_n \vec{w}_n \quad \text{by}$$

$$T(\vec{w}) := [\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n] [L]^{-1} \begin{bmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{bmatrix}, \text{ i.e. } [T] = [L]^{-1}$$

Isomorphism

Pf (cont'd): \Leftarrow if $[L]$ is invertible, then $m=n$
and we have $[L]^{-1}$

- We define $T: W \longrightarrow V$ for

$$\vec{w} = a_1 \vec{w}_1 + a_2 \vec{w}_2 + \dots + a_n \vec{w}_n \quad \text{by}$$

$$T(\vec{w}) := [\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n] [L]^{-1} \begin{bmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{bmatrix}, \text{ i.e. } [T] = [L]^{-1}$$

- Then $\vec{v}_i [T \circ L]_{\vec{v}_i} = I_{n \times n}$
 $\vec{w}_i [L \circ T]_{\vec{w}_i} = I_{n \times n} \Rightarrow T \text{ is the inverse of } L.$

Summary



Summary

linear transformation $L: V \rightarrow W$
(with a choice of basis). | $m \times n$ matrix . [L]

Summary

linear transformation $L: V \rightarrow W$
(with a choice of basis).

$m \times n$ matrix . [L]

- Composition
- matrix multiplication.

Summary

linear transformation $L: V \rightarrow W$
(with a choice of basis).

$m \times n$ matrix. $[L]$

- Composition
- Isomorphism
- matrix multiplication.
- Invertible .

Summary

linear transformation $L: V \rightarrow W$
(with a choice of basis).

- Composition
- Isomorphism

$m \times n$ matrix. $[L]$

- matrix multiplication.
- Invertible
transition matrix is an example

Summary

linear transformation $L: V \rightarrow W$
(with a choice of basis).

- Composition
- Isomorphism
- $\text{Im}(L)$

$m \times n$ matrix. $[L]$

- matrix multiplication.
- Invertible
transition matrix is an example
- $C([L])$

Summary

linear transformation $L: V \rightarrow W$
(with a choice of basis).

- Composition
- Isomorphism
- $\text{Im}(L)$
- $\text{Ker}(L)$

$m \times n$ matrix . $[L]$

- matrix multiplication.
- Invertible
transition matrix is an example
- $C([L])$
- $N([L])$

Summary

linear transformation $L: V \rightarrow W$
(with a choice of basis).

- Composition
- Isomorphism
- $\text{Im}(L)$
- $\text{Ker}(L)$
- $\dim(\text{Im}(L))$

$m \times n$ matrix . $[L]$

- matrix multiplication.
- Invertible
transition matrix is an example
- $C([L])$
- $N([L])$
- $\text{rk}([L]).$