

# **Lecture 11**

## **Linear transformation**

# Linear transformation

# Linear transformation

- Recall that we can think of a  $m \times n$  matrix  $A$  as an assignment/map:

$$L_A : \mathbb{R}^n \longrightarrow \mathbb{R}^m$$

# Linear transformation

- Recall that we can think of a  $m \times n$  matrix  $A$  as an assignment/map:

$$L_A : \mathbb{R}^n \longrightarrow \mathbb{R}^m$$

Sending a vector  $\vec{x} \in \mathbb{R}^n$  to the vector  $A\vec{x}$  in  $\mathbb{R}^m$ .

# Linear transformation

- Recall that we can think of a  $m \times n$  matrix  $A$  as an assignment/map:

$$L_A : \mathbb{R}^n \longrightarrow \mathbb{R}^m$$

Sending a vector  $\vec{x} \in \mathbb{R}^n$  to the vector  $A\vec{x}$  in  $\mathbb{R}^m$ .

- We write  $L_A(\vec{x}) = A\vec{x}$

# Linear transformation

- Recall that we can think of a  $m \times n$  matrix  $A$  as an assignment/map:

$$L_A : \mathbb{R}^n \longrightarrow \mathbb{R}^m$$

Sending a vector  $\vec{x} \in \mathbb{R}^n$  to the vector  $A\vec{x}$  in  $\mathbb{R}^m$ .

- We write  $L_A(\vec{x}) = A\vec{x}$

e.g.: 1.  $A = cI$      $L_I(\vec{x}) = c\vec{x}$ .

# Linear transformation

- Recall that we can think of a  $m \times n$  matrix  $A$  as an assignment/map:

$$L_A : \mathbb{R}^n \longrightarrow \mathbb{R}^m$$

Sending a vector  $\vec{x} \in \mathbb{R}^n$  to the vector  $A\vec{x}$  in  $\mathbb{R}^m$ .

- We write  $L_A(\vec{x}) = A\vec{x}$

e.g.: 1.  $A = cI$      $L_I(\vec{x}) = c\vec{x}$ .

2.  $A = 0_{m \times n}$ ,     $L_0(\vec{x}) = \vec{0}$

# Linear transformation

- Recall that we can think of a  $m \times n$  matrix  $A$  as an assignment/map:

$$L_A : \mathbb{R}^n \longrightarrow \mathbb{R}^m$$

Sending a vector  $\vec{x} \in \mathbb{R}^n$  to the vector  $A\vec{x}$  in  $\mathbb{R}^m$ .

- We write  $L_A(\vec{x}) = A\vec{x}$

e.g.: 1.  $A = cI$      $L_I(\vec{x}) = c\vec{x}$ .

2.  $A = 0_{m \times n}$ ,  $L_0(\vec{x}) = \vec{0}$

3.  $A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ ,  $L_A(e_1) = e_2$ ,  $L_A(e_2) = e_1$

# Linear transformation


# Linear transformation

- $A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$      $L_A \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{bmatrix} x_2 \\ x_1 \end{bmatrix}$

# Linear transformation

- $A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$      $L_A \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{bmatrix} x_2 \\ x_1 \end{bmatrix}$


Picture:



# Linear transformation

- $A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$      $L_A\begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{bmatrix} x_2 \\ x_1 \end{bmatrix}$

Picture:



# Linear transformation

- $A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$      $L_A\begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{bmatrix} x_2 \\ x_1 \end{bmatrix}$

Picture:




It gives a reflection along the  $45^\circ$  line.

# Linear transformation

- $A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$      $L_A \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{bmatrix} x_2 \\ x_1 \end{bmatrix}$

Picture:



It gives a reflection along the  $45^\circ$  line.

- $A = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$  ,  $L_A(e_1) = \begin{bmatrix} \cos \theta \\ \sin \theta \end{bmatrix}$  ,  $L_A(e_2) = \begin{bmatrix} -\sin \theta \\ \cos \theta \end{bmatrix}$

# Linear transformation


- $A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \quad L_A \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{bmatrix} x_2 \\ x_1 \end{bmatrix}$

Picture:



It gives a reflection along the  $45^\circ$  line.


- $A = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}, L_A(e_1) = \begin{bmatrix} \cos \theta \\ \sin \theta \end{bmatrix}, L_A(e_2) = \begin{bmatrix} -\sin \theta \\ \cos \theta \end{bmatrix}$



# Linear transformation


- $A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \quad L_A \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} x_2 \\ x_1 \end{pmatrix}$

Picture:



It gives a reflection along the  $45^\circ$  line.

- $A = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}, L_A(e_1) = \begin{bmatrix} \cos \theta \\ \sin \theta \end{bmatrix}, L_A(e_2) = \begin{bmatrix} -\sin \theta \\ \cos \theta \end{bmatrix}$




# Linear transformation

# Linear transformation

- $A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, \quad L_A(e_1) = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad L_A(e_2) = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$


# Linear transformation

- $A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$ ,  $L_A(e_1) = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ ,  $L_A(e_2) = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$



# Linear transformation


- $A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, \quad L_A(e_1) = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad L_A(e_2) = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$




- $A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \quad L_A(e_1) = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad L_A(e_2) = \begin{bmatrix} 0 \\ 0 \end{bmatrix}.$

# Linear transformation

- $A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$ ,  $L_A(e_1) = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ ,  $L_A(e_2) = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$



- $A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ ,  $L_A(e_1) = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ ,  $L_A(e_2) = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$ .



is the projection along y-axis.

# Linear transformation

# Linear transformation

- Observe that For  $A = [\vec{a}_1, \vec{a}_2, \dots, \vec{a}_n]$ , by definition

# Linear transformation

- Observe that For  $A = [\vec{a}_1, \vec{a}_2, \dots, \vec{a}_n]$ , by definition
  1.  $L_A(\vec{e}_i) = \vec{a}_i$  is the  $i$ -th column of  $A$ .

# Linear transformation

- Observe that For  $A = [\vec{a}_1, \vec{a}_2, \dots, \vec{a}_n]$ , by definition
  - 1.  $L_A(\vec{e}_i) = \vec{a}_i$  is the  $i$ -th column of  $A$ .
  - 2.  $L_A$  is determined by its value at  $\vec{e}_i$ ; more explicitly

# Linear transformation

- Observe that For  $A = [\vec{a}_1, \vec{a}_2, \dots, \vec{a}_n]$ , by definition
  1.  $L_A(\vec{e}_i) = \vec{a}_i$  is the  $i$ -th column of  $A$ .
  2.  $L_A$  is determined by its value at  $\vec{e}_i$ 's, more explicitly

$$L_A \left( \begin{bmatrix} c_1 \\ c_2 \\ \vdots \\ c_n \end{bmatrix} \right) = A \begin{bmatrix} c_1 \\ \vdots \\ c_n \end{bmatrix}$$

# Linear transformation

- Observe that For  $A = [\vec{a}_1, \vec{a}_2, \dots, \vec{a}_n]$ , by definition
  - $L_A(\vec{e}_i) = \vec{a}_i$  is the  $i$ -th column of  $A$ .
  - $L_A$  is determined by its value at  $\vec{e}_i$ 's, more explicitly

$$L_A \left( \begin{bmatrix} c_1 \\ c_2 \\ \vdots \\ c_n \end{bmatrix} \right) = A \begin{bmatrix} c_1 \\ \vdots \\ c_n \end{bmatrix} = c_1 \vec{a}_1 + c_2 \vec{a}_2 + \dots + c_n \vec{a}_n \\ = c_1 L_A(\vec{e}_1) + c_2 L_A(\vec{e}_2) + \dots + c_n L_A(\vec{e}_n)$$

# Linear transformation

- Observe that For  $A = [\vec{a}_1, \vec{a}_2, \dots, \vec{a}_n]$ , by definition
  - $L_A(\vec{e}_i) = \vec{a}_i$  is the  $i$ -th column of  $A$ .
  - $L_A$  is determined by its value at  $\vec{e}_i$ 's, more explicitly

$$L_A\left(\begin{bmatrix} c_1 \\ c_2 \\ \vdots \\ c_n \end{bmatrix}\right) = A \begin{bmatrix} c_1 \\ \vdots \\ c_n \end{bmatrix} = c_1 \vec{a}_1 + c_2 \vec{a}_2 + \dots + c_n \vec{a}_n$$
$$= c_1 L_A(\vec{e}_1) + c_2 L_A(\vec{e}_2) + \dots + c_n L_A(\vec{e}_n)$$

$\parallel$

$$L_A(c_1 \vec{e}_1 + \dots + c_n \vec{e}_n).$$

# Linear transformation

- Observe that For  $A = [\vec{a}_1, \vec{a}_2, \dots, \vec{a}_n]$ , by definition


1.  $L_A(\vec{e}_i) = \vec{a}_i$  is the  $i$ -th column of  $A$ .

2.  $L_A$  is determined by its value at  $\vec{e}_i$ 's, more explicitly

$$L_A \left( \begin{bmatrix} c_1 \\ c_2 \\ \vdots \\ c_n \end{bmatrix} \right) = A \begin{bmatrix} c_1 \\ \vdots \\ c_n \end{bmatrix} = c_1 \vec{a}_1 + c_2 \vec{a}_2 + \dots + c_n \vec{a}_n \\ = c_1 L_A(\vec{e}_1) + c_2 L_A(\vec{e}_2) + \dots + c_n L_A(\vec{e}_n)$$

||

$$L_A(c_1 \vec{e}_1 + \dots + c_n \vec{e}_n),$$



this characterize a linear transform.


# Linear transformation

- Observe that For  $A = [\vec{a}_1, \vec{a}_2, \dots, \vec{a}_n]$ , by definition
  - 1.  $L_A(\vec{e}_i) = \vec{a}_i$  is the  $i$ -th column of  $A$ .
  - 2.  $L_A$  is determined by its value at  $\vec{e}_i$ 's, more explicitly

$$L_A \left( \begin{bmatrix} c_1 \\ c_2 \\ \vdots \\ c_n \end{bmatrix} \right) = A \begin{bmatrix} c_1 \\ \vdots \\ c_n \end{bmatrix} = c_1 \vec{a}_1 + c_2 \vec{a}_2 + \dots + c_n \vec{a}_n$$
$$= c_1 L_A(\vec{e}_1) + c_2 L_A(\vec{e}_2) + \dots + c_n L_A(\vec{e}_n)$$

||

$$L_A(c_1 \vec{e}_1 + \dots + c_n \vec{e}_n).$$



this characterize a linear transform.

- 3.  $L_A(\vec{c}) \in C(A)$  for any  $\vec{c} \in \mathbb{R}^n$ .

$C(A)$  is also called the image of  $L_A$

# Linear transformation

# Linear transformation

Def: • A map  $f$  from a set  $S_1$  to another set  $S_2$  is an assignment of element  $x$  in  $S_1$  to an element  $f(x)$  in  $S_2$ .

# Linear transformation

- Def: • A **map**  $f$  from a set  $S_1$  to another set  $S_2$  is an assignment of element  $x$  in  $S_1$  to **an element**  $f(x)$  in  $S_2$ .
- We write  $f: S_1 \rightarrow S_2$  for a map.

# Linear transformation

Def: • A **map**  $f$  from a set  $S_1$  to another set  $S_2$  is an assignment of element  $x$  in  $S_1$  to **an element**  $f(x)$  in  $S_2$ .

• We write  $f: S_1 \rightarrow S_2$  for a map.

e.g. • Let  $S_1$  be the set of students in Linear algebra A-8 class.  $S_2 = \{0, 1, \dots, 50\}$ .

# Linear transformation

Def:

- A **map**  $f$  from a set  $S_1$  to another set  $S_2$  is an assignment of element  $x$  in  $S_1$  to **an element**  $f(x)$  in  $S_2$ .
- We write  $f: S_1 \rightarrow S_2$  for a map.

e.g.

- Let  $S_1$  be the set of students in Linear algebra A-8 class.  $S_2 = \{0, 1, \dots, 50\}$ .
- Let  $f: S_1 \rightarrow S_2$  sending student  $x$  to his / her score in quiz 1.

# Linear transformation

Def:

- A **map**  $f$  from a set  $S_1$  to another set  $S_2$  is an assignment of element  $x$  in  $S_1$  to **an element**  $f(x)$  in  $S_2$ .
- We write  $f: S_1 \rightarrow S_2$  for a map.

e.g.

- Let  $S_1$  be the set of students in Linear algebra A-8 class.  $S_2 = \{0, 1, \dots, 50\}$ .
- Let  $f: S_1 \rightarrow S_2$  sending student  $x$  to his / her score in quiz 1.
- $f$  is a map.

# Linear transformation

# Linear transformation

Def. • Given two vector space  $V, W$ , a linear map/transformation is a map

$$L : V \longrightarrow W$$

# Linear transformation

Def. • Given two vector space  $V, W$ , a linear map/transformation is a map

$$L : V \longrightarrow W$$

Satisfying:  $L(a\vec{x} + b\vec{y}) = aL(\vec{x}) + bL(\vec{y}).$

# Linear transformation

Def. • Given two vector space  $V, W$ , a linear map/transformation is a map

$$L : V \longrightarrow W$$

satisfying:  $L(a\vec{x} + b\vec{y}) = aL(\vec{x}) + bL(\vec{y}).$

e.g.:  $L_A : \mathbb{R}^n \longrightarrow \mathbb{R}^m$  is linear

# Linear transformation

Def. • Given two vector space  $V, W$ , a linear map/transformation is a map

$$L : V \longrightarrow W$$

satisfying:  $L(a\vec{x} + b\vec{y}) = aL(\vec{x}) + bL(\vec{y}).$

e.g.:  $L_A : \mathbb{R}^n \longrightarrow \mathbb{R}^m$  is linear

$$L_A(a\vec{x} + b\vec{y}) = A(a\vec{x} + b\vec{y})$$

# Linear transformation

Def. • Given two vector space  $V, W$ , a linear map/transformation is a map

$$L : V \longrightarrow W$$

Satisfying:  $L(a\vec{x} + b\vec{y}) = aL(\vec{x}) + bL(\vec{y}).$

e.g.:  $L_A : \mathbb{R}^n \longrightarrow \mathbb{R}^m$  is linear

$$\begin{aligned} L_A(a\vec{x} + b\vec{y}) &= A(a\vec{x} + b\vec{y}) \\ &= aA\vec{x} + bA\vec{y} \end{aligned}$$

# Linear transformation

Def. • Given two vector space  $V, W$ , a linear map/transformation is a map

$$L : V \longrightarrow W$$

Satisfying:  $L(a\vec{x} + b\vec{y}) = aL(\vec{x}) + bL(\vec{y}).$

e.g.:  $L_A : \mathbb{R}^n \longrightarrow \mathbb{R}^m$  is linear

$$\begin{aligned} L_A(a\vec{x} + b\vec{y}) &= A(a\vec{x} + b\vec{y}) \\ &= aA\vec{x} + bA\vec{y} \\ &= aL_A(\vec{x}) + bL_A(\vec{y}). \end{aligned}$$

# Examples of Linear transformation

## Examples of Linear transformation

e.g.: • We learn that  $P_3 = \{a_0 + a_1x + a_2x^2 + a_3x^3 \mid a_i \in \mathbb{R}\}$ , the space of degree 3 polynomials, is a vector space.

# Examples of Linear transformation

- e.g.:
- We learn that  $P_3 = \{a_0 + a_1x + a_2x^2 + a_3x^3 \mid a_i \in \mathbb{R}\}$ , the space of degree 3 polynomials, is a vector space.
  - Define  $D: P_3 \longrightarrow P_3$  be

# Examples of Linear transformation

- e.g.: • We learn that  $P_3 = \{a_0 + a_1x + a_2x^2 + a_3x^3 \mid a_i \in \mathbb{R}\}$ , the space of degree 3 polynomials, is a vector space.
- Define  $D: P_3 \longrightarrow P_3$  be

$$D(a_0 + a_1x + a_2x^2 + a_3x^3) = \frac{d}{dx}(a_0 + a_1x + a_2x^2 + a_3x^3)$$
$$= a_1 + 2a_2x + 3a_3x^2.$$

# Examples of Linear transformation

e.g.: • We learn that  $P_3 = \{a_0 + a_1x + a_2x^2 + a_3x^3 \mid a_i \in \mathbb{R}\}$

, the space of degree 3 polynomials, is a vector space.

- Define  $D: P_3 \longrightarrow P_3$  be

$$D(a_0 + a_1x + a_2x^2 + a_3x^3) = \frac{d}{dx}(a_0 + a_1x + a_2x^2 + a_3x^3)$$

$$= a_1 + 2a_2x + 3a_3x^2.$$

- It is a linear transformation :

$$\begin{cases} D(cf(x)) = cD(f(x)) \\ D(f(x) + g(x)) = D(f(x)) + D(g(x)). \end{cases}$$

# Examples of Linear transformation

## Examples of Linear transformation

- We let  $m: P_3 \rightarrow P_4$  given by multiplication  
 $m(f(x)) := x f(x).$

## Examples of Linear transformation

- We let  $m: P_3 \rightarrow P_4$  given by multiplication  
 $m(f(x)) := x f(x).$

then explicitly :  $m(a_0 + a_1 x + a_2 x^2 + a_3 x^3)$   
 $= a_0 x + a_1 x^2 + a_2 x^3 + a_3 x^4.$

## Examples of Linear transformation

- We let  $m: P_3 \rightarrow P_4$  given by multiplication  
 $m(f(x)) := x f(x).$

then explicitly :  $m(a_0 + a_1 x + a_2 x^2 + a_3 x^3)$   
 $= a_0 x + a_1 x^2 + a_2 x^3 + a_3 x^4.$

- Let  $D = x \frac{d^2}{dx^2}$ ,  $D(f(x)) := x \frac{d^2 f(x)}{dx^2}$

## Examples of Linear transformation

- We let  $m: P_3 \rightarrow P_4$  given by multiplication  
 $m(f(x)) := x f(x).$

then explicitly :  $m(a_0 + a_1 x + a_2 x^2 + a_3 x^3)$   
 $= a_0 x + a_1 x^2 + a_2 x^3 + a_3 x^4.$

- Let  $D = x \frac{d^2}{dx^2}$ ,  $D(f(x)) := x \frac{d^2 f(x)}{dx^2}$

$$D(a_0 + a_1 x + a_2 x^2 + a_3 x^3) = x \frac{d}{dx}(a_1 + 2a_2 x + 3a_3 x^2)$$

## Examples of Linear transformation

- We let  $m: P_3 \rightarrow P_4$  given by multiplication  
 $m(f(x)) := x f(x).$

then explicitly :  $m(a_0 + a_1 x + a_2 x^2 + a_3 x^3)$   
 $= a_0 x + a_1 x^2 + a_2 x^3 + a_3 x^4.$

- Let  $D = x \frac{d^2}{dx^2}$ ,  $D(f(x)) := x \frac{d^2 f(x)}{dx^2}$

$$\begin{aligned} D(a_0 + a_1 x + a_2 x^2 + a_3 x^3) &= x \frac{d}{dx}(a_1 + 2a_2 x + 3a_3 x^2) \\ &= x(2a_2 + 6a_3 x) = 2a_2 x + 6a_3 x^2. \end{aligned}$$

# Examples of Linear transformation

## Examples of Linear transformation

- Integration:  $I : P_3 \longrightarrow P_4$  defined by

$$I(f(x)) := \int_0^x f(t) dt.$$

## Examples of Linear transformation

- Integration:  $I : P_3 \longrightarrow P_4$  defined by

$$I(f(x)) := \int_0^x f(t) dt.$$

explicity:  $I(a_0 + a_1 x + a_2 x^2 + a_3 x^3)$

$$= a_0 x + \left(\frac{a_1}{2}\right) x^2 + \left(\frac{a_2}{3}\right) x^3 + \left(\frac{a_3}{4}\right) x^4.$$

# Examples of Linear transformation

- Integration:  $I: P_3 \longrightarrow P_4$  defined by

$$I(f(x)) := \int_0^x f(t) dt.$$

explicity:  $I(a_0 + a_1 x + a_2 x^2 + a_3 x^3)$

$$= a_0 x + \left(\frac{a_1}{2}\right) x^2 + \left(\frac{a_2}{3}\right) x^3 + \left(\frac{a_3}{4}\right) x^4.$$

- $I(cf(x)) = \int_0^x cf(t) dt = c \int_0^x f(t) dt = c I(f(x))$

# Examples of Linear transformation

- Integration:  $I: P_3 \longrightarrow P_4$  defined by

$$I(f(x)) := \int_0^x f(t) dt.$$

explicity:  $I(a_0 + a_1 x + a_2 x^2 + a_3 x^3)$

$$= a_0 x + \left(\frac{a_1}{2}\right) x^2 + \left(\frac{a_2}{3}\right) x^3 + \left(\frac{a_3}{4}\right) x^4.$$

- $I(cf(x)) = \int_0^x cf(t) dt = c \int_0^x f(t) dt = c I(f(x))$

$$I(f(x) + g(x)) = \int_0^x (f(t) + g(t)) dt$$

# Examples of Linear transformation

- Integration:  $I: P_3 \rightarrow P_4$  defined by

$$I(f(x)) := \int_0^x f(t) dt.$$

explicity:  $I(a_0 + a_1 x + a_2 x^2 + a_3 x^3)$

$$= a_0 x + \left(\frac{a_1}{2}\right) x^2 + \left(\frac{a_2}{3}\right) x^3 + \left(\frac{a_3}{4}\right) x^4.$$

- $I(cf(x)) = \int_0^x cf(t) dt = c \int_0^x f(t) dt = c I(f(x))$

$$I(f(x) + g(x)) = \int_0^x (f(t) + g(t)) dt = \int_0^x f(t) dt + \int_0^x g(t) dt$$

# Examples of Linear transformation

- Integration:  $I: P_3 \rightarrow P_4$  defined by

$$I(f(x)) := \int_0^x f(t) dt.$$

explicity:  $I(a_0 + a_1 x + a_2 x^2 + a_3 x^3)$

$$= a_0 x + \left(\frac{a_1}{2}\right) x^2 + \left(\frac{a_2}{3}\right) x^3 + \left(\frac{a_3}{4}\right) x^4.$$

- $I(cf(x)) = \int_0^x cf(t) dt = c \int_0^x f(t) dt = c I(f(x))$

$$\begin{aligned} I(f(x) + g(x)) &= \int_0^x (f(t) + g(t)) dt = \int_0^x f(t) dt + \int_0^x g(t) dt \\ &= I(f(x)) + I(g(x)). \end{aligned}$$

# Composition of Linear transformation

# Composition of Linear transformation

prop (composition): Let  $f: V \rightarrow W$ ,  $g: W \rightarrow U$  be linear Transformation, then  $gof: V \rightarrow U$  is a linear transformation

# Composition of Linear transformation

- prop (composition): • Let  $f: V \rightarrow W$ ,  $g: W \rightarrow U$  be linear Transformation, then  $gof: V \rightarrow U$  is a linear transformation
- $gof$  is defined as  $gof(\vec{v}) = g(f(\vec{v}))$ .

# Composition of Linear transformation

- prop (composition): • Let  $f: V \rightarrow W$ ,  $g: W \rightarrow U$  be linear Transformation, then  $gof: V \rightarrow U$  is a linear transformation
- $gof$  is defined as  $gof(\vec{v}) = g(f(\vec{v}))$ .

Pf:  $gof(a\vec{v}_1 + b\vec{v}_2) = g(f(a\vec{v}_1 + b\vec{v}_2))$

# Composition of Linear transformation

- prop (composition): • Let  $f: V \rightarrow W$ ,  $g: W \rightarrow U$  be linear Transformation, then  $gof: V \rightarrow U$  is a linear transformation
- $gof$  is defined as  $gof(\vec{v}) = g(f(\vec{v}))$ .

Pf:

$$\begin{aligned}gof(a\vec{v}_1 + b\vec{v}_2) &= g(f(a\vec{v}_1 + b\vec{v}_2)) \\&= g(a f(\vec{v}_1) + b f(\vec{v}_2))\end{aligned}$$

# Composition of Linear transformation

- prop (composition): • Let  $f: V \rightarrow W$ ,  $g: W \rightarrow U$  be linear Transformation, then  $gof: V \rightarrow U$  is a linear transformation
- $gof$  is defined as  $gof(\vec{v}) = g(f(\vec{v}))$ .

Pf:

$$\begin{aligned} g \circ f (a\vec{v}_1 + b\vec{v}_2) &= g(f(a\vec{v}_1 + b\vec{v}_2)) \\ &= g(a\underbrace{f(\vec{v}_1)}_{\vec{w}_1} + b\underbrace{f(\vec{v}_2)}_{\vec{w}_2}) \end{aligned}$$

# Composition of Linear transformation

- prop (composition): • Let  $f: V \rightarrow W$ ,  $g: W \rightarrow U$  be linear Transformation, then  $gof: V \rightarrow U$  is a linear transformation
- $gof$  is defined as  $gof(\vec{v}) = g(f(\vec{v}))$ .

Pf:

$$\begin{aligned} g \circ f (a\vec{v}_1 + b\vec{v}_2) &= g(f(a\vec{v}_1 + b\vec{v}_2)) \\ &= g\left(a\underbrace{f(\vec{v}_1)}_{\vec{w}_1} + b\underbrace{f(\vec{v}_2)}_{\vec{w}_2}\right) \\ &= g(a\vec{w}_1 + b\vec{w}_2) \end{aligned}$$

# Composition of Linear transformation

- prop (composition): • Let  $f: V \rightarrow W$ ,  $g: W \rightarrow U$  be linear Transformation, then  $gof: V \rightarrow U$  is a linear transformation
- $gof$  is defined as  $gof(\vec{v}) = g(f(\vec{v}))$ .

Pf:

$$\begin{aligned} g \circ f (a\vec{v}_1 + b\vec{v}_2) &= g(f(a\vec{v}_1 + b\vec{v}_2)) \\ &= g(a\underbrace{f(\vec{v}_1)}_{\vec{w}_1} + b\underbrace{f(\vec{v}_2)}_{\vec{w}_2}) \\ &= g(a\vec{w}_1 + b\vec{w}_2) = ag(\vec{w}_1) + bg(\vec{w}_2) \end{aligned}$$

# Composition of Linear transformation

- prop (composition): • Let  $f: V \rightarrow W$ ,  $g: W \rightarrow U$  be linear Transformation, then  $gof: V \rightarrow U$  is a linear transformation
- $gof$  is defined as  $gof(\vec{v}) = g(f(\vec{v}))$ .

Pf:

$$\begin{aligned} g \circ f (a\vec{v}_1 + b\vec{v}_2) &= g(f(a\vec{v}_1 + b\vec{v}_2)) \\ &= g\left(a\underbrace{f(\vec{v}_1)}_{\vec{w}_1} + b\underbrace{f(\vec{v}_2)}_{\vec{w}_2}\right) \\ &= g(a\vec{w}_1 + b\vec{w}_2) = ag(\vec{w}_1) + bg(\vec{w}_2) \\ &= ag(f(\vec{v}_1)) + bg(f(\vec{v}_2)) \end{aligned}$$

# Composition of Linear transformation

- prop (composition): • Let  $f: V \rightarrow W$ ,  $g: W \rightarrow U$  be linear Transformation, then  $gof: V \rightarrow U$  is a linear transformation
- $gof$  is defined as  $gof(\vec{v}) = g(f(\vec{v}))$ .

Pf:

$$\begin{aligned}gof(a\vec{v}_1 + b\vec{v}_2) &= g(f(a\vec{v}_1 + b\vec{v}_2)) \\&= g\left(a\underbrace{f(\vec{v}_1)}_{\vec{w}_1} + b\underbrace{f(\vec{v}_2)}_{\vec{w}_2}\right) \\&= g(a\vec{w}_1 + b\vec{w}_2) = ag(\vec{w}_1) + bg(\vec{w}_2) \\&= ag(f(\vec{v}_1)) + bg(f(\vec{v}_2)) = a(gof)(\vec{v}_1) + b(gof)(\vec{v}_2)\end{aligned}$$

# Composition of Linear transformation

# Composition of Linear transformation

e.g.: We consider  $\mathcal{D} = x \frac{d^2}{dx^2}$ ,

# Composition of Linear transformation

e.g.: We consider  $D = x \frac{d^2}{dx^2}$ , we can think of

$$f = \frac{d^2}{dx^2} : P_3 \longrightarrow P_1, \text{ and } g : P_1 \longrightarrow P_2$$

# Composition of Linear transformation

e.g.: We consider  $D = x \frac{d^2}{dx^2}$ , we can think of

$f = \frac{d^2}{dx^2} : P_3 \rightarrow P_1$ , and  $g : P_1 \rightarrow P_2$

$\nwarrow$  multiplication by  $x$ .

# Composition of Linear transformation

e.g.: We consider  $D = x \frac{d^2}{dx^2}$ , we can think of

$f = \frac{d^2}{dx^2} : P_3 \rightarrow P_1$ , and  $g : P_1 \rightarrow P_2$

$\nwarrow$  multiplication by  $x$ .

then we have  $(g \circ f)(a_0 + a_1x + a_2x^2 + a_3x^3)$

# Composition of Linear transformation

e.g.: We consider  $D = x \frac{d^2}{dx^2}$ , we can think of

$f = \frac{d^2}{dx^2} : P_3 \rightarrow P_1$ , and  $g : P_1 \rightarrow P_2$

$\nwarrow$  multiplication by  $x$ .

then we have  $(gof)(a_0 + a_1x + a_2x^2 + a_3x^3)$

$$= g(f(a_0 + a_1x + a_2x^2 + a_3x^3))$$

# Composition of Linear transformation

e.g.: We consider  $D = x \frac{d^2}{dx^2}$ , we can think of

$f = \frac{d^2}{dx^2} : P_3 \rightarrow P_1$ , and  $g : P_1 \rightarrow P_2$

$\nwarrow$  multiplication by  $x$ .

then we have  $(g \circ f)(a_0 + a_1x + a_2x^2 + a_3x^3)$

$$= g(f(a_0 + a_1x + a_2x^2 + a_3x^3))$$

$$= g(2a_2 + 6a_3x)$$

# Composition of Linear transformation

e.g.: We consider  $D = x \frac{d^2}{dx^2}$ , we can think of

$f = \frac{d^2}{dx^2} : P_3 \rightarrow P_1$ , and  $g : P_1 \rightarrow P_2$

$\nwarrow$  multiplication by  $x$ .

$$\begin{aligned} \text{then we have } & (g \circ f)(a_0 + a_1x + a_2x^2 + a_3x^3) \\ &= g(f(a_0 + a_1x + a_2x^2 + a_3x^3)) \\ &= g(2a_2 + 6a_3x) = 2a_2x + 6a_3x^2 \\ &= D(a_0 + a_1x + a_2x^2 + a_3x^3) \end{aligned}$$

# Composition of Linear transformation

e.g.: We consider  $D = x \frac{d^2}{dx^2}$ , we can think of

$f = \frac{d^2}{dx^2} : P_3 \rightarrow P_1$ , and  $g : P_1 \rightarrow P_2$

$\nwarrow$  multiplication by  $x$ .

$$\begin{aligned} \text{then we have } & (g \circ f)(a_0 + a_1x + a_2x^2 + a_3x^3) \\ &= g(f(a_0 + a_1x + a_2x^2 + a_3x^3)) \\ &= g(2a_2 + 6a_3x) = 2a_2x + 6a_3x^2 \\ &= D(a_0 + a_1x + a_2x^2 + a_3x^3) \end{aligned}$$

then from the above proposition, we can deduce  
linearity of  $D$  from that of  $\frac{d}{dx}$  and  $g$ .

# Composition of Linear transformation

# Composition of Linear transformation

- Back to matrices:  $A$ :  $m \times n$  matrix  
 $B$ :  $n \times r$  matrix.

# Composition of Linear transformation

- Back to matrices:  $A$ :  $m \times n$  matrix  
 $B$ :  $n \times r$  matrix.
- $L_B: \mathbb{R}^r \rightarrow \mathbb{R}^n$ ,  $L_A: \mathbb{R}^n \rightarrow \mathbb{R}^m$

# Composition of Linear transformation

- Back to matrices:  $A$ :  $m \times n$  matrix  
 $B$ :  $n \times r$  matrix.
- $L_B : \mathbb{R}^r \rightarrow \mathbb{R}^n$ ,  $L_A : \mathbb{R}^n \rightarrow \mathbb{R}^m$
- $L_A \circ L_B : \mathbb{R}^r \rightarrow \mathbb{R}^m$  which is given by

# Composition of Linear transformation

- Back to matrices:  $A$ :  $m \times n$  matrix  
 $B$ :  $n \times r$  matrix.
- $L_B : \mathbb{R}^r \rightarrow \mathbb{R}^n$ ,  $L_A : \mathbb{R}^n \rightarrow \mathbb{R}^m$
- $L_A \circ L_B : \mathbb{R}^r \rightarrow \mathbb{R}^m$  which is given by  
$$L_A \circ L_B (\vec{x}) = L_A(L_B(\vec{x}))$$

# Composition of Linear transformation

- Back to matrices:  $A$ :  $m \times n$  matrix  
 $B$ :  $n \times r$  matrix.
- $L_B : \mathbb{R}^r \rightarrow \mathbb{R}^n$ ,  $L_A : \mathbb{R}^n \rightarrow \mathbb{R}^m$
- $L_A \circ L_B : \mathbb{R}^r \rightarrow \mathbb{R}^m$  which is given by  
$$L_A \circ L_B (\vec{x}) = L_A(L_B(\vec{x}))$$
 \in \mathbb{R}^m

# Composition of Linear transformation

- Back to matrices:  $A$ :  $m \times n$  matrix  
 $B$ :  $n \times r$  matrix.
- $L_B : \mathbb{R}^r \rightarrow \mathbb{R}^n$ ,  $L_A : \mathbb{R}^n \rightarrow \mathbb{R}^m$
- $L_A \circ L_B : \mathbb{R}^r \rightarrow \mathbb{R}^m$  which is given by  
$$L_A \circ L_B (\vec{x}) = L_A(L_B(\vec{x})) = L_A(B\vec{x})$$
 \in \mathbb{R}^r

# Composition of Linear transformation

- Back to matrices:  $A$ :  $m \times n$  matrix  
 $B$ :  $n \times r$  matrix.
- $L_B : \mathbb{R}^r \rightarrow \mathbb{R}^n$ ,  $L_A : \mathbb{R}^n \rightarrow \mathbb{R}^m$
- $L_A \circ L_B : \mathbb{R}^r \rightarrow \mathbb{R}^m$  which is given by  
$$L_A \circ L_B (\vec{x}) = L_A(L_B(\vec{x})) = L_A(B\vec{x})$$

# Composition of Linear transformation

- Back to matrices:  $A$ :  $m \times n$  matrix  
 $B$ :  $n \times r$  matrix.
- $L_B : \mathbb{R}^r \rightarrow \mathbb{R}^n$ ,  $L_A : \mathbb{R}^n \rightarrow \mathbb{R}^m$
- $L_A \circ L_B : \mathbb{R}^r \rightarrow \mathbb{R}^m$  which is given by  
$$L_A \circ L_B (\vec{x}) = L_A(L_B(\vec{x})) = L_A(B\vec{x})$$
  
$$= A(B\vec{x}) = (AB)(\vec{x})$$

# Composition of Linear transformation

- Back to matrices:  $A$ :  $m \times n$  matrix  
 $B$ :  $n \times r$  matrix.
- $L_B : \mathbb{R}^r \rightarrow \mathbb{R}^n$ ,  $L_A : \mathbb{R}^n \rightarrow \mathbb{R}^m$
- $L_A \circ L_B : \mathbb{R}^r \rightarrow \mathbb{R}^m$  which is given by  
$$L_A \circ L_B (\vec{x}) = L_A(L_B(\vec{x})) = L_A(B\vec{x})$$
$$= A(B\vec{x}) = (AB)(\vec{x})$$
$$= L_{AB}(\vec{x}).$$

# Composition of Linear transformation

- Back to matrices:  $A$ :  $m \times n$  matrix  
 $B$ :  $n \times r$  matrix.

- $L_B : \mathbb{R}^r \rightarrow \mathbb{R}^n$ ,  $L_A : \mathbb{R}^n \rightarrow \mathbb{R}^m$

- $L_A \circ L_B : \mathbb{R}^r \rightarrow \mathbb{R}^m$  which is given by

$$L_A \circ L_B (\vec{x}) = L_A(L_B(\vec{x})) = L_A(B\vec{x})$$

$$= A(B\vec{x}) = (AB)(\vec{x})$$

$$= L_{AB}(\vec{x}).$$

# Matrix representation of Linear transformation

# Matrix representation of Linear transformation

- If we look at  $L_A : \mathbb{R}^n \longrightarrow \mathbb{R}^m$  associated to a matrix, it is determined by  $L_A(\bar{e}_i)$

# Matrix representation of Linear transformation

- If we look at  $L_A : \mathbb{R}^n \longrightarrow \mathbb{R}^m$  associated to a matrix, it is determined by  $L_A(\vec{e}_i)$   
i.e. its action on a basis  $\vec{e}_1, \dots, \vec{e}_n$

# Matrix representation of Linear transformation

- If we look at  $L_A : \mathbb{R}^n \rightarrow \mathbb{R}^m$  associated to a matrix, it is determined by  $L_A(\vec{e}_i)$   
i.e. its action on a basis  $\vec{e}_1, \dots, \vec{e}_n$

Prop: Any linear transformation  $L : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is given by  $L = L_A$  for some  $m \times n$  matrix  $A$ .

# Matrix representation of Linear transformation

- If we look at  $L_A : \mathbb{R}^n \rightarrow \mathbb{R}^m$  associated to a matrix, it is determined by  $L_A(\vec{e}_i)$ i.e. its action on a basis  $\vec{e}_1, \dots, \vec{e}_n$

Prop: Any linear transformation  $L : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is given by  $L = L_A$  for some  $m \times n$  matrix  $A$ .

Pf: (say  $n=3, m=2$ )

# Matrix representation of Linear transformation

- If we look at  $L_A : \mathbb{R}^n \rightarrow \mathbb{R}^m$  associated to a matrix, it is determined by  $L_A(\vec{e}_i)$   
i.e. its action on a basis  $\vec{e}_1, \dots, \vec{e}_n$

Prop: Any linear transformation  $L : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is given by  $L = L_A$  for some  $m \times n$  matrix  $A$ .

Pf: (say  $n=3, m=2$ ) Write  $\begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix} = c_1 \vec{e}_1 + c_2 \vec{e}_2 + c_3 \vec{e}_3$ .

# Matrix representation of Linear transformation

- If we look at  $L_A : \mathbb{R}^n \rightarrow \mathbb{R}^m$  associated to a matrix, it is determined by  $L_A(\vec{e}_i)$   
i.e. its action on a basis  $\vec{e}_1, \dots, \vec{e}_n$

Prop: Any linear transformation  $L : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is given by  $L = L_A$  for some  $m \times n$  matrix  $A$ .

Pf: (say  $n=3, m=2$ ) Write  $\begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix} = c_1 \vec{e}_1 + c_2 \vec{e}_2 + c_3 \vec{e}_3$ .

$$L(c_1 \vec{e}_1 + c_2 \vec{e}_2 + c_3 \vec{e}_3) = c_1 L(\vec{e}_1) + c_2 L(\vec{e}_2) + c_3 L(\vec{e}_3)$$

# Matrix representation of Linear transformation

- If we look at  $L_A : \mathbb{R}^n \rightarrow \mathbb{R}^m$  associated to a matrix, it is determined by  $L_A(\vec{e}_i)$   
i.e. its action on a basis  $\vec{e}_1, \dots, \vec{e}_n$

Prop: Any linear transformation  $L : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is given by  $L = L_A$  for some  $m \times n$  matrix  $A$ .

Pf: (say  $n=3, m=2$ ) Write  $\begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix} = c_1 \vec{e}_1 + c_2 \vec{e}_2 + c_3 \vec{e}_3$ .

$$L(c_1 \vec{e}_1 + c_2 \vec{e}_2 + c_3 \vec{e}_3) = c_1 \underbrace{L(\vec{e}_1)}_{\text{Vectors in } \mathbb{R}^2} + c_2 \underbrace{L(\vec{e}_2)}_{\text{Vectors in } \mathbb{R}^2} + c_3 \underbrace{L(\vec{e}_3)}_{\text{Vectors in } \mathbb{R}^2}$$

Vectors in  $\mathbb{R}^2$

# Matrix representation of Linear transformation

- If we look at  $L_A : \mathbb{R}^n \rightarrow \mathbb{R}^m$  associated to a matrix, it is determined by  $L_A(\vec{e}_i)$   
i.e. its action on a basis  $\vec{e}_1, \dots, \vec{e}_n$

Prop: Any linear transformation  $L : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is given by  $L = L_A$  for some  $m \times n$  matrix  $A$ .

Pf: (say  $n=3, m=2$ ) Write  $\begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix} = c_1 \vec{e}_1 + c_2 \vec{e}_2 + c_3 \vec{e}_3$ .

$$L(c_1 \vec{e}_1 + c_2 \vec{e}_2 + c_3 \vec{e}_3) = c_1 \underbrace{L(\vec{e}_1)}_{\text{Vectors in } \mathbb{R}^2} + c_2 \underbrace{L(\vec{e}_2)}_{\text{Vectors in } \mathbb{R}^2} + c_3 \underbrace{L(\vec{e}_3)}_{\text{Vectors in } \mathbb{R}^2}$$

Set  $A = [L(\vec{e}_1), L(\vec{e}_2), L(\vec{e}_3)]$  be the matrix.

# Matrix representation of Linear transformation

# Matrix representation of Linear transformation

- For general  $L : V \rightarrow W$  between two vector space

# Matrix representation of Linear transformation

- For general  $L : V \rightarrow W$  between two vector space

Let say  $\vec{v}_1, \dots, \vec{v}_n$  a basis for  $V$ .

$\vec{w}_1, \dots, \vec{w}_m$  a basis for  $W$ .

# Matrix representation of Linear transformation

- For general  $L : V \rightarrow W$  between two vector space
  - Let say  $\vec{v}_1, \dots, \vec{v}_n$  a basis for  $V$ .
  - $\vec{w}_1, \dots, \vec{w}_m$  a basis for  $W$ .
- (Say  $n=3, m=2$ ):

# Matrix representation of Linear transformation

- For general  $L : V \rightarrow W$  between two vector space

Let say  $\vec{v}_1, \dots, \vec{v}_n$  a basis for  $V$ .

$\vec{w}_1, \dots, \vec{w}_m$  a basis for  $W$ .

- (Say  $n=3, m=2$ ):

We can write every vector  $\vec{v} = a_1 \vec{v}_1 + a_2 \vec{v}_2 + a_3 \vec{v}_3$

# Matrix representation of Linear transformation

- For general  $L : V \rightarrow W$  between two vector space

Let say  $\vec{v}_1, \dots, \vec{v}_n$  a basis for  $V$ .

$\vec{w}_1, \dots, \vec{w}_m$  a basis for  $W$ .

- (Say  $n=3, m=2$ ):

We can write every vector  $\vec{v} = a_1 \vec{v}_1 + a_2 \vec{v}_2 + a_3 \vec{v}_3$

$$L(\vec{v}) = L(a_1 \vec{v}_1 + a_2 \vec{v}_2 + a_3 \vec{v}_3)$$

# Matrix representation of Linear transformation

- For general  $L : V \rightarrow W$  between two vector space

Let say  $\vec{v}_1, \dots, \vec{v}_n$  a basis for  $V$ .

$\vec{w}_1, \dots, \vec{w}_m$  a basis for  $W$ .

- (Say  $n=3, m=2$ ):

We can write every vector  $\vec{v} = a_1 \vec{v}_1 + a_2 \vec{v}_2 + a_3 \vec{v}_3$

$$L(\vec{v}) = L(a_1 \vec{v}_1 + a_2 \vec{v}_2 + a_3 \vec{v}_3)$$

$$= a_1 L(\vec{v}_1) + a_2 L(\vec{v}_2) + a_3 L(\vec{v}_3)$$

# Matrix representation of Linear transformation

- For general  $L : V \rightarrow W$  between two vector space

Let say  $\vec{v}_1, \dots, \vec{v}_n$  a basis for  $V$ .

$\vec{w}_1, \dots, \vec{w}_m$  a basis for  $W$ .

- (Say  $n=3, m=2$ ):

We can write every vector  $\vec{v} = a_1 \vec{v}_1 + a_2 \vec{v}_2 + a_3 \vec{v}_3$

$$L(\vec{v}) = L(a_1 \vec{v}_1 + a_2 \vec{v}_2 + a_3 \vec{v}_3)$$

$$= a_1 L(\vec{v}_1) + a_2 L(\vec{v}_2) + a_3 L(\vec{v}_3)$$

- $L$  is uniquely determined by its value on a basis  $\vec{v}_1, \vec{v}_2, \vec{v}_3$ .

# Matrix representation of Linear transformation

# Matrix representation of Linear transformation

- Let's go back to the situation that  $L: V \rightarrow W$  with basis  $\vec{v}_1, \vec{v}_2, \vec{v}_3$  for  $V$ , and  $\vec{w}_1, \vec{w}_2$  for  $W$ .

# Matrix representation of Linear transformation

- Let's go back to the situation that  $L: V \rightarrow W$  with basis  $\vec{v}_1, \vec{v}_2, \vec{v}_3$  for  $V$ , and  $\vec{w}_1, \vec{w}_2$  for  $W$ .
- We know  $L$  is determined by  $L(\vec{v}_1), L(\vec{v}_2), L(\vec{v}_3)$ .

# Matrix representation of Linear transformation

- Let's go back to the situation that  $L: V \rightarrow W$  with basis  $\vec{v}_1, \vec{v}_2, \vec{v}_3$  for  $V$ , and  $\vec{w}_1, \vec{w}_2$  for  $W$ .
- We know  $L$  is determined by  $L(\vec{v}_1), L(\vec{v}_2), L(\vec{v}_3)$ .
- Let's further write

$$L(\vec{v}_1) = a_{11} \vec{w}_1 + a_{21} \vec{w}_2$$

$$L(\vec{v}_2) = a_{12} \vec{w}_1 + a_{22} \vec{w}_2$$

$$L(\vec{v}_3) = a_{13} \vec{w}_1 + a_{23} \vec{w}_2.$$

# Matrix representation of Linear transformation

- Let's go back to the situation that  $L: V \rightarrow W$  with basis  $\vec{v}_1, \vec{v}_2, \vec{v}_3$  for  $V$ , and  $\vec{w}_1, \vec{w}_2$  for  $W$ .
- We know  $L$  is determined by  $L(\vec{v}_1), L(\vec{v}_2), L(\vec{v}_3)$ .
- Let's further write

$$L(\vec{v}_1) = a_{11} \vec{w}_1 + a_{21} \vec{w}_2$$

$$L(\vec{v}_2) = a_{12} \vec{w}_1 + a_{22} \vec{w}_2$$

$$L(\vec{v}_3) = a_{13} \vec{w}_1 + a_{23} \vec{w}_2.$$

or

$$[L(\vec{v}_1), L(\vec{v}_2), L(\vec{v}_3)] = [\vec{w}_1, \vec{w}_2] \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \end{bmatrix}$$

# Matrix representation of Linear transformation

- Let's go back to the situation that  $L: V \rightarrow W$  with basis  $\vec{v}_1, \vec{v}_2, \vec{v}_3$  for  $V$ , and  $\vec{w}_1, \vec{w}_2$  for  $W$ .
- We know  $L$  is determined by  $L(\vec{v}_1), L(\vec{v}_2), L(\vec{v}_3)$ .
- Let's further write

$$L(\vec{v}_1) = a_{11} \vec{w}_1 + a_{21} \vec{w}_2$$

$$L(\vec{v}_2) = a_{12} \vec{w}_1 + a_{22} \vec{w}_2$$

$$L(\vec{v}_3) = a_{13} \vec{w}_1 + a_{23} \vec{w}_2.$$

this we call  
it the matrix  
representing  $L$ .  
↓

or

$$[L(\vec{v}_1), L(\vec{v}_2), L(\vec{v}_3)] = [\vec{w}_1, \vec{w}_2]$$

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \end{bmatrix}$$

# Matrix representation of Linear transformation

# Matrix representation of Linear transformation

- Fixing basis  $\vec{v}_i$ 's for  $V$ , and  $\vec{w}_i$ 's for  $W$ .  
The matrix  $[L]$  encode all information about  $L$ .

# Matrix representation of Linear transformation

- Fixing basis  $\vec{v}_i$ 's for  $V$ , and  $\vec{w}_i$ 's for  $W$ .  
The matrix  $[L]$  encode all information about  $L$ .
- If  $\vec{v} = c_1 \vec{v}_1 + c_2 \vec{v}_2 + c_3 \vec{v}_3$ .  
 $L(\vec{v}) = c_1 L(\vec{v}_1) + c_2 L(\vec{v}_2) + c_3 L(\vec{v}_3)$

# Matrix representation of Linear transformation

- Fixing basis  $\vec{v}_i$ 's for  $V$ , and  $\vec{w}_i$ 's for  $W$ .  
The matrix  $[L]$  encode all information about  $L$ .
- If  $\vec{v} = c_1 \vec{v}_1 + c_2 \vec{v}_2 + c_3 \vec{v}_3$ .  
$$L(\vec{v}) = c_1 L(\vec{v}_1) + c_2 L(\vec{v}_2) + c_3 L(\vec{v}_3)$$
$$= [L(\vec{v}_1), L(\vec{v}_2), L(\vec{v}_3)] \begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix}$$

# Matrix representation of Linear transformation

- Fixing basis  $\vec{v}_i$ 's for  $V$ , and  $\vec{w}_i$ 's for  $W$ .  
The matrix  $[L]$  encode all information about  $L$ .
- If  $\vec{v} = c_1 \vec{v}_1 + c_2 \vec{v}_2 + c_3 \vec{v}_3$ .  
 $L(\vec{v}) = c_1 L(\vec{v}_1) + c_2 L(\vec{v}_2) + c_3 L(\vec{v}_3)$

$$= [L(\vec{v}_1), L(\vec{v}_2), L(\vec{v}_3)] \begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix}$$

$$= [\vec{w}_1, \vec{w}_2] [L] \begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix}.$$

# Matrix representation of Linear transformation

- Fixing basis  $\vec{v}_i$ 's for  $V$ , and  $\vec{w}_i$ 's for  $W$ .  
The matrix  $[L]$  encode all information about  $L$ .

- If  $\vec{v} = c_1 \vec{v}_1 + c_2 \vec{v}_2 + c_3 \vec{v}_3$ .

$$L(\vec{v}) = c_1 L(\vec{v}_1) + c_2 L(\vec{v}_2) + c_3 L(\vec{v}_3)$$

$$= [L(\vec{v}_1), L(\vec{v}_2), L(\vec{v}_3)] \begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix}$$

$$= [\vec{w}_1, \vec{w}_2] \begin{bmatrix} L \\ \vdots \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix}.$$

this usual matrix multiplication tell us how to find  $L(\vec{v})$ .

# Matrix representation of Linear transformation

# Matrix representation of Linear transformation

e.g. • Let's consider  $D = \frac{d}{dx} : P_3 \rightarrow P_3$  be the linear transformation.

# Matrix representation of Linear transformation

- e.g. • Let's consider  $D = \frac{d}{dx} : P_3 \rightarrow P_3$  be the linear transformation.
- We fix a basis  $1, x, x^2, x^3$  for  $P_3$ .

# Matrix representation of Linear transformation

- e.g. • Let's consider  $D = \frac{d}{dx} : P_3 \rightarrow P_3$  be the linear transformation.
- We fix a basis  $1, x, x^2, x^3$  for  $P_3$ .
- 

# Matrix representation of Linear transformation

e.g. • Let's consider  $D = \frac{d}{dx} : P_3 \rightarrow P_3$  be the linear transformation.

$$\begin{matrix} \vec{v}_1 & \vec{v}_2 & \vec{v}_3 & \vec{v}_4 \\ \vec{w}_1 & \vec{w}_2 & \vec{w}_3 & \vec{w}_4 \end{matrix}$$

• We fix a basis  $1, x, x^2, x^3$  for  $P_3$ .

( both  $\vec{v}_i$ 's and  $\vec{w}_i$ 's are chosen to be this )

# Matrix representation of Linear transformation

e.g. • Let's consider  $D = \frac{d}{dx} : P_3 \rightarrow P_3$  be the linear transformation.

$$\begin{matrix} \vec{v}_1 & \vec{v}_2 & \vec{v}_3 & \vec{v}_4 \\ \vec{w}_1 & \vec{w}_2 & \vec{w}_3 & \vec{w}_4 \end{matrix}$$

- We fix a basis  $1, x, x^2, x^3$  for  $P_3$ .

( both  $\vec{v}_i$ 's and  $\vec{w}_i$ 's are chosen to be this )

$$D(\vec{v}_1) = 0 = 0 \cdot \vec{w}_1 + 0 \cdot \vec{w}_2 + 0 \cdot \vec{w}_3 + 0 \cdot \vec{w}_4$$

# Matrix representation of Linear transformation

- e.g.: • Let's consider  $D = \frac{d}{dx} : P_3 \rightarrow P_3$  be the linear transformation.
- We fix a basis  $1, x, x^2, x^3$  for  $P_3$ .
- ( both  $\vec{v}_i$ 's and  $\vec{w}_i$ 's are chosen to be this )
- $D(\vec{v}_1) = 0 = 0 \cdot \vec{w}_1 + 0 \cdot \vec{w}_2 + 0 \cdot \vec{w}_3 + 0 \cdot \vec{w}_4 = [\vec{w}_1 \dots \vec{w}_4] \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$

# Matrix representation of Linear transformation

e.g. • Let's consider  $D = \frac{d}{dx} : P_3 \rightarrow P_3$  be the linear transformation.

$$\begin{matrix} \vec{v}_1 & \vec{v}_2 & \vec{v}_3 & \vec{v}_4 \\ \vec{w}_1 & \vec{w}_2 & \vec{w}_3 & \vec{w}_4 \end{matrix}$$

• We fix a basis  $1, x, x^2, x^3$  for  $P_3$ .

( both  $\vec{v}_i$ 's and  $\vec{w}_i$ 's are chosen to be this )

$$D(\vec{v}_1) = 0 = 0 \cdot \vec{w}_1 + 0 \cdot \vec{w}_2 + 0 \cdot \vec{w}_3 + 0 \cdot \vec{w}_4 = [\vec{w}_1, \dots, \vec{w}_4] \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

$$D(\vec{v}_2) = 1 = 1 \cdot \vec{w}_1 + 0 \cdot \vec{w}_2 + 0 \cdot \vec{w}_3 + 0 \cdot \vec{w}_4 = [\vec{w}_1, \dots, \vec{w}_4] \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

# Matrix representation of Linear transformation

e.g. • Let's consider  $D = \frac{d}{dx} : P_3 \rightarrow P_3$  be the linear transformation.

$$\begin{array}{c} \vec{v}_1 \quad \vec{v}_2 \quad \vec{v}_3 \quad \vec{v}_4 \\ \vec{w}_1 \quad \vec{w}_2 \quad \vec{w}_3 \quad \vec{w}_4 \end{array}$$

• We fix a basis  $1, x, x^2, x^3$  for  $P_3$ .

( both  $\vec{v}_i$ 's and  $\vec{w}_i$ 's are chosen to be this )

$$D(\vec{v}_1) = 0 = 0 \cdot \vec{w}_1 + 0 \cdot \vec{w}_2 + 0 \cdot \vec{w}_3 + 0 \cdot \vec{w}_4 = [\vec{w}_1, \dots, \vec{w}_4] \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

$$D(\vec{v}_2) = 1 = 1 \cdot \vec{w}_1 + 0 \cdot \vec{w}_2 + 0 \cdot \vec{w}_3 + 0 \cdot \vec{w}_4 = [\vec{w}_1, \dots, \vec{w}_4] \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

$$D(\vec{v}_3) = 2x = 0 \cdot \vec{w}_1 + 2 \cdot \vec{w}_2 + 0 \cdot \vec{w}_3 + 0 \cdot \vec{w}_4 = [\vec{w}_1, \dots, \vec{w}_4] \begin{bmatrix} 0 \\ 2 \\ 0 \\ 0 \end{bmatrix}$$

# Matrix representation of Linear transformation

- e.g. • Let's consider  $D = \frac{d}{dx} : P_3 \rightarrow P_3$  be the linear transformation.

$$\begin{array}{c} \vec{v}_1 \quad \vec{v}_2 \quad \vec{v}_3 \quad \vec{v}_4 \\ \vec{w}_1 \quad \vec{w}_2 \quad \vec{w}_3 \quad \vec{w}_4 \end{array}$$

- We fix a basis  $1, x, x^2, x^3$  for  $P_3$ .

( both  $\vec{v}_i$ 's and  $\vec{w}_i$ 's are chosen to be this )

$$D(\vec{v}_1) = 0 = 0 \cdot \vec{w}_1 + 0 \cdot \vec{w}_2 + 0 \cdot \vec{w}_3 + 0 \cdot \vec{w}_4 = [\vec{w}_1, \dots, \vec{w}_4] \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

$$D(\vec{v}_2) = 1 = 1 \cdot \vec{w}_1 + 0 \cdot \vec{w}_2 + 0 \cdot \vec{w}_3 + 0 \cdot \vec{w}_4 = [\vec{w}_1, \dots, \vec{w}_4] \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

$$D(\vec{v}_3) = 2x = 0 \cdot \vec{w}_1 + 2 \cdot \vec{w}_2 + 0 \cdot \vec{w}_3 + 0 \cdot \vec{w}_4 = [\vec{w}_1, \dots, \vec{w}_4] \begin{bmatrix} 0 \\ 2 \\ 0 \\ 0 \end{bmatrix}$$

$$D(\vec{v}_4) = 3x^2 = 0 \cdot \vec{w}_1 + 0 \cdot \vec{w}_2 + 3 \cdot \vec{w}_3 + 0 \cdot \vec{w}_4 = [\vec{w}_1, \dots, \vec{w}_4] \begin{bmatrix} 0 \\ 0 \\ 3 \\ 0 \end{bmatrix}$$

# Matrix representation of Linear transformation

# Matrix representation of Linear transformation

- We collect the column vector representing

$$L(\vec{v}_i) = [\vec{w}_1, \vec{w}_2, \vec{w}_3, \vec{w}_4] \begin{bmatrix} 1 \\ a_i \\ 1 \end{bmatrix}$$

# Matrix representation of Linear transformation

- We collect the column vector representing

$$L(\vec{v}_i) = [\vec{w}_1, \vec{w}_2, \vec{w}_3, \vec{w}_4] \begin{bmatrix} | \\ a_i \\ | \end{bmatrix}$$

- We put  $\begin{bmatrix} | \\ a_i \\ | \end{bmatrix}$  as columns of  $[L]$ :  $[L] = \begin{bmatrix} | & | & | & | \\ a_1 & a_2 & a_3 & a_4 \\ | & | & | & | \end{bmatrix}$

# Matrix representation of Linear transformation

- We collect the column vector representing

$$L(\vec{v}_i) = [\vec{w}_1, \vec{w}_2, \vec{w}_3, \vec{w}_4] \begin{bmatrix} | \\ a_i \\ | \end{bmatrix}$$

- We put  $\begin{bmatrix} | \\ a_i \\ | \end{bmatrix}$  as columns of  $[L]$ :  $[L] = \begin{bmatrix} | & | & | & | \\ a_1, a_2, a_3, a_4 \\ | & | & | & | \end{bmatrix}$

- In the example:  $[L] = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 3 \\ 0 & 0 & 0 & 0 \end{bmatrix}$

# Matrix representation of Linear transformation

# Matrix representation of Linear transformation

- Let say we want to find  $D(1+x-3x^2+5x^3)$

# Matrix representation of Linear transformation

- Let say we want to find  $D(1+x-3x^2+5x^3)$

We can write  $1+x-3x^2+5x^3 = [\vec{v}_1, \vec{v}_2, \vec{v}_3, \vec{v}_4]$

$$\begin{bmatrix} 1 \\ -1 \\ -3 \\ 5 \end{bmatrix}$$

# Matrix representation of Linear transformation

- Let say we want to find  $D(1+x-3x^2+5x^3)$   
we can write  $1+x-3x^2+5x^3 = [\vec{v}_1, \vec{v}_2, \vec{v}_3, \vec{v}_4]$   $\begin{bmatrix} 1 \\ -1 \\ -3 \\ 5 \end{bmatrix}$
- We multiple  $\begin{bmatrix} 1 \\ -1 \\ -3 \\ 5 \end{bmatrix}$  to  $[L]$  from R.H.S :

# Matrix representation of Linear transformation

- Let say we want to find  $D(1+x-3x^2+5x^3)$

We can write  $1+x-3x^2+5x^3 = [\vec{v}_1, \vec{v}_2, \vec{v}_3, \vec{v}_4]$

$$\begin{bmatrix} 1 \\ -1 \\ -3 \\ 5 \end{bmatrix}$$

- We multiple  $\begin{bmatrix} 1 \\ -1 \\ -3 \\ 5 \end{bmatrix}$  to  $[L]$  from R.H.S :

$$\begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 3 \\ 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ -1 \\ -3 \\ 5 \end{bmatrix} = \begin{bmatrix} 1 \\ -6 \\ 15 \\ 0 \end{bmatrix}$$

# Matrix representation of Linear transformation

- Let say we want to find  $D(1+x-3x^2+5x^3)$

We can write  $1+x-3x^2+5x^3 = [\vec{v}_1, \vec{v}_2, \vec{v}_3, \vec{v}_4]$

$$\begin{bmatrix} 1 \\ -1 \\ -3 \\ 5 \end{bmatrix}$$

- We multiple  $\begin{bmatrix} 1 \\ -1 \\ -3 \\ 5 \end{bmatrix}$  to  $[L]$  from R.H.S :

$$\begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 3 \\ 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ -1 \\ -3 \\ 5 \end{bmatrix} = \begin{bmatrix} 1 \\ -6 \\ 15 \\ 0 \end{bmatrix}$$

- Then  $[\vec{w}_1, \vec{w}_2, \vec{w}_3, \vec{w}_4] \begin{bmatrix} 1 \\ -6 \\ 15 \\ 0 \end{bmatrix} = 1-6x+15x^2$  is the resulting vector.