


Chapter 3: Characterizing Running Times

Introduction

When we look at input sizes large enough to make only the order of growth of the running time relevant, we are studying **asymptotic** efficiency of algorithms. That is, we are concerned with how the running time increases with the size of the input *in the limit*, as the size of the input increases without bound.

Usually, an algorithm that is asymptotically more efficient will be the best choice for all but very small inputs.



Experiment on your own with an online graphing calculator to see how the growth of different functions compare. For example, try <http://desmos.com/calculator>.

Asymptotic Notation: informal introduction


- **O -notation** (big-Oh): The O -notation characterizes an upper bound on the asymptotic behavior of a function. It says that a function grows no faster than a certain rate, based upon its highest order term. Example: $7n^3 + 100n^2 - 20n + 6$ would be $O(n^3)$. It is also $O(n^4)$.
- **Ω -notation** (big-Omega): The Ω -notation characterizes a lower bound on the asymptotic behavior of a function. It says that a function grows at least as fast as a certain rate, based (again) upon its highest order term. Example: $7n^3 + 100n^2 - 20n + 6$ would be $\Omega(n^3)$. It is also $\Omega(n^2)$, and $\Omega(n)$.

- **Θ -notation (big-Theta)** The Θ -notation characterizes a tight bound on the asymptotic behavior of a function. It says that a function grows precisely at a certain rate, based on its highest order term. Example: $7n^3 + 100n^2 - 20n + 6$ would be $\Theta(n^3)$. However, it isn't $\Theta(n^2)$ or $\Theta(n^4)$.

Note: If a function is both $O(f(n))$ and $\Omega(f(n))$ for some function $f(n)$, then we have shown that the function is $\Theta(f(n))$.

Example: Do an informal asymptotic analysis of insertion sort to show that its worst-case running time is $O(n^2)$ and $\Omega(n^2)$ and thus is $\Theta(n^2)$.

Asymptotic Notation: formal definition



- **O -notation: asymptotically upper bound**

– *Definition:* For a given function $g(n)$, we denote by $O(g(n))$ to be the *set of functions*:

$$O(g(n)) = \{f(n): \text{there exist positive constants } c, \text{ and } n_0 \text{ such that } 0 \leq f(n) \leq cg(n) \text{ for all } n \geq n_0\}$$

– *Alternative Definition:* for two given functions $f(n)$ and $g(n)$:

$$f(n) = O(g(n)) \iff \lim_{n \rightarrow \infty} \frac{g(n)}{f(n)} = c \text{ or } \infty$$

– We can use the alternative definition to show

$$\begin{aligned} f(n) = \frac{1}{2}n^2 - 4n + 100 &= O(n^2) \quad \text{since } \lim_{n \rightarrow \infty} \frac{n^2}{\frac{1}{2}n^2 - 4n + 100} = 2 \\ &= O(n^3) \quad \text{since } \lim_{n \rightarrow \infty} \frac{n^3}{\frac{1}{2}n^2 - 4n + 100} = \infty \\ &= O(\infty) \quad \text{since } \lim_{n \rightarrow \infty} \frac{\infty}{\frac{1}{2}n^2 - 4n + 100} = \infty \end{aligned}$$

– Another example:

$$f(n) = 10 + \frac{1}{n} = O(1) \text{ since } \lim_{n \rightarrow \infty} \frac{1}{10 + \frac{1}{n}} = \frac{1}{10} \quad (1)$$

(2)

- **Ω -notation: asymptotically lower bound**

- *Definition:* For a given function $g(n)$, we denote by $\Omega(g(n))$ to be the *set of functions*:

$$\Omega(g(n)) = \{f(n): \text{there exist positive constants } c, \text{ and } n_0 \text{ such that } 0 \leq cg(n) \leq f(n) \text{ for all } n \geq n_0\}$$

- *Alternative Definition:* for two given functions $f(n)$ and $g(n)$:

$$f(n) = \Omega(g(n)) \iff \lim_{n \rightarrow \infty} \frac{g(n)}{f(n)} = c \text{ or } 0$$

- We can use the definition to show

$$\begin{aligned} f(n) = \frac{1}{2}n^2 - 4n + 100 &= \Omega(n^2) \quad \text{since } \lim_{n \rightarrow \infty} \frac{n^2}{\frac{1}{2}n^2 - 4n + 100} = 2 \\ &= \Omega(n) \quad \text{since } \lim_{n \rightarrow \infty} \frac{n}{\frac{1}{2}n^2 - 4n + 100} = 0 \\ &= \Omega(1) \quad \text{since } \lim_{n \rightarrow \infty} \frac{1}{\frac{1}{2}n^2 - 4n + 100} = 0 \end{aligned}$$

- For example:

$$\begin{aligned} \text{INSERTION-SORT}(n) &= \Omega(1) \\ &= \Omega(n) \\ &\neq \Omega(n^2), \text{ since best-case takes time proportional to } n \end{aligned}$$

- **Θ -notation: asymptotically tight bound**

- *Definition:* For a given function $g(n)$, we denote by $\Theta(g(n))$ to be the *set of functions*:

$$\Theta(g(n)) = \{f(n): \text{there exist positive constants } c_1, c_2, \text{ and } n_0 \text{ such that } 0 \leq c_1g(n) \leq f(n) \leq c_2g(n) \text{ for all } n \geq n_0\}$$

- *Alternative Definition:* For two given functions $f(n)$ and $g(n)$:

$$f(n) = \Theta(g(n)) \iff \lim_{n \rightarrow \infty} \frac{g(n)}{f(n)} = c$$

- We can use the definition to show $f(n) = \frac{1}{2}n^2 - 4n + 100 = \Theta(n^2)$, since

$$\lim_{n \rightarrow \infty} \frac{n^2}{\frac{1}{2}n^2 - 4n + 100} = 2$$

- Without using definition, we can determine the Θ -notation (as well as other notations) of a function $f(n)$ by throwing away lower-order terms and ignores the leading coefficient of the highest-order term.

For example,

$$f(n) = \frac{1}{2}n^2 - 4n + 100 = \Theta(n^2)$$

- People often use big-O notation to describe the running time of an algorithm rather than using Θ -notation (but they often mean to use Θ). However, Θ -notation tells people more exact running time of an algorithm. For example:

INSERTION-SORT(A) = $O(n^2)$ for all input n

INSERTION-SORT(A) $\neq \Theta(n^2)$
 $\neq \Theta(n)$

The worst case of INSERTION-SORT(A) = $\Theta(n^2)$

- *Theorem: for any two functions $f(n)$ and $g(n)$,*
 $f(n) = \Theta(g(n)) \implies f(n) = O(g(n))$ and $f(n) = \Omega(g(n))$

- **Properties of asymptotics**

- **Transitivity**

- $f(n) = \Theta(g(n))$ and $g(n) = \Theta(h(n))$ imply $f(n) = \Theta(h(n))$

- $f(n) = O(g(n))$ and $g(n) = O(h(n))$ imply $f(n) = O(h(n))$

- $f(n) = \Omega(g(n))$ and $g(n) = \Omega(h(n))$ imply $f(n) = \Omega(h(n))$

- **Reflexivity**

- $f(n) = \Theta(f(n))$

- $f(n) = O(f(n))$

- $f(n) = \Omega(f(n))$

- **Symmetry**

- $f(n) = \Theta(g(n))$ if and only if $g(n) = \Theta(f(n))$

- **Transpose Symmetry**

- $f(n) = O(g(n))$ if and only if $g(n) = \Omega(f(n))$

- **Trichotomy:** For any two numbers a and b , exactly one of the following must hold:
 $a < b, a = b, a > b$. Not all functions are asymptotically comparable. For example: n and $n^{1+\sin n}$ cannot be asymptotically compared.

- Based on the above properties, we can draw an analogy to comparing two numbers a and b :

- $f(n) = O(g(n))$ is similar to $a \leq b$

- $f(n) = \Omega(g(n))$ is similar to $a \geq b$

- $f(n) = \Theta(g(n))$ is similar to $a = b$

- **Asymptotic notations in equations**

- $2n^2 + 3n + 1 = 2n^2 + \Theta(n) \implies 2n^2 + 3n + 1 = 2n^2 + f(n)$, where $f(n) = \Theta(n)$. In this case, $f(n) = 3n + 1$

This example shows that we can mix asymptotics in equations. For example, we replaced lower order terms above by $\Theta(n)$ so we could focus on the more important terms.

– $T(n) = O(n) + \Theta(n) + \Omega(n) \implies T(n) = \Omega(n)$

This example shows how to combine three asymptotic components in an equation. $T(n)$ consists of the sum of the following three terms:

- * $O(n)$ which is growing at a rate $\leq n$,
- * $\Theta(n)$ so growing at the same rate as n , and
- * $\Omega(n)$ so growing at a rate greater than n .

If we combine those three, the best we can say is $\Omega(n)$ since that will be the dominant term for $T(n)$. This is because something growing faster than n will be growing bigger than two other terms growing at a rate equal to or less than n .

Categories of functions

growth rate	slowest	$\rightarrow$	$\rightarrow$	$\rightarrow$	fastest
run time	fastest	$\leftarrow$	$\leftarrow$	$\leftarrow$	slowest
categories	constant	logarithms	polynomials	exponentials	super exponentials
examples	5 1 10000	$\log_2 n$ $\log_{10} n$ $100 \log_e n$	n^2 n^3 $n^{0.1}$	$2^{n/2}$ 2^n 3^n	$(\log n)^n$ $n!$ n^n

- For logarithm functions, the base does not matter. $\log_{10} n = \Theta(\log_2 n)$ or $\log_e n = \Theta(\log_2 n)$
- For exponential functions, the base matters. $2^{n/2} = \sqrt{2^n} = O(2^n)$, but $2^{n/2} \neq \Theta(2^n)$
- Comparison between polynomials and exponentials: n^a and b^n
No matter how big the a is and how small the $b > 1$ is, the exponential b^n will always out-grow the polynomial n^a if n approaches ∞ .
For example, 1.000001^n will out-grow $n^{1,000,000}$ when $n \rightarrow \infty$
- **Review Section 3.3 from the textbook:** Discrete mathematics that is useful for analysis. Floors and ceilings, Modular arithmetic, Polynomials, Exponentials, Logarithms, Factorials (pay attention to Stirling's Approximation), and Fibonacci numbers.

– **Exponentials:** For all real $a > 0$, m , and n

$$\begin{aligned}
 a^0 &= 1 \\
 a^1 &= a \\
 a^{-1} &= \frac{1}{a} \\
 (a^m)^n &= a^{mn} \\
 (a^m)^n &= (a^n)^m
 \end{aligned}$$

$$a^m a^n = a^{m+n}$$

- **Logarithms:** Useful properties to remember:

$$\begin{aligned}\log(ab) &= \log a + \log b \\ \log\left(\frac{a}{b}\right) &= \log a - \log b \\ \log(a^b) &= b \log a\end{aligned}$$

- Logarithms and Exponentials are inverse functions:

$$\begin{aligned}b^{\log_b a} &= a \\ \log_b(b^a) &= a\end{aligned}$$

- **Change of base formula:**

$$\log_b a = \frac{\log_k a}{\log_k b}$$

This shows that logarithms in different bases differ only by a constant factor.

- We often use the base-2 logarithm in computer science since computers are binary.

$$\begin{aligned}\lg n &= \log_2 n \\ \ln n &= \log_e n \\ \log n &= \log_{10} n \\ \log^k n &= (\log n)^k \\ \lg \lg n &= \lg(\lg n)\end{aligned}$$

- **Factorials and Stirling's Approximation:**

$$n! = \sqrt{2\pi n} \left(\frac{n}{e}\right)^n \left(1 + \Theta\left(\frac{1}{n}\right)\right)$$

- This allows us to show that $\lg(n!) = \Theta(n \lg n)$, which is useful in the analysis of several algorithms.
- **Review Calculus:** The derivative rule and *L'Hôpital's rule* are useful for obtaining limits of more complex functions.

- **Summations:** Arithmetic series and Geometric series sums are commonly found during analysis of algorithms.

– **Simple Arithmetic series:**

$$1 + 2 + 3 + \dots + n = \frac{n(n+1)}{2}$$

Here is a common variation:

$$1 + 2 + 3 + \dots + n - 1 = \frac{n(n-1)}{2}$$

- **General Arithmetic Series:** Each term a_i is separated from the previous term a_{i-1} by a constant.

$$a_1 + a_2 + \dots + a_n = \frac{(a_1 + a_n)}{2}n$$

– **Simple Geometric Series:**

$$1 + 2 + 2^2 + 2^3 + \dots + 2^n = 2^{n+1} - 1$$

Here is a common variation:

$$1 + 2 + 2^2 + 2^3 + \dots + 2^{n-1} = 2^n - 1$$

– **General Geometric Series:**

For $x \neq 1$, we have:

$$1 + x + x^2 + x^3 + \dots + x^n = \frac{x^{n+1} - 1}{x - 1}$$

If $|x| < 1$, then the sum is $\frac{1}{x-1}$.

Practice Problems

- Exercise 3.2-3. Is $2^{n+1} = O(2^n)$? Is $2^{2n} = O(2^n)$?
- Order the following functions by growth rate: n , n^2 , $\sqrt{n}$, $n^{1.5}$, 2^n , $n!$, $(\log n)^2$, 3^n , $\log n^2$.

Math review

- Derivative rules: <https://www.mathsisfun.com/calculus/derivatives-rules.html>
- *L'Hôpital's rule*: <https://www.mathsisfun.com/calculus/l-hopitals-rule.html>