



UPPSALA  
UNIVERSITET

# Bayesian Statistics and Data Analysis

## Lecture 5

Måns Magnusson

Department of Statistics, Uppsala University  
Thanks to Aki Vehtari, Aalto University

- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation



UPPSALA  
UNIVERSITET

# It's all about expectations

---

$$E_{p(\theta|y)}[f(\theta)] = \int f(\theta) p(\theta|y) d\theta,$$

$$\text{where } p(\theta|y) = \frac{p(y|\theta)p(\theta)}{\int p(y|\theta)p(\theta)d\theta}$$

- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation



# It's all about expectations

$$E_{p(\theta|y)}[f(\theta)] = \int f(\theta) p(\theta|y) d\theta,$$

$$\text{where } p(\theta|y) = \frac{p(y|\theta)p(\theta)}{\int p(y|\theta)p(\theta)d\theta}$$

We can easily evaluate  $p(y|\theta)p(\theta)$  for any  $\theta$ , but the integral  $\int p(y|\theta)p(\theta)d\theta$  is usually difficult.

- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation



# It's all about expectations

$$E_{p(\theta|y)}[f(\theta)] = \int f(\theta) p(\theta|y) d\theta,$$

$$\text{where } p(\theta|y) = \frac{p(y|\theta)p(\theta)}{\int p(y|\theta)p(\theta)d\theta}$$

We can easily evaluate  $p(y|\theta)p(\theta)$  for any  $\theta$ , but the integral  $\int p(y|\theta)p(\theta)d\theta$  is usually difficult.

We can use the unnormalized posterior  $q(\theta|y) = p(y|\theta)p(\theta)$ , for example, in

- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

## It's all about expectations

$$E_{p(\theta|y)}[f(\theta)] = \int f(\theta)p(\theta|y)d\theta,$$

$$\text{where } p(\theta|y) = \frac{p(y|\theta)p(\theta)}{\int p(y|\theta)p(\theta)d\theta}$$

We can easily evaluate  $p(y|\theta)p(\theta)$  for any  $\theta$ , but the integral  $\int p(y|\theta)p(\theta)d\theta$  is usually difficult.

We can use the unnormalized posterior  $q(\theta|y) = p(y|\theta)p(\theta)$ , for example, in

- Monte Carlo methods which can sample from  $p(\theta^{(s)}|y)$  using only  $q(\theta^{(s)}|y)$

$$E_{p(\theta|y)}[f(\theta)] \approx \frac{1}{S} \sum_{s=1}^S f(\theta^{(s)})$$



UPPSALA  
UNIVERSITET

# Monte Carlo

---

- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation
- Monte Carlo methods we have discussed so far
  - Inverse CDF works for 1D



- Monte Carlo recap
  - Markov Chain Monte Carlo (MCMC)
    - Gibbs sampling
    - Metropolis-Hastings
  - Diagnostics
    - Warm-up
    - Convergence
    - $S_{\text{eff}}$ , MCSE, and autocorrelation
- Monte Carlo methods we have discussed so far
    - Inverse CDF works for 1D
    - Analytic transformations work for only certain distributions



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

- Monte Carlo methods we have discussed so far
  - Inverse CDF works for 1D
  - Analytic transformations work for only certain distributions
  - Grid methods works in less than a few dimensions





- Monte Carlo recap

- Markov Chain Monte Carlo (MCMC)

- Gibbs sampling
- Metropolis-Hastings

- Diagnostics

- Warm-up
- Convergence
- $S_{\text{eff}}$ , MCSE, and autocorrelation

- Monte Carlo methods we have discussed so far
  - Inverse CDF works for 1D
  - Analytic transformations work for only certain distributions
  - Grid methods works in less than a few dimensions
  - Rejection sampling works mostly in 1D (truncation is a special case)



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

- Monte Carlo methods we have discussed so far
  - Inverse CDF works for 1D
  - Analytic transformations work for only certain distributions
  - Grid methods works in less than a few dimensions
  - Rejection sampling works mostly in 1D (truncation is a special case)
  - Importance sampling is reliable only in special cases



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

- Monte Carlo methods we have discussed so far
  - Inverse CDF works for 1D
  - Analytic transformations work for only certain distributions
  - Grid methods works in less than a few dimensions
  - Rejection sampling works mostly in 1D (truncation is a special case)
  - Importance sampling is reliable only in special cases



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

- Monte Carlo methods we have discussed so far
  - Inverse CDF works for 1D
  - Analytic transformations work for only certain distributions
  - Grid methods works in less than a few dimensions
  - Rejection sampling works mostly in 1D (truncation is a special case)
  - Importance sampling is reliable only in special cases
- What to do in high dimensions?



- Monte Carlo recap

- Markov Chain Monte Carlo (MCMC)

- Gibbs sampling
- Metropolis-Hastings

- Diagnostics

- Warm-up
- Convergence
- $S_{\text{eff}}$ , MCSE, and autocorrelation

- Monte Carlo methods we have discussed so far
  - Inverse CDF works for 1D
  - Analytic transformations work for only certain distributions
  - Grid methods works in less than a few dimensions
  - Rejection sampling works mostly in 1D (truncation is a special case)
  - Importance sampling is reliable only in special cases
- What to do in high dimensions?
  - Markov chain Monte Carlo (Ch 11-12)



- Monte Carlo recap

- Markov Chain Monte Carlo (MCMC)

- Gibbs sampling
- Metropolis-Hastings

- Diagnostics

- Warm-up
- Convergence
- $S_{\text{eff}}$ , MCSE, and autocorrelation

- Monte Carlo methods we have discussed so far
  - Inverse CDF works for 1D
  - Analytic transformations work for only certain distributions
  - Grid methods works in less than a few dimensions
  - Rejection sampling works mostly in 1D (truncation is a special case)
  - Importance sampling is reliable only in special cases
- What to do in high dimensions?
  - Markov chain Monte Carlo (Ch 11-12)
  - Laplace, Variational\*, EP\* (Ch 4, 13\*, next course)



- Monte Carlo recap
  - Markov Chain Monte Carlo (MCMC)
    - Gibbs sampling
    - Metropolis-Hastings
  - Diagnostics
    - Warm-up
    - Convergence
    - $S_{\text{eff}}$ , MCSE, and autocorrelation
- Andrey Markov proved weak law of large numbers and central limit theorem for certain dependent-random sequences, which were later named Markov chains



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

# Markov chains

---

- Andrey Markov proved weak law of large numbers and central limit theorem for certain dependent-random sequences, which were later named Markov chains
- The probability of each event depends only on the state attained in the previous event (or finite number of previous events)

$$p(\theta_t | \theta_{t-1}, \theta_{t-2}, \dots) = p(\theta_t | \theta_{t-1})$$





- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

# Markov chains

---

- Andrey Markov proved weak law of large numbers and central limit theorem for certain dependent-random sequences, which were later named Markov chains
- The probability of each event depends only on the state attained in the previous event (or finite number of previous events)

$$p(\theta_t | \theta_{t-1}, \theta_{t-2}, \dots) = p(\theta_t | \theta_{t-1})$$

- $T_t(\theta_t | \theta_{t-1}) \equiv p(\theta_t | \theta_{t-1})$  is usually referred to as the **transition distribution**



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

# Markov chains

- Andrey Markov proved weak law of large numbers and central limit theorem for certain dependent-random sequences, which were later named Markov chains
- The probability of each event depends only on the state attained in the previous event (or finite number of previous events)

$$p(\theta_t | \theta_{t-1}, \theta_{t-2}, \dots) = p(\theta_t | \theta_{t-1})$$

- $T_t(\theta_t | \theta_{t-1}) \equiv p(\theta_t | \theta_{t-1})$  is usually referred to as the **transition distribution**
- Under some assumptions  $p(\theta_t | \theta_{t-1})$  will converge (in total variation) to *one* **stationary distribution**  $p(\theta)$



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

# Markov chains

- Andrey Markov proved weak law of large numbers and central limit theorem for certain dependent-random sequences, which were later named Markov chains
- The probability of each event depends only on the state attained in the previous event (or finite number of previous events)

$$p(\theta_t | \theta_{t-1}, \theta_{t-2}, \dots) = p(\theta_t | \theta_{t-1})$$

- $T_t(\theta_t | \theta_{t-1}) \equiv p(\theta_t | \theta_{t-1})$  is usually referred to as the **transition distribution**
- Under some assumptions  $p(\theta_t | \theta_{t-1})$  will converge (in total variation) to *one* **stationary distribution**  $p(\theta)$
- Goal in MCMC: Construct a **transition distribution** with  $p(\theta|y)$  as the **stationary distribution**



UPPSALA  
UNIVERSITET

# Markov chain Monte Carlo (MCMC)

---

- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation
- Produce draws  $\theta^{(t)}$  given  $\theta^{(t-1)}$  from a Markov chain, with **stationary distribution**  $p(\theta|y)$



# Markov chain Monte Carlo (MCMC)

---

- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation
- Produce draws  $\theta^{(t)}$  given  $\theta^{(t-1)}$  from a Markov chain, with **stationary distribution**  $p(\theta|y)$ 
  - + generic
  - + combine sequence of easier Monte Carlo draws to form a Markov chain



# Markov chain Monte Carlo (MCMC)

---

- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation
- Produce draws  $\theta^{(t)}$  given  $\theta^{(t-1)}$  from a Markov chain, with **stationary distribution**  $p(\theta|y)$ 
  - + generic
  - + combine sequence of easier Monte Carlo draws to form a Markov chain
  - + chain goes where most of the posterior mass is
  - + asymptotically chain spends the  $\alpha\%$  of time where  $\alpha\%$  posterior mass is



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation
- Produce draws  $\theta^{(t)}$  given  $\theta^{(t-1)}$  from a Markov chain, with **stationary distribution**  $p(\theta|y)$ 
  - + generic
  - + combine sequence of easier Monte Carlo draws to form a Markov chain
  - + chain goes where most of the posterior mass is
  - + asymptotically chain spends the  $\alpha\%$  of time where  $\alpha\%$  posterior mass is
  - + central limit theorem holds for expectations



# Markov chain Monte Carlo (MCMC)

---

- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation
- Produce draws  $\theta^{(t)}$  given  $\theta^{(t-1)}$  from a Markov chain, with **stationary distribution**  $p(\theta|y)$ 
  - + generic
  - + combine sequence of easier Monte Carlo draws to form a Markov chain
  - + chain goes where most of the posterior mass is
  - + asymptotically chain spends the  $\alpha\%$  of time where  $\alpha\%$  posterior mass is
  - + central limit theorem holds for expectations
    - draws are dependent
    - construction of efficient Markov chains is not always easy





# Markov chain Monte Carlo (MCMC)

---

- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

- Set of random variables  $\theta_1, \theta_2, \dots$ , so that with all values of  $t$ ,  $\theta_t$  depends only on the previous  $\theta_{(t-1)}$

$$p(\theta_t | \theta_1, \dots, \theta_{(t-1)}) = p(\theta_t | \theta_{(t-1)})$$



# Markov chain Monte Carlo (MCMC)

---

- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

- Set of random variables  $\theta_1, \theta_2, \dots$ , so that with all values of  $t$ ,  $\theta_t$  depends only on the previous  $\theta_{(t-1)}$

$$p(\theta_t | \theta_1, \dots, \theta_{(t-1)}) = p(\theta_t | \theta_{(t-1)})$$

- Chain has to be initialized with some starting point  $\theta_0$



# Markov chain Monte Carlo (MCMC)

---

- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

- Set of random variables  $\theta_1, \theta_2, \dots$ , so that with all values of  $t$ ,  $\theta_t$  depends only on the previous  $\theta_{(t-1)}$

$$p(\theta_t | \theta_1, \dots, \theta_{(t-1)}) = p(\theta_t | \theta_{(t-1)})$$

- Chain has to be initialized with some starting point  $\theta_0$
- Transition distribution  $T_t(\theta_t | \theta_{t-1})$  (may depend on  $t$ )



# Markov chain Monte Carlo (MCMC)

---

- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

- Set of random variables  $\theta_1, \theta_2, \dots$ , so that with all values of  $t$ ,  $\theta_t$  depends only on the previous  $\theta_{(t-1)}$

$$p(\theta_t | \theta_1, \dots, \theta_{(t-1)}) = p(\theta_t | \theta_{(t-1)})$$

- Chain has to be initialized with some starting point  $\theta_0$
- Transition distribution  $T_t(\theta_t | \theta_{t-1})$  (may depend on  $t$ )
- Choose a transition distribution so the stationary distribution of the Markov chain is  $p(\theta | y)$



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

- Alternate sampling from conditional distributions
- Basic algorithm, for  $j \in \{1, \dots, J\}$

sample  $\theta_{j,t}$  from  $p(\theta_j | \theta_{-j,t-1}, y)$ ,

where  $\theta_{j,t-1} = (\theta_{1,J}, \dots, \theta_{j-1,t}, \theta_{j+1,t-1}, \dots, \theta_{t-1,J})$

- Will converge (in total variation) to  $p(\theta|y)$  as  $T \rightarrow \infty$



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

- Alternate sampling from conditional distributions
- Basic algorithm, for  $j \in \{1, \dots, J\}$

sample  $\theta_{j,t}$  from  $p(\theta_j | \theta_{-j,t-1}, y)$ ,

where  $\theta_{j,t-1} = (\theta_{1,J}, \dots, \theta_{j-1,t}, \theta_{j+1,t-1}, \dots, \theta_{t-1,J})$

- Will converge (in total variation) to  $p(\theta|y)$  as  $T \rightarrow \infty$
- $j$  can be multiple (blocked) parameters
- 1D sampling ( $|j| = 1$ ) is generally easy



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

- Alternate sampling from conditional distributions
- Basic algorithm, for  $j \in \{1, \dots, J\}$


sample  $\theta_{j,t}$  from  $p(\theta_j | \theta_{-j,t-1}, y)$ ,  
where  $\theta_{j,t-1} = (\theta_{1,J}, \dots, \theta_{j-1,t}, \theta_{j+1,t-1}, \dots, \theta_{t-1,J})$

- Will converge (in total variation) to  $p(\theta|y)$  as  $T \rightarrow \infty$
- $j$  can be multiple (blocked) parameters
- 1D sampling ( $|j| = 1$ ) is generally easy
- Related to the (stochastic) EM algorithm



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

# Gibbs sampling



demo





UPPSALA  
UNIVERSITET

# Gibbs sampling

---

- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation
- With *conditionally* conjugate priors, the sampling from the conditional distributions is easy for wide range of models



UPPSALA  
UNIVERSITET

# Gibbs sampling

---

- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation
- With *conditionally* conjugate priors, the sampling from the conditional distributions is easy for wide range of models
- BUGS / WinBUGS / OpenBUGS / JAGS



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation
- With *conditionally* conjugate priors, the sampling from the conditional distributions is easy for wide range of models
- BUGS / WinBUGS / OpenBUGS / JAGS
- No algorithm parameters to tune



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation
- With *conditionally* conjugate priors, the sampling from the conditional distributions is easy for wide range of models
- BUGS / WinBUGS / OpenBUGS / JAGS
- No algorithm parameters to tune
- For not so easy conditionals, use e.g. inverse-CDF



- Monte Carlo recap
  - Markov Chain Monte Carlo (MCMC)
    - Gibbs sampling
    - Metropolis-Hastings
  - Diagnostics
    - Warm-up
    - Convergence
    - $S_{\text{eff}}$ , MCSE, and autocorrelation
- With *conditionally* conjugate priors, the sampling from the conditional distributions is easy for wide range of models
  - BUGS / WinBUGS / OpenBUGS / JAGS
  - No algorithm parameters to tune
  - For not so easy conditionals, use e.g. inverse-CDF
  - Several parameters can be updated in blocks (*blocking*)




- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation
- With *conditionally* conjugate priors, the sampling from the conditional distributions is easy for wide range of models
- BUGS / WinBUGS / OpenBUGS / JAGS
- No algorithm parameters to tune
- For not so easy conditionals, use e.g. inverse-CDF
- Several parameters can be updated in blocks (*blocking*)
- Slow if parameters are highly dependent in the posterior...



UPPSALA  
UNIVERSITET

- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

# Gibbs sampling



demo



# Sampling conditional vs joint

---

- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation
- How about sampling  $\theta$  jointly?
  - e.g. it is easy to sample from multivariate normal





- Monte Carlo recap
  - Markov Chain Monte Carlo (MCMC)
    - Gibbs sampling
    - Metropolis-Hastings
  - Diagnostics
    - Warm-up
    - Convergence
    - $S_{\text{eff}}$ , MCSE, and autocorrelation
- How about sampling  $\theta$  jointly?
    - e.g. it is easy to sample from multivariate normal
  - Can we use that to form a Markov chain?



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

# The Metropolis algorithm

---

- Algorithm

1. starting point  $\theta^0$

2.  $t = 1, 2, \dots$

- (a) pick a proposal  $\theta^*$  from a **proposal distribution**  $J_t(\theta^*|\theta_{t-1})$ .

Proposal distribution has to be symmetric, i.e.

$$J_t(\theta_a|\theta_b) = J_t(\theta_b|\theta_a), \text{ for all } \theta_a, \theta_b$$



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

# The Metropolis algorithm

- Algorithm

1. starting point  $\theta^0$

2.  $t = 1, 2, \dots$

- (a) pick a proposal  $\theta^*$  from a **proposal distribution**  $J_t(\theta^*|\theta_{t-1})$ .

Proposal distribution has to be symmetric, i.e.

$J_t(\theta_a|\theta_b) = J_t(\theta_b|\theta_a)$ , for all  $\theta_a, \theta_b$

- (b) calculate acceptance ratio

$$r = \frac{p(\theta^*|y)}{p(\theta_{t-1}|y)}$$



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

# The Metropolis algorithm

- Algorithm

1. starting point  $\theta^0$

2.  $t = 1, 2, \dots$

- (a) pick a proposal  $\theta^*$  from a **proposal distribution**  $J_t(\theta^*|\theta_{t-1})$ .

Proposal distribution has to be symmetric, i.e.

$J_t(\theta_a|\theta_b) = J_t(\theta_b|\theta_a)$ , for all  $\theta_a, \theta_b$

- (b) calculate acceptance ratio

- (c) set

$$r = \frac{p(\theta^*|y)}{p(\theta_{t-1}|y)}$$

$$\theta_t = \begin{cases} \theta^* & \text{with probability } \min(r, 1) \\ \theta_{t-1} & \text{otherwise} \end{cases}$$



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

# The Metropolis algorithm

- Algorithm

1. starting point  $\theta^0$
2.  $t = 1, 2, \dots$

(a) pick a proposal  $\theta^*$  from a **proposal distribution**  $J_t(\theta^*|\theta_{t-1})$ .

Proposal distribution has to be symmetric, i.e.  
 $J_t(\theta_a|\theta_b) = J_t(\theta_b|\theta_a)$ , for all  $\theta_a, \theta_b$

(b) calculate acceptance ratio

$$r = \frac{p(\theta^*|y)}{p(\theta_{t-1}|y)}$$

(c) set

$$\theta_t = \begin{cases} \theta^* & \text{with probability } \min(r, 1) \\ \theta_{t-1} & \text{otherwise} \end{cases}$$

ie, if  $p(\theta^*|y) > p(\theta_{t-1}|y)$  accept the proposal always  
and otherwise accept the proposal with probability  $r$



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

# The Metropolis algorithm

- Algorithm

1. starting point  $\theta^0$

2.  $t = 1, 2, \dots$

- (a) pick a proposal  $\theta^*$  from a **proposal distribution**  $J_t(\theta^*|\theta_{t-1})$ .

Proposal distribution has to be symmetric, i.e.

$J_t(\theta_a|\theta_b) = J_t(\theta_b|\theta_a)$ , for all  $\theta_a, \theta_b$

- (b) calculate acceptance ratio

- (c) set
$$r = \frac{p(\theta^*|y)}{p(\theta_{t-1}|y)}$$
$$\theta_t = \begin{cases} \theta^* & \text{with probability } \min(r, 1) \\ \theta_{t-1} & \text{otherwise} \end{cases}$$

- rejection of a proposal increments the time  $t$  also by one ie, the new state is the same as previous



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

# The Metropolis algorithm

- Algorithm

1. starting point  $\theta^0$

2.  $t = 1, 2, \dots$

- (a) pick a proposal  $\theta^*$  from a **proposal distribution**  $J_t(\theta^*|\theta_{t-1})$ .

Proposal distribution has to be symmetric, i.e.

$J_t(\theta_a|\theta_b) = J_t(\theta_b|\theta_a)$ , for all  $\theta_a, \theta_b$

- (b) calculate acceptance ratio

- (c) set
$$r = \frac{p(\theta^*|y)}{p(\theta_{t-1}|y)}$$
$$\theta_t = \begin{cases} \theta^* & \text{with probability } \min(r, 1) \\ \theta_{t-1} & \text{otherwise} \end{cases}$$

- rejection of a proposal increments the time  $t$  also by one ie, the new state is the same as previous
- step c is executed by generating a random number from  $\mathcal{U}(0, 1)$



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

# The Metropolis algorithm

- Algorithm

1. starting point  $\theta^0$

2.  $t = 1, 2, \dots$

- (a) pick a proposal  $\theta^*$  from a **proposal distribution**  $J_t(\theta^*|\theta_{t-1})$ .

Proposal distribution has to be symmetric, i.e.

$$J_t(\theta_a|\theta_b) = J_t(\theta_b|\theta_a), \text{ for all } \theta_a, \theta_b$$

- (b) calculate acceptance ratio

$$r = \frac{p(\theta^*|y)}{p(\theta_{t-1}|y)}$$

- (c) set

$$\theta_t = \begin{cases} \theta^* & \text{with probability } \min(r, 1) \\ \theta_{t-1} & \text{otherwise} \end{cases}$$

- rejection of a proposal increments the time  $t$  also by one ie, the new state is the same as previous
- step c is executed by generating a random number from  $\mathcal{U}(0, 1)$
- $p(\theta^*|y)$  and  $p(\theta_{t-1}|y)$  have the same normalization terms, and thus instead of  $p(\cdot|y)$ , unnormalized  $q(\cdot|y)$  can be used, **as the normalization terms cancel out!**





- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

- Example: one bivariate observation  $(y_1, y_2)$ 
  - bivariate normal distribution with unknown mean and known covariance

$$\begin{pmatrix} \theta_1 \\ \theta_2 \end{pmatrix} \Big| y \sim \mathcal{N} \left( \begin{pmatrix} y_1 \\ y_2 \end{pmatrix}, \begin{pmatrix} 1 & \rho \\ \rho & 1 \end{pmatrix} \right)$$

- proposal distribution  $J_t(\theta^* | \theta_{t-1}) = \mathcal{N}(\theta^* | \theta_{t-1}, \sigma_p^2)$

demo



UPPSALA  
UNIVERSITET

# Why Metropolis algorithm works

---

- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation
- Intuitively more draws from the higher density areas as jumps to higher density are always accepted and only some of the jumps to the lower density are accepted



# Why Metropolis algorithm works

---

- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation
- Intuitively more draws from the higher density areas as jumps to higher density are always accepted and only some of the jumps to the lower density are accepted
- Theoretically
  1. Prove that simulated series is a Markov chain which has unique stationary distribution
  2. Prove that this stationary distribution is the desired target distribution



# Why Metropolis algorithm works

---

- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

1. Prove that simulated series is a Markov chain which has unique stationary distribution
  - a) irreducible
  - b) aperiodic
  - c) recurrent / not transient



# Why Metropolis algorithm works

---

- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

1. Prove that simulated series is a Markov chain which has unique stationary distribution
  - a) irreducible
    - = positive probability of eventually reaching any state from any other state
  - b) aperiodic
  - c) recurrent / not transient



# Why Metropolis algorithm works

---

- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

1. Prove that simulated series is a Markov chain which has unique stationary distribution
  - a) irreducible
    - = positive probability of eventually reaching any state from any other state
  - b) aperiodic
    - = aperiodic (return times are not periodic)
    - holds for a random walk on any proper distribution (except for trivial exceptions)
  - c) recurrent / not transient



# Why Metropolis algorithm works

---

- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

1. Prove that simulated series is a Markov chain which has unique stationary distribution
  - a) irreducible
    - = positive probability of eventually reaching any state from any other state
  - b) aperiodic
    - = aperiodic (return times are not periodic)
    - holds for a random walk on any proper distribution (except for trivial exceptions)
  - c) recurrent / not transient
    - = probability to return to a state  $i$  is 1 as  $T \rightarrow \infty$
    - holds for a random walk on any proper distribution (except for trivial exceptions)



# Why Metropolis algorithm works

2. Prove that this stationary distribution is the desired target distribution  $p(\theta|y)$

- consider starting algorithm at time  $t - 1$  with a draw  $\theta_{t-1} \sim p(\theta|y)$

- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation





- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

## Why Metropolis algorithm works

2. Prove that this stationary distribution is the desired target distribution  $p(\theta|y)$ 
  - consider starting algorithm at time  $t - 1$  with a draw  $\theta_{t-1} \sim p(\theta|y)$
  - consider any two such points  $\theta_a$  and  $\theta_b$  drawn from  $p(\theta|y)$  and labeled so that  $p(\theta_b|y) \geq p(\theta_a|y)$



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

## Why Metropolis algorithm works

2. Prove that this stationary distribution is the desired target distribution  $p(\theta|y)$ 
  - consider starting algorithm at time  $t - 1$  with a draw  $\theta_{t-1} \sim p(\theta|y)$
  - consider any two such points  $\theta_a$  and  $\theta_b$  drawn from  $p(\theta|y)$  and labeled so that  $p(\theta_b|y) \geq p(\theta_a|y)$
  - the unconditional probability density of a transition from  $\theta_a$  to  $\theta_b$  is

$$p(\theta_{t-1} = \theta_a, \theta_t = \theta_b) = p(\theta_a|y)J_t(\theta_b|\theta_a),$$



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

## Why Metropolis algorithm works

### 2. Prove that this stationary distribution is the desired target distribution $p(\theta|y)$

- consider starting algorithm at time  $t - 1$  with a draw  $\theta_{t-1} \sim p(\theta|y)$
- consider any two such points  $\theta_a$  and  $\theta_b$  drawn from  $p(\theta|y)$  and labeled so that  $p(\theta_b|y) \geq p(\theta_a|y)$
- the unconditional probability density of a transition from  $\theta_a$  to  $\theta_b$  is

$$p(\theta_{t-1} = \theta_a, \theta_t = \theta_b) = p(\theta_a|y)J_t(\theta_b|\theta_a),$$

- the unconditional probability density of a transition from  $\theta_b$  to  $\theta_a$  is

$$\begin{aligned} p(\theta_t = \theta_a, \theta_{t-1} = \theta_b) &= p(\theta_b|y)J_t(\theta_a|\theta_b) \left( \frac{p(\theta_a|y)}{p(\theta_b|y)} \right) \\ &= p(\theta_a|y)J_t(\theta_a|\theta_b), \end{aligned}$$



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

## Why Metropolis algorithm works

### 2. Prove that this stationary distribution is the desired target distribution $p(\theta|y)$

- consider starting algorithm at time  $t - 1$  with a draw  $\theta_{t-1} \sim p(\theta|y)$
- consider any two such points  $\theta_a$  and  $\theta_b$  drawn from  $p(\theta|y)$  and labeled so that  $p(\theta_b|y) \geq p(\theta_a|y)$
- the unconditional probability density of a transition from  $\theta_a$  to  $\theta_b$  is

$$p(\theta_{t-1} = \theta_a, \theta_t = \theta_b) = p(\theta_a|y)J_t(\theta_b|\theta_a),$$

- the unconditional probability density of a transition from  $\theta_b$  to  $\theta_a$  is

$$\begin{aligned} p(\theta_t = \theta_a, \theta_{t-1} = \theta_b) &= p(\theta_b|y)J_t(\theta_a|\theta_b) \left( \frac{p(\theta_a|y)}{p(\theta_b|y)} \right) \\ &= p(\theta_a|y)J_t(\theta_a|\theta_b), \end{aligned}$$

which is the same as the probability of transition from  $\theta_a$  to  $\theta_b$ , since we have required that  $J_t(\cdot|\cdot)$  is symmetric



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

## Why Metropolis algorithm works

### 2. Prove that this stationary distribution is the desired target distribution $p(\theta|y)$

- consider starting algorithm at time  $t - 1$  with a draw  $\theta_{t-1} \sim p(\theta|y)$
- consider any two such points  $\theta_a$  and  $\theta_b$  drawn from  $p(\theta|y)$  and labeled so that  $p(\theta_b|y) \geq p(\theta_a|y)$
- the unconditional probability density of a transition from  $\theta_a$  to  $\theta_b$  is

$$p(\theta_{t-1} = \theta_a, \theta_t = \theta_b) = p(\theta_a|y)J_t(\theta_b|\theta_a),$$

- the unconditional probability density of a transition from  $\theta_b$  to  $\theta_a$  is

$$\begin{aligned} p(\theta_t = \theta_a, \theta_{t-1} = \theta_b) &= p(\theta_b|y)J_t(\theta_a|\theta_b) \left( \frac{p(\theta_a|y)}{p(\theta_b|y)} \right) \\ &= p(\theta_a|y)J_t(\theta_a|\theta_b), \end{aligned}$$

which is the same as the probability of transition from  $\theta_a$  to  $\theta_b$ , since we have required that  $J_t(\cdot|\cdot)$  is symmetric

- since their joint distribution is symmetric,  $\theta_t$  and  $\theta_{t-1}$  have the same marginal distributions, and so  $p(\theta|y)$  is the stationary distribution of the Markov chain of  $\theta$



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation
- Generalization of Metropolis algorithm for non-symmetric proposal distributions
  - acceptance ratio includes ratio of proposal distributions

$$r = \frac{p(\theta^*|y)/J_t(\theta^*|\theta_{t-1})}{p(\theta_{t-1}|y)/J_t(\theta_{t-1}|\theta^*)}$$



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

- Generalization of Metropolis algorithm for non-symmetric proposal distributions
  - acceptance ratio includes ratio of proposal distributions

$$r = \frac{p(\theta^*|y)/J_t(\theta^*|\theta_{t-1})}{p(\theta_{t-1}|y)/J_t(\theta_{t-1}|\theta^*)} = \frac{p(\theta^*|y)J_t(\theta_{t-1}|\theta^*)}{p(\theta_{t-1}|y)J_t(\theta^*|\theta_{t-1})}$$



# Metropolis-Hastings algorithm

---

- Ideal proposal distribution is the distribution itself
  - $J(\theta^*|\theta) \equiv p(\theta^*|y)$  for all  $\theta$
  - acceptance probability is 1
  - independent draws
  - not usually feasible

- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation





- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

# Metropolis-Hastings algorithm

---

- Ideal proposal distribution is the distribution itself
  - $J(\theta^*|\theta) \equiv p(\theta^*|y)$  for all  $\theta$
  - acceptance probability is 1
  - independent draws
  - not usually feasible
- Good proposal distribution resembles the target distribution
  - if the shape of the target distribution is unknown, usually normal or  $t$  distribution is used



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

# Metropolis-Hastings algorithm

- Ideal proposal distribution is the distribution itself
  - $J(\theta^*|\theta) \equiv p(\theta^*|y)$  for all  $\theta$
  - acceptance probability is 1
  - independent draws
  - not usually feasible
- Good proposal distribution resembles the target distribution
  - if the shape of the target distribution is unknown, usually normal or  $t$  distribution is used
- After the proposal distribution shape has been selected, it is important to select the scale
  - small scale
    - many steps accepted, but the chain moves slowly due to small steps
  - big scale
    - long steps proposed, but many of those rejected and again chain moves slowly

demo



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

## Metropolis-Hastings algorithm

- Ideal proposal distribution is the distribution itself
  - $J(\theta^*|\theta) \equiv p(\theta^*|y)$  for all  $\theta$
  - acceptance probability is 1
  - independent draws
  - not usually feasible
- Good proposal distribution resembles the target distribution
  - if the shape of the target distribution is unknown, usually normal or  $t$  distribution is used
- After the proposal distribution shape has been selected, it is important to select the scale
  - small scale
    - many steps accepted, but the chain moves slowly due to small steps
  - big scale
    - long steps proposed, but many of those rejected and again chain moves slowly

demo

- Generic rule for rejection rate is 60-90%




- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation
- Specific case of Metropolis-Hastings algorithm
  - single updated (or blocked)
  - proposal distribution is the conditional distribution
    - proposal and target distributions are same
    - acceptance probability is 1




- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

- Usually doesn't scale well to high dimensions
  - if the shape doesn't match the whole distribution, the efficiency drops



• Draws — Steps of the sampler — 90% HPI



• Draws — Steps of the sampler — 90% HPI



## Warm-up

---


- Asymptotically chain spends the  $\alpha\%$  of time where  $\alpha\%$  posterior mass is

- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation



## Warm-up

- Asymptotically chain spends the  $\alpha\%$  of time where  $\alpha\%$  posterior mass is
  - but in finite time the initial part of the Markov chain may be non-representative



• Draws — Steps of the sampler — 90% HPD


- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

## Warm-up

- Asymptotically chain spends the  $\alpha\%$  of time where  $\alpha\%$  posterior mass is
  - but in finite time the initial part of the Markov chain may be non-representative



• Draws — Steps of the sampler — 90% HPD

- Warm-up period = (non-representative) draws from the beginning of the Markov chain
  - remove warm-up before using samples for inference
  - warm-up may include also phase for adapting algorithm parameters






- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

## Warm-up

- Asymptotically chain spends the  $\alpha\%$  of time where  $\alpha\%$  posterior mass is
  - but in finite time the initial part of the Markov chain may be non-representative



• Draws — Steps of the sampler — 90% HPD

- Warm-up period = (non-representative) draws from the beginning of the Markov chain
  - remove warm-up before using samples for inference
  - warm-up may include also phase for adapting algorithm parameters
- Also called **burn-in**



UPPSALA  
UNIVERSITET

# Assesing convergence

---

- Several Markov chains make convergence diagnostics easier

- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - **Convergence**
  - $S_{\text{eff}}$ , MCSE, and autocorrelation




- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

## Assesing convergence

- Several Markov chains make convergence diagnostics easier
- Start chains from different starting points – preferably overdispersed


No convergence



- Remove warm-up draws and run chains long enough




- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - **Convergence**
  - $S_{\text{eff}}$ , MCSE, and autocorrelation





- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - **Convergence**
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

## Visually converged






- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - **Convergence**
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

## Several chains

### Visually converged



Visual convergence check is not sufficient



# $\hat{R}$ : comparison of within and between variances of the chains

---

- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation


- $\hat{R}$  or potential scale reduction factor (PSRF)
- Compare means and variances of the chains




# $\hat{R}$ : comparison of within and between variances of the chains

- $\hat{R}$  or **potential scale reduction factor** (PSRF)
- Compare means and variances of the chains  
 $W$  = within chain variance estimate  
 $\text{var\_hat\_plus}$  = total variance estimate

50 warmup, 50 post warmup iterations



Rhat = 1.64









# $\hat{R}$ : comparison of within and between variances of the chains

- $\hat{R}$  or **potential scale reduction factor** (PSRF)
- Compare means and variances of the chains  
 $W$  = within chain variance estimate  
 $\text{var\_hat\_plus}$  = total variance estimate

500 warmup, 500 post warmup iterations



Rhat = 1.08






# $\hat{R}$ : comparison of within and between variances of the chains

- $\hat{R}$  or **potential scale reduction factor** (PSRF)
- Compare means and variances of the chains  
 $W$  = within chain variance estimate  
 $\text{var\_hat\_plus}$  = total variance estimate

5000 warmup, 5000 post warmup iterations



Rhat = 1





UPPSALA  
UNIVERSITET

$\hat{R}$

---

- $M$  chains, each having  $N$  draws

- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - **Convergence**
  - $S_{\text{eff}}$ , MCSE, and autocorrelation



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - **Convergence**
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

- $M$  chains, each having  $N$  draws
- Within chains variance  $W$

$$W = \frac{1}{M} \sum_{m=1}^M s_m^2, \text{ where } s_m^2 = \frac{1}{N-1} \sum_{n=1}^N (\theta_{nm} - \bar{\theta}_{.m})^2$$



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - **Convergence**
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

- $M$  chains, each having  $N$  draws
- Within chains variance  $W$

$$W = \frac{1}{M} \sum_{m=1}^M s_m^2, \text{ where } s_m^2 = \frac{1}{N-1} \sum_{n=1}^N (\theta_{nm} - \bar{\theta}_{.m})^2$$

- Between chains variance  $B$

$$B = \frac{N}{M-1} \sum_{m=1}^M (\bar{\theta}_{.m} - \bar{\theta}_{..})^2,$$

$$\text{where } \bar{\theta}_{.m} = \frac{1}{N} \sum_{n=1}^N \theta_{nm}, \bar{\theta}_{..} = \frac{1}{M} \sum_{m=1}^M \bar{\theta}_{.m}$$



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - **Convergence**
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

- $M$  chains, each having  $N$  draws
- Within chains variance  $W$

$$W = \frac{1}{M} \sum_{m=1}^M s_m^2, \text{ where } s_m^2 = \frac{1}{N-1} \sum_{n=1}^N (\theta_{nm} - \bar{\theta}_{.m})^2$$

- Between chains variance  $B$

$$B = \frac{N}{M-1} \sum_{m=1}^M (\bar{\theta}_{.m} - \bar{\theta}_{..})^2,$$

$$\text{where } \bar{\theta}_{.m} = \frac{1}{N} \sum_{n=1}^N \theta_{nm}, \bar{\theta}_{..} = \frac{1}{M} \sum_{m=1}^M \bar{\theta}_{.m}$$

- $B/N$  is variance of the means of the chains



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - **Convergence**
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

- $M$  chains, each having  $N$  draws
- Within chains variance  $W$

$$W = \frac{1}{M} \sum_{m=1}^M s_m^2, \text{ where } s_m^2 = \frac{1}{N-1} \sum_{n=1}^N (\theta_{nm} - \bar{\theta}_{.m})^2$$

- Between chains variance  $B$

$$B = \frac{N}{M-1} \sum_{m=1}^M (\bar{\theta}_{.m} - \bar{\theta}_{..})^2,$$

$$\text{where } \bar{\theta}_{.m} = \frac{1}{N} \sum_{n=1}^N \theta_{nm}, \bar{\theta}_{..} = \frac{1}{M} \sum_{m=1}^M \bar{\theta}_{.m}$$

- $B/N$  is variance of the means of the chains
- Estimate total variance  $\text{var}(\theta|y)$  as a weighted mean of  $W$  and  $B$

$$\widehat{\text{var}}^+(\theta|y) = \frac{N-1}{N} W + \frac{1}{N} B$$



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - **Convergence**
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

- Estimate total variance  $\text{var}(\theta|y)$  as a weighted mean of  $W$  and  $B$

$$\widehat{\text{var}}^+(\theta|y) = \frac{N-1}{N}W + \frac{1}{N}B$$

- this *overestimates* marginal posterior variance if the starting points are overdispersed





- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - **Convergence**
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

- Estimate total variance  $\text{var}(\theta|y)$  as a weighted mean of  $W$  and  $B$

$$\widehat{\text{var}}^+(\theta|y) = \frac{N-1}{N}W + \frac{1}{N}B$$

- this *overestimates* marginal posterior variance if the starting points are overdispersed
- Given finite  $N$ ,  $W$  *underestimates* marginal posterior variance
  - single chains have not yet visited all points in the distribution
  - when  $N \rightarrow \infty$ ,  $E(W) \rightarrow \text{var}(\theta|y)$



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - **Convergence**
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

- Estimate total variance  $\text{var}(\theta|y)$  as a weighted mean of  $W$  and  $B$

$$\widehat{\text{var}}^+(\theta|y) = \frac{N-1}{N}W + \frac{1}{N}B$$

- this *overestimates* marginal posterior variance if the starting points are overdispersed
- Given finite  $N$ ,  $W$  *underestimates* marginal posterior variance
  - single chains have not yet visited all points in the distribution
  - when  $N \rightarrow \infty$ ,  $E(W) \rightarrow \text{var}(\theta|y)$
- As  $\widehat{\text{var}}^+(\theta|y)$  overestimates and  $W$  underestimates, compute

$$\hat{R} = \sqrt{\frac{\widehat{\text{var}}^+}{W}}$$




# UPPSALA UNIVERSITET

- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation


$\hat{R}$

- BDA3:  $\hat{R}$  aka *potential scale reduction factor* (PSRF)
- Compare means and variances of the chains  
 $W$  = within chain variance estimate  
 $\text{var\_hat\_plus}$  = total variance estimate

50 warmup, 50 post warmup iterations



Rhat = 1.64







- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

- BDA3:  $\hat{R}$  aka *potential scale reduction factor* (PSRF)
- Compare means and variances of the chains  
W = within chain variance estimate  
var\_hat\_plus = total variance estimate

500 warmup, 500 post warmup iterations



Rhat = 1.08






# UPPSALA UNIVERSITET

- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

$\hat{R}$

- BDA3:  $\hat{R}$  aka *potential scale reduction factor* (PSRF)
- Compare means and variances of the chains  
W = within chain variance estimate  
var\_hat\_plus = total variance estimate

5000 warmup, 5000 post warmup iterations



Rhat = 1





# UPPSALA UNIVERSITET

$\hat{R}$

- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - **Convergence**
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

$$\hat{R} = \sqrt{\frac{\widehat{\text{var}}^+}{W}}$$

- Estimates how much the scale could reduce if  $N \rightarrow \infty$
- $\hat{R} \rightarrow 1$ , when  $N \rightarrow \infty$
- If  $\hat{R}$  is big (e.g.,  $R > 1.01$ ), keep sampling



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - **Convergence**
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

$$\hat{R} = \sqrt{\frac{\widehat{\text{var}}^+}{W}}$$

- Estimates how much the scale could reduce if  $N \rightarrow \infty$
- $\hat{R} \rightarrow 1$ , when  $N \rightarrow \infty$
- If  $\hat{R}$  is big (e.g.,  $R > 1.01$ ), keep sampling
- If  $\hat{R}$  close to 1, it is still possible that chains have not converged
  - if starting points were not overdispersed
  - distribution far from normal (especially if infinite variance)
  - just by chance when  $N$  is finite



## UPPSALA UNIVERSITET

$\hat{R}$

---

- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - **Convergence**
  - $S_{\text{eff}}$ , MCSE, and autocorrelation
- Additional  $\hat{R}$  methods to assess convergence
- Split- $\hat{R}$ 
  - Examines *mixing* and *stationarity* of chains
  - To examine stationarity chains are split to two parts: compare means and variances of the split chains





- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - **Convergence**
  - $S_{\text{eff}}$ , MCSE, and autocorrelation
- Additional  $\hat{R}$  methods to assess convergence
- Split- $\hat{R}$ 
  - Examines *mixing* and *stationarity* of chains
  - To examine stationarity chains are split to two parts: compare means and variances of the split chains
- Rank normalized  $\hat{R}$ 
  - Does not requires that the target distribution has finite mean and variance



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

- Monte Carlo estimates still valid (central limit theorem)

$$E_{p(\theta|y)}[f(\theta)] \approx \frac{1}{S} \sum_{s=1}^S f(\theta^{(s)})$$

- Estimation of Monte Carlo error is more difficult
  - evaluation of *effective* sample size,  $S_{\text{eff}}$ .




- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation
- Auto correlation function
  - describes the correlation given a certain lag
  - can be used to compare efficiency of MCMC methods



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

# Autocorrelation




• Draws — Steps of the sampler — 90% HPD




- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

# Autocorrelation




• Draws — Steps of the sampler — 90% HPD






- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation


# Autocorrelation



• Draws — Steps of the sampler — 90% HPD



—  $\theta_1$  —  $\theta_2$




—  $\theta_1$  —  $\theta_2$




# UPPSALA UNIVERSITET

- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation


## Autocorrelation



• Draws — Steps of the sampler — 90% HPD



— theta1 — theta2




— theta1 — theta2




- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation


# Autocorrelation



• Draws — Steps of the sampler — 90% HPI



—  $\theta_1$  —  $\theta_2$







- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

- The autocorrelation can be used to estimate Monte Carlo error in case of MCMC
- For expectation  $\bar{\theta}$

$$\text{Var}[\bar{\theta}] = \frac{\sigma_{\theta}^2}{S_{\text{eff}}}$$

where  $S_{\text{eff}} = S/\tau$ , and  $\tau$  is sum of autocorrelations



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

- The autocorrelation can be used to estimate Monte Carlo error in case of MCMC
- For expectation  $\bar{\theta}$

$$\text{Var}[\bar{\theta}] = \frac{\sigma_{\theta}^2}{S_{\text{eff}}}$$

where  $S_{\text{eff}} = S/\tau$ , and  $\tau$  is sum of autocorrelations

- $\tau$  describes how many dependent draws correspond to one independent draw



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

- The autocorrelation can be used to estimate Monte Carlo error in case of MCMC
- For expectation  $\bar{\theta}$

$$\text{Var}[\bar{\theta}] = \frac{\sigma_{\theta}^2}{S_{\text{eff}}}$$

where  $S_{\text{eff}} = S/\tau$ , and  $\tau$  is sum of autocorrelations

- $\tau$  describes how many dependent draws correspond to one independent draw
- Here  $S = NM$  (in BDA3  $N = nm$  and  $n_{\text{eff}} = N/\tau$ )



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

- The autocorrelation can be used to estimate Monte Carlo error in case of MCMC
- For expectation  $\bar{\theta}$

$$\text{Var}[\bar{\theta}] = \frac{\sigma_{\theta}^2}{S_{\text{eff}}}$$

where  $S_{\text{eff}} = S/\tau$ , and  $\tau$  is sum of autocorrelations

- $\tau$  describes how many dependent draws correspond to one independent draw
- Here  $S = NM$  (in BDA3  $N = nm$  and  $n_{\text{eff}} = N/\tau$ )
- BDA3 focuses on  $S_{\text{eff}}$  and not the Monte Carlo error directly



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

- Estimation of the autocorrelation using several chains

$$\hat{\rho}_n = 1 - \frac{W - \frac{1}{M} \sum_{m=1}^M \hat{\rho}_{n,m}}{2\widehat{\text{var}}^+}$$

where  $\hat{\rho}_{n,m}$  is autocorrelation at lag  $n$  for chain  $m$



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

- Estimation of the autocorrelation using several chains

$$\hat{\rho}_n = 1 - \frac{W - \frac{1}{M} \sum_{m=1}^M \hat{\rho}_{n,m}}{2\widehat{\text{var}}^+}$$

where  $\hat{\rho}_{n,m}$  is autocorrelation at lag  $n$  for chain  $m$

- BDA3 has slightly different and less accurate equation. The above equation is used in Stan 2.18+



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

- Estimation of the autocorrelation using several chains

$$\hat{\rho}_n = 1 - \frac{W - \frac{1}{M} \sum_{m=1}^M \hat{\rho}_{n,m}}{2\widehat{\text{var}}^+}$$

where  $\hat{\rho}_{n,m}$  is autocorrelation at lag  $n$  for chain  $m$

- BDA3 has slightly different and less accurate equation. The above equation is used in Stan 2.18+
- Compared to a method which computes the autocorrelation from a single chain, the multi-chain estimate has smaller variance



- Estimation of  $\tau$

$$\tau = 1 + 2 \sum_{t=1}^{\infty} \hat{\rho}_t$$

where  $\hat{\rho}_t$  is empirical autocorrelation

- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation





- Estimation of  $\tau$

$$\tau = 1 + 2 \sum_{t=1}^{\infty} \hat{\rho}_t$$

where  $\hat{\rho}_t$  is empirical autocorrelation

- empirical autocorrelation function is noisy
- noise is larger for longer lags (less observations)

- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation



- Estimation of  $\tau$

$$\tau = 1 + 2 \sum_{t=1}^{\infty} \hat{\rho}_t$$

where  $\hat{\rho}_t$  is empirical autocorrelation

- empirical autocorrelation function is noisy
- noise is larger for longer lags (less observations)
- less noisy estimate is obtained by truncating

$$\hat{\tau} = 1 + 2 \sum_{t=1}^T \hat{\rho}_t$$

- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

## Geyer's adaptive window estimator of $\tau$

---


- Truncation  $T$  can be decided adaptively
  - for stationary, irreducible, recurrent Markov chain
  - let  $\Gamma_m = \rho_{2m} + \rho_{2m+1}$ , which is sum of two consequent autocorrelations
  - $\Gamma_m$  is positive, decreasing and convex function of  $m$



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

## Geyer's adaptive window estimator of $\tau$

- Truncation  $T$  can be decided adaptively
  - for stationary, irreducible, recurrent Markov chain
  - let  $\Gamma_m = \rho_{2m} + \rho_{2m+1}$ , which is sum of two consequent autocorrelations
  - $\Gamma_m$  is positive, decreasing and convex function of  $m$
- Initial positive sequence estimator (Geyer's IPSE)
  - Choose the largest  $m$  so, that all values of the sequence  $\hat{\Gamma}_1, \dots, \hat{\Gamma}_m$  are positive





UPPSALA  
UNIVERSITET

# Effective sample size, $S_{\text{eff}}$

---

Effective sample size  $\text{ESS} = S_{\text{eff}} \approx S/\hat{\tau}$




- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

# Effective sample size, $S_{\text{eff}}$

$$\text{Effective sample size ESS} = S_{\text{eff}} \approx S / \hat{\tau}$$



$$\hat{\tau} = 1 + 2 \sum_{t=1}^T \hat{\rho}_t$$




$$\approx 24$$



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

# Effective sample size

$$\text{Effective sample size } ESS = S_{\text{eff}} \approx S / \hat{\tau}$$



$$\hat{\tau} = 1 + 2 \sum_{t=1}^T \hat{\rho}_t$$




$$\approx 104$$



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation

# Effective sample size

$$\text{Effective sample size } ESS = S_{\text{eff}} \approx S / \hat{\tau}$$



$$\hat{\tau} = 1 + 2 \sum_{t=1}^T \hat{\rho}_t$$

$$\approx 63$$





- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation
- Nonlinear dependencies
  - optimal proposal depends on location



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation
- Nonlinear dependencies
  - optimal proposal depends on location
- Funnels
  - optimal proposal depends on location



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation
- Nonlinear dependencies
  - optimal proposal depends on location
- Funnels
  - optimal proposal depends on location
- Multimodal
  - difficult to move from one mode to another



- Monte Carlo recap
- Markov Chain Monte Carlo (MCMC)
  - Gibbs sampling
  - Metropolis-Hastings
- Diagnostics
  - Warm-up
  - Convergence
  - $S_{\text{eff}}$ , MCSE, and autocorrelation
- Nonlinear dependencies
  - optimal proposal depends on location
- Funnels
  - optimal proposal depends on location
- Multimodal
  - difficult to move from one mode to another
- Long-tailed with non-finite variance and mean
  - central limit theorem for expectations does not hold