

COMP527  
Data Mining and Visualisation  
Problem Set 3

Danushka Bollegala




Figure 1: Projecting three points A, B, C onto the line  $y = mx + c$ .

**Question** Consider the problem of projecting a two-dimensional dataset consisting of three points  $A = (-1, 0)$ ,  $B = (0, 1)$ , and  $C = (1, 0)$  onto the one-dimensional line given by  $y = mx + c$ . The dataset and the line is shown in Figure 1. Answer the following questions.

- A. Compute the co-ordinates of the base of the perpendicular from point  $(\alpha, \beta)$  to line  $y = mx + c$ . **(10 marks)**

Assume the base of the coordinates to be  $(p, q)$ . Then it must satisfy the following equations as it is on the line  $y = mx + c$  and the line section connecting base to  $(\alpha, \beta)$  is perpendicular (hence gradient of  $-1/m$ ).

$$q = mp + c \quad (1)$$

$$\frac{q - \beta}{p - \alpha} = -\frac{1}{m} \quad (2)$$

Solving these equations we get,

$$p = \frac{\alpha + m\beta - mc}{1 + m^2}$$

and

$$q = \frac{m\alpha + m^2\beta + c}{1 + m^2}.$$

- B. Compute the perpendicular distance to the line  $y = mx + c$  from point  $(\alpha, \beta)$ . **(10 marks)**

The distance  $d$  is given by,

$$d^2 = (p - \alpha)^2 + (q - \beta)^2.$$

By substituting for  $(p, q)$  from the previous question we get,

$$d = \frac{|\alpha m - \beta + c|}{\sqrt{1 + m^2}}$$

- C. Show that if  $y = mx + c$  is a solution to the one dimensional PCA projection, then  $y = mx + c'$  is also a solution. Here,  $c \neq c'$ . **(10 marks)**

Any line parallel to  $y = mx + c$  will have the same variations between the corresponding pairs of projected points. Therefore, any parallel line would be a solution. Therefore, we can set  $c = 0$  for the rest of the question and find the solution for  $m$ . Then any line  $y = mx + c$  for the found  $m$  will be a solution for any  $c \in \mathbb{R}$ .

- D. Find  $m$  such that the variance of the projected points on to the straight line is maximised. **(20 marks)**

Let us assume the projections of  $A, B, C$  onto  $y = mx + c$  are given by  $A', B', C'$ . Then the variation  $v$  is given by,

$$v = A'B'^2 + A'B'^2 + B'C'^2$$

Substituting for the base points

$$A' = \left( \frac{-1 - mc}{1 + m^2}, \frac{-m + c}{1 + m^2} \right), \quad (3)$$

$$B' = \left( \frac{1 - mc}{1 + m^2}, \frac{m^2 + c}{1 + m^2} \right), \quad (4)$$

$$C' = \left( \frac{1 - mc}{1 + m^2}, \frac{m + c}{1 + m^2} \right). \quad (5)$$

we get,  $v = \frac{2m^2+6}{m^2+1} = 2 + \frac{4}{m^2}$  Therefore,  $v$  is maximised when  $m = 0$ , giving the line  $y = c$ .

- E. Find  $m$  such that the projection error is minimised. **(20 marks)**

$$d^2 = AA'^2 + BB'^2 + CC'^2 \quad (6)$$

$$= \frac{m^2 + 1 + m^2}{m^2 + 1} \quad (7)$$

$$= 2 - \frac{1}{m^2 + 1}. \quad (8)$$

Note that we have set  $c = 0$  following question 3. Therefore,  $d$  is minimised by setting  $m = 0$ , giving the solution  $y = c$ .

- F. Compute the covariance matrix for this dataset. **(10 marks)**

Let  $\mathbf{x}_1 = (-1, 0)^\top$ ,  $\mathbf{x}_2 = (1, 0)^\top$ ,  $\mathbf{x}_3 = (0, 1)^\top$ . Then their mean  $\boldsymbol{\mu} = (0, 1/3)^\top$ . Variances are computed as,

$$(\mathbf{x}_1 - \boldsymbol{\mu})(\mathbf{x}_1 - \boldsymbol{\mu})^\top = \begin{bmatrix} 1 & 1/3 \\ 1/3 & 1/9 \end{bmatrix}, \quad (9)$$

$$(\mathbf{x}_2 - \boldsymbol{\mu})(\mathbf{x}_2 - \boldsymbol{\mu})^\top = \begin{bmatrix} 1 & -1/3 \\ -1/3 & 1/9 \end{bmatrix}, \quad (10)$$

$$(\mathbf{x}_3 - \boldsymbol{\mu})(\mathbf{x}_3 - \boldsymbol{\mu})^\top = \begin{bmatrix} 0 & 0 \\ 0 & 4/9 \end{bmatrix}. \quad (11)$$

Adding those three matrices and dividing by 3 we get the covariance matrix  $\mathbf{S}$  as follows:

$$\mathbf{S} = \begin{bmatrix} 2 & 0 \\ 0 & 2/3 \end{bmatrix}$$

- G. Find the eigenvalues and eigenvectors of the covariance matrix. **(10 marks)**

*Eigenvalue equation for  $\mathbf{S}$  is:*

$$\mathbf{S}\boldsymbol{\theta} = \lambda\boldsymbol{\theta} \quad (12)$$

$$|\mathbf{S} - \lambda\mathbf{I}| = 0 \quad (13)$$

*From which we get  $\lambda = 2, 2/3$ . The corresponding eigenvectors are respectively  $(1, 0)^\top$  and  $(0, 1)^\top$ .*

- H. Find the PCA projection using the eigenvalue decomposition of the covariance matrix. **(10 marks)**

*For PCA we must select the eigenvector corresponding to the largest eigenvalue as it maximises the variance of the projected data points. Therefore, we select  $(1, 0)^\top$ , which means that we select the  $x$ -axis, giving the solution  $y = c$ .*