

Proofs of some technical results

Sayantan Khan

July 2017

Contents

1	Blakers-Massey Theorem	2
2	Comparison theorem for cohomology theories	2
3	Brown's representability theorem	3
3.1	Spectra and cohomology theories	3
3.2	Proof of Brown's representability theorem	4
A	Definitions and notation	7
B	Some useful lemmas and theorems	7

1 Blakers-Massey Theorem

Fill in later

2 Comparison theorem for cohomology theories

Theorem 2.1 (Comparison Theorem). *If h and k are two reduced cohomology theories satisfying the (DV) axiom, such that there exists a natural transformation χ from h^* to k^* and χ induces a natural isomorphism from $h^n(S^0)$ to $k^n(S^0)$ for all n , then χ is a natural isomorphism of cohomology theories.*

Proof. First, we'll use a simple reduction. For any pointed space X , we'll show if χ is an isomorphism from $h^n(X_+)$ to $k^n(X_+)$, then χ is an isomorphism between $h^n(X)$ and $k^n(X)$. Here, X_+ is the space $X \amalg +$, with the basepoint being $+$. Consider the following cofiber sequence.

$$S^0 \xrightarrow{i} X_+ \rightarrow X$$

The map from S^0 to X_+ sends the basepoint of S^0 to $+$, and the other point to the original basepoint of X . This cofiber sequence splits, which means an isomorphism from $h^n(X_+)$ to $k^n(X_+)$ is equivalent to an isomorphism from $h(X)$ to $k(X)$ (by the five lemma).

The next step in the proof is to show the isomorphism for finite CW complexes. This will be done by showing the isomorphism for S^n and D^n for all n , and then using the fact that all finite CW complexes are pushouts of disks and spheres. We have an isomorphism on all spheres from the suspension isomorphism, and that extends to wedges of spheres. And since all disks are contractible, we have an isomorphism there as well. This also tells us that χ is an isomorphism from 0-dimensional CW complexes. We'll proceed by induction at this point. An n skeleton is defined by the following pushout.

$$\begin{array}{ccc} \coprod S_+^n & \longrightarrow & X_+^{(n-1)} \\ \downarrow & & \downarrow \\ \coprod D_+^n & \dashrightarrow & X_+^{(n)} \end{array}$$

We then apply the Mayer-Vietoris sequence to the subspaces $\coprod D_+^n$ and $X_+^{(n-1)}$. Call these subspaces T_1 and T_2 . Their intersection T_0 is the wedge of spheres.

$$\begin{array}{ccccccccc} h^{n+1}(T_1) \oplus h^{n+1}(T_2) & \longleftarrow & h^{n+1}(T_0) & \longleftarrow & h^n(X^{(n)}) & \longleftarrow & h^{n+1}(T_1) \oplus h^{n+1}(T_2) & \longleftarrow & h^{n+1}(T_0) \\ \downarrow \cong & & \downarrow \cong & & \downarrow \chi & & \downarrow \cong & & \downarrow \cong \\ k^{n+1}(T_1) \oplus k^{n+1}(T_2) & \longleftarrow & k^{n+1}(T_0) & \longleftarrow & k^n(X^{(n)}) & \longleftarrow & k^{n+1}(T_1) \oplus k^{n+1}(T_2) & \longleftarrow & k^{n+1}(T_0) \end{array}$$

By the induction hypothesis, all but the middle arrows are isomorphisms, hence by the five lemma, χ is an isomorphism too. This shows χ is a natural isomorphism for finite dimensional CW complexes.

The final step is to show this for infinite dimensional CW complexes. This will require the use of Milnor's theorem B.5. For any infinite dimensional CW complex X , take the filtration consisting of its finite dimensional skeletons. Then we have maps between the following short exact sequences.

$$\begin{array}{ccccccccc}
0 & \longrightarrow & \lim_n h^{q-1}(X^{(n)}) & \longrightarrow & h^q(X) & \longrightarrow & \lim_n h^q(X^{(n)}) & \longrightarrow & 0 \\
\downarrow \cong & & \downarrow \cong & & \downarrow \chi & & \downarrow \cong & & \downarrow \cong \\
0 & \longrightarrow & \lim_n k^{q-1}(X^{(n)}) & \longrightarrow & k^q(X) & \longrightarrow & \lim_n k^q(X^{(n)}) & \longrightarrow & 0
\end{array}$$

By the proof for the finite dimensional case, we have the all but the middle arrow are isomorphisms. By the five lemma, we get the middle arrow is an isomorphism too, which concludes the proof. \square

3 Brown's representability theorem

In this section, we shall see that all reduced cohomology theories that satisfy the wedge sum (DV) axiom are representable functors, i.e. they are naturally isomorphic to the hom functor in the homotopy category \mathbf{hCW}_* . In particular, for a given reduced cohomology theory h^* , we'll construct a sequence of spaces $\mathcal{Z}(n)$, which we'll call a spectrum, such that $\tilde{h}^n(X)$ is naturally isomorphic to $[X, \mathcal{Z}(n)]$.

3.1 Spectra and cohomology theories

Definition 3.1 (Ω -Spectrum). A spectrum is a \mathbb{Z} indexed sequence of pointed spaces $\mathcal{Z}(n)$ together with structure maps $\sigma_n : \Sigma \mathcal{Z}(n) \rightarrow \mathcal{Z}(n+1)$. If the adjoints of the structure maps, i.e. the maps $\tilde{\sigma}_n : \mathcal{Z}(n) \rightarrow \Omega \mathcal{Z}(n+1)$ are homotopy equivalences, then the spectrum is called an Ω -spectrum.

Proposition 3.1. Given a Ω -spectrum \mathcal{Z} , one can define the following functor.

$$\tilde{h}^n(X; \mathcal{Z}) = [X, \mathcal{Z}(n)]$$

This is a contravariant functor which satisfies the homotopy invariance (H), suspension (S), exactness (E), and the wedge sum (DV) axiom. It is therefore a reduced cohomology theory.

Proof. We'll deal with the axioms one at a time.

Homotopy invariance (H): This is obvious, because we are looking at homotopy classes of maps.

Suspension (S): We need to show there is a natural isomorphism from $\tilde{h}^n(X)$ to $\tilde{h}^{n+1}(\Sigma X)$. Note that the adjoint of the structure maps are homotopy equivalences. We therefore have a natural isomorphism.

$$[X, \mathcal{Z}(n)] \cong [X, \Omega \mathcal{Z}(n+1)]$$

On the other hand, since Σ and Ω are adjoints, we have the following natural isomorphism.

$$[X, \Omega \mathcal{Z}(n+1)] \cong [\Sigma X, \mathcal{Z}(n+1)]$$

Composing the two natural isomorphisms, we get our required isomorphism.

Exactness (E): We need to show for any cofibration $i : A \hookrightarrow X$, the following sequence is exact.

$$\tilde{h}^n(A) \leftarrow \tilde{h}^n(X) \leftarrow \tilde{h}^n(X/A)$$

Using the cofiber sequence, we get that following sequence is exact.

$$[A, \mathcal{Z}(n)] \leftarrow [X, \mathcal{Z}(n)] \leftarrow [(X/A), \mathcal{Z}(n)]$$

Wedge sum (DV): The functor $[\cdot, \mathcal{Z}(n)]$ satisfies (DV) axiom. This is fairly easy to check. That means \tilde{h}^* satisfies (DV) axiom.

□

3.2 Proof of Brown's representability theorem

In the previous section, we saw that if we are given an Ω -spectrum, we can construct a reduced cohomology theory using the spectrum. Brown's representability theorem is the converse of the previous theorem, i.e. given a reduced cohomology theory which satisfies the (DV) axiom, it can be represented by an Ω -spectrum, which is unique up to homotopy. This theorem is fairly technical, and will require the use of the theorem on Milnor exact sequence (theorem B.5).

Theorem 3.2 (Brown's representability theorem). *Let \tilde{h}^* be a reduced cohomology theory satisfying the (DV) axiom. Then there is an Ω -spectrum \mathcal{Z} such that \tilde{h}^n is naturally isomorphic to $[\cdot, \mathcal{Z}(n)]$.*

Proof. The proof will have two main parts. The first part will involve constructing the spaces $\mathcal{Z}(n)$ for each n such that there is a natural isomorphism from $\tilde{h}^n(X)$ to $[X, \mathcal{Z}(n)]$ for all CW complexes X . The second part will involve constructing the structure maps from $\Sigma \mathcal{Z}(n) \rightarrow \mathcal{Z}(n+1)$.

Fix an $n \in \mathbb{Z}$. We will construct the space $\mathcal{Z}(n)$ as a CW complex, using finite dimensional skeletons $\mathcal{Z}(n)_k$. For each k , we will also pick a cohomology class $c_n(k)$ in $\tilde{h}^n(\mathcal{Z}(n)_k)$ such that the map $d_n^m(k) : [S^m, \mathcal{Z}(n)_k] \rightarrow \tilde{h}^n(S^m)$ is an isomorphism for $m < k$ and surjection for $m = k$.

$$\begin{aligned} d_n^m(k) &: [S^m, \mathcal{Z}(n)_k] \rightarrow \tilde{h}^n(S^m) \\ d_n^m(k) &: [f] \mapsto f^*(c_n(k)) \end{aligned}$$

To construct easy examples of spectra, one needs to check that the filtered colimits commute with the loop space functor, at least for nice enough spaces.

Show that this is a group homomorphism for $m \geq 1$

For $k = 0$, we define $\mathcal{Z}(n)_0$ as follows.

$$\mathcal{Z}(n)_0 := \bigvee_{\alpha \in \tilde{h}^n(S^0)} S_\alpha^0$$

The cohomology group of $\mathcal{Z}(n)_0$ is given by a direct product, since \tilde{h}^n satisfies the (DV) axiom.

$$\tilde{h}^n(\mathcal{Z}(n)_0) \cong \prod_{\alpha \in \tilde{h}^n(S^0)} \tilde{h}^n(S_\alpha^0)$$

Pick the following element as $c_n(0)$.

$$c_n(0) := \prod_{\alpha \in \tilde{h}^n(S^0)} \alpha$$

Since $k = 0$, we only need to show that $d_n^0(0)$ is a surjection. Pick any $\alpha \in \tilde{h}^n(S^0)$. Corresponding to this α , there's a copy of S^0 sitting inside $\mathcal{Z}(n)_0$. Let f be the inclusion map of this copy of S^0 into $\mathcal{Z}(n)_0$. Then the induced map on cohomology is the projection map on the α^{th} coordinate, since the cohomology theory satisfies the (DV) axiom. Applying this induced map on $c_n(0)$, we see that in the α^{th} coordinate, it has α , because of the way we defined it. This shows the map is surjective.

To prove the induction step, suppose we have defined the space $\mathcal{Z}(n)_k$ and $c_n(k)$ that satisfy the required properties. Let $K_k \trianglelefteq [S^k, \mathcal{Z}(n)_k]$ be the kernel of the map $d_n^k(k)$. We construct the following map.

$$\phi_n(k) : \bigvee_{x \in K_k} S^k \rightarrow \mathcal{Z}(n)_k \vee \bigvee_{y \in \tilde{h}^n(S^{k+1})} S^{k+1}$$

The cofibration probably works if you take a subset of K_k that does not contain the constant map

This map is obtained by taking the wedge of maps from S^k to $\mathcal{Z}(n)_k$ which are contained in K_k . This is a cofibration. By the (DV) axiom, we have the following cohomology groups.

$$\tilde{h}^n \left(\mathcal{Z}(n)_k \vee \bigvee_{y \in \tilde{h}^n(S^{k+1})} S^{k+1} \right) = \tilde{h}^n(\mathcal{Z}(n)_k) \times \prod_{y \in \tilde{h}^n(S^{k+1})} \tilde{h}^n(S^{k+1})$$

Not sure how to show this, or whether this is entirely correct. Need to check later

From this, we can immediately see that the elements of $\tilde{h}^n \left(\mathcal{Z}(n)_k \vee \bigvee_{y \in \tilde{h}^n(S^{k+1})} S^{k+1} \right)$ of the form $(c_n(k), \bullet)$ (where \bullet is any arbitrary element) is in the kernel of $\phi_n^*(k)$. Define $\mathcal{Z}(n)_{k+1}$ to be the cofiber of the map $\phi_n(k)$, and let the map to the cofiber be $b_n(k)$. By the exactness axiom, we have that the following sequence is exact.

$$\tilde{h}^n(\mathcal{Z}(n)_{k+1}) \xrightarrow{b_n^*(k)} \tilde{h}^n(\mathcal{Z}(n)_k) \times \prod_{y \in \tilde{h}^n(S^{k+1})} \tilde{h}^n(S^{k+1}) \xrightarrow{\phi_n^*(k)} \prod_{x \in K_k} \tilde{h}^n(S^k)$$

Pick the following element $A \in \prod_{y \in \tilde{h}^n(S^{k+1})} \tilde{h}^n(S^{k+1})$.

$$A := \prod_{\alpha \in \tilde{h}^n(S^{k+1})} \alpha$$

The element $(c_n(k), A)$ lies in the kernel of $\phi_n^*(k)$, which means it lies in the image of $b_n^*(k)$. We define $c_n(k+1)$ to be a pre-image of $(c_n(k), A)$. Seeing that the associated map $d_n^m(k+1)$ is surjective for $m = k+1$ is easy enough. The proof is the same as that in the case of $d_n^0(0)$. The trickier part is showing injectivity for $m < k+1$. Since $d_m^n(k)$ is a group homomorphism (for $M \geq 1$), for $m \geq 1$, it will suffice to show the kernel is trivial. Pick an element, say $[f]$ in the kernel. We need to show that f is a nullhomotopic map. But at each step, we coned off the kernel of $d_n^m(m)$. That means f is nullhomotopic. This shows the injectivity and hence the isomorphism for $m < k+1$.

Next, we define $\mathcal{Z}(n)$ the colimit of the following diagram.

$$\mathcal{Z}(n)_0 \xrightarrow{b_n(0)} \mathcal{Z}(n)_1 \xrightarrow{b_n(1)} \mathcal{Z}(n)_2 \xrightarrow{b_n(2)} \dots$$

Note that $\mathcal{Z}(n)_k$ are CW subcomplexes of $\mathcal{Z}(n)$, in particular, we can appeal to Milnor's theorem B.5, i.e. the following sequence is exact.

$$0 \rightarrow \lim_k^1 \tilde{h}^{n-1}(\mathcal{Z}(n)_k) \rightarrow \tilde{h}^n(\mathcal{Z}(n)) \rightarrow \lim_k \tilde{h}^n(\mathcal{Z}(n)_k) \rightarrow 0$$

Furthermore the element $(c(n)_0, c(n)_1, c(n)_2, \dots)$ lies in $\lim_k \tilde{h}^n(\mathcal{Z}(n)_k)$ since we pick $c(n)_{k+1}$ as a preimage of $c(n)_k$. By exactness, we get a preimage c_n in $\tilde{h}^n(\mathcal{Z}(n))$. We define a map $d_n^m : [S^m, \mathcal{Z}(n)] \rightarrow \tilde{h}^n(S^m)$ which sends $[f]$ to $f^*(c_n)$. Because of the inductive construction, we know this is an isomorphism for all $m \geq 0$ (To see this, observe that a map from a compact space like S^m factors through a finite stage in the colimit $\mathcal{Z}(n)$). This can be extended to a natural isomorphism for all finite CW complexes. Now that we know how to represent the individual functors \tilde{h}^n , we need to construct the structure maps from the suspension homomorphism of the cohomology theory. Let T_n be the suspension homomorphism from $\tilde{h}^n(X)$ to $\tilde{h}^{n+1}(\Sigma X)$. If we set X to be $\mathcal{Z}(n)$, we have the following homomorphism.

$$T_n : \tilde{h}^n(\mathcal{Z}(n)) \rightarrow \tilde{h}^{n+1}(\Sigma \mathcal{Z}(n+1))$$

But this is equivalent to the following homomorphism.

$$\tilde{T}_n : [\mathcal{Z}(n), \mathcal{Z}(n)] \rightarrow [\Sigma \mathcal{Z}(n), \mathcal{Z}(n+1)]$$

We do the most obvious thing, i.e. apply \tilde{T}_n to the homotopy class of the identity map, and we pick a map to be our structure map from the resulting homotopy class. □

This result enables us to study any reduced cohomology theory by studying its associated spectrum. This lets us study many cohomology theories that were intractable by the usual methods, e.g. cobordism, which is represented by the Thom spectrum.

Maybe write the proof. I'm skipping the proof of the case when $m = 0$. I think it should be fairly easy once I figure out why the map is a cofibration.

Need to prove this, but can't think of a proof yet. The five lemma might help

Show that this forms an Ω -spectrum

A Definitions and notation

Definition A.1 (Suspension of a pointed space). The suspension ΣX of a pointed space X is the smash product $S^1 \wedge X$.

Definition A.2 (Loop space of a pointed space). The loop spaces ΩX of a pointed space X is the set of all pointed maps from S^1 to X with the compact-open topology.

Definition A.3 (\lim^1). Let T be the category of towers of abelian groups, i.e. \mathbb{N} indexed set of abelian groups G_i with maps $f_i : G_i \rightarrow G_{i-1}$, and maps are set of arrows that make the whole thing commute. Then \lim is a left exact functor from T to AbGrp , and we define \lim^1 to be the first right derived functor of \lim .

Check that this category has enough injectives

B Some useful lemmas and theorems

Note: Although we state many of the lemmas here for TOP , they are also true for TOP_* , and the proof is similar.

Lemma B.1. If $i : A \hookrightarrow X$ is a cofibration (in the category TOP), then the mapping cone $C(i)$ is homotopy equivalent to X/A .

Proof. We will first construct the maps to and from $C(i)$ to X/A . The maps from $C(i)$ to X/A is the map that collapses the cone of A to a point corresponding to A in X/A . Now consider a map from $H : A \times I$ to $C(i)$, such that H contracts A to a point in $C(i)$, starting from the inclusion of A in X . Let the map j from X to $C(i)$ be the inclusion map. Since i is a cofibration, we can extend H with the initial condition j to a map $J : X \times I \rightarrow C(i)$. But $J(\cdot, 1)$ collapses A to a point. That means it factors through a X/A . This gives us a map k from X/A to $C(i)$.

The fact that these maps are homotopy inverses can be verified using the homotopy J . □

Not sure of this. Verify later.

Lemma B.2. In the category TOP , the following sequence is h -coexact.

$$A \xrightarrow{f} B \xrightarrow{i} C(f)$$

That means for any space Z , the following sequence of abelian groups is exact.

$$[A, Z] \leftarrow [B, Z] \leftarrow [C(f), Z]$$

Proof. If an element $[c] \in [B, Z]$ goes to 0 in $[A, Z]$, that means $c \circ f : A \rightarrow Z$ is nullhomotopic, where c is a representative of $[c]$. But that means there is some function $d \in C(f)$ such that $c = d \circ i$. This shows the exactness of the sequence. □

Lemma B.3. If K is a compact space, let A_i be a sequence of spaces where points are closed, and A is the colimit of the following diagram:

$$A_0 \hookrightarrow A_1 \hookrightarrow A_2 \hookrightarrow \dots$$

where all the embeddings are closed, then a map from K to A factors finitely through some A_i .

Proof. Let $J = f(K)$ be the compact image of K in A . For each set $A_i \setminus A_{i-1}$, pick an element c_i of J in the set, if J intersects $A_i \setminus A_{i-1}$. Since A_i 's are closed, that means the subset c_i has the discrete topology. Furthermore, since points are closed, the set $\{\cup c_i\}$ is a closed subset of J , hence compact. And compact spaces with discrete topology are finite. That means only finitely many $A_i \setminus A_{i-1}$ intersect J . This means the map factors through at some finite stage. \square

Theorem B.4 (Alternative characterization of \lim^1). *If F is an object in the tower category, then $\lim^1(F)$ is the cokernel of the following map.*

$$\alpha_F : \prod_{i \in \mathbb{N}} F_i \rightarrow \prod_{i \in \mathbb{N}} F_i$$

$$\alpha_F : (g_0, g_1, g_2, \dots) \mapsto (g_0 - f_1(g_1), g_1 - f_2(g_2), \dots)$$

Proof. The first step in characterizing \lim^1 in the following manner is to pick an appropriate injective resolution. Let F be a tower of abelian groups and I an injective tower it maps into via a monomorphism m .

$$\begin{array}{ccccccc} F_0 & \xleftarrow{f_1} & F_1 & \xleftarrow{f_2} & F_2 & \xleftarrow{f_3} & F_3 & \xleftarrow{f_4} & \dots \\ \downarrow m_0 & & \downarrow m_1 & & \downarrow m_2 & & \downarrow m_3 & & \\ I_0 & \xleftarrow{i_1} & I_1 & \xleftarrow{i_2} & I_2 & \xleftarrow{i_3} & I_3 & \xleftarrow{i_4} & \dots \end{array}$$

Without losing any generality, we can assume all the maps i_k in I are surjective. Otherwise, we just replace I_k by $\bigoplus_{j=0}^k I_j$, and have the maps on all but the last coordinate be the identity. This is important, because we'll need surjectivity of the maps later. We can now construct an injective resolution of F as the following exact sequence.

$$0 \rightarrow F \xrightarrow{m} I \xrightarrow{q} \text{coker}(m) \rightarrow 0$$

The first derived functor is the homology at $\lim(\text{coker}(m))$ of the following sequence.

$$0 \rightarrow \lim(F) \xrightarrow{\lim(m)} \lim(I) \xrightarrow{\lim(q)} \lim(\text{coker}(m)) \rightarrow 0$$

Now, just like α_F was defined in the statement of the theorem, we define α_I and $\alpha_{\text{coker}(m)}$. Then we get the following short exact sequence of chain complexes (the rows are exact).

$$\begin{array}{ccccccc} & & 0 & & 0 & & 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \prod_i F_i & \xrightarrow{m} & \prod_i I_i & \xrightarrow{q} & \prod_i \text{coker}(m) \longrightarrow 0 \\ & & \alpha_F \downarrow & & \alpha_I \downarrow & & \alpha_{\text{coker}(m)} \downarrow \\ 0 & \longrightarrow & \prod_i F_i & \xrightarrow{m} & \prod_i I_i & \xrightarrow{q} & \prod_i \text{coker}(m) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ & & 0 & & 0 & & 0 \end{array}$$

We can apply the snake lemma to get the following long exact sequence.

$$\begin{array}{ccccccc}
0 & \longrightarrow & \ker(\alpha_F) & \xrightarrow{m_*} & \ker(\alpha_I) & \xrightarrow{q_*} & \ker(\alpha_{\text{coker}(m)}) \\
& & & & \searrow \partial & & \nearrow \\
& & \text{coker}(\alpha_F) & \xrightarrow{m^*} & \text{coker}(\alpha_I) & \xrightarrow{q^*} & \text{coker}(\alpha_{\text{coker}(m)}) \longrightarrow 0
\end{array}$$

But we see from the definition of \lim that the kernels of α are precisely the \lim . Thus we have the following long exact sequence.

$$\begin{array}{ccccccc}
0 & \longrightarrow & \lim(F) & \xrightarrow{\lim(m)} & \lim(I) & \xrightarrow{\lim(q)} & \lim(\text{coker}(m)) \\
& & & & \searrow \partial & & \nearrow \\
& & \text{coker}(\alpha_F) & \xrightarrow{m^*} & \text{coker}(\alpha_I) & \xrightarrow{q^*} & \text{coker}(\alpha_{\text{coker}(m)}) \longrightarrow 0
\end{array}$$

The last step in the proof will be to show that $\text{coker}(\alpha_I)$ is 0, in which case $\lim^1(F)$ is isomorphic to $\text{coker}(\alpha_F)$. Showing that $\text{coker}(\alpha_I)$ is 0 is equivalent to showing that α_I is surjective. To see this, pick any element $(j_0, j_1, j_2, \dots) \in \prod_i I_i$. We need to find an element $(k_0, k_1, k_2, k_3, \dots)$ such that we have the following equalities.

$$\begin{aligned}
j_0 &= k_0 - i_1(k_1) \\
j_1 &= k_1 - i_2(k_2) \\
j_2 &= k_2 - i_3(k_3) \\
&\vdots
\end{aligned}$$

But notice that we constructed I such that all the i_k are surjective. That means this system of equations can be solved simultaneously and $\text{coker}(\alpha_I)$ is 0. This shows the result. \square

Theorem B.5 (Milnor exact sequence). *If $\{i_n : X_n \hookrightarrow X_{n+1}\}$ for $n \in \mathbb{N}$ are a sequence of nested (pointed) CW subcomplexes such that $X = \bigcup_n X_n$, and \tilde{h}^* is a reduced cohomology theory, then we have the following exact sequence for all $i \geq 1$.*

$$0 \rightarrow \lim_n^1 \tilde{h}^{i-1}(X_n) \rightarrow \tilde{h}^i(X) \rightarrow \lim_n \tilde{h}^i(X_n) \rightarrow 0$$

Proof. We first construct the mapping telescope of the inclusions. We start with the (unpointed) space $X \times \mathbb{R}_+$. We then consider the following subspace S .

$$S = \bigcup_{i \in \mathbb{N}} X_i \times [i, i+1]$$

We then quotient S by the subspace $* \times \mathbb{R}_+$ and we use the quotient out space as the base-point of the pointed mapping telescope T .

We define two subspaces of the mapping telescope; we'll apply Mayer-Vietoris to these subspaces.

$$\begin{aligned}
T_1 &:= \bigcup_{i \in \mathbb{N}} X_{2i+1} \times [2i+1, 2i+2] \\
T_2 &:= \bigcup_{i \in \mathbb{N}} X_{2i} \times [2i, 2i+1]
\end{aligned}$$

To be more precise, we should be taking open neighbourhoods of these spaces to apply Mayer-Vietoris, but the open neighbourhoods deformation retract to these spaces anyways, so it's alright. Observe that the intersection T_0 of T_1 and T_2 is homeomorphic to the wedge of all the X_i . By similar reasoning (although we replace homeomorphism by homotopy equivalence), we get that T_1 is homotopy equivalent to the wedge of all the X_i for odd i and T_2 the wedge of X_i for all even i . We consider the Mayer-Vietoris sequence, and use the (DV) axiom to get isomorphisms.

$$\begin{array}{ccccccc}
\tilde{h}^{q+1}(T) & \longleftarrow & \tilde{h}^q(T_0) & \xleftarrow{\phi_q} & \tilde{h}^q(T_1) \oplus \tilde{h}^q(T_2) & \longleftarrow & \tilde{h}^{q+1}(T) \\
& & \downarrow \cong & & \downarrow \cong & & \\
& & \prod_{i \in \mathbb{N}} \tilde{h}^q(X_i) & & \prod_{i \in \mathbb{N}} \tilde{h}^q(X_{2i}) \oplus \prod_{i \in \mathbb{N}} \tilde{h}^q(X_{2i+1}) & & \\
& \swarrow \cdot (-1)^i & & \searrow m_q & & & \\
\prod_{i \in \mathbb{N}} \tilde{h}^q(X_i) & & & & & &
\end{array}$$

We need to determine how the map m_q should be defined to make the diagram commute. We do this by seeing how the map behaves on the basis. First, we take $x_0 \in \tilde{h}^q(X_0)$. The first isomorphism takes it to $(x, 0)$, the map ϕ_q takes it to $x - 0$ and finally it goes to x . Next, we see how $x_1 \in \tilde{h}^q(X_1)$ behaves. It first gets sent to $(i_0^* x_1, x_1)$ and then to $i_0^* x_1 - x_1$. Finally, the $(-1)^i$ sends it to $x_1 - i_0^* x_1$. Similarly, $x_2 \in \tilde{h}^q(X_2)$ gets sent to $x_2 - i_1^* x_2$, and so go on the rest of the basis elements. We now have the following long exact sequence.

$$\tilde{h}^{q+1}(T) \longleftarrow \prod_{i \in \mathbb{N}} \tilde{h}^q(X_i) \xleftarrow{m_q} \prod_{i \in \mathbb{N}} \tilde{h}^q(X_i) \longleftarrow \tilde{h}^q(T)$$

From this, we get the following short exact sequence.

$$0 \longleftarrow \ker(m_q) \longleftarrow \tilde{h}^q(T) \longleftarrow \operatorname{coker}(m_{q-1}) \longleftarrow 0$$

From theorem B.4, we see that $\ker(m_q)$ is $\lim_n (\tilde{h}^q(X_n))$ and $\operatorname{coker}(m_{q-1})$ is $\lim_n^1 (\tilde{h}^{q-1}(X_n))$. This proves the theorem. \square