# THE VARIABILITY CLIMATE CHANGE IS RESPONSIBLE FOR IN VEGETATION LOSS IN GHANA

Quantifying The Status of Galamsey With Time Series Analysis

# KALONG BONIFACE FUGAH SELETEY MITCHELL SUPERVISER: MR JUSTIES AMAYOR KESSIE

Department of Mathematics and Statistics University Of Energy and Natural Resource, Sunyani

October 9, 2022



# Outline Of Presentation

- Introduction
- Problem Statement
- Objective
- Methodology
- Reference





### INTRODUCTION

One would anticipate that the majority of emerging nations, which are still in the early stages of economic development and growth, would have a high forest cover and little deforestation. This, however, has not been the case. Ghana is a lower-middle-income nation that is still working toward middle-income classification. However, it has already begun to see a deforestation rate that is comparable to that of middle-income countries. The rapid population expansion, clearing of field for Galamsey operation, increased domestic need of wood for things like fuel, furniture, construction, and timber exports have all contributed to this trend, Bush fires in the 1980s, climate change, and lax law enforcement have all had an impact.



3/14

#### Introduction Con't

The purpose of this paper is to establish an understanding in time series analysis on remotely sensed data. Which will introduced us to the fundamentals of time series modeling, including decomposition, autocorrelation and modeling historical changes in Galamsey Operation in Ghana, the Cause, Dangers and it's Environmental impact



#### Introduction Con't

Galamsey also known as "gather them and sell", Owusu-Nimo2018 is the term given by local Ghanaian for illegal small-scale gold mining in Ghana . The major cause of Galamsey is unemployment among the youth in Ghana Gracia2018. Young university graduates rarely find work and when they do it hardly sustains them. The result is that these youth go the extra mile to earn a living for themselves and their family.



#### PROBLEM STATEMENT

The Footprint of Galamsey is Spreading at a very faster rate, causing vegetation loss. Other factors accounting to vegetation loss may largely include climate change, urban and exurban development, bush fires. But not much works or research has been done to tell the extent to which Galamsey causes vegetation loss. This research attempts to segregate the variability climate is responsible for in vegetation loss so as to attribute the residual variability to Galamsey and other related activities such as bush-fires etc.





#### Research Questions

To address the challenge of the vegetation variability in this work, the following several statements were formed:

- Are there any changes in vegetation cause by Galamsey and Climate change in Ghana?
- Is there any relationship between vegetation and land surface temperature in Ghana?





Group Nine (UENR) Proposal October 9, 2022 7 / 14

#### **OBJECTIVE**

The purpose is to establish an understanding in time series analysis on remotely sensed data. We will be introduced to the fundamentals of time series modeling, including decomposition, auto-correlation and modeling historical changes.

- Perform time series analysis on satellite derived vegetation indices
- Estimate the extent to which Galamsey causes vegetation loss in Ghana.
- Dissociate or single out the variability climate is responsible for in vegetation loss





## Significance Of The Study

In this study, we want to examine the effects of climatic change on Ghana's vegetation during these thirty years.

- This study allows us to explore climatic differences and climate-related drivers.
- It offers a chance to research how climatic variability affects the ecosystem and human health.
- This study explores historical and projected vegetation and climate data, by sector, impacts, key vulnerabilities and what adaptation measures can be taken.
- It also explores the overview for a general context of how climate change is affecting Ghana.



# Limitation Of The Study

The goal of time series modeling is to employ the simplest model feasible to account for as much data as possible while still developing an explanatory model of the data that does not over-fit the issue set.

- Missing Data
- It is almost certain that data from distant sensing will not provide the same level of precision.
- Atmospheric factors can distort the visual findings, causing the vegetation's color to shift dramatically from image to image as a result of atmospheric factors (fog, ground moisture, cloud cover)





#### Data

The Satellite (MOD13) provide consistent, and spatial time series comparisons of global vegetation conditions that can be used to monitor the Earth's photosynthetic vegetation activity in support of **change detection**, and biophysical interpretations. For this analysis, we will be using the MOD13Q1 V6 product which contains the Enhanced Vegetation Index (EVI). Calculated as: and;










Figure:

Instead of analyzing the imagery directly, we will summarize the mean EVI values. We will apply a smoothing strategy using an ARIMA function to fix the situation where some cells may not have EVI for a particular month. Once NA values have been eliminated, the time series will be divided to eliminate seasonality before the normalized data is fitted using a linear model. We will go to classify our data on the map and map it after we have extracted the linear trend.

# Methodology

we will first collect data from Google Earth Engine in order to choose EVI values and Climate Change data.




Figure:



13 / 14

# Methodology Cont'

- \* To help machine learning classifiers works better with time series data .
- \* Local features or patterns in time series can be found and combined to address challenges involving time-series categorization.
- \* Then, a method to discover patterns that are helpful for classification is suggested.
- \* combine these patterns to create computable categorization rules.

#### **Exponential Family**

It is of the form  $f(y) = c(y, \phi) exp \frac{y\theta - a(\theta)}{\phi}$  where:

- $\theta$  and  $\phi$  are parameters. The parameters  $\theta$  is called the canonical parameter and  $\phi$  the dispersion parameter.
- The choice of a(t) and  $c(y, \phi)$  determine the actual probability function such as the binomial, normal, gamma, etc.

Group Nine (UENR) Proposal October 9, 2022 14/14