



UNIVERSIDAD DE CHILE
FACULTAD DE CIENCIAS FÍSICAS Y MATEMÁTICAS
DEPARTAMENTO DE ASTRONOMÍA

**STUDY OF THE ROLE OF GAS IN THE FORMATION OF SMBH SEED, THROUGH
SIMULATIONS OF NSC WITH GAS EMBEDDED**

TESIS PARA OPTAR AL GRADO DE MAGÍSTER EN CIENCIAS, MENCIÓN ASTRONOMÍA

BORIS CUEVAS GOMEZ

PROFESOR GUÍA:
ANDRÉS ESCALA

MIEMBROS DE LA COMISIÓN:
BASTIÁN REINOSO
RICARDO MUÑOZ
WALTER MAX-MOERBECK

Este trabajo ha sido parcialmente financiado por:
CENTRO DE ASTROFÍSICA Y TECNOLOGÍAS AFINES

SANTIAGO DE CHILE

2026

RESUMEN DE LA TESIS PARA OPTAR
AL GRADO DE MAGÍSTER EN CIENCIAS,
MENCIÓN ASTRONOMÍA
POR: BORIS CUEVAS GOMEZ
FECHA: 2026
PROF. GUÍA: ANDRÉS ESCALA

STUDY OF THE ROLE OF GAS IN THE FORMATION OF SMBH SEED, THROUGH SIMULATIONS OF NSC WITH GAS EMBEDDED

Observaciones de cuásares a alto redshift imponen fuertes restricciones sobre los mecanismos de formación de agujeros negros supermasivos (SMBHs). En consecuencia, varias vías han sido propuestas para explicar la formación de estos objetos masivos (con masas del orden de $\geq 10^6 M_\odot$) en el Universo temprano. Entre ellas, el escenario de colisiones descontroladas en cúmulos estelares nucleares parece ser prometedor para generar semillas de agujeros negros, que posteriormente podrían crecer hasta el rango de supermasivo. En este trabajo, presentamos un conjunto de simulaciones de cúmulos estelares nucleares con una componente gaseosa, para estudiar el efecto del gas en la formación de semillas de agujeros negros supermasivos. Exploramos un escenario que cuantifica el papel de las colisiones y de los procesos de relajación. Acoplamos un código SPH con un código de N cuerpos, ambos del proyecto AMUSE, usando el método BRIDGE. El código SPH resuelve las ecuaciones hidrodinámicas del gas, mientras que el código de N cuerpos resuelve la dinámica de las estrellas. Para una escala de tiempo característica t_H , la comparación con las escalas de tiempo de colisión t_{coll} y relajación t_{relax} define tres regímenes dinámicos, basados en la estabilidad frente a las colisiones. Mostramos que para simulaciones en el régimen intermedio, definido por $t_{\text{relax}} \leq t_{\text{coll}} \leq t_H$, la eficiencia de formación de agujeros negros es $\gtrsim 20\%$, y en nuestro modelo más masivo, se forma un agujero negro de masa intermedia con $\gtrsim 10^4 M_\odot$. Nuestros hallazgos muestran que el gas mejora la formación de semillas de agujeros negros supermasivos, ya que incrementa la masa final del objeto más masivo formado, además de acortar el tiempo de colapso del núcleo del cúmulo.

RESUMEN DE LA TESIS PARA OPTAR
AL GRADO DE MAGÍSTER EN CIENCIAS,
MENCIÓN ASTRONOMÍA
POR: BORIS CUEVAS GOMEZ
FECHA: 2026
PROF. GUÍA: ANDRÉS ESCALA

STUDY OF THE ROLE OF GAS IN THE FORMATION OF SMBH SEED, THROUGH SIMULATIONS OF NSC WITH GAS EMBEDDED

Observations of quasars at high redshift impose strong constraints on the formation mechanisms of supermassive black holes (SMBHs). Consequently, several pathways have been proposed to explain the formation of such massive objects ($\geq 10^6 M_\odot$) in the early Universe. Among them, runaway collisions in nuclear star clusters (NSCs) are a promising channel for SMBH seed formation. In this work, we present a set of simulations of gas-embedded nuclear star clusters to study the effect of gas on the formation of a supermassive black hole seed. We explore a scenario that quantifies the role of collisions and the interplay with relaxation processes. We couple an SPH code with a pure N-body code, both from the AMUSE framework, using the BRIDGE method. The SPH code solves the hydrodynamics equations for the gas, whereas the N-body code is used to solve the equations for the dynamics of the stars. For a characteristic timescale t_H , the interplay between the collision timescale t_{coll} and the relaxation timescale t_{relax} defines three dynamical regimes based on the stability under collisions. We show that for simulations in the intermediate regime, defined by $t_{\text{relax}} \leq t_{\text{coll}} \leq t_H$, the black hole formation efficiency is $\gtrsim 20\%$, and in our most massive models, an IMBH of $\gtrsim 10^4 M_\odot$ is formed. Our findings show that gas enhances the formation of a SMBH seeds by increasing the mass of the final most massive object and shortening the timescale for core collapse of the star cluster.

Dedicado a mis padres, y mis dos hermanas, por su amor incondicional.

Agradecimientos

Quiero agradecer a mi profesor guía, Andrés Escala, por su apoyo en este proyecto. También a los coautores del documento científico producto de este trabajo, que sumados a mi profesor guía son Bastián Reinoso, Marcelo Vergara, Dominik Schleicher, y Sebastián Valdebenito, por su participación y comentarios de dicho documento. En particular Bastián, quien ha participado activamente durante todo mi proyecto de tesis.

También quisiera agradecer a los profesores Ricardo Muñoz y Walter Max-Moerbeck por aceptar ser parte de mi comité.

También quisiera agradecer a mis grupos de amigos, tanto del colegio, del pregrado y posteriormente dentro del magíster, por ofrecerme su desinteresada amistad, y su apoyo en momentos no tan buenos.

También quisiera agradecer a mi familia. A mis padres, Nelson y Jacqueline, por su amor incondicional y paciencia infinita por todo este tiempo. A mis hermanas, Daniela y Jacqueline, que al igual que mis padres me han apoyado de manera incondicional y me han abierto las puertas de sus hogares. A mis cuñados, Pablo y Lucas, a quienes considero mis hermanos postizos, y que me han regalado hermosos sobrinos.

Finalmente agradecer al soporte financiero del CATA, que a través del proyecto BASAL ANID FB21003 han financiado parte de esta investigación.

Table of Contents

1. Introduction	1
1.1. Brief history: from Quasars to SMBHs	2
1.2. Formation channels for SMBHs	4
1.2.1. Population III stars	6
1.2.2. Direct collapse of primordial gas clouds	6
1.2.3. Runaway collisions in dense star clusters	7
1.3. Research Objectives and Thesis Outline	7
2. A scenario for SMBH formation	9
2.1. Critical mass	10
2.2. Relaxation mass	11
2.3. Global instability scenario	12
3. Numerical Methods	15
3.1. The AMUSE Framework	15
3.2. N-body code: PH4	16
3.2.1. Hermite integrator	17
3.2.2. Block time steps	19
3.2.3. Collisions and mergers	20
3.2.4. Escaping stars	21
3.3. SPH code: Fi	23
3.3.1. Barnes-Hut tree	23
3.3.2. Smoothed Particle Hydrodynamics	25
3.4. Coupling Strategy: The BRIDGE Method	28
3.4.1. Symplectic Integrator	28
3.4.2. BRIDGE	30
3.4.3. Adaptive timestep	32
4. Initial Conditions and Setup	34
4.1. Plummer distribution	34
4.1.1. N-body units	36
4.2. Stars and Gas Initial Conditions	38

4.2.1. Particles generation	40
4.3. Numerical Parameters	41
4.3.1. Ph4	43
4.3.2. Fi	44
4.3.3. Bridge	45
4.3.4. Code validation	45
5. Results	48
5.1. Statement about my contribution	48
6. Conclusions	67
6.1. Caveats	68
References	69
Appendices	76
A. Derivation of Critical Masses for Collisions and Relaxation	76
A.1. Critical Mass in the Simplified Case	76
A.2. Effect of an External Gravitational Potential	77
A.3. Generalized Critical Mass Including Gravitational Focusing	79
A.4. Derivation of the Relaxation Mass	81
B. Fourth-Order Hermite Integration Scheme	86
B.1. Step 1: Particle selection	86
B.2. Step 2: Calculate the predicted position and velocities for all particles . . .	87
B.3. Step 3: Calculate acceleration and jerk for particle i	88
B.4. Step 4: Second and third time derivative of acceleration, using Hermite interpolation	88
B.5. Step 5: Add corrections	90
B.6. Step 6: Time step update	91
C. SPH equations	91
C.1. Solving λ_i	92
C.2. Equation of motion for SPH particles	94
C.3. Equation of energy for SPH particles	96

List of Tables

4.1.	Initial conditions for the stellar systems. R_v , M , N , m_s , r_s , and σ denote the virial radius, total stellar mass, number of stars, stellar mass, stellar radius, and velocity dispersion, respectively.	38
4.2.	Initial conditions for gas systems. q , M_{SPH} , N_{SPH} , m_{SPH} , and c_s denote the gas mass fraction, total gas mass, number of SPH particles, SPH particle mass and mean sound speed, respectively.	39
4.3.	Key runtime parameters for the PH4 N-body code. Parameters shown with their default and the values used in this work.	44
4.4.	Key runtime parameters for the Fi SPH code. Parameters shown with their default and the values used in this work.	45
4.5.	Key runtime parameters for the BRIDGE coupling scheme. GPU acceleration is enabled via <code>mode=1</code> for the FASTKICK force calculations. The N-body and SPH components run on CPU (PH4 with 4 workers, Fi with 8 workers). The parameter <code>do_accretion</code> is set to <code>False</code> in our simulations, as we do not include gas accretion.	46

List of Figures

1.1.	Image of 3C 273 from Hubble data. Credits: https://science.nasa.gov/asset/hubble/quasar-3c-273-2/	2
1.2.	Simulated images of Sgr A [*] . The left panel is a snapshot of a numerical simulation of Sgr A [*] , and middle panel is the average of snapshots of the simulation. On the right panel, a representative image reconstruction, using the synthetic visibilities from the simulation. Credits: EHT Collaboration et al. (2022).	4
1.3.	BH abundance (number density) in cMpc ⁻³ (y-axis) vs redshift (x-axis). The y-axis, from top to bottom, goes from very rare ($\sim 10^{-9}$) to common ~ 1 . Credits: Regan and Volonteri (2024).	5
2.1.	Collision and relaxation timescales in a mass versus radius diagram. The blue dashed line correspond to the critical mass M_{crit} , while blue solid line correspond to the relaxation mass M_{relax} , both calculated for clusters formed by solar mass stars, characteristic time $t_{\text{H}} = 10^{10}$ yr and $q = 0$. The black dash-dotted line represents $t_{\text{coll}} = t_{\text{relax}}$, with t_{H} varying from 10^2 yr to 10^{10} yr and $q = 0$. Clusters above the black dash-dotted line satisfy $t_{\text{coll}} \leq t_{\text{relax}}$. The unstable regime (UR) is defined by $t_{\text{coll}} \leq t_{\text{H}} \leq t_{\text{relax}}$ (red colored zone); the stable regime is where $t_{\text{relax}} \leq t_{\text{H}} \leq t_{\text{coll}}$ (gray colored zone); and the intermediate regime (IR) fulfills $t_{\text{relax}} \leq t_{\text{coll}} \leq t_{\text{H}}$ (orange colored zone). The arrows point toward the predicted movements in the M–R diagram.	13
3.1.	Principal domains in AMUSE framework. Multiple physical problems can be solved coupling codes from different domains. Credits: Portegies Zwart and McMillan (2018).	16
3.2.	Schematic representation of the three-layer architecture of the AMUSE framework. The user interacts with the system via a Python script (top layer), which communicates through interfaces (middle layer) with the community codes (bottom layer). Credits: Portegies Zwart and McMillan (2018).	17
3.3.	Representation of an oct tree	24
3.4.	Comparison of force calculation normalized (masses and gravitational constant set to 1). In dashed green, the exact gravitational force. In blue, the force with plummer softening, and in red, the force calculated with cubic spline softening. Credits: Reinoso (2024)	31

3.5.	Scheme of the BRIDGE coupling strategy. In the kick steps, FASTKICK computes the mutual gravitational forces between stars and gas, and the velocities of both components are updated. In the drift step, each code evolves its particles independently: Ph4 handles stellar dynamics (including collision detection and mergers), while Fi evolves the gas. After the kick-drift-kick cycle, the global BRIDGE time step is updated based on the gas free-fall time and the coupling timescales (t_{ff} , $\Delta t_{\text{coup,gas}}$, $\Delta t_{\text{coup,stars}}$). This updated timestep is used for both the next BRIDGE step and the Fi integration. Additionally, escaped N-body particles are identified and tagged in the post-processing phase.	33
4.1.	Energy conservation and virial equilibrium for a test simulation of a gas-embedded star cluster. Top panel: relative energy error $\Delta E/E$. Bottom panel: virial ratio $2K/ U $. The system remains in virial equilibrium with an energy error of $\lesssim 10^{-4}$ over 50 crossing times.	47
A.1.	Lambert function. The black line shows the $W_0(z)$ branch, and the blue line shows the $W_{-1}(z)$ branch.	85

Chapter 1

Introduction

One of the most enigmatic phenomena in astronomy is the existence of black holes (BHs). Since our understanding of the Universe is based primarily on the observation of light, the discovery of objects with no emission is, at least, challenging. This lack of emission can be understood from a fundamental property of astronomical objects, the escape velocity. For any body sufficiently massive, it is possible to measure its gravitational acceleration:

$$\mathbf{a} = \frac{GM}{r^2} , \quad (1.1)$$

where M is the mass and r the distance from the object. With this data is possible to calculate the velocity for escaping from their gravity, the escape velocity:

$$v_{\text{esc}} = \sqrt{\frac{2GM}{r}} . \quad (1.2)$$

When an object is dense enough that the escape velocity at its surface exceeds the speed of light, it is classified as a black hole. However, black holes are not totally invisible, as their effect on the surroundings is detected.

Black holes are commonly classified according to their mass M_{BH} . Stellar-mass black holes, typically with masses up to $\sim 100 M_{\odot}$, are thought to form from the collapse of massive stars, either through supernova explosions or via direct collapse (Belczynski et al., 2010; O’Connor and Ott, 2011). On the other side are supermassive black holes (SMBHs), exhibiting masses in the range 10^6 – $10^{10} M_{\odot}$ (Natarajan and Treister, 2009; Woods et al., 2019). However, the mechanism of their formation remains an open question in astrophysics.

1.1. Brief history: from Quasars to SMBHs

By the 1960s, there was evidence accounting for star-like objects with radio sources. In 1962, three of the brightest radio sources were identified in the northern sky by Thomas Matthews and Allan Sandage: 3C 48, 3C 196 and 3C 286 (Garwin and Lincoln, 2019). The spectrum of 3C 48 was characterized by broad emission lines, and from photometry the object showed variability on timescales of months, and also an excess of ultraviolet emission, compared with normal stars (Garwin and Lincoln, 2019; Shields, 1999). Those objects became known as quasi-stellar radio sources (QSRS), quasi-stellar sources (QSO), or quasars (Shields, 1999). In 1963, Greenstein and Matthews (1963) identified emission lines in 3C 48 of Mg II, [Ne III], [Ne V], and [O II], with a redshift of $z = 0.3675$. Interpreting this object as a galaxy core, they estimated a distance of 1.1×10^9 pc, and absolute magnitude (after redshift correction) of -26.3 , making it at least ten times brighter than the brightest known elliptical galaxies. By the same year, Schmidt (1963) reported the observation of 3C 273 (see Figure 1.1 for a contemporary image from Hubble data). Schmidt noticed that the spectrum presented emission lines from hydrogen, shifted to longer wavelengths. He identified four emission lines as Balmer lines, $H\beta$, $H\gamma$, $H\delta$ and $H\epsilon$, with a redshift of $z = 0.158$.




Figure 1.1: Image of 3C 273 from Hubble data. Credits: <https://science.nasa.gov/asset/hubble/quasar-3c-273-2/>.

If these objects were stars in our Galaxy, the large redshift would have to be gravitational in origin, requiring extremely high densities. Schmidt (1963) estimated a radius on the order of ~ 10 km

for such a compact object. The second explanation, which Schmidt considered the most direct and least objectionable, was that 3C 273 was the nucleus of a distant galaxy with a cosmological redshift, with a relative velocity of 47400 km s^{-1} , at a distance of $\sim 500 \text{ Mpc}$, and with a diameter of $\lesssim 1 \text{ kpc}$.

The first suggestion that quasars are fueled by SMBHs came with the work of Salpeter (1965). Salpeter proposed that extremely massive objects could accrete interstellar gas. Given an object of mass M , velocity v relative to a gas of density n (H atoms per cm^3), he defined a characteristic radius $s_0 = GM/v^2$ (from Hoyle and Lyttleton (1939)). Then, an accretion rate is defined:

$$\frac{dM}{dt} = 2\pi\alpha s_0^2 n v = \frac{\alpha M}{t_0}, \quad (1.3)$$

where α is a parameter that depends on the specific heat ratio of the gas. Salpeter also estimated that for the total mass being accreted per unit time, a fraction f of 0.05–0.2 can escape as energy to infinity, while a fraction $1 - f > 0.8$ contribute to the increase in mass of the object. The optical luminosity is estimated as:

$$L_{\text{opt}} = \left(\frac{25}{v}\right)^3 n \frac{\alpha}{0.25} \frac{f_{\text{opt}}}{0.03} \left(\frac{M}{10^7}\right)^2 \times 2 \times 10^9 L_{\odot}, \quad (1.4)$$

where $L_{\odot}$ is the solar luminosity and v is in km s^{-1} . In solar units, L_{opt}/M can grow as large as 3×10^4 , then radiation pressure lowers α , such that L_{opt}/M remains approximately constant.

The model was further developed and popularized by Lynden-Bell (1969), who argued that the remnants of quasars should be common in the centres of nearby galaxies. He proposed that most quasars should have consumed their fuel, on a timescale of 10^{10} years (the age of the universe). Those dead quasars should then be common in galactic centers. This is explained by the fact that as quasars consume the accreted gas, the Schwarzschild radius (the event horizon, a spherical boundary of radius $r = 2GM/c^2$ where the escape velocity equals the speed of light) becomes larger, consuming larger amounts of material. Also, in the past galaxies had larger amounts of gas than now.

One of the most important confirmations came with the discovery of an SMBH in the center of the Milky Way. In 1974, a compact radio source at the center of our galaxy, Sagittarius A* (Sgr A*), was discovered (Balick and Brown, 1974). In the late 1990s, Ghez et al. (1999) studied the proper motions of 90 stars near the center. They found that the movements indicated that a dark mass of $2.6(\pm 0.2) \times 10^6 M_{\odot}$ was confined to a volume of 10^{-6} pc^3 , coincident with Sgr A*. They concluded that our Galaxy harbours an SMBH at its center. The definitive proof came years later. In 2017, the Event Horizon Telescope (EHT; Doeleman et al. (2009)), an interferometer based on Very Long Baseline Interferometry (VLBI), observed Sgr A* (EHT Collaboration et al., 2022). They found that images of Sgr A* were consistent with a Kerr black hole, with a mass

of $\sim 4 \times 10^6 M_\odot$, and a bright ring with a diameter of $51.8 \pm 2.3 \mu\text{as}$, and a dim interior (see Figure 1.2).




Figure 1.2: Simulated images of Sgr A^{*}. The left panel is a snapshot of a numerical simulation of Sgr A^{*}, and middle panel is the average of snapshots of the simulation. On the right panel, a representative image reconstruction, using the synthetic visibilities from the simulation. Credits: EHT Collaboration et al. (2022).

1.2. Formation channels for SMBHs

SMBHs are conventionally defined as black holes with masses in the range $\sim 10^6$ – $10^{10} M_\odot$. There is accumulated evidence that support their existence (Gillessen et al., 2009; Latif and Ferrara, 2016; Natarajan and Treister, 2009; Volonteri, 2010; Woods et al., 2019). However, the origin of such dense and massive BHs is an active research topic in astronomy, as their main formation mechanism remains unknown (Woods et al., 2019).

The challenge intensified with the discovery of quasars at redshifts $z > 6$. Early surveys, such as the Canada-France High- z Quasar Survey (CFHQS), identified the first quasars beyond $z = 6$ (Willott et al., 2007). Similarly, Jiang et al. (2015) reported eight quasars at $z \sim 6$ from the Sloan Digital Sky Survey (SDSS). More recently, Yang et al. (2026) reported the discovery of 31 new quasars at $6.6 < z < 7.8$ (Yang et al., 2026) from the Euclid Wide Survey. Among them, EUCL J1729 is the most distant quasar discovered to date, with a redshift of $z \sim 7.77$, surpassing the quasar reported by Mortlock et al. (2011) at $z \sim 7.085$, and J0313-1806 at $z \sim 7.6$ (Wang et al., 2021), which hosts an SMBH of $\sim 1.6 \times 10^9 M_\odot$. At $z \sim 7.77$, the age of the Universe was ~ 662 Myr (Yang et al., 2026), raising the natural question: how could SMBHs have formed at such early times?

To understand the formation pathways of SMBHs, it is important to know how they are seeded. In this context, a BH seed is defined as the progenitor object (i.e., a stellar-mass BH or a very massive star) that forms in the early Universe and subsequently grows into a SMBH. Regan and Volonteri (2024) make the distinction between light and heavy seeds, where the former are objects

with mass $< 1000 M_{\odot}$, and the latter are objects with mass $> 1000 M_{\odot}$. They estimate that light seeds accreting gas at a 'standard' radiative efficiency of $\epsilon_r \sim 0.1$, takes ~ 1 Gyr to grow up to a mass of $\sim 10^9 M_{\odot}$. However, those seeds are unlikely to grow efficiently enough. Alvarez et al. (2009) showed that for black hole remnants of Population III stars (Pop III), radiation emitted by the accreting BH produces photoionization and heating that suppress the inflow of gas, resulting in negligible mass growth of the seed, making them unlikely to produce the observed quasars at $z > 6$. In agreement, another study of a simulation of 15000 black holes remnants of Population III stars, over 300 Myr, showed that fewer than 100 BHs grew by more than 10^{-3} of their initial masses, most of them coming from dense regions (Smith et al., 2018). Given the inefficiency of light seed growth, heavier seeds have been proposed as an alternative. Heavy seeds can be formed by runaway collisions of dense star clusters (Escala, 2021; Portegies Zwart et al., 2004), or the formation of a supermassive star (SMS) of $\gtrsim 10^5 M_{\odot}$ (Woods et al., 2017).

Although heavy seeds are more likely to grow into SMBHs, they require rare environmental conditions, resulting in a lower number density. For a redshift of $z \sim 10$, the heaviest seeds, with masses up to $\sim 10^5 M_{\odot}$, have number densities of $\sim 10^{-10}$ – 10^{-5} cMpc^{-3} , while other heavy seeds have number densities of $\sim 10^{-10}$ – 10^{-1} cMpc^{-3} (see Figure 1.3). In contrast, light seeds are more common, with a number density of $\sim 10^{-1}$ – 10^1 cMpc^{-3} (Regan and Volonteri, 2024).




Figure 1.3: BH abundance (number density) in cMpc^{-3} (y-axis) vs redshift (x-axis). The y-axis, from top to bottom, goes from very rare ($\sim 10^{-9}$) to common ~ 1 . Credits: Regan and Volonteri (2024).

There are many pathways for the formation of SMBHs (Volonteri, 2010). In the following,

we describe three channels: (i) remnants of Pop III stars, (ii) direct collapse black hole, and (iii) runaway stellar collisions in dense star clusters.

1.2.1. Population III stars

One of the pathways of SMBH formation is through remnants of Population III stars. Simulations show that Population III stars form in halos of 10^5 – $10^6 M_\odot$ at $z \sim 17$ (O’Shea and Norman, 2007, 2008). While early studies suggested that fragmentation was suppressed, recent high-resolution simulations have demonstrated that primordial protostellar disks are unstable to gravitational fragmentation, producing binary and multiple systems (Clark et al., 2011). After ~ 3 Myr, those stars can end in a supernova explosion or via direct collapse, leaving a BH remnant of $\sim 10^2 M_\odot$ (Fryer et al., 2001). However, their subsequent growth into the SMBH mass range seems unlikely. In an optimistic scenario, Vesperini et al. (2010) showed that the gas ejected by first-generation (Population III) stars can explain the chemical abundances in second generation (SG) stars, and that this dense gas can lead to significant accretion onto a pre-existing BH at the Eddington rate, increasing the mass by up to a factor of 100 ($\sim 10^4 M_\odot$). They estimated that a BH mass of $\gtrsim 10^4 M_\odot$ is required for the radiation-driven wind from the accreting BH to expel the surrounding gas. However, simulations show that a $100 M_\odot$ BH accreting metal-free gas exhibits intermittent accretion, which is on average reduced to $\sim 32\%$ of the Eddington rate (Milosavljević et al., 2009). In consequence, the mass increase is reduced to a factor of ~ 2 – 5 (Vesperini et al., 2010). Other simulations also report inefficient accretion onto Pop III BH remnants, due to the erratic dynamics of these BHs within their host clusters (Pfister et al., 2019; Smith et al., 2018). However, if Pop III stars form in dense clusters rather than in isolation, runaway collisions can lead to the formation of heavier seeds ($\sim 10^3$ – $10^4 M_\odot$), as we discuss later.

1.2.2. Direct collapse of primordial gas clouds

Another pathway is the formation of a heavy seed of $\sim 10^5 M_\odot$, via the direct collapse of primordial gas clouds. A set of simulations performed by Bromm and Loeb (2003) studied the evolution of an atomic-cooling halo with a total mass of $10^8 M_\odot$ ($1.5 \times 10^7 M_\odot$ in baryons), with a virial temperature of $\sim 10^4$ K. They varied the spin parameter λ , the inclusion of H_2 cooling, and the presence of a photo-dissociating Lyman-Werner (LW) background. They found that when H_2 cooling is suppressed, the gas collapses towards the centre via atomic hydrogen cooling, without fragmenting. Their simulations showed that $\gtrsim 10^6 M_\odot$ of the gas condenses into a region of $\lesssim 1$ pc, after which a supermassive star is likely to form. This SMS could subsequently collapse into a BH, retaining approximately 90% of its mass (Baumgarte and Shapiro, 1999; Shibata and Shapiro, 2002). However, the presence of H_2 cools the gas efficiently, leading to the formation of Population III stars. Their short lives (~ 3 Myr) produce supernova events (SNe) that enrich the gas, favouring subsequent star formation. To suppress H_2 cooling, the collapsing cloud must be exposed to a

strong Lyman-Werner flux, typically provided by a neighbouring star-forming halo. This seeding mechanism requires very specific conditions, making it difficult to explain the observed number density of quasars at $z > 6$. Although Shang et al. (2010) found that the number of halos in which H_2 formation can be suppressed may be larger, the subsequent mass growth of a $\sim 10^5 M_\odot$ seed faces significant challenges in sustaining the necessary accretion rates, as demonstrated by Chon et al. (2021).

1.2.3. Runaway collisions in dense star clusters

As discussed in the previous section, in low-metallicity gas, efficient H_2 cooling can lead to fragmentation, leading to the formation of Population III stars (Abel et al., 2000; Liao and Turk, 2019). As more massive halos form, metal enrichment from Population III stars can lead to efficient fragmentation and the formation of low-mass stars (Omukai et al., 2008; Volonteri, 2010). This opens the possibility of another scenario for the formation of SMBH seeds: runaway collisions in dense stellar clusters.

As the cluster evolves, mass segregation drives heavy stars to the center and light stars to the outer regions, through dynamical friction and two-body relaxation. This tendency towards equipartition of kinetic energy can lead to the gravothermal instability (Spitzer, 1987). In this scenario, the core may undergo a runaway collision process, leading to the formation of a massive black hole (Escala, 2021; Gürkan et al., 2004; Rasio et al., 2003). This scenario has been tested through N-body simulations of nuclear star clusters (NSCs) (Das et al., 2021; Portegies Zwart et al., 2004; Portegies Zwart and McMillan, 2002; Sakurai et al., 2017; Vergara et al., 2024, 2025, 2023), as well as young Population III star clusters (Freitag et al., 2006a,b; Reinoso et al., 2023, 2020; Vergara et al., 2021), showing the formation of a BH with mass $\sim 10^3\text{--}10^4 M_\odot$.

Most of these simulations consider purely gravitational interactions in gas-free clusters. However, gas can significantly enhance the formation of an SMBH seed. Gas damping can significantly accelerate mass segregation and drive overall cluster contraction (Leigh et al., 2014). Gas accretion also contributes to cluster contraction (Bonnell and Bate, 2002; Reinoso, 2024), and can subsequently fuel growth at the Eddington rate (Chung et al., 2026). Dynamical friction from star-gas interactions can also drive massive stars towards the centre (Leigh et al., 2014; Ostriker, 1999), however, this requires sufficient resolution to be adequately captured in simulations.

1.3. Research Objectives and Thesis Outline

Dynamical friction and mass segregation funnel material towards galactic centres. Also, objects that lose their dynamical support fall to the centre. The resulting high densities make NSCs likely sites for SMBH formation (Escala, 2006; Naiman et al., 2015; Shlosman et al., 1990). This thesis

investigates the role of gas in the runaway collision process within gas-embedded NSCs. The main goal is to quantify how the presence of gas affects this process: whether it increases the mass of the resulting BH seed, and by how much. To this end, we perform coupled N-body and SPH simulations, extending the global instability scenario of Escala (2021) to the smaller, denser systems explored by Vergara et al. (2023).

The outline of the thesis is as follows: In Chapter 2, we describe the scenario for the SMBH seed formation. The numerical methods are described in Chapter 3, and the initial conditions are presented in Chapter 4. The main results are presented in Chapter 5, which contains a scientific paper. Finally, we summarize our conclusions in Chapter 6.

Chapter 2

A scenario for SMBH formation

In this section, we describe the scenario proposed by Escala (2021) extended to include an external potential to mimic the effects of gas. The scenario explores the role of collisional and relaxation timescales (t_{coll} and t_{relax}) in determining the global stability of stellar systems.

As stars move in a cluster, gravitational interactions redistribute kinetic and potential energy through two-body encounters. Strong encounters can lead to the formation or disruption of binaries, the ejection of stars, or stellar mergers when stars physically overlap (Binney and Tremaine, 2008; Spitzer, 1987). Mass segregation drives massive stars toward the centre, increasing the central density and accelerating core collapse. When the density becomes sufficiently high, stellar collisions become frequent, potentially triggering a runaway phase that can lead to a global instability (Escala, 2021).

Weak encounters produce small, gradual changes in stellar velocities, but their cumulative effect leads to two-body relaxation, which drives mass segregation and can bring the system toward a state of partial energy equipartition (Bianchini et al., 2016; Trenti and van der Marel, 2013). The relevant timescales for describing these processes are the collisional timescale (t_{coll}) and the relaxation timescale (t_{relax}).

The collisional timescale t_{coll} provides an estimate of the collision rate between particles in a system. For a stellar system of identical stars, a first estimate of the collision rate $1/t_{\text{coll}}$ is given by Binney and Tremaine (2008):

$$\frac{1}{t_{\text{coll}}} = 16\sqrt{\pi}n\sigma r_{\star}^2(1 + \Theta) , \quad (2.1)$$

where $r_{\star}$ is the stellar radius, σ is the velocity dispersion of the system, n is the number density of stars, and

$$\Theta \equiv \frac{v_\star^2}{4\sigma^2} = 9.54 \frac{m}{M_\odot} \frac{R_\odot}{r_\star} \left(\frac{100 \text{ km s}^{-1}}{\sigma} \right)^2, \quad (2.2)$$

is the Safronov number, with m the mass of each star and $v_\star = \sqrt{2Gm/r_\star}$ the escape speed from the stellar surface. This derivation assumes a Maxwellian velocity distribution with dispersion σ . Equation (2.1) gives the global collision rate for the system, from typical stellar properties (m , $r_\star$, and σ).

The other relevant timescale is the relaxation time. As a system evolves, two-body relaxation drives it away from its initial state. Close encounters and mass segregation redistribute energy and mass, modifying the system's global properties (e.g., radius and velocity dispersion). Although strong encounters can significantly change a star's velocity, they are less frequent than weak encounters. The relaxation timescale t_{relax} is the time required for a typical star to change its velocity significantly through the cumulative effect of weak encounters. From Binney and Tremaine (2008),

$$t_{\text{relax}} = n_{\text{relax}} t_{\text{cross}}, \quad (2.3)$$

where t_{cross} is the crossing time of a typical star, and n_{relax} is the number of crossings needed for the cumulative change in velocity δv to become comparable to the star's velocity ($\Delta v = \sum \delta v \sim v$). For a system of N identical stars, $n_{\text{relax}} \sim 0.1N/\log N$ (see Appendix A.4). Thus,

$$t_{\text{relax}} = \frac{0.1N}{\log(N)} t_{\text{cross}}. \quad (2.4)$$

2.1. Critical mass

The relevance of the relaxation and collision timescales depends on the characteristic timescale of the system, t_{H} , typically taken as the age of the system. If $t_{\text{coll}} \lesssim t_{\text{H}}$, collisions are expected to be dynamically relevant. For a virialized system of total mass M and radius R , Equation (2.1) can be rearranged (Escala, 2021):

$$t_{\text{coll}} = \frac{1}{n\Sigma_0} \sqrt{\frac{R}{GM}}, \quad (2.5)$$

where $\Sigma_0 = 16\sqrt{\pi}r_\star^2(1 + \Theta)$ is the effective cross-section, and $\sigma = \sqrt{GM/R}$ for a virialized system. The Safronov number can be simplified to $\Theta \sim 9.54(m/M_\odot)(R_\odot/r_\star)$ if we assume $\sigma \sim 100 \text{ km s}^{-1}$. Assuming a homogeneous density distribution as a first approximation, the number density is $n = 3M/(4\pi R^3 m)$. Imposing the condition $t_{\text{coll}} \leq t_{\text{H}}$ and solving for the mass we obtain the critical mass (Escala, 2021; Vergara et al., 2023):

$$M \geq \left(\frac{4\pi m}{3\Sigma_0 t_H G^{1/2}} \right)^{2/3} R^{7/3} = M_{\text{crit}} . \quad (2.6)$$

The condition $t_{\text{coll}} \leq t_H$ is equivalent to $M \geq M_{\text{crit}}$. Thus, M_{crit} is the mass of a virialized system of radius R and equal-mass stars whose collision timescale equals $t_{\text{coll}} = t_H$. Including gravitational focusing and an external potential of mass $M_{\text{ext}} = qM$, the critical mass can be written as (see Appendix A.3):

$$M_{\text{crit}} = \left(\frac{\beta R^{7/2}}{2} + \sqrt{\frac{\beta^2 R^7}{4} + \frac{\alpha^3 R^3}{27}} \right)^{2/3} + \left(\frac{\beta R^{7/2}}{2} - \sqrt{\frac{\beta^2 R^7}{4} + \frac{\alpha^3 R^3}{27}} \right)^{2/3} - \frac{2\alpha R}{3} , \quad (2.7)$$

with

$$\alpha = \frac{9.54 m R_\odot v_{100}^2}{r_\star G M_\odot (1+q)^2} , \quad (2.8)$$

$$\beta = \frac{\sqrt{\pi} m}{12 G^{1/2} t_H r_\star^2 (1+q)} . \quad (2.9)$$

2.2. Relaxation mass

Similarly, we can obtain a condition using the relaxation time. From Appendix A.4, including an external potential, the relaxation timescale (Equation (A.41)) yields

$$t_{\text{relax}} = \frac{0.1 N (1+q)^3}{\ln(N(1+q)^2)} \frac{R^{3/2}}{\sqrt{GM}} . \quad (2.10)$$

Replacing $N = M/m$, and solving $t_{\text{relax}} \leq t_H$ for the characteristic radius R :

$$R \leq \left(\frac{t_H m}{0.1(1+q)^3} \ln \left(\frac{M(1+q)^2}{m} \right) \right)^{2/3} \left(\frac{G}{M} \right)^{1/3} . \quad (2.11)$$

This equation gives a condition on the radius such that the system evolves significantly due to two-body relaxation over t_H . We can rewrite Equation (2.11) by solving for the mass,

$$M \leq M_{\text{relax}} = \frac{m}{(1+q)^2} \exp\left(-2W_{-1}\left(\frac{-1}{2\lambda}\right)\right), \quad (2.12)$$

where $W_{-1}(z)$ is the lower branch of the Lambert function, and $\lambda = (t_{\text{H}}\sqrt{Gm})/(0.1(1+q)^2R^{3/2})$. In such systems, two-body relaxation does not significantly alter the system's properties over the characteristic timescale t_{H} . If, in addition, $t_{\text{coll}} \gg t_{\text{H}}$, the system remains approximately in virial equilibrium. The two mass scales defined above, M_{crit} and M_{relax} , delimit different dynamical regimes. We discuss their properties in the following section.

2.3. Global instability scenario

In this section, we analyze the scenario for supermassive black hole (SMBH) formation from NSCs proposed by Escala (2021). The critical mass M_{crit} from Equation (2.7) gives a threshold for the condition $t_{\text{coll}} = t_{\text{H}}$. For stellar clusters where $t_{\text{coll}} \leq t_{\text{H}}$ ($M \geq M_{\text{crit}}$), the onset of runaway collisions is expected, eventually forming a BH at the centre. On the other hand, when $t_{\text{relax}} \leq t_{\text{H}}$ ($M \leq M_{\text{relax}}$), two-body interactions could expand the cluster, altering its global properties (e.g., radius and velocity dispersion). Escala (2021) proposed a new regime of global instability, given by $t_{\text{coll}} \leq t_{\text{H}} \leq t_{\text{relax}}$. This regime has two key aspects. First, since $t_{\text{coll}} \leq t_{\text{H}}$, collisions occur within the characteristic timescale t_{H} . Second, the condition $t_{\text{coll}} \leq t_{\text{relax}}$ implies that two-body relaxation is unable to reverse the global collapse. As a result, the onset of runaway collisions leads to the formation of a massive black hole (MBH).

A second scenario can also lead to the formation of a central BH. We define an intermediate regime, given by clusters where $t_{\text{relax}} \leq t_{\text{coll}} \leq t_{\text{H}}$. In this regime, the system could begin to expand on a timescale comparable to t_{relax} . However, this does not necessarily prevent the onset of a collisional runaway in the core. Since the central density is significantly higher than the global average, the local collision time $t_{\text{coll,core}}$ is considerably shorter than the global collision time t_{coll} . If $t_{\text{coll,core}} \leq t_{\text{relax,core}}$, the core could contract and decouple dynamically from the expanding envelope, which could be compatible with a gravothermal instability scenario (Cohn, 1980; Spitzer, 1987). Then, the core could undergo runaway collisions, forming a central BH coexisting with a surrounding star cluster, while the outer stars, heated by two-body relaxation and dynamical encounters, are driven outward, expanding the outer regions of the cluster. Finally, a stable regime is defined by $t_{\text{relax}} \leq t_{\text{H}} \leq t_{\text{coll}}$. In this stable regime, clusters can expand due to two-body relaxation, which further decreases the collision rate. Although a BH can form via collisions in the core, most of the mass could remain in the stellar cluster.

The final fate of the central massive object (CMO) is conveniently explained in terms of the black hole formation efficiency, defined as $\epsilon_{\text{BH}} = M_{\text{BH}}/M_{\text{CMO}}$ (Escala, 2021; Vergara et al., 2023), where M_{BH} is the mass of the central black hole, and $M_{\text{CMO}} = M_{\text{BH}} + M_{\text{NSC}}$ is the mass of the central massive object, composed of a central BH and/or a nuclear stellar cluster. In Figure 2.1, the




Figure 2.1: Collision and relaxation timescales in a mass versus radius diagram. The blue dashed line correspond to the critical mass M_{crit} , while blue solid line correspond to the relaxation mass M_{relax} , both calculated for clusters formed by solar mass stars, characteristic time $t_H = 10^{10}$ yr and $q = 0$. The black dash-dotted line represents $t_{\text{coll}} = t_{\text{relax}}$, with t_H varying from 10^2 yr to 10^{10} yr and $q = 0$. Clusters above the black dash-dotted line satisfy $t_{\text{coll}} \leq t_{\text{relax}}$. The unstable regime (UR) is defined by $t_{\text{coll}} \leq t_H \leq t_{\text{relax}}$ (red colored zone); the stable regime is where $t_{\text{relax}} \leq t_H \leq t_{\text{coll}}$ (gray colored zone); and the intermediate regime (IR) fulfills $t_{\text{relax}} \leq t_{\text{coll}} \leq t_H$ (orange colored zone). The arrows point toward the predicted movements in the M–R diagram.

critical mass and relaxation mass are displayed (dashed and solid curves, respectively), for clusters formed by solar-mass stars, with $t_H = 10^{10}$ yr (the age of the Universe). The boundaries between the three regimes are defined by the curves M_{crit} , M_{relax} , and $t_{\text{coll}} = t_{\text{relax}}$, where the first two correspond to the conditions $t_{\text{coll}} = t_H$ and $t_{\text{relax}} = t_H$, respectively.

The unstable regime (UR), characterized by $t_{\text{coll}} \leq t_H \leq t_{\text{relax}}$, defines the zone where the mass of the system is simultaneously larger than the critical and the relaxation mass. In this regime, the system is unable to expand via two-body relaxation before collisions become relevant. Although binary heating can, in principle, oppose core contraction, in systems with masses larger than $10^7 M_\odot$ binary heating cannot reverse core collapse, and the system can suffer a merger instability, which may lead to the formation of a central BH (Lee, 1987). The dissipative nature of collisions thus drives a net contraction of the cluster. Then, most of the mass could remain in the central BH,

with efficiencies $\epsilon_{\text{BH}} \sim 1$. In this regime, we expect systems to move towards the bottom-left of the diagram, as the core contracts and stars are ejected from the core. In contrast, in the stable regime, defined by $t_{\text{relax}} \leq t_{\text{H}} \leq t_{\text{coll}}$, the dynamics are dominated by two-body relaxation over collisions. Since relaxation processes are relevant, the expansion of the cluster moves it to the right in the M–R diagram, potentially crossing the solid line. Although a central black hole could form, most of the mass remains in the stellar cluster, resulting in an NSC. Consequently, efficiencies are $\epsilon_{\text{BH}} \ll 1$. In addition, we define an intermediate regime (IR), which corresponds to the region where $t_{\text{relax}} \leq t_{\text{coll}} \leq t_{\text{H}}$. In this regime, relaxation processes and collisions can occur during the evolution: the system could expand while the core undergoes a runaway collision process, forming a central BH. These systems eventually move to the right as their radius increases, and downwards, as strong encounters can eject stars at high velocities. These systems may eventually cross into the stable regime (right side of the dashed curve), with efficiencies spanning a wider range $\epsilon_{\text{BH}} \leq 1$. While the classification above is general and valid for any characteristic timescale the figure illustrates the case $t_{\text{H}} = 10^{10}$ yr (the age of the Universe). In our simulations, we adopt $t_{\text{age}} = 5$ Myr, the duration of the simulations, following the approach of Vergara et al. (2023).

As noted by Escala (2021), the observed BH formation efficiency as a function of M_{CMO} shows a step-like form that clearly separates NSCs from SMBHs. A possible explanation for this observational gap is that systems in the intermediate regime may form IMBHs; however, an important fraction of the mass could remain in the surrounding stellar cluster. Consequently, their gravitational sphere of influence is small and challenging to resolve (Trenti, 2006), so these systems are likely to be classified as NSCs rather than as IMBH hosts. This naturally contributes to the lack of IMBH detections in NSC-dominated systems.

Chapter 3

Numerical Methods

In this chapter, we describe the numerical methods employed to model the coupled evolution of stars and gas in dense stellar clusters. We require three principal components: (i) an N-body code (ph4) to model stellar dynamics, (ii) a smoothed particle hydrodynamics (SPH) code (Fi) to simulate the gas, and (iii) a coupling scheme (BRIDGE) to handle their mutual gravitational interaction. The N-body and SPH codes used are accessed through the Astrophysical Multipurpose Software Environment (AMUSE; Portegies Zwart and McMillan (2018)), which provides a modular framework for solving multi-physics problems. We do not include gas accretion onto stars, as our focus is exclusively on the gravitational interaction between the two components.

3.1. The AMUSE Framework

The numerical codes used in this work are accessed through the Astrophysical Multipurpose Software Environment (AMUSE; Portegies Zwart and McMillan (2018)). The AMUSE project aims to provide a flexible and modular framework for solving multi-physics problems in astrophysics. It offers a variety of community codes, classified into four main domains: gravitational dynamics, hydrodynamics, stellar evolution, and radiative transfer (Figure 3.1). Some codes, such as GADGET-2 (Springel, 2005), can cover more than one domain.

The structure of AMUSE is composed of three principal layers (Figure 3.2). The top layer is the user interface, where a Python script is written to set up and run a specific scientific problem. In this script, codes are initialized, and additional routines (e.g., for handling stellar collisions) can be implemented. This layer also contains the manager, whose primary functions are to provide data structures and methods for communication between the user script and the community codes, as well as to handle unit conversions. The middle layer provides the interfaces to the community codes. It facilitates bidirectional communication between the manager and the underlying code




Figure 3.1: Principal domains in AMUSE framework. Multiple physical problems can be solved coupling codes from different domains. Credits: Portegies Zwart and McMillan (2018).

layer. Each interface consists of two components: a proxy, written in Python, which initializes the necessary functions and parameters, and a partner, written in the code’s native language (e.g., C++, CUDA, or Fortran). The proxy and partner communicate via the Message Passing Interface (MPI). Finally, the bottom layer contains the community codes themselves.

3.2. N-body code: PH4

The proposed scenario presented in Section 2 requires high numerical precision to accurately model close encounters and direct collisions. We therefore employ the ph4 N-body code from the AMUSE framework (Portegies Zwart and McMillan, 2018). In this section, we include a description of the code, and the external routines implemented to handle collisions.




Figure 3.2: Schematic representation of the three-layer architecture of the AMUSE framework. The user interacts with the system via a Python script (top layer), which communicates through interfaces (middle layer) with the community codes (bottom layer). Credits: Portegies Zwart and McMillan (2018).

3.2.1. Hermite integrator

The ph4 code uses a fourth-order Hermite integrator (Makino and Aarseth, 1992) to compute the positions and velocities of stars under gravity. This predictor-corrector method employs adaptive individual time steps, ensuring high accuracy in regions of strong gravitational interactions. Considering a particle i with its own time t_i , time step Δt_i , position $\mathbf{x}_i$, velocity $\mathbf{v}_i$, acceleration $\mathbf{a}_i$ and

jerk $\dot{\mathbf{a}}_i$, all calculated at time t_i . The integration algorithm proceeds as follows:

1. Select the particle with the minimum $t_i + \Delta t_i$. Set the global time to $t = t_i + \Delta t_i$.
2. Calculate the predicted positions ($\mathbf{x}_p$) and velocities ($\mathbf{v}_p$) for all particles, using the current values for $\mathbf{x}$, $\mathbf{v}$, $\mathbf{a}$, $\dot{\mathbf{a}}$.
3. Calculate the acceleration ($\mathbf{a}_i$) and jerk ($\dot{\mathbf{a}}_i$) for particle i at time $t_i + \Delta t_i$ using the predicted positions and velocities.
4. Calculate the second and third time derivatives of acceleration ($\mathbf{a}_i^{(2)}$ and $\mathbf{a}_i^{(3)}$) using Hermite interpolation.
5. Apply corrections to the position and velocity of particle i .
6. Calculate and update the time step Δt_i .
7. Repeat the algorithm.

The interpolation uses the acceleration and jerk at the beginning and end of the time step, allowing the scheme to achieve fourth-order accuracy without explicitly computing higher-order derivatives. A detailed derivation of the algorithm and the full set of equations is provided in Appendix B. From Eqs. (B.2) and (B.3), the predicted positions and velocities are:

$$\mathbf{x}_{p,j} = \mathbf{x}_j + \mathbf{v}_j(t - t_j) + \frac{\mathbf{a}_{0,j}}{2}(t - t_j)^2 + \frac{\dot{\mathbf{a}}_{0,j}}{3!}(t - t_j)^3, \quad (3.1)$$

$$\mathbf{v}_{p,j} = \mathbf{v}_j + \mathbf{a}_{0,j}(t - t_j) + \frac{\dot{\mathbf{a}}_{0,j}}{2!}(t - t_j)^2. \quad (3.2)$$

Here, $\mathbf{x}_j$ and $\mathbf{v}_j$ are the position and velocity of particle j at time t_j . The current acceleration $\mathbf{a}_{0,j}$ and jerk $\dot{\mathbf{a}}_{0,j}$ are computed as:

$$\mathbf{a}_{0,j} = \sum_{k \neq j} Gm_k \frac{\mathbf{r}_{0,jk}}{(\mathbf{r}_{0,jk}^2 + \epsilon^2)^{3/2}}, \quad (3.3)$$

$$\dot{\mathbf{a}}_{0,j} = \sum_{k \neq j} Gm_k \left[\frac{\mathbf{v}_{0,jk}}{(\mathbf{r}_{0,jk}^2 + \epsilon^2)^{3/2}} - \frac{3(\mathbf{v}_{0,jk} \cdot \mathbf{r}_{0,jk})\mathbf{r}_{0,jk}}{(\mathbf{r}_{0,jk}^2 + \epsilon^2)^{5/2}} \right], \quad (3.4)$$

$$\mathbf{r}_{0,jk} = \mathbf{r}_k - \mathbf{r}_j, \quad (3.5)$$

$$\mathbf{v}_{0,jk} = \mathbf{v}_k - \mathbf{v}_j. \quad (3.6)$$

Using Hermite interpolation, the second and third time derivatives of the acceleration can be estimated. This estimation is subsequently used to correct the positions and velocities. The relevant expressions, derived in the appendix, are:

$$\mathbf{a}_{0,i}^{(2)} = \frac{-6(\mathbf{a}_{0,i} - \mathbf{a}_{1,i}) - \Delta t_i(4\dot{\mathbf{a}}_{0,i} + 2\dot{\mathbf{a}}_{1,i})}{\Delta t_i^2}, \quad (3.7)$$

$$\mathbf{a}_{0,i}^{(3)} = \frac{12(\mathbf{a}_{0,i} - \mathbf{a}_{1,i}) + 6(\dot{\mathbf{a}}_{0,i} + \dot{\mathbf{a}}_{1,i})\Delta t_i}{\Delta t_i^3}. \quad (3.8)$$

The final corrected position and velocity of particle i are then given by:

$$\mathbf{x}_i(t_i + \Delta t_i) = \mathbf{x}_{p,i} + \frac{\mathbf{a}_{0,i}^{(2)} \Delta t_i^4}{24} + \frac{\mathbf{a}_{0,i}^{(3)} \Delta t_i^5}{120}, \quad (3.9)$$

$$\mathbf{v}_i(t_i + \Delta t_i) = \mathbf{v}_{p,i} + \frac{\mathbf{a}_{0,i}^{(2)} \Delta t_i^3}{6} + \frac{\mathbf{a}_{0,i}^{(3)} \Delta t_i^4}{24}. \quad (3.10)$$

These corrections incorporate terms up to fourth order. The main advantage of this method is that it only requires the explicit calculation of the acceleration and jerk at each step, with higher-order derivatives being obtained through the interpolation scheme.

3.2.2. Block time steps

While the Hermite integrator computes individual time steps for each particle, evolving particles one by one is computationally expensive. `ph4` therefore groups particles into block time steps, which are powers of two of the N-body time unit. This allows particles with the same time step to be integrated simultaneously, improving efficiency while maintaining synchronization. The choice of time step directly affects the accuracy of the integration, the conservation of energy, and the total computational cost.

In the block scheme, time steps are set in two steps. First, a dynamic time step is calculated. Then, the time step is assigned using a block scheme. For the dynamical time step, the standard formula (B.35) is used:

$$\Delta t_i = \sqrt{\eta \frac{|\mathbf{a}_{1,i}| |\mathbf{a}_{1,i}^{(2)}| + |\dot{\mathbf{a}}_{1,i}|^2}{|\dot{\mathbf{a}}_{1,i}| |\mathbf{a}_{1,i}^{(3)}| + |\mathbf{a}_{1,i}^{(2)}|^2}} . \quad (3.11)$$

As can be noted, $\mathbf{a}_{1,i}$ and $\dot{\mathbf{a}}_{1,i}$ are calculated in each step, while $\mathbf{a}_{1,i}^{(3)}$ is constant. Only $\mathbf{a}_{1,i}^{(2)}$ needs to be estimated:

$$\mathbf{a}_{1,i}^{(2)} = \mathbf{a}_{0,i}^{(2)} + \Delta t_i \mathbf{a}_{1,i}^{(3)} . \quad (3.12)$$

When the simulation begins, the initial time steps are calculated using (B.37):

$$\Delta t = \eta_s \frac{|\mathbf{a}|}{|\dot{\mathbf{a}}|} . \quad (3.13)$$

To avoid the computational cost of evolving particles individually, ph4 assigns particles to discrete block time steps that are powers of two of the N-body time unit. Given the initial time step $\Delta t_{0,j}$ (calculated from Equation (3.13)), and the N-body time τ , an integer n is adjusted to satisfy the condition:

$$0.5 < \frac{\tau 2^n}{\Delta t_{0,j}} < 1 . \quad (3.14)$$

The initial time step of particles is then set to $\tau 2^n$, where n can be positive or negative. As the model evolves, the new time step is computed using (3.11). The time step is halved if the calculated time step is smaller than the current time step, and doubled if it is larger than twice the current time step. The advantage of this time step criterion is that the time step in the code is equal to or lower than that required by the dynamics. Moreover, since particles share the same time steps, they can be integrated simultaneously. Additionally, all particles can be synchronized, as their times are multiples of powers of two of the same time unit.

3.2.3. Collisions and mergers

Stellar collisions are detected through a two-stage filter. First, ph4 checks whether two particles are closer than the sum of their radii: $|\Delta \mathbf{x}| < r_i + r_j$. A second filter then verifies close encoun-

ters using the criterion $\varepsilon|\Delta\mathbf{x}||\Delta\mathbf{v}| > \Delta\mathbf{x} \cdot \Delta\mathbf{v}$, where $\varepsilon = 0.001$. This avoids false detections from close encounters detected after the stars have already passed their point of closest approach and are receding, which can occur due to the finite timestep. Once a collision is detected, the flag `collision_detection` in the code is activated, and a custom handling routine is executed. In our implementation, the two particles involved in the collision are identified as particles i and j , corresponding to the most and less massive between them, respectively. The properties of particle i are then modified: its mass is set to the sum of the masses, its position to the centre of mass, its velocity to the centre-of-mass velocity, and its radius is updated to conserve mass density:

$$m_{i,\text{new}} = m_i + m_j , \quad (3.15)$$

$$\mathbf{x}_{i,\text{new}} = \frac{m_i \mathbf{x}_i + m_j \mathbf{x}_j}{m_i + m_j} , \quad (3.16)$$

$$\mathbf{v}_{i,\text{new}} = \frac{m_i \mathbf{v}_i + m_j \mathbf{v}_j}{m_i + m_j} , \quad (3.17)$$

$$r_{i,\text{new}} = r_i \left(\frac{m_i + m_j}{m_i} \right)^{1/3} . \quad (3.18)$$

$$(3.19)$$

Finally, particle j is removed from the simulation.

3.2.4. Escaping stars

After each time step, potential escapers are identified through a multi-stage detection algorithm. To maintain mass and energy conservation, escaped stars are not removed from the simulation. Instead, we add an extra attribute, `escaped` (all stars begin with `particles.escaped = False`), to track bound and unbound stars for subsequent analysis.

To detect escaping stars, the first step is to define an appropriate length scale. In the initialization, the escape scale is defined as the minimum between ten times the virial radius and the Lagrangian radius containing the total mass:

$$r_{0,\text{esc}} = \min(10 \times r_{\text{vir}}, r_{100}) , \quad (3.20)$$

where r_{100} represents the Lagrangian radius containing the total stellar mass. In subsequent iterations, the escape distance is updated as:

$$r_{\text{esc}} = \max(r_{\text{esc}}, 10 \times r_{b,\text{vir}}) , \quad (3.21)$$

where $r_{b,\text{vir}}$ denotes the virial radius of bound stars. Stars ejected from the core can have positive total energy while still interacting with their neighbours, and stars in marginally bound binaries may also be misclassified as unbound. To avoid tagging stars that are not genuinely escaping, we apply a multi-stage filter at the end of each time step:

1. Find the core center and identify stars beyond the escape radius.
2. Select gravitationally unbound stars.
3. Identify stars with outward radial motion.
4. Verify minimal gravitational influence on the nearest neighbour, to discard candidates that are still bound to another star (e.g., in a binary).

The core centre location is determined using a density-weighted scheme following Sweatman (1993). For each star, the distance to its 7th nearest neighbor ($r_{i,7}$) is computed, and the density center is calculated as:

$$\mathbf{x}_c = \frac{\sum_{i=1}^N \mathbf{x}_i / r_{i,7}^3}{\sum_{i=1}^N 1 / r_{i,7}^3} . \quad (3.22)$$

This formulation resembles a center of mass calculation but uses local stellar density instead of mass. Stars satisfying the condition

$$|\mathbf{x}_i - \mathbf{x}_c| > r_{\text{esc}} , \quad (3.23)$$

are identified as potential escapers. Gravitational unbinding is evaluated using the total energy $E_i = E_{p,i} + E_{k,i}$, where $E_{p,i}$ is the gravitational potential energy (including contributions from both stars and gas) and $E_{k,i}$ is the kinetic energy. Stars with $E_i \leq 0$ are considered bound, while those with $E_i > 0$ are classified as unbound. Among unbound stars, outward-moving candidates are identified through the condition:

$$\Delta \mathbf{x}_i \cdot \Delta \mathbf{v}_i > 0 , \quad (3.24)$$

where $\Delta \mathbf{x}_i = \mathbf{x}_i - \mathbf{x}_c$ and $\Delta \mathbf{v}_i = \mathbf{v}_i - \mathbf{v}_c$ represent position and velocity relative to the cluster center. Similar to (3.22), the velocity of the core center is computed as

$$\mathbf{v}_c = \frac{\sum_{i=1}^N \mathbf{v}_i m_i / r_{i,7}^3}{\sum_{i=1}^N m_i / r_{i,7}^3} . \quad (3.25)$$

The final validation step quantifies the gravitational influence on neighbouring stars, ensuring that the escaping candidate is not bound to a nearby star (e.g., in a binary), which would mean it is not genuinely escaping the cluster. For each candidate escaper, the relative potential energy ($e_{p,i}$) and kinetic energy ($e_{k,i}$) with respect to its nearest neighbor are computed. The energy ratio is defined as:

$$X_E := \frac{|e_{p,i}|}{e_{k,i}}. \quad (3.26)$$

Stars satisfying $X_E < 0.01$ are confirmed as escapers, ensuring minimal gravitational perturbation to neighboring stars. These stars are tagged as escapers (`particles.escaped = True`). Tagged escapers are excluded from subsequent dynamical calculations (e.g., core radius, density profiles), but they are not removed from the simulation, ensuring mass and energy conservation.

3.3. SPH code: Fi

The gas dynamics are modelled using the Fi code (Gerritsen et al., 1997), an MPI-parallel smoothed particle hydrodynamics (SPH) implementation within the AMUSE framework. We focus on the aspects most relevant to our simulations; a comprehensive description of the code can be found in Pelupessy (2005).

3.3.1. Barnes-Hut tree

A key feature of Fi is its ability to compute the self-gravity of the gaseous component. Particle-based codes offer a range of methods for gravitational force calculation. The most accurate approach is the direct summation of all pairwise interactions, which scales as $\mathcal{O}(N^2)$ and is computationally expensive for large N . While such precision is often necessary for collisional stellar dynamics (e.g., in our N-body component), the gas system in our simulations is treated as a collisionless fluid for the purpose of gravity calculation, allowing for faster, approximate methods, sacrificing accuracy for computational efficiency.

Fi employs the Barnes-Hut (BH) tree algorithm (Barnes and Hut, 1986). This method uses a hierarchical oct-tree spatial decomposition and a geometric opening criterion to approximate gravitational forces.

The tree is constructed by enclosing all particles within a root cell. If this cell contains more than one particle, it is subdivided into eight daughter cells of equal volume. This subdivision process repeats recursively for each cell containing multiple particles, until every cell holds at most one

particle (a leaf node). Figure 3.3 illustrates this structure in two dimensions.




Figure 3.3: Representation of an oct tree

Once the tree is built, the gravitational force on a given particle is calculated by going through the tree from the root. For each node encountered (linked to a cell), a decision is made: either open the node to examine its children, or accept the node as a single interacting entity. The classic Barnes-Hut opening criterion is:

$$l/d < \theta, \quad (3.27)$$

where l is the side length of the cubic cell, d is the distance from the target particle to the node's center of mass, and θ is an accuracy parameter. If this condition is met, the entire mass of the node is treated as a single point mass at its center of mass, providing an acceptable approximation when the node is sufficiently distant (i.e., appears small in angular size). The recursion stops upon reaching a leaf node, which by definition represents a single particle and is interacted with directly. This approximation reduces the computational cost of the force calculation from $\mathcal{O}(N^2)$ to $\mathcal{O}(N \log N)$ and adapts efficiently to clustered mass distributions, making it suitable for collisionless systems (Barnes and Hut, 1986).

The standard geometric criterion (Equation 3.27), however, can fail when the mass distribution within a cell is highly asymmetric. In those cases, the criterion may be satisfied even though a significant fraction of the mass of the cell lies close to the body, leading to large errors in the estimated force (Salmon and Warren, 1994). Therefore, Fi implements a modified criterion following Springel and Hernquist (2002):

$$M_n l^4 < \alpha |\mathbf{a}| d^6, \quad (3.28)$$

where M_n is the total mass contained in the node, $\mathbf{a}$ is an estimate of the gravitational acceleration (typically from the previous timestep), and α is a parameter to control accuracy.

3.3.2. Smoothed Particle Hydrodynamics

The gas dynamics are solved using the Smoothed Particle Hydrodynamics (SPH) method (Gingold and Monaghan, 1977; Lucy, 1977). In SPH, the hydrodynamic equations are solved from a Lagrangian perspective, representing the fluid with a discrete set of particles rather than a continuous medium. For an inviscid fluid described by density ρ , pressure p , and specific internal energy u , the Lagrangian takes the form:

$$L = \int \left[\frac{1}{2} v^2 - u(\rho) \right] \rho d^3 \mathbf{x}, \quad (3.29)$$

where the specific internal energy satisfies $\partial u / \partial \rho = p / \rho^2$, and $\mathbf{r} = (x_1, x_2, x_3)$ is the position vector. From this Lagrangian, the Euler equations of motion can be derived via a variational principle (Rasio, 2000):

$$\frac{\partial \rho}{\partial t} + \nabla \cdot (\rho \mathbf{v}) = 0, \quad (3.30)$$

$$\frac{\partial \mathbf{v}}{\partial t} + (\mathbf{v} \cdot \nabla) \mathbf{v} = -\frac{\nabla p}{\rho}, \quad (3.31)$$

$$\frac{\partial u}{\partial t} + (\mathbf{v} \cdot \nabla) u = -\frac{p}{\rho} (\nabla \cdot \mathbf{v}). \quad (3.32)$$

Equations (3.30), (3.31), and (3.32) are the continuity, momentum, and energy equations, respectively.

To obtain a discrete representation, the continuous Lagrangian is replaced by a particle-based analogue:

$$L_{\text{SPH}} = \sum_{i=1}^N m_i \left[\frac{1}{2} \mathbf{v}_i^2 - u_i \right], \quad (3.33)$$

where m_i , $\mathbf{v}_i$, and ρ_i are the mass, velocity, and density of each particle (representing a fluid element), and $u_i = u(\rho_i)$ is the specific internal energy. To use Eq. (3.33) as the system's Lagrangian, a suitable expression for the density in terms of the particle coordinates is required.

A key ingredient of the method is the density estimate. In SPH, the density at any point $\mathbf{r}$ is estimated by interpolating over neighboring particles:

$$\rho(\mathbf{r}) = \sum_j m_j W(\mathbf{r} - \mathbf{r}_j; h), \quad (3.34)$$

where $W(\mathbf{r} - \mathbf{r}_j; h)$ is a smoothing kernel of characteristic width h (the smoothing length). The kernel must satisfy two conditions: its integral over all space must be unity, and it should approach a Dirac delta function in the limit $h \rightarrow 0$. The first condition ensures that integrating Eq. (3.34) over space recovers the total mass of the system.

In the case of Fi, the cubic spline kernel is adopted (Monaghan, 1992; Pelupessy, 2005):

$$W(r, h) = \frac{1}{\pi h^3} \begin{cases} 1 - \frac{3}{2} \left(\frac{r}{h}\right)^2 + \frac{3}{4} \left(\frac{r}{h}\right)^3, & 0 \leq \frac{r}{h} \leq 1, \\ \frac{1}{4} \left(2 - \frac{r}{h}\right)^3, & 1 \leq \frac{r}{h} \leq 2, \\ 0, & \text{otherwise.} \end{cases} \quad (3.35)$$

With the density estimation and kernel defined, we now turn to the treatment of the smoothing length h . In modern SPH implementations, it is common to use an adaptive smoothing length h_i for each particle, adjusted to keep the number of neighboring particles approximately constant. However, the use of adaptive smoothing lengths introduces ∇h terms in the equations, which can lead to violations of energy and entropy conservation if not properly accounted for (Nelson and Papaloizou, 1994). A rigorous way to incorporate variable smoothing lengths is to treat h_i as additional variables in the Lagrangian formulation, subject to constraints (Springel and Hernquist, 2002). In Fi, the following constraint is imposed:

$$\frac{4\pi}{3} h_i^3 (\rho_i + \tilde{\rho}) = M_{\text{SPH}}, \quad (3.36)$$

where $M_{\text{SPH}} = \bar{m} N_{\text{nb}}$, with $\bar{m}$ the mean particle mass and N_{nb} the target number of neighbors. The term $\tilde{\rho}$ is an extra density introduced to prevent h_i from diverging as $\rho_i \rightarrow 0$ (Pelupessy, 2005). This constraint can be incorporated into the variational principle via Lagrangian multipliers, yielding correction factors in the equations of motion (see Appendix C).

With the notation $W_{ij}(h_i) \equiv W(|\mathbf{r}_i - \mathbf{r}_j|, h_i)$ and the symmetrized kernel $\bar{W}_{ij} = [W_{ij}(h_i) + W_{ij}(h_j)]/2$, the equation of motion for particle i becomes:

$$\frac{d\mathbf{v}_i}{dt} = - \sum_{j=1}^N m_j \left[f_i \frac{P_i}{\rho_i^2} \nabla_i W_{ij}(h_i) + f_j \frac{P_j}{\rho_j^2} \nabla_i W_{ij}(h_j) + \Pi_{ij} \nabla_i \bar{W}_{ij} \right], \quad (3.37)$$

where the correction factor f_i accounts for the variability of the smoothing length:

$$f_i = \left(1 + \frac{h_i}{3(\rho_i + \bar{\rho})} \frac{\partial \rho_i}{\partial h_i} \right)^{-1}. \quad (3.38)$$

The first two terms come from a conservative derivation of SPH equations. To handle discontinuities such as shocks, an artificial viscosity (AV) term Π_{ij} is included. In SPH, entropy generation is confined to shocks through this viscous term, which also prevents unphysical particle interpenetration.

Fi implements the standard Monaghan artificial viscosity (Monaghan, 1992; Pelupessy, 2005):

$$\Pi_{ij} = \frac{-\alpha \mu_{ij} \bar{c}_{ij} + \beta \mu_{ij}^2}{\bar{\rho}_{ij}}, \quad (3.39)$$

with

$$\mu_{ij} = \begin{cases} \frac{\mathbf{v}_{ij} \cdot \mathbf{r}_{ij}}{\bar{h}_{ij}(\mathbf{r}_{ij}^2/h_{ij}^2 + \eta^2)}, & \text{for } \mathbf{v}_{ij} \cdot \mathbf{r}_{ij} < 0, \\ 0, & \text{for } \mathbf{v}_{ij} \cdot \mathbf{r}_{ij} \geq 0, \end{cases} \quad (3.40)$$

where $\mathbf{v}_{ij} = \mathbf{v}_i - \mathbf{v}_j$ and $\mathbf{r}_{ij} = \mathbf{r}_i - \mathbf{r}_j$. The dimensionless parameters α and β control the strength of the viscosity (default values are $\alpha = 0.5$, $\beta = 1.0$), and $\eta \ll 1$ prevents numerical divergences. Quantities overlaid with a bar are symmetrized averages, e.g., $\bar{c}_{ij} = (c_i + c_j)/2$.

The AV is only activated when particles are approaching each other ($\mathbf{v}_{ij} \cdot \mathbf{r}_{ij} < 0$), ensuring that $\mu_{ij} \leq 0$ and therefore $\Pi_{ij} \geq 0$. As the term $-\nabla_i W_{ij}(h_i)$ is proportional to $(\mathbf{r}_i - \mathbf{r}_j) = \mathbf{r}_{ij}$, the AV acts to decrease kinetic energy, with the dissipated energy converted into thermal energy, thereby increasing the internal energy (or, equivalently, the entropic function, depending on the thermodynamic formulation chosen) of the particles.

Finally, the thermodynamic evolution of the gas must be specified. Fi offers two formulations: evolving either the specific internal energy u_i or the entropic function A_i (related to the specific entropy). For an ideal gas with adiabatic index γ , these quantities are related through

$$P_i = A_i \rho_i^\gamma, \quad (3.41)$$

$$u_i = \frac{A_i}{\gamma - 1} \rho_i^{\gamma-1} = \frac{P_i}{(\gamma - 1) \rho_i}, \quad (3.42)$$

where A_i is the entropic function, which remains constant for adiabatic processes in the absence of

shocks. The evolution equation for the entropic function is

$$\frac{dA_i}{dt} = -\frac{\gamma-1}{\rho_i^\gamma}(\Gamma - \Lambda) + \frac{1}{2} \frac{\gamma-1}{\rho_i^{\gamma-1}} \sum_{j=1}^N m_j \Pi_{ij} \mathbf{v}_{ij} \cdot \nabla_i \bar{W}_{ij}, \quad (3.43)$$

where Γ and Λ represent external heating and cooling rates (both set to zero in our simulations). Alternatively, the internal energy equation can be integrated:

$$\frac{du_i}{dt} = \sum_{j=1}^N m_j \left[\frac{f_i}{2} \frac{p_i}{\rho_i^2} \nabla_i W_{ij}(h_i) + \frac{f_j}{2} \frac{p_j}{\rho_j^2} \nabla_i W_{ij}(h_j) + \Pi_{ij} \nabla_i \bar{W}_{ij} \right] \mathbf{v}_{ij}. \quad (3.44)$$

For our simulations, we adopt the energy equation (Eq. (3.44)), as it provides better numerical stability in the coupled stellar-gas framework (see Appendix C).

3.4. Coupling Strategy: The BRIDGE Method

The N-body and SPH codes operate as independent modules and must be coupled for mixed star-gas simulations. A key advantage of using SPH for the gas component is that both systems are represented by particles, which simplifies the implementation of gravitational interactions. In this section, we describe the coupling scheme employed in this work, based on the BRIDGE method (Fujii et al., 2007) and following the implementation of Reinoso (2024).

3.4.1. Symplectic Integrator

The BRIDGE method (Fujii et al., 2007) employs a symplectic integrator, specifically the second-order leapfrog scheme. The evolution of a dynamical system can be described by Hamilton's equations:

$$\frac{df}{dt} = \{f, H\}, \quad (3.45)$$

where f, H denotes the Poisson bracket:

$$\{f, H\} = \sum_i \left(\frac{\partial f}{\partial q_i} \frac{\partial H}{\partial p_i} - \frac{\partial f}{\partial p_i} \frac{\partial H}{\partial q_i} \right) = D_H f(t), \quad (3.46)$$

with (q_i, p_i) being the canonical coordinates. The formal solution of this equation is given by the exponential operator:

$$f(t) = e^{tD_H} f(0) . \quad (3.47)$$

For a gravitational Hamiltonian of the form

$$H(\mathbf{x}, \mathbf{p}) = \frac{\mathbf{p}^2}{2m} + \Phi(\mathbf{x}) \equiv H_{\text{kin}}(\mathbf{p}) + H_{\text{pot}}(\mathbf{x}), \quad (3.48)$$

the Liouville operator D_H can be decomposed as:

$$D_H = \frac{\partial H}{\partial \mathbf{p}} \cdot \frac{\partial}{\partial \mathbf{x}} - \frac{\partial H}{\partial \mathbf{x}} \cdot \frac{\partial}{\partial \mathbf{p}}, \quad (3.49)$$

$$= \frac{\mathbf{p}}{m} \cdot \frac{\partial}{\partial \mathbf{x}} + (-\nabla \Phi(\mathbf{x})) \cdot \frac{\partial}{\partial \mathbf{p}}, \quad (3.50)$$

$$\equiv D_{\text{drift}} + D_{\text{kick}} . \quad (3.51)$$

The solution after a time step Δt is then:

$$f(t + \delta t) = e^{\delta t D_H} f(t) . \quad (3.52)$$

When applied to the phase-space coordinates $(\mathbf{x}, \mathbf{v})$, the exponential operator $e^{\delta t D_H}$ has no closed-form expression due to the coupling between position and velocity (Portegies Zwart and McMillan, 2018). However, the individual operators D_{drift} and D_{kick} have simple actions:

$$D_{\text{drift}}(\mathbf{x}, \mathbf{v}) = (\mathbf{v}, \mathbf{0}) , \quad (3.53)$$

$$D_{\text{kick}}(\mathbf{x}, \mathbf{v}) = (\mathbf{0}, -\nabla \Phi(\mathbf{x})) . \quad (3.54)$$

The drift operator advances positions at constant velocity, while the kick operator updates velocities at constant position. Importantly, these operators do not commute: $[D_{\text{drift}}, D_{\text{kick}}] \neq 0$. However, the exponential operator can be approximated using an operator splitting technique (Fujii et al., 2007):

$$e^{\Delta t D_H} = e^{\delta t (D_{\text{kick}} + D_{\text{drift}})} = \prod_{i=1}^k e^{a_i \delta t D_{\text{kick}}} e^{b_i \delta t D_{\text{drift}}} + \mathcal{O}(\Delta t^{k+1}) . \quad (3.55)$$

For $k = 2$ with coefficients $a_1 = a_2 = 1/2$, $b_1 = 1$, $b_2 = 0$, we obtain the second-order accurate approximation:

$$e^{\delta t(D_{\text{kick}} + D_{\text{drift}})} \approx e^{(\delta t/2)D_{\text{kick}}} e^{\delta t D_{\text{drift}}} e^{(\delta t/2)D_{\text{kick}}} + \mathcal{O}(\delta t^3) . \quad (3.56)$$

This leads to the kick-drift-kick leapfrog integrator, which proceeds as follows for a time step Δt :

1. Kick: Advance velocities by $\Delta t/2$ using accelerations computed at the current positions:

$$\mathbf{v}_0 \rightarrow \mathbf{v}_{1/2} = \mathbf{v}_0 + \frac{\Delta t}{2} \mathbf{a}_0 ,$$

2. Drift: Advance positions by Δt using the new half-step velocities:

$$\mathbf{x}_0 \rightarrow \mathbf{x}_1 = \mathbf{x}_0 + \Delta t \mathbf{v}_{1/2} ,$$

3. Kick: Advance velocities by another $\Delta t/2$ using accelerations computed at the updated positions:

$$\mathbf{v}_{1/2} \rightarrow \mathbf{v}_1 = \mathbf{v}_{1/2} + \frac{\Delta t}{2} \mathbf{a}_1 ,$$

where $\mathbf{a}_0 = \mathbf{a}(\mathbf{x}_0)$ and $\mathbf{a}_1 = \mathbf{a}(\mathbf{x}_1)$ are the accelerations due to the gravitational potential. The subscripts 0, 1/2, and 1 denote quantities evaluated at times t , $t + \Delta t/2$, and $t + \Delta t$, respectively.

3.4.2. BRIDGE

AMUSE includes an implementation of the original BRIDGE scheme. An example coupling an N-body code with an SPH code can be found in the AMUSE distribution at:

`amuse/examples/simple/gas_in_cluster.py`

This example models a star cluster with an embedded gas component, coupling GADGET-2 (SPH) and HERMITE (N-body). In this implementation, each code performs its own drift step internally. The kick step, however, is handled externally as follows:

- Accelerations $\mathbf{a}_{\text{NB},i}$ for each N-body particle due to all SPH particles are computed using the SPH code's `get_gravity_at_point(eps, x, y, z)` method, which returns the gravitational acceleration at a given point (x, y, z) with softening length eps . The velocities of N-body particles are then kicked using these accelerations.

- Similarly, accelerations $\mathbf{a}_{\text{SPH},i}$ for each SPH particle due to all N-body particles are computed using the `get_gravity_at_point(eps, x, y, z)` method of a tree code (BHTree in the example), with the softening length set to the SPH smoothing length h_i . The velocities of SPH particles are then kicked accordingly.

As identified by Reinoso (2024), this implementation has two main issues, both related to the kick steps. First, SPH and N-body particles use different gravitational softening lengths (h and ϵ , respectively), leading to asymmetric forces:

$$F_{\text{NB}} = \frac{Gm_{\text{NB}}m_{\text{SPH}}}{(r^2 + \epsilon^2)^{3/2}}r \neq F_{\text{SPH}} = \frac{Gm_{\text{NB}}m_{\text{SPH}}}{(r^2 + h^2)^{3/2}}r, \quad (3.57)$$

where $r = |\mathbf{r}_{\text{NB}} - \mathbf{r}_{\text{SPH}}|$. This asymmetry can be resolved by introducing an effective softening length $\bar{\epsilon}$ for the mutual gravitational interaction during the kick steps.

Second, the force calculation may use different softening kernels: SPH codes often employ a cubic spline softening, while N-body codes typically use a Plummer (or similar) softening. The difference between these softening formulations is illustrated in Figure 3.4.




Figure 3.4: Comparison of force calculation normalized (masses and gravitational constant set to 1). In dashed green, the exact gravitational force. In blue, the force with plummer softening, and in red, the force calculated with cubic spline softening. Credits: Reinoso (2024)

To handle these issues, Reinoso (2024) developed a modified BRIDGE implementation that uses the FASTKICK code to compute gravitational interactions between SPH and N-body particles. FASTKICK computes forces using a Plummer softening with a constant softening length $\bar{\epsilon}$, ensuring symmetry and consistency. The code provides a Python interface, making it compatible with AMUSE simulation scripts, with the computationally intensive force calculations performed in compiled languages (C/Fortran), and a GPU-accelerated version (C++/CUDA) is also available.

The modified BRIDGE algorithm proceeds as follows for each global time step Δt :

1. At time t , compute accelerations $\mathbf{a}_{\text{NB},i}$ for all N-body particles due to SPH particles, and $\mathbf{a}_{\text{SPH},i}$ for all SPH particles due to N-body particles, using FASTKICK with softening length $\bar{\epsilon}$. Kick velocities by $\Delta t/2$.
2. Evolve each code independently for a full time step Δt (drift), updating positions via their own internal integrators.
3. Recompute accelerations using FASTKICK with the updated positions, and apply a second velocity kick of $\Delta t/2$.

3.4.3. Adaptive timestep

The evolution of the coupled system is controlled by the BRIDGE time step. At the end of each BRIDGE step, the time step is adjusted to accurately resolve the physics of the gas and the star-gas interaction. For the gas, the minimum free-fall time is calculated:

$$t_{\text{ff}} = \sqrt{\frac{3\pi}{32G\rho_{\text{max}}}}, \quad (3.58)$$

where ρ_{max} is the maximum density among all SPH particles. For the gas-stars interaction, Courant-type time steps that depend on the acceleration between gas and star particles are calculated:

$$\Delta t_{\text{coup,gas}} = \min_i \left(f_{\text{coup}} \sqrt{\frac{2h_i}{\mathbf{a}_{\text{SPH},i}}} \right), \quad (3.59)$$

$$\Delta t_{\text{coup,stars}} = \min_i \left(f_{\text{coup}} \sqrt{\frac{\epsilon_{\text{coup}}}{\mathbf{a}_{\text{NB},i}}} \right), \quad (3.60)$$

where h_i is the softening length of SPH particles, $\mathbf{a}_{\text{SPH},i}$ and $\mathbf{a}_{\text{NB},i}$ are the forces calculated by FASTKICK in the kick step, and ϵ_{coup} is the softening length used by FASTKICK in the calculation of gravitational interactions. For $f_{\text{coup}} = 1$, Equations (3.59) and (3.60) set the minimum

resolution for gas and stars interactions to be solved in the BRIDGE steps. Since f_{coup} is an accuracy parameter, it is recommended to use values < 1 .

Taking into account the minimum gas free-fall time (Equation (3.60)) and the Courant-type time steps (Equations (3.59) and (3.60)), the timestep is decided as

$$\tau_{\text{step}} = \min(t_{\text{ff}}, \Delta t_{\text{coup,gas}}, \Delta t_{\text{coup,stars}}) . \quad (3.61)$$

Finally, the time step of BRIDGE is adjusted by the condition

$$0.5\tau_{\text{step}} \leq \frac{\tau}{2^n} \leq \tau_{\text{step}} , \quad (3.62)$$

with τ the crossing time of the stellar cluster and n an integer. The time step of the BRIDGE $\Delta t_{\text{BRIDGE}} = \tau/2^n$ is adjusted at the end of each BRIDGE step, and is also set as the timestep for the SPH code. Note that the adjusted time step is a power of two of the initial N-body time unit, consistent with the block time steps used for the N-body particles. Consequently, the three interacting codes are in synchrony. Figure 3.5 shows a schematic of the coupling strategy.




Figure 3.5: Scheme of the BRIDGE coupling strategy. In the kick steps, FAST-KICK computes the mutual gravitational forces between stars and gas, and the velocities of both components are updated. In the drift step, each code evolves its particles independently: Ph4 handles stellar dynamics (including collision detection and mergers), while Fi evolves the gas. After the kick-drift-kick cycle, the global BRIDGE time step is updated based on the gas free-fall time and the coupling timescales (t_{ff} , $\Delta t_{\text{coup,gas}}$, $\Delta t_{\text{coup,stars}}$). This updated timestep is used for both the next BRIDGE step and the Fi integration. Additionally, escaped N-body particles are identified and tagged in the post-processing phase.

Chapter 4

Initial Conditions and Setup

Chapter 3 described the numerical methods used in this work, including the SPH code for the gas, the N-body code for the stars, and the coupling scheme. This chapter provides a complete specification of the simulations, detailing the initial conditions for both the stellar and gaseous systems, as well as the numerical parameters used in the code initialization.

4.1. Plummer distribution

The scenario presented in Chapter 2 and proposed by Escala (2021) was previously explored by Vergara et al. (2023) through simulations sampling different collisional regimes. These models represent stellar systems characterized by their initial mass and size scale. The stellar clusters are initialized using a Plummer distribution (Plummer, 1911). The Plummer model is characterized by the potential (Binney and Tremaine, 2008):

$$\Phi_p(r) = -\frac{GM}{\sqrt{r^2 + b^2}}, \quad \text{with } r = |\mathbf{r}|, \quad (4.1)$$

where b is the Plummer scale. From the Poisson equation, the associated density is

$$\rho_p(r) = \frac{3M}{4\pi b^3} \left(1 + \frac{r^2}{b^2}\right)^{-5/2}. \quad (4.2)$$

The potential energy of a Plummer model is:

$$U_p = -\frac{3\pi GM^2}{32b}. \quad (4.3)$$

We now require a description for the velocity. The velocity dispersion σ for an isotropic spherical stellar system can be obtained through the Jeans equation (Binney and Tremaine, 2008):

$$\frac{d(\rho\sigma^2)}{dr} = -\rho \frac{d\Phi}{dr} . \quad (4.4)$$

Solving for the Plummer model gives the velocity dispersion:

$$\sigma_p(r)^2 = \frac{GM}{6\sqrt{r^2 + b^2}} . \quad (4.5)$$

We can relate the Plummer scale parameter b to a physical scale. First, we use the definition of virial radius as the characteristic system radius R via the potential energy U :

$$-\frac{GM^2}{2R} = U \quad (4.6)$$

$$\frac{1}{R} = -\frac{2}{GM^2}U , \quad (4.7)$$

where U is the total gravitational potential energy. In the case of a system of N particles, this radius is defined as:

$$R = \left(\frac{1}{M^2} \sum_{i,j \neq i}^N \frac{m_i m_j}{|\mathbf{r}_i - \mathbf{r}_j|} \right)^{-1} , \quad (4.8)$$

In the case of a Plummer model, replacing Eq. 4.3 into Eq. 4.7, this radius can be expressed in function of the Plummer scale parameter b :

$$\frac{1}{R} = \left(-\frac{2}{GM^2} \right) \left(-\frac{3\pi GM^2}{32b} \right) , \quad (4.9)$$

$$R = \frac{16b}{3\pi} . \quad (4.10)$$

Once the physical scale of the system is defined via b (or equivalently R), we need to generate the actual particle distribution. This is typically done in a dimensionless system of units, which we now describe.

4.1.1. N-body units

Most gravitational N-body codes employ a dimensionless system of units, known as N-body units or Hénon units (see Heggie and Mathieu (1986)). In this system, physical quantities are scaled to be of order unity, which brings two main benefits: (i) it eliminates the need to multiply by G in every force calculation, and (ii) it improves numerical precision by keeping variables close to unity, thereby minimizing round-off errors that would otherwise arise when working with astrophysical scales spanning many orders of magnitude. Additionally, N-body units facilitate comparison between results obtained with different codes.

N-body units are defined by setting the gravitational constant, total mass, and total energy to convenient values:

$$G = 1 , \tag{4.11}$$

$$M = 1 , \tag{4.12}$$

$$E = -\frac{1}{4} . \tag{4.13}$$

With this, the units of mass, length, and time can be defined. The mass unit is simply the total mass:

$$u_m = M . \tag{4.14}$$

Assuming virial equilibrium ($2T + U = 0$), the kinetic and potential energies become:

$$T = \frac{1}{4} = -E , \tag{4.15}$$

$$U = -\frac{1}{2} = 2E . \tag{4.16}$$

Using the virial radius definition (Eq. (4.7)) and substituting the potential energy from Eq. (4.16), we obtain:

$$R = -\frac{GM^2}{4E} . \tag{4.17}$$

Using $G = 1$, $M = 1$ and $E = -1/4$, we find that $R = 1$, so the length unit u_l is:

$$u_l = R = -\frac{GM^2}{4E} . \quad (4.18)$$

Finally, we derive the time unit. A characteristic velocity follows from the kinetic energy:

$$T = \frac{M\sigma^2}{2} , \quad (4.19)$$

$$\sigma^2 = \frac{2T}{M} = \frac{-2E}{M} . \quad (4.20)$$

The crossing time is defined as:

$$t_{cross} = \frac{2R}{\sigma} = \frac{GM^{5/2}}{(-2E)^{3/2}} . \quad (4.21)$$

With $G = 1$, $M = 1$, and $E = -1/4$, we obtain $t_{cross} = 2^{3/2}$. The standard N-body time unit is then defined as:

$$u_t = t_{cross} 2^{-2/3} = \frac{GM^{5/2}}{(-4E)^{3/2}} , \quad (4.22)$$

which indeed yields $u_t = 1$ for $G = 1$, $M = 1$ and $E = -1/4$. Note that defining the unit system requires only a scale radius and the total mass of the system. AMUSE provides its own physical units system, which can be converted to N-body units using a converter. In AMUSE, N-body units are handled through the `nbody_system` class, which requires at least two physical quantities, typically the total mass and the virial radius. The following example illustrates how to initialize a converter:

```

1 from amuse.units import units, nbody_system
2
3 R_vir = 1 | units.pc          # Define a scale radius
4 M_tot = 1e4 | units.MSun     # Define the total mass
5
6 converter = nbody_system.nbody_to_si(R_vir, M_tot)
```

Code 4.1: How to define a converter in AMUSE

The converter serves a bidirectional role: it translates physical quantities (e.g., positions, ve-

locities, masses) into dimensionless N-body units for the numerical calculations, and converts the simulation output back to dimensional physical units for subsequent analysis.

4.2. Stars and Gas Initial Conditions

For the results presented in Chapter 5, we adopt the same initial conditions as the models in Vergara et al. (2023). For each model, we perform four simulations with different gas fractions, using the same stellar initial conditions (same as in Vergara et al. (2023)) for all gas fractions. We use the notation of a family model to reference the four variations of a specific model. For example, model family A is composed of the four models A₀, A₅, A₁₀ and A₂₀, where the capital letter indicates that they share the same initial stellar cluster conditions, and the subscript indicates the percentage of gas mass added. Table 4.1 summarizes the initial conditions of the stellar components of each model.

Table 4.1: Initial conditions for the stellar systems. R_v , M , N , m_s , r_s , and σ denote the virial radius, total stellar mass, number of stars, stellar mass, stellar radius, and velocity dispersion, respectively.

Model ID ¹	R_v [pc]	M [$M_\odot$]	N	m_s [$M_\odot$]	r_s [$R_\odot$]	σ [km/s]
A	0.005	5×10^4	10^3	50	11.7	146.66
B	0.005	2.5×10^4	5×10^2	50	11.7	103.71
C	0.005	10^4	10^3	10	4.7	65.59
D	0.005	5×10^3	5×10^2	10	4.7	46.38
F	0.005	10^3	10^3	1	1.0	20.74
H	0.01	5×10^4	5×10^3	10	4.7	103.71
I	0.01	10^4	10^3	10	4.7	46.38
J	0.01	5×10^3	5×10^3	1	1.0	32.79
K	0.01	10^3	10^3	1	1.0	14.66
L	0.1	5×10^5	10^4	50	11.7	103.71
M	0.1	10^5	10^4	10	4.7	46.38
N	0.1	10^4	10^4	1	1.0	14.67

¹ The "Model ID" column identifies the family of models. All members of a given family (e.g., A₀, A₅, A₁₀, and A₂₀) share the same stellar initial conditions, which are listed here with capital letters with no suffix (A, B, C, etc.).

In Table 4.1, R_v , M , N , and σ denote the virial radius, total mass, number of stars, and velocity dispersion of the stellar system, respectively. The initial mass and radius of each star are given by m_s and r_s . All variants of a family model have the same initial conditions for the stellar system. For the gas-free models, the stellar clusters are initialized in virial equilibrium ($2K + U = 0$). For models with gas, the stellar component retains the same initial velocities and is therefore not in virial equilibrium with the total mass (stars and gas). The total system is subvirial, with $2K < |U_{\text{total}}|$, and will evolve dynamically toward a new equilibrium. The gas is initialized in hydrostatic

equilibrium, supported by pressure in the gravitational potential of the stars. The initial conditions of gas components are presented in Table 4.2, where $q = M_{\text{SPH}}/M$, M_{SPH} is the total mass of SPH particles, N_{SPH} and m_{SPH} are the total number and mass per SPH particle, and c_s is the sound speed of the gas component, calculated as the mass-weighted average within the virial radius.

Table 4.2: Initial conditions for gas systems. q , M_{SPH} , N_{SPH} , m_{SPH} , and c_s denote the gas mass fraction, total gas mass, number of SPH particles, SPH particle mass and mean sound speed, respectively.

Model ID	q	$M_{\text{SPH}} [M_\odot]$	N_{SPH}	$m_{\text{SPH}} [M_\odot]$	$c_s [km/s]$
A ₅	0.05	2.5×10^3	5×10^3	0.5	95.97
A ₁₀	0.1	5×10^3	10^4	0.5	96.67
A ₂₀	0.2	10^4	2×10^4	0.5	96.90
B ₅	0.05	1.25×10^3	2.5×10^3	0.5	66.35
B ₁₀	0.1	2.5×10^3	5×10^3	0.5	67.17
B ₂₀	0.2	5×10^3	10^4	0.5	68.11
C ₅	0.05	5×10^2	5×10^3	0.1	43.10
C ₁₀	0.1	10^3	10^4	0.1	43.24
C ₂₀	0.2	2×10^3	2×10^4	0.1	43.35
D ₅	0.05	2.5×10^2	2.5×10^3	0.1	29.77
D ₁₀	0.1	5×10^2	5×10^3	0.1	30.51
D ₂₀	0.2	10^3	10^4	0.1	30.46
F ₅	0.05	5×10^1	5×10^3	0.01	13.63
F ₁₀	0.1	10^2	10^4	0.01	13.64
F ₂₀	0.2	2×10^2	2×10^4	0.01	13.72
H ₅	0.05	2.5×10^3	2.5×10^4	0.1	68.50
H ₁₀	0.1	5×10^3	5×10^4	0.1	68.57
H ₂₀	0.2	10^4	10^5	0.1	69.04
I ₅	0.05	5×10^2	5×10^3	0.1	30.25
I ₁₀	0.1	10^3	10^4	0.1	30.24
I ₂₀	0.2	2×10^3	2×10^4	0.1	30.72
J ₅	0.05	2.5×10^2	2.5×10^4	0.01	21.64
J ₁₀	0.1	5×10^2	5×10^4	0.01	21.85
J ₂₀	0.2	10^3	10^5	0.01	21.78
K ₅	0.05	5×10^1	5×10^3	0.01	9.65
K ₁₀	0.1	10^2	10^4	0.01	9.62
K ₂₀	0.2	2×10^2	2×10^4	0.01	9.63
L ₅	0.05	2.5×10^4	5×10^4	0.5	68.84
L ₁₀	0.1	5×10^4	10^5	0.5	68.85
L ₂₀	0.2	10^5	2×10^5	0.5	69.15
M ₅	0.05	5×10^3	5×10^4	0.1	30.76
M ₁₀	0.1	10^4	10^5	0.1	30.84
M ₂₀	0.2	2×10^4	2×10^5	0.1	30.90
N ₅	0.05	5×10^2	5×10^4	0.01	9.76
N ₁₀	0.1	10^3	10^5	0.01	9.71
N ₂₀	0.2	2×10^3	2×10^5	0.01	9.78

4.2.1. Particles generation

Both the stellar and gaseous systems are generated using the AMUSE implementation of the Plummer model. To generate a stellar system with the initial conditions of model A, for example, we first initialize a converter:

```
1 from amuse.units import units, nbody_system
2
3 M_stars = 5e4 | units.MSun
4 R_vir = 0.005 | units.pc
5 converter = nbody_system.nbody_to_si(R_vir, M_stars)
```

Code 4.2: Converter example for model A

After importing the **plummer** module, the stellar particles are generated by calling the function **new_plummer_model** with the converter and number of particles:

```
1 from amuse.ic import plummer
2
3 N = 1000
4 stars = plummer.new_plummer_model(N, converter, radius_cutoff=5., do_scale=True)
5 stars.radius = 11.7 | units.RSun
```

Code 4.3: Stars particles in model A

The **radius_cutoff** parameter truncates the distribution at $5b$, where b is the Plummer scale parameter (here corresponding to a virial radius $R_v = 0.005$ pc). Setting **do_scale=True** ensures that the generated particles are scaled to N-body units, satisfying the virial conditions described in Section 4.1.1 (total mass $M = 1$, total energy $E = -1/4$, kinetic energy $T = 1/4$, and potential energy $U = -1/2$).

For the gaseous component, we use the **gasplummer** function from the **amuse.ic** module to generate the SPH particles. The number of gas particles is chosen such that the mass ratio $m_{\text{SPH}}/m_{\text{NB}} = 0.01$, where m_{NB} is the mass of each N-body particle. For example, for the variation with 20%, A_{20} , the gas particles are generated as follows:

```
1 from amuse.ic import gasplummer
2
3 f_gas = 0.2
4 N_gas = N*100
5 N_gas = int(N_gas*f_gas)
6
7 gas = gasplummer.new_plummer_gas_model(N_gas,
8                                         convert_nbody=converter,
```

```

9                                     do_scale=True,
10                                     mass_cutoff=0.95535)
11 gas.mass = (M_stars*f_gas / N_gas)

```

Code 4.4: Gas particles in model A

While the stellar component is initialized in virial equilibrium with respect to its own mass, the gas particles are initially at rest. To prevent immediate collapse, their internal energy is set to achieve hydrostatic equilibrium in the gravitational potential of the stellar cluster. For a fluid in hydrostatic equilibrium,

$$\nabla p = -\rho \nabla \Phi, \quad (4.23)$$

where p , ρ , and Φ stand for the pressure, density, and gravitational potential, respectively. Assuming a polytropic equation of state (Eqs. (3.41) and (3.42)), this condition gives a relation for the specific internal energy u :

$$u = -\frac{\Phi}{\gamma}, \quad (4.24)$$

where γ is the polytropic index.

The initialization described in Code 4.4 raises a potential issue. The gas particles are generated using a converter scaled to the total stellar mass ($M = 5 \times 10^4 M_\odot$), and their internal energy is set accordingly to achieve hydrostatic equilibrium in the combined potential. However, after generation, we explicitly reduce the gas mass to $f_{\text{gas}}M$ (in this case, 20% of the stellar mass). In isolation, this reduction would lead to an imbalance: the gravitational force from the gas diminishes, while the internal pressure remains unchanged, causing the gaseous component to expand. However, because the gas is gravitationally coupled to the stellar system, the dominant contribution to the potential comes from the stars. We therefore expect the gas to remain near dynamical equilibrium, with any minor adjustments occurring on a dynamical timescale as the simulation evolves.

4.3. Numerical Parameters

Having described the numerical methods in the previous chapter, we now detail the specific parameter configurations used for each code. As discussed in Section 3.1, the AMUSE framework provides a unified interface that connects Python scripts with compiled codes through class structures. Each code is accessible via the community module, e.g., `amuse.community.ph4.interface` for Ph4. The following example illustrates how to create an instance of Ph4:

```

1 from amuse.community.ph4.interface import ph4

```

```
2
3 grav = ph4()
```

Code 4.5: Instance creation for Ph4

In Code 4.5, **grav** is an instance of the class **amuse.community.ph4.interface.ph4** (shortened **ph4**). The code classes have a variety of parameters and internal methods that control gravitational softening, integration accuracy, and the timestep, among other features. These are used to set up the particles before running the code. The community codes run internally in N-body units; however, AMUSE provides converters that translate between these internal units and physical units, allowing the user to work with physical quantities throughout the simulation. In the following, we show how to correctly initialize and run an instance of **ph4**, including particle creation using a Plummer distribution and a converter:

```
1 from amuse.community.ph4.interface import ph4
2 from amuse.units import units, nbody_system
3 from amuse.ic.plummer import new_plummer_model
4
5 # Step 1, creation of a converter and particles
6 virial_radius = 1 | units.pc
7 total_mass = 1000 | units.MSun
8 N_stars = 1000
9
10 converter = nbody_system.nbody_to_si(total_mass, virial_radius)
11
12 bodies = new_plummer_model(N_stars,
13                             convert_nbody=converter,
14                             do_scale=True)
15 bodies.radius = 1 | units.RSun
16
17 # Step 2, creation of an instance of the code
18 grav = ph4(converter,
19             mode = 'cpu',
20             number_of_workers=1)
21
22 grav.initialize_code()
23 grav.parameters.epsilon_squared = (1 | units.RSun) ** 2
24
25 # Step 3, adding particles
26 grav.particles.add_particles(bodies)
27 grav.commit_particles()
28
29 # Step 4, enable collision detection
30 sc_col_det = grav.stopping_conditions.collision_detection
```

Code 4.6: Example setup for Ph4 with particles

In Code 4.6, we have created a converter. The principal function of the converter is to translate units from the N-body system of units (see Section 4.1.1) to the International System of Units (SI). The converter needs the virial radius and total mass, then escalates quantities such that one N-body unit of length is equal to the virial radius. When the instance of `ph4` is created, we provide the converter, the `mode` (where options are `'gpu'` and `'cpu'`), and the `number_of_workers`, that controls the number of CPUs used in parallel calculations. Since the stopping condition for detection of collisions is enabled, it is necessary to provide a function to handle collisions. We can then evolve the instance of `grav`. The next listing shows a minimal evolution loop:

```

1  from amuse.community.ph4.interface import ph4
2  from amuse.units import units, nbody_system
3  from amuse.ic.plummer import new_plummer_model
4
5  t_end = 1000 | units.yr
6
7  # Principal loop
8  while grav.model_time < t_end:
9      grav.evolve_model(t_end)
10
11     if sc_col_det.is_set():
12         solve_collision(grav, sc_col_det)

```

Code 4.7: Loop with evolution for Ph4

where `solve_collision` is a function that handles the collisions. Note that when a collision is detected, the code stops before reaching the final time. Then, if there is no function to correctly solve the collisions, the loop could be infinite. Since the structures provided by AMUSE are generic, the initialization of most codes is very similar.

4.3.1. Ph4

The principal parameters of `ph4` are summarized in Table 4.3.

In our simulations we use a gravitational softening of $0.05 R_{\odot}$. Since the smallest stars are solar-mass stars, the gravitational softening introduces only minor deviations from the real force while preventing singularities. To use the GPU, it is necessary to set `mode='gpu'` during initialization. By default, the `ph4` instance is created with `mode='cpu'`. All our simulations run on the CPU, with `number_of_workers=4` for parallel calculations.

Table 4.3: Key runtime parameters for the PH4 N-body code. Parameters shown with their default and the values used in this work.

Parameter	Description	Default	Used
epsilon_squared	Gravitational softening length squared	0.0	$(0.05 R_{\odot})^2$
timestep_parameter	Accuracy parameter for Hermite integrator	0.14	0.01
force_sync	Force synchronization of particles	0 (False)	1 (True)
manage_encounters	Close encounter management	4	4
initial_timestep_fac	Initial timestep factor	0.0625	0.0625
initial_timestep_limit	Initial timestep limit	0.03125	0.03125

The smaller timestep parameter (0.01) increases the accuracy of the Hermite integrator, which is necessary to resolve close encounters and collisions accurately. The gravitational softening is set to a small fraction $((0.05 R_{\odot})^2)$ of the stellar radii to avoid singularities while minimally affecting the dynamics.

4.3.2. Fi

To initialize Fi, we create an instance of `amuse.community.fi.interface.Fi`. However, SPH particles are stored as `gas_particles`. The following example shows how to initialize Fi with SPH particles following a Plummer distribution:

```

1  from amuse.community.fi.interface import Fi
2  from amuse.units import units, nbody_system
3  from amuse.ic.gasplummer import new_plummer_gas_model
4
5  # Creation of SPH particles
6  gas_pl = gasplummer.new_plummer_gas_model
7  SPH_bodies = gas_pl(N_gas,
8                      convert_nbody=converter,
9                      do_scale=True)
10
11 # Creation of Fi instance
12 SPH = Fi(converter, mode='openmp', number_of_workers=8)
13 SPH.initialize_code()
14
15 # Adding gas particles
16 SPH.gas_particles.add_particles(SPH_bodies)

```

Code 4.8: Example setup for FI with gas particles

Now, we present the main parameters used in Fi, summarized in Table 4.4. In our simulations,

Fi runs with 8 parallel processes, specified by `Fi(number_of_workers = 8)` during initialization.

Table 4.4: Key runtime parameters for the Fi SPH code. Parameters shown with their default and the values used in this work.

Parameter	Description	Default	Used
<code>integrate_entropy_flag</code>	Integrate entropy (True) or internal energy (False)	True	False
<code>radiation_flag</code>	Include radiative cooling/heating	False	False
<code>star_formation_flag</code>	Include star formation	False	False
<code>use_hydro_flag</code>	Enable SPH hydrodynamics	True	True
<code>eps_is_h_flag</code>	Gravitational softening = SPH smoothing length	True	True
<code>gamma</code>	Adiabatic index (polytropic)	1.6666667	1.6666667
<code>n_smooth</code>	Target number of SPH neighbours	64	64
<code>alpha</code>	Artificial viscosity parameter (Monaghan)	0.5	0.5
<code>beta</code>	Artificial viscosity parameter (Monaghan)	1.0	1.0
<code>courant</code>	Courant condition parameter	0.3	0.3
<code>acc_timestep_flag</code>	Use acceleration-based timestep criterion	True	True
<code>timestep</code>	Initial system timestep	1.0	1.0

The only parameter modified from its default is `integrate_entropy_flag`, which is set to False to evolve the specific internal energy instead of the entropic function. This choice provides better numerical stability for energy conservation (see Section 3.3). All other parameters are left at their default values.

4.3.3. Bridge

Finally, we present the parameters used in our BRIDGE implementation. Although the implementation¹ runs in Python and is independent of AMUSE, it is designed to work with AMUSE data structures. The principal parameters used to initialize BRIDGE are summarized in Table 4.5.

4.3.4. Code validation

To verify the accuracy and stability of our numerical setup, we present the results of a test simulation. Figure 4.1 shows the relative energy error $\Delta E/E$ (top panel) and the virial ratio $2K/|U|$ (bottom panel) as a function of time over 50 crossing times. The test system consists of a star

¹ This implementation was developed by B. Reinoso, a co-author of the paper presented in Chapter 5.

Table 4.5: Key runtime parameters for the BRIDGE coupling scheme. GPU acceleration is enabled via `mode=1` for the FASTKICK force calculations. The N-body and SPH components run on CPU (PH4 with 4 workers, Fi with 8 workers). The parameter `do_accretion` is set to False in our simulations, as we do not include gas accretion.

Parameter	Description	Value
<code>mode</code>	Use GPU-accelerated kicks (1) or CPU (0)	1 (GPU enabled)
<code>f_coup</code>	Coupling safety factor for timestep calculation	0.5
<code>eps_nb</code>	Softening length for gravitational coupling	5 AU
<code>dt_bridge</code>	Maximum BRIDGE timestep (N-body units) ¹	1.0
<code>do_accretion</code>	Use accretion implementation	False

¹ The BRIDGE timestep is adaptive and determined by the gas free-fall time and the coupling timescales (t_{ff} , $\Delta t_{\text{coup,gas}}$, $\Delta t_{\text{coup,stars}}$). The value listed here is the maximum allowed timestep in N-body units, which is adjusted downward as needed.

cluster of 5000 stars of $1 M_{\odot}$ (total stellar mass $M = 5000 M_{\odot}$), with virial radius $R_v = 1.0$ pc, embedded in a gas cloud of mass $5000 M_{\odot}$ represented by 5000 SPH particles. The initial crossing time of the system is 0.1491 Myr.

The relative energy error remains below $\sim 10^{-4}$ throughout the simulation, confirming that the integration scheme conserves energy to high accuracy. The virial ratio fluctuates around unity, indicating that the system remains in virial equilibrium. These results validate the numerical setup described in this chapter.




Figure 4.1: Energy conservation and virial equilibrium for a test simulation of a gas-embedded star cluster. Top panel: relative energy error $\Delta E/E$. Bottom panel: virial ratio $2K/|U|$. The system remains in virial equilibrium with an energy error of $\lesssim 10^{-4}$ over 50 crossing times.

Chapter 5

Results

5.1. Statement about my contribution

This section presents the research article entitled "The role of a gaseous component on the collisional runaway path to supermassive black hole seeds". My contributions include the derivation of the generalized critical mass, which incorporates gravitational focusing and an external potential, as well as the definition of the relaxation mass scale. I performed all the simulations, carried out the data analysis, and produced all the tables and figures shown in the article. All the authors contributed to the discussion and to the revision of the manuscript. In particular, Bastián Reinoso developed the BRIDGE implementation used for the coupling strategy.

The role of gas on the collisional runaway path to supermassive black hole seeds

B. Cuevas¹, B. Reinoso², A. Escala¹, M. C. Vergara^{1,3}, S. Valdebenito¹, and D. R. G. Schleicher⁴

¹ Departamento de Astronomía, Universidad de Chile, Casilla 36-D, Santiago, Chile

² Department of Physics, Gustaf Hållströmin katu 2, FI-00014, University of Helsinki, Finland
e-mail: bastian.reinoso@helsinki.fi

³ Departamento de Astronomía, Facultad de Ciencias Físicas y Matemáticas, Universidad de Concepción, Concepción, Chile

⁴ Dipartimento di Fisica, Sapienza, Università di Roma, Piazza le Aldo Moro 5, 00185 Roma, Italy

Received August 27, 2026

ABSTRACT

Context. The origin of supermassive black holes (SMBHs) remains unclear, despite observational evidence for their rapid formation in the early Universe, as evidenced by active galactic nuclei (AGN) at high redshift.

Aims. In this paper, we investigate stellar dynamics in dense, gas-embedded clusters. We explore how gas influences the runaway collision process and the subsequent formation of SMBH seeds, in a global instability scenario.

Methods. We perform a series of coupled N-body and SPH simulations using the AMUSE framework, with the stellar and gaseous components interacting gravitationally via the BRIDGE method. We use 12 initial condition models with equal-mass stars following Plummer distributions, with total masses ranging from 10^3 to $5 \times 10^5 M_\odot$, virial radii from 0.005 to 0.1 pc, and stellar masses of 50, 10 and $1 M_\odot$. For each model, we perform four simulations with gas fractions ranging from 0% to 20%, resulting in a total of 48 simulations. The gas is sampled with the same spatial distribution as the stars, with a mass ratio per particle of $m_{\text{SPH}}/m = 0.01$, where m is the stellar mass.

Results. We find that in most of our simulations, higher gas fractions lead to the formation of more massive black holes, as the core collapses earlier, triggering collisions through gas-star interactions. Family N models, which lie in the stable regime, form only a negligible BH even at high gas fractions. Our results show increases in BH mass of $\sim 8\%$ for low gas fractions, to $\sim 44\%$ at 20% gas fraction. The core collapse time is also shortened, with the strongest reduction at 20% gas fraction (reduction of $\sim 27\%$).

Conclusions. The transonic regime of our initial conditions is where dynamical friction is most efficient in principle. Although our simulations do not fully resolve this effect, and in the absence of other dissipation mechanisms (e.g., supernovae, stellar feedback, cooling), gas-star interactions are the principal mechanism driving the enhancement in BH formation efficiency. This implies that the presence of gas during runaway collisions can lead to the formation of more massive SMBH seeds.

Key words. Galaxies: star clusters: general – Gas dynamics – Hydrodynamics – Methods: numerical – Stars: black holes – Stars: kinematics and dynamics

1. Introduction

The origin of supermassive black holes (SMBHs) is currently a topic of active research, and the formation mechanism remains unclear (Woods et al. 2019). There is evidence of SMBHs with masses in the range $\sim 10^6$ – $10^{10} M_\odot$ (Natarajan & Treister 2009; Volonteri 2010; Woods et al. 2019; Latif & Ferrara 2016; King 2015). However, the origin of SMBHs becomes more enigmatic with the observations of quasars at $z \gtrsim 6$ (Willott et al. 2007; Jiang et al. 2015), and the more recent detection of 31 quasars at $6.6 < z < 7.8$ from the Euclid Wide Survey (Yang et al. 2026). In particular, Mortlock et al. (2011) reported a quasar at $z \sim 7.085$ (0.77 Gyr after the Big Bang) hosting a black hole of $2 \times 10^9 M_\odot$.

These observations imply that the seeds of the first SMBHs must have formed and grown on very short timescales (Latif & Ferrara 2016). Several formation channels have been proposed to produce SMBH seeds, each with distinct initial masses and associated challenges. The remnants of Population III stars are expected to form light seeds of $\sim 10^2 M_\odot$ (Fryer et al. 2001). These seeds can grow into intermediate-mass black holes (IMBHs) in dense early globular clusters through gas accretion (Vesperini et al. 2010); collisions between Pop

III stars can also contribute to building more massive seeds (Vergara et al. 2021; Reinoso et al. 2018, 2020), and both mechanisms may occur simultaneously (Boekholt et al. 2018). However, their growth to SMBH masses requires sustained Eddington accretion or super-Eddington bursts, which are difficult to achieve (Regan & Volonteri 2024; Smith et al. 2018). The direct collapse of primordial gas clouds can form heavier seeds of $\sim 10^5 M_\odot$ (Bromm & Loeb 2003; Lodato & Natarajan 2006; Latif & Schleicher 2015; van Dokkum et al. 2025); but this pathway requires specific conditions to suppress fragmentation and maintain effective cooling (Shang et al. 2010; Suazo et al. 2019). Moreover, direct collapse black holes (DCBHs) typically form in metal-free environments, and subsequent accretion can be severely suppressed by supernovae (SNe) feedback and by the large relative velocity that the BH acquires due to its formation at ~ 1 kpc from the galactic centre (Chon et al. 2021). Runaway collisions in dense stellar clusters provide another pathway, efficiently forming IMBHs with masses of $\sim 10^3$ – $10^4 M_\odot$, which can serve as seeds for SMBHs (Sakurai et al. 2017; Chung et al. 2026). However, current simulations of the collisional runaway typically produce IMBHs rather than SMBHs (Portegies Zwart & McMillan

2002; Vergara et al. 2023; Vergara et al. 2026; Reinoso et al. 2018), partly due to the computational cost of simulating the large number of stars required to model more massive clusters. Once formed, these IMBHs could grow through subsequent gas accretion and mergers with other BHs (Inayoshi et al. 2022; Kritos et al. 2023; Inayoshi et al. 2016; Chung et al. 2026; Pelupessy et al. 2007).

Galactic centers are one of the most likely places where SMBHs can form, since the process of mass segregation and dynamical friction (DF) can funnel material to the center (Shlosman et al. 1990; Escala 2006; Naiman et al. 2015). Besides SMBHs, nuclear stellar clusters (NSCs) can also be found in galactic centers (Neumayer et al. 2020; Schiavi et al. 2021), with both referred to under the nomenclature of central massive objects (CMOs; Ferrarese et al. (2006)). There are correlations between properties of the host galaxy and the CMO it contains, including the M – σ relation between BH mass and bulge velocity dispersion (Gültekin et al. 2009; Tremaine et al. 2002), the $M_{\text{BH}}-M_{\text{bulge}}$ relation (Häring & Rix 2004; Marconi & Hunt 2003), as well as correlations between the host galaxy mass and the mass of the SMBH or NSC it harbors (Ferrarese et al. 2006). These correlations suggest that there could be a link in the origin of SMBHs and NSCs (Escala 2021). NSCs dominate in galaxies with masses $\lesssim 10^{10} M_{\odot}$, while SMBHs dominate in those with masses $\gtrsim 10^{12} M_{\odot}$; in the intermediate-mass regime, both types coexist (Georgiev et al. 2016). Runaway collisions are the primary candidate for forming a massive BH in the center (Escala 2021; Chung et al. 2026; Das et al. 2021). Mass segregation and the Spitzer instability drive the migration of the most massive stars toward the center. This increase in the core density triggers a runaway collision process, forming a massive BH as a consequence (Gürkan et al. 2004; Rasio et al. 2003).

Observations with the James Webb Space Telescope (JWST; Gardner et al. (2023)) have discovered a considerable population of SMBHs at $5 \lesssim z \lesssim 11$ (Schneider et al. 2023; Natarajan et al. 2024; Eilers et al. 2023; Harikane 2025), as well as young massive clusters (YMCs) that could eventually form IMBHs through runaway collisions (Schleicher et al. 2026). The presence of such SMBHs challenges the light seed scenario, since Pop III remnants would require sustained super-Eddington accretion to grow to these masses (Kiyuna 2026). In contrast, heavy seeds can reach SMBH masses through Eddington-limited accretion (Jeon et al. 2025; Silk et al. 2024). Recent observations of extremely red, point-like objects, known as little red dots (LRDs), may help elucidate the origins of early SMBHs, as they could represent an intermediate population between seeds and the observed SMBHs (Vaida & Farber 2026). Observations of LRDs span a redshift range of $3 \lesssim z \lesssim 10$. LRDs were initially identified as extremely red galaxies (Endsley et al. 2023; Labbé et al. 2023). There is no consensus on their nature. Evidence supporting an AGN interpretation includes the detection of high-ionization coronal lines, such as [FeX], and broad H α emission (Kocevski et al. 2023; Furtak et al. 2024). Although many LRDs are radio and X-ray quiet, this could be expected from super-Eddington accretion models, where thick disks suppress high-energy emission (Secunda et al. 2026). From a stellar interpretation, LRDs are thought to be dusty, intensely star-forming galaxies (Escala et al. 2025; Kritos & Silk 2026; Rantala 2026). If this is the case, these objects would be among the densest stellar systems ever observed, with core densities as high as $\sim 10^8 M_{\odot} \text{ pc}^{-3}$, largely exceeding the threshold for runaway collisions, and may include tidal disruption events (TDEs) in some models (Bellovary 2025).

In this paper, we investigate the role of gas in the formation of SMBH seeds. We test the scenario of global instability in NSCs proposed by Escala (2021) through coupled N-body/SPH simulations, by selecting a set of initial conditions from Vergara et al. (2023) and adding a gas component. We first describe the scenario in Section 2 and then present the numerical methods in Section 3. We present the main results in Section 4. We discuss the results and their implications for SMBH seed formation, along with caveats, in Section 5. Finally, we summarize the main results and outline considerations for future work in Section 6.

2. The collisional runaway scenario for SMBH seeds

A promising channel for the formation of massive black hole seeds involves runaway stellar collisions in dense stellar clusters Escala (2021). This scenario is motivated by the observation that NSCs and SMBHs are found in galactic centers (Georgiev et al. 2016), suggesting a potential evolutionary link. While stellar collisions are negligible in typical galactic fields (Binney & Tremaine 2008), dense star clusters could have reached densities high enough for the collision timescale (t_{coll}) to become shorter than a characteristic timescale t_{H} (e.g., the age of the system or the Hubble time), allowing collisions to become a dominant dynamical process. Escala (2021) established a critical density criterion, derived from $t_{\text{coll}} < t_{\text{H}}$, to define this unstable, collision-dominated regime. This work showed that observed NSCs generally avoid this parameter space, whereas systems with massive black holes (MBHs) are consistent with having undergone such a phase.

2.1. Timescale criteria and the instability regime

The viability of the collisional runaway scenario depends on the interplay between three fundamental timescales: the collision time (t_{coll}), the relaxation time (t_{relax}), and a characteristic timescale (t_{H}). Following the scenario proposed by Escala (2021), the collisional runaway predicts two global collisional instabilities, both defined by the condition that the collision time is comparable to or shorter than the age of the system ($t_{\text{coll}} \leq t_{\text{H}}$). In the first regime, defined by systems where $t_{\text{coll}} < t_{\text{relax}} (\leq t_{\text{H}})$, collisions dominate before relaxation can act, leading to runaway collisions that form a massive remnant without the system being able to expand (Escala et al. 2025). In the second regime, defined by systems where $t_{\text{relax}} < t_{\text{coll}} (\leq t_{\text{H}})$, the system is initially in the collisional regime but expands before collisions are triggered, as two-body relaxation operates on shorter timescales. However, core collapse could be present in this scenario (Portegies Zwart & McMillan 2002), enhancing collisions within the core. We now derive the relations that quantify these regimes, following the approach of Vergara et al. (2023).

Consider a stellar system of total mass M and characteristic radius R , composed of N stars of mass m and radius $r_{\star}$. We assume the system is virialized, with a velocity dispersion $\sigma = (GM/R)^{1/2}$. The collision time can be estimated as $t_{\text{coll}} = (n\Sigma\sigma)^{-1}$, where $n = 3M/(4\pi R^3 m)$ is the number density and Σ is the effective cross-section (Binney & Tremaine 2008). Identifying terms in the collisional rate $1/t_{\text{coll}} = 16\sqrt{\pi}n\sigma r_{\star}^2(1 + \Theta)$, the cross-section can be defined as $\Sigma_0 = 16\sqrt{\pi}r_{\star}^2(1 + \Theta)$, where $\Theta = Gm/(2r_{\star}\sigma^2)$ is the Safronov number. Substituting these expressions yields $t_{\text{coll}} = (4\pi m R^{7/2})/(3\Sigma_0 G^{1/2} M^{3/2})$.

For a typical velocity dispersion of $\sim 100 \text{ km s}^{-1}$, the Safronov number becomes $9.54(m/M_{\odot})(R_{\odot}/r_{\star})$ (Binney & Tremaine 2008). Consequently, the effective

cross-section is determined primarily by the stellar radii, corresponding to the limit where the Safronov number is small. Then, the condition for collisions to be significant over the age of the system, $t_{\text{coll}} \leq t_H$, leads to a mass-radius relation that serves as the first boundary in an M - R diagram:

$$M \geq \left(\frac{4\pi m}{3\Sigma_0 t_H G^{1/2}} \right)^{2/3} R^{7/3}. \quad (1)$$

If, instead, we include the virialized velocity dispersion in the Safronov number, the dependence of Σ_0 on σ translates into a dependence on the total mass M through $\sigma = \sqrt{GM/R}$. This distinguishes our derivation from that of Vergara et al. (2023), where the cross-section is dominated by stellar radii and the Safronov number is neglected. Moreover, in the absence of an external potential, M denotes the stellar mass. When an external potential of mass M_{ext} is present, the total mass is $M_{\text{tot}} = M + M_{\text{ext}}$, and the velocity dispersion increases to $\sigma = \sqrt{GM_{\text{tot}}/R(1+q)}$ (Reinoso et al. 2020), where $q = M_{\text{ext}}/M$. This makes Eq. (1) an implicit relation for M . Solving this relation for M (see Appendix A) yields:

$$M_{\text{crit}} = \left(\frac{\beta R^{7/2}}{2} + \sqrt{\frac{\beta^2 R^7}{4} + \frac{\alpha^3 R^3}{27}} \right)^{2/3} + \left(\frac{\beta R^{7/2}}{2} - \sqrt{\frac{\beta^2 R^7}{4} + \frac{\alpha^3 R^3}{27}} \right)^{2/3} - \frac{2\alpha R}{3}, \quad (2)$$

where

$$\alpha = \frac{9.54 m R_{\odot} v_{100}^2}{r_{\star} G M_{\odot} (1+q)^2}, \quad \beta = \frac{\sqrt{\pi} m}{12 G^{1/2} r_{\star}^2 (1+q) t_H}. \quad (3)$$

Here, $v_{100} = 100 \text{ km s}^{-1}$. The external potential is assumed to have the same virial radius and mass distribution as the stellar system. Eq. (2) is the marginal condition for $t_{\text{coll}} = t_H$. Systems that exceed this mass ($M > M_{\text{crit}}$) have $t_{\text{coll}} < t_H$.

The relaxation time, which characterizes the timescale for stellar orbits to be significantly altered by cumulative weak encounters, is given by Binney & Tremaine (2008): $t_{\text{relax}} = t_{\text{cross}} \frac{0.1N}{\ln N}$, where $t_{\text{cross}} = R/\sigma$, and $\sigma = \sqrt{GM/R}$ for virialized system. Replacing the number of stars $N = M/m$, the relaxation time can be written as $t_{\text{relax}} = (0.1 R^{3/2} M^{1/2}) / (m G^{1/2} \ln(M/m))$.

The condition $t_{\text{relax}} \leq t_H$ defines a second relation:

$$R \leq \left(\frac{t_H m \ln(M/m)}{0.1} \right)^{2/3} \left(\frac{G}{M} \right)^{1/3}. \quad (4)$$

Solving $t_{\text{relax}} = t_H$ for M including the effects of an external potential ($M = M_{\text{star}}(1+q)$) yields the mass scale (Appendix B):

$$M_{\text{relax}} = \frac{m}{(1+q)^2} \exp\left(-2W_{-1}\left(-\frac{1}{2\lambda}\right)\right), \quad (5)$$

where W_{-1} is the lower branch of the Lambert W function evaluated at $z = -1/(2\lambda)$, and $\lambda = t_H \sqrt{mG}/(0.1 R^{3/2} (1+q)^2)$. We use the lower branch of the Lambert function since it is physically more relevant (see Appendix B.1). The relation in Eq. (5) defines the characteristic mass scale above which a system of radius R can evolve over a time t_H without two-body relaxation processes significantly altering its initial conditions.

The timescales t_{coll} and t_{relax} define three distinct dynamical regimes for a stellar system of mass M , radius R and external potential of mass ratio q . The collisional regime (CR) occurs when $t_{\text{coll}} \leq t_H \leq t_{\text{relax}}$ (i.e., $M \geq M_{\text{crit}}$ and $M \geq M_{\text{relax}}$),

where collisions dominate before two-body relaxation can expand the cluster, potentially leading to a global collapse and the formation of a massive BH. The stable regime (SR) is defined by $t_{\text{relax}} \leq t_H \leq t_{\text{coll}}$ ($M \leq M_{\text{crit}}$ and $M \leq M_{\text{relax}}$), where relaxation expands the cluster before collisions become relevant. Finally, we define an intermediate regime (IR) by $t_{\text{relax}} \leq t_{\text{coll}} \leq t_H$ ($M_{\text{crit}} \leq M \leq M_{\text{relax}}$), where two-body relaxation begins to expand the cluster before global collisions become frequent; however, the core may still undergo collisional collapse while the outer envelope expands. The described regimes characterize the stability of the system against a global collisional collapse. In the CR, we expect that most of the mass ends up in a central BH, resulting in a BH-dominated system. In the SR, the gravitational potential remains dominated by the stellar cluster. The IR represents an intermediate case, where a broad range of BH masses may form embedded in a surrounding NSC.

3. Numerical methods and initial conditions

We model the coupled evolution of stars and gas using the Astrophysical Multipurpose Software Environment, AMUSE¹ (Portegies Zwart et al. 2009, 2013; Pelupessy et al. 2013; Portegies Zwart & McMillan 2018). The stellar component is evolved with a pure N-body code, while the gas is modeled with the smoothed particle hydrodynamics (SPH) method (Gingold & Monaghan 1977). The two components interact gravitationally via the BRIDGE method (Fujii et al. 2007). We provide more details below.

3.1. Stellar dynamics and collision treatment

The stellar dynamics are computed with the pure N-body code PH4 (McMillan et al. 2012), which implements a 4th-order Hermite integrator (Makino & Aarseth 1992), with adaptive block time steps. Stars are treated as point masses with an associated physical radius.

A collision between two stars is detected when the distance between their centers is less than the sum of their radii, $|\mathbf{x}_i - \mathbf{x}_j| < R_i + R_j$. When a collision is detected, the integration is interrupted and the two stars are replaced by a new single star. The merger product inherits the total mass, the center-of-mass position, and center-of-mass velocity of the progenitors. The new radius is calculated under the assumption of constant density: $R_{\text{coll}} = R_i((m_1 + m_2)/m_i)^{1/3}$. This simple treatment conserves linear momentum and mass.

3.2. Gas dynamics

The gas is evolved using the SPH code FI (Gerritsen & Icke 1997), which solves the hydrodynamic equations in Lagrangian form. The code utilizes a cubic spline kernel for interpolation (Monaghan 1992), with an adaptive smoothing length to keep a total of 64 neighbors. Self-gravity of the gas is computed using a Barnes-Hut tree code (Barnes & Hut 1986), with a softening length that is equal to the smoothing length. The gas is modeled as an ideal, adiabatic fluid without radiative cooling or external heating.

Although fully conservative equations for entropy and energy are implemented in some SPH codes (Springel & Hernquist 2002), we set up the FI so that it evolves the internal energy, which consequently becomes the relevant thermodynamic quantity. For an ideal adiabatic gas with adiabatic index γ , the equa-

¹ <https://github.com/amusecode/amuse>

tion of state is $P = A\rho^\gamma$, where P is the pressure and ρ the density. In SPH, the equation of state is written in terms of the specific internal energy u :

$$P = (\gamma - 1)\rho u. \quad (6)$$

The sound speed, which is the relevant dynamical quantity in the absence of radiative cooling and heating, is given by (Horedt 2004):

$$c_s = \sqrt{\frac{\partial P}{\partial \rho}} = \sqrt{\frac{\gamma P}{\rho}} = \sqrt{\gamma(\gamma - 1)u}. \quad (7)$$

3.3. Gravitational coupling

The gravitational interaction between the stellar (PH4) and gaseous (FI) systems is handled by the BRIDGE coupling scheme (Fujii et al. 2007). This method is necessary to efficiently integrate systems with very different intrinsic time scales. BRIDGE uses a symplectic, kick-drift-kick (Verlet) integrator to couple the subsystems. In this scheme, the gravitational force from one subsystem on the other is applied as impulses (kicks) for a half timestep, updating the velocity. Then, each subsystem is evolved independently for a full timestep (drift). Finally, a kick is applied for a half timestep. The gravitational force between gas and stars is calculated with a Plummer softening kernel and with a softening length given by

$$\epsilon_{g,s} = 0.5(h + \epsilon_{\text{coup}}), \quad (8)$$

where h is the smoothing length of the gas particle and $\epsilon_{\text{coup}} = 5$ AU is the coupling softening length. The initial BRIDGE timestep is chosen to be the crossing time of the stellar component, and we implement an adaptive timestep Δt_{BRIDGE} . After every BRIDGE timestep, Δt_{BRIDGE} is set to be the minimum between the free-fall time in the densest SPH particle and a Courant-type timestep that depends on the acceleration between gas and star particles. These timesteps are calculated as

$$t_{\text{ff}} = \sqrt{\frac{3\pi}{32G\rho_{\text{max}}}}, \quad (9)$$

$$\Delta t_{\text{coup,gas}} = \min\left(f_{\text{coup}} \sqrt{\frac{2h}{a_{g,s}}}\right) \quad (10)$$

$$\Delta t_{\text{coup,stars}} = \min\left(f_{\text{coup}} \sqrt{\frac{\epsilon_{\text{coup}}}{a_{s,g}}}\right) \quad (11)$$

$$\tau = \min(t_{\text{ff}}, \Delta t_{\text{coup,gas}}, \Delta t_{\text{coup,stars}}), \quad (12)$$

where ρ_{max} is the maximum density among SPH particles. h and $a_{g,s}$ are the softening length and the highest acceleration among all SPH particles due to the stars. ϵ_{coup} is set to be 5 AU, and $a_{g,s}$ is the highest acceleration among all stars due to the gas. $f_{\text{coup}} = 0.5$ is a free parameter that controls the time-step accuracy.

Finally, the BRIDGE timestep is set to fulfill the condition

$$0.5\tau \leq \Delta t_{\text{BRIDGE}} \leq \tau. \quad (13)$$

This adaptive timestep ensures stability against strong gas-stars gravitational interactions.

3.4. Initial conditions

In order to test the scenario proposed by Escala (2021) we simulated the same initial conditions as Vergara et al. (2023), keeping the same model IDs to facilitate comparison. For each model, we perform four simulations with different gas fractions, using the same stellar initial conditions for all gas fractions. Table D.2 summarizes the initial conditions for the stellar component, where ID is the model identifier, R_v is the initial virial radius, M is the total mass, N is the number of stars, m is the initial star mass, r is the initial star radius and σ is the velocity dispersion. Each model is a Plummer distribution (Plummer 1911) with equal-mass stars. The initial virial radii are $R_v = 0.005, 0.01, 0.1$ pc, and the number of stars $N = 500, 1000, 5000, 10000$. The stellar masses and radii are the same as in Vergara et al. (2023): for $m = 1, 10$ and $50 M_\odot$, we adopt $r = 1, 4.7$ and $11.7 R_\odot$, respectively, following the mass-radius relations of Bond et al. (1984) and Demircan & Kahraman (1991). Due to computational limitations, we simulate clusters that are less massive but more compact than real NSCs. This allows us to explore the instability on shorter timescales ($t_H = 5$ Myr) while maintaining the same physical regime.

Fig. 1 shows the initial conditions of models in Table D.2 (see Appendix D.1), in a mass-radius space. The solid lines represent the relaxation mass (Eq. (5)), and the dashed lines are for the critical mass (Eq. (2)), calculated for the final simulation time ($t_H = 5$ Myr) for $q = 0$; the colored areas indicate the range of gas fractions, from $q = 0$ to $q = 0.2$. Black dot-dashed line stands for the condition $t_{\text{coll}} = t_{\text{relax}}$, for $q = 0$. Colors indicate the initial star mass, with yellow for $m = 1 M_\odot$, red for $m = 10 M_\odot$ and blue for $m = 50 M_\odot$. Each model is represented by a different symbol, with the same color convention. We note that family models A, B, C, D, H and I lie in the IR, while family models F, J, K, L, M and N are in the SR, with no models in the CR.




Fig. 1: SMBH formation scenario represented in a mass-radius space. For all symbols and lines, the colors represent different stellar masses, with yellow for $m = 1 M_\odot$, red for $m = 10 M_\odot$ and blue for $m = 50 M_\odot$. Solid and dashed lines are for the relaxation and critical mass, with $t_H = 5$ Myr and $q = 0$, while colored areas indicate the range of gas fractions from $q = 0$ to $q = 0.2$. Black dot-dashed line represents the condition $t_{\text{coll}} = t_{\text{relax}}$ for $q = 0$. The symbols represent the initial conditions of Table D.2.

For each star cluster model, we first generate a Plummer distribution of SPH particles with the same virial radius and total mass as the stellar system (so that the gas would be in virial equilibrium if considered alone). We then scale the gas mass to 5%, 10% and 20% of the stellar mass by adjusting the particle masses while keeping the same spatial distribution. For all models, the SPH particle mass is $m_{\text{SPH}} = m/100$, where m is the stellar mass. The number of SPH particles for a given gas fraction f is then $N_{\text{SPH}} = f \times N \times 100$, with N the number of stars. Since the added gas mass is small compared to the stellar mass, we assume that the overall virial equilibrium of the stellar component is not significantly disturbed.

To characterize the gas further, we consider the initial sound speed profiles (Eq. (7)) for the gas variants of the A family (A_5 , A_{10} and A_{20}) and the N family (N_5 , N_{10} and N_{20}), which are the most and least collisional models families in our sample (in their respective regimes), respectively. For the A gas variants, the stellar velocity dispersion decreases from 186.80 km s^{-1} in the central region (r_5) to 101.60 km s^{-1} in the outer regions (r_{95}), while the gas sound speed falls from 129.89 km s^{-1} at the center to 65.51 km s^{-1} in the outermost layers (see Appendix E).

The fact that $\sigma/c_s > 1$ throughout the stellar cluster has important consequences. The transonic motion of stars through the gas produces an efficient DF, as described by Ostriker (1999), which leads to stars losing orbital energy and sinking toward the center, enhancing the occurrence of collisions in the central region. Although our simulations do not fully resolve DF (see Appendix D.3), the dissipation of kinetic energy due to gas-star interactions is still present. For the N gas variants (N_5 , N_{10} and N_{20}), the velocities are an order of magnitude lower ($\sigma \sim 15 \text{ km s}^{-1}$, $c_s \sim 10 \text{ km s}^{-1}$), but the condition $\sigma/c_s > 1$ still holds.

4. Results

Here we present the main results from our simulations. We use the black hole efficiency defined in Escala (2021):

$$\epsilon_{\text{BH}} = (1 + M_{\text{NSC}}/M_{\text{BH}})^{-1}, \quad (14)$$

where M_{BH} corresponds to the most massive object, and M_{NSC} is the mass of gravitationally bound stars. Considering M_{ESC} as the escaped stars mass, the total initial mass of the stellar system is $M_s = M_{\text{ESC}} + M_{\text{NSC}} + M_{\text{BH}}$. The central massive object is defined as $M_{\text{CMO}} = M_{\text{NSC}} + M_{\text{BH}}$, i.e., the total bound mass of the stellar system. In terms of the efficiency, the BH candidate mass is then $M_{\text{BH}} = \epsilon_{\text{BH}} M_{\text{CMO}}$.

4.1. Dynamical evolution

In this section we present the time evolution of the mass components of the system. We select two model families from different regimes, A and N. For each family, we consider two variants: a stars-only simulation (denoted A_0 and N_0), and simulations with gas masses of 20% of the initial stellar mass (denoted A_{20} , N_{20}). The A variants have an initial stellar mass of $M = 5 \times 10^4 M_\odot$, virial radius $R_{\text{vir}} = 0.005 \text{ pc}$, and is composed of 10^3 stars. The N variants have an initial mass of $M = 10^4 M_\odot$, virial radius $R_{\text{vir}} = 0.1 \text{ pc}$ and 10^4 stars. As shown in Fig. 1, the A family is in the IR, where the collision time is shorter than the simulation time. In contrast, the N family is in the SR where collisions are not globally relevant.

Each row in Fig. 2 corresponds to A_0 and A_{20} , from top to bottom. The left column shows the evolution over the full 5 Myr,

while the right column zooms in to 40 relaxation timescales. The dashed blue line marks the initial collision time, the dot-dashed green line indicates the initial relaxation time, and the dashed black line shows the estimated core collapse time (see Appendix C). We observe that the inner 10% and 30% Lagrangian radii rapidly contract towards the center, on scales of the collision time. After 1 Myr, A_0 and A_{20} show that the Lagrangian radii remain stable, with A_0 showing a slight increase in the 90% Lagrangian radii compared with A_{20} . In the second column we can observe slight differences in the core collapse (slightly shorter for A_{20}), and no considerable changes during $1 t_{\text{relax}}$.




Fig. 2: Lagrangian radii for stars, from 10% (red) to 90% (blue). The top panel corresponds to A_0 , and bottom panel to A_{20} . The left column shows the full simulation time (5 Myr), while the right column shows the evolution up to 40 relaxation times. Blue dashed and green dot-dashed lines are the initial collision and relaxation timescales, respectively. Black dot-dashed lines are the core collapse.

The time evolution of the mass components and ϵ_{BH} of stars for A_0 and A_{20} is displayed in Fig. 3. The top panel shows the escaped mass normalized by the initial stellar mass M_s . We find that approximately 40% of the initial stellar mass escapes during the first 1 Myr of evolution for both variants, and the total escaped mass reaches the range of 50-60% by 5 Myr. The A_0 variant shows only 57.3% of escaped mass vs 50.2% for A_{20} . The second panel shows collisions normalized by the initial number of stars, with solid lines for total collisions and dashed lines for collisions involving the most massive object. We can see that most collisions happen during the first Myr, augmenting as the gas mass increases. During the first Myr, A_0 shows 27.2% of collisions, reaching 27.5% at 5 Myr. A_{20} shows 32.5% of collisions at 1 Myr, and 33.5% of collisions at 5 Myr. The influence of gas is stronger during the early evolution of the cluster, increasing the number of collisions by 19.49% relative to A_0 . The gas also increases the number of collisions at late times. The third panel shows the evolution of the black hole efficiency (Eq. (14)), which is slightly higher for A_{20} . As described in Vergara et al. (2023), the increase in efficiency in the late stage of the system is dominated by the escaped mass, as collisions are less common. The bottom panel shows the mass evolution of the black hole candidate, exhibiting a similar behavior to the collision panel, with a rapid increase at early times, and a negligible increase in late evolution. The final mass of the most massive object is $13\,300 M_\odot$ for A_0 , and $16\,400 M_\odot$ for A_{20} , representing 62.30% and a 65.86% of the CMO mass, respectively. This represents, respectively, 26.6 and 32.8% of the initial stellar mass, which means that most of the collisions end in a single object. From the right column, the black dash-dotted line is the mean core collapse time, where we note that it coincides with the beginning of the onset of runaway collisions, as expected. By the collision




Fig. 3: Mass components and ϵ_{BH} evolution over time of stars. Blue curves are for A_0 , and purple for A_{20} . Top panel is for escaped mass normalized by the total stellar mass. Second panel shows collisions normalized by the initial number of stars, where solid lines are for the total and dashed lines for collisions involving the most massive object. Third panel shows ϵ_{BH} (Eq. 14), and bottom panel shows the mass of the most massive object. Left column is for 5 Myr, with the vertical blue dashed line representing t_{coll} . Right column is a zoom to 40 relaxation times (~ 0.014 Myr), where the dot-dashed black line is the mean core collapse time.

time (vertical dashed blue line), an important part of collisions have occurred, while ϵ_{BH} has reached approximately half of its final value.

Fig. 4 shows the Lagrangian radii of N_0 and N_{20} . For N_0 , the 90% and 70% radii expand slowly, while the 30% and 10% radii contract slightly. N_{20} exhibits qualitatively the same behavior, with no significant differences compared with N_0 .




Fig. 4: Lagrangian radii for stars, from 10% (red) to 90% (blue). The top panel corresponds to N_0 , and bottom panel to N_{20} . Green and black dot-dashed line are the initial relaxation timescales and core collapse, respectively.

The mass and ϵ_{BH} evolution of N_0 and N_{20} is presented in Fig. 5. The top panel shows that the escaped mass is lower than 1%, which means that most of the initial stellar mass remains gravitationally bound. From the second panel, collisions are near zero, which may be related to a long collisional timescale. As there are no significant collisions and escaped mass, the efficiency is near zero, as can be observed in the third panel. Con-

sequently, as can be observed from the bottom panel, the most massive object shows only 4 collisions.




Fig. 5: Mass components evolution over time of stars for four variants of model N. Equivalent to Fig. 3 with no zoom.

Models of family N lie in the regime where $t_{\text{coll}} > t_{\text{H}}$, i.e., collisions are not globally relevant. The results shown in Figs. 4 and 5 are therefore expected: no significant mass loss, negligible collisions, near-zero BH efficiency, and no formation of a massive object.

4.2. Effect of gas on core collapse timescale

For each model, we estimated the core collapse time t_{cc} using the method described in Appendix C. Table 1 shows the ratio $x_{\text{cc}} = t_{\text{cc,gas}}/t_{\text{cc,0}}$ averaged over models, where the columns, from left to right are for models in the IR, SR and all, respectively. Each row corresponds to a gas fraction. In the case of models in the IR, for a gas fraction of 5%, the mean ratio is 0.8765 ± 0.1818 ; for 10%, 0.8345 ± 0.1345 ; and for 20%, 0.7410 ± 0.1094 . All ratios are below 1, indicating a reduced t_{cc} in the presence of gas. The core collapse time tends to be shorter as the gas fraction increases, while the dispersion decreases. The decrease in dispersion with higher gas fraction suggests that the effect becomes more predictable when more gas is present.

For models in the SR, the mean ratio is 1.0111 ± 0.1570 for 5% gas fraction, 0.8571 ± 0.1068 for 10% gas fraction, and 0.7082 ± 0.0533 for 20% gas fraction. The reduction of t_{cc} is less pronounced for SR models at 5% and 10% gas fractions, and at 5% it is almost negligible. The dispersions are smaller for all gas fractions compared with models in the IR. For a gas fraction of 20%, the reduction on t_{cc} is slightly more pronounced in the SR models (0.7082 ± 0.0533) than in the IR models (0.7410 ± 0.1094). Despite these small differences between IR and SR models, the general tendency is to shorten the core collapse time as the gas mass is larger.

Fig. 6 shows the core collapse time as a function of the relaxation time and collision time, on left and right panels respectively. In addition to the described behavior, we observe correlations between the t_{cc} and both the relaxation and collision times:

Table 1: Mean ratio $x_{cc} = t_{cc,gas}/t_{cc,0}$ of core collapse times for gas-embedded models relative to their gas-free counterparts, and standard deviation for IR family models (A, B, C, D, H, I), SR family models (F, J, K, L, M), and the combined sample.

$X_{gas} (%)$	IR		SR		Total	
	$\bar{x}_{cc}$	σ	$\bar{x}_{cc}$	σ	$\bar{x}_{cc}$	σ
5	0.8765	0.1818	1.0111	0.1570	0.9377	0.1836
10	0.8345	0.1345	0.8571	0.1068	0.8448	0.1232
20	0.7410	0.1094	0.7082	0.0533	0.7261	0.0899

t_{cc} decreases with shorter t_{relax} (left panel) and with shorter t_{coll} (right panel).




Fig. 6: Core collapse time as a function of t_{relax} (left panel) and t_{coll} (right panel).

4.3. Critical mass and BH formation efficiency

In this subsection, we analyze the black hole formation efficiency of our models. First, we compare our stars-only results with those of Vergara et al. (2023) to validate our setup. Then, we show how the presence of gas modifies the efficiency as a function of the critical mass, testing the scenario proposed by Escala (2021).

To make a first comparison, Table 2 presents results for the same models as Vergara et al. (2023). From left to right, the columns are: model identifier, escaped mass (in $M_\odot$), black hole mass (in $M_\odot$) and formation efficiency (ϵ_{BH}). For each quantity, two sub-columns show our stars-only results (left) and those of Vergara et al. (2023) (right). For the escaped mass, the differences are small: the ratio of our values to those of Vergara et al. (2023) ranges from less than 1% to approximately 10%, corresponding to less than 5% of the initial stellar mass. In the case of the BH mass, the largest differences occur for the most collisional models (A₀, B₀, C₀, in the IR), with relative differences of 34.11%, 36.36%, and 15.66%, respectively. These correspond to differences of 6.77%, 5.60%, and 2.60% of the initial stellar mass.

These differences probably arise from the different numerical codes used. Our simulations employ PH4 (via AMUSE), while Vergara et al. (2023) use NBODY6++GPU (Wang et al. 2015; Aarseth & Aarseth 2003). Both use a fourth-order Hermite integrator with hierarchical block time-steps. However, NBODY6++GPU additionally implements the Kustaanheimo–Stiefel algorithm (Kustaanheimo et al. 1965) and chain regularization (Mikkola & Aarseth 1993) as special treatments for close encounters, binaries, and multiple systems.

The impact on BH mass translates into efficiency differences. Again, the most collisional models (A₀, B₀, C₀) show a considerable enhancement in BH formation efficiency, with differences of 15.97, 8.87, and 8.17 percentage points, respectively. For the remaining models, the differences are below 4%.

Significant differences are observed for N₀. In our simulations, our values are lower by 82.48% and 95.77% for escaped mass and BH mass, respectively. This discrepancy arises because the relaxation time of model N is relatively long (~ 0.5 Myr) and our simulation time is limited to 5 Myr. In contrast, models L₀ and M₀, despite also having large relaxation times, reach core collapse within 5 Myr (at ~ 0.9 and ~ 2.3 Myr, respectively). Since the simulations of Vergara et al. (2023) extend to 10 Myr, model N₀ has evolved considerably further in their case. Assuming core collapse occurs on a timescale of $\sim 20t_{relax}$ (Quinlan 1996), and given that $t_{relax} \sim 0.5$ Myr for N₀, the core collapse is expected at ~ 10 Myr.

Next, we show results for all models, including their gas variants. Fig. 7 shows, ϵ_{BH} (top panel), and normalized escaped mass (bottom panel). In the left column, quantities are plotted as a function of M/M_{relax} , while in the right column they are plotted as a function of M/M_{crit} . From the bottom panels, we observe a tendency for the escaped mass to grow as M/M_{relax} decreases and M/M_{crit} increases, with no clear distinction between gas variants. There are two processes that produce escaping stars: diffusive or small-angle relaxation evaporates stars slowly from the cluster; on the other hand, two-body encounters produce ejections of high-energy stars, more likely in dense cores (Khalisi et al. 2007). Family models in the IR that are more collisional (A, B, and C) have shorter core collapse and collision times, showing an important fraction of escapers (probably due to ejections). In the other extreme, the N family models, where core collapse is not reached and collisions are not globally relevant, exhibit a small fraction of escapers.

In the top-right panel, we observe a clear tendency for efficiency to grow as M/M_{crit} increases. All models follow the same overall trend, but with a systematic vertical shift: the stars-only variants (blue) lie systematically below the fitted curves, while the 20% gas variants lie above. This is confirmed by the fits. Near the first regime ($M/M_{crit} < 10^{-4}$) and the third regime ($M/M_{crit} > 10^5$), where the efficiency approaches 0 and 1 respectively, these differences become smaller, as all curves converge. To model this effect, we performed a unified fit where the shape of the curve is determined by a common set of parameters (x_0 , k , a), while the gas fraction q introduces a horizontal scaling of the mass ratio. We define an effective mass ratio $x_{eff} = (M/M_{crit})(1 + \alpha_q q)$, where α_q is a free parameter that depends on q . Similar to Vergara et al. (2026), the efficiency is then modeled by the form:

$$\epsilon_{BH} = f(x_{eff}) = (1 + \exp[-k(\log_{10}(x_{eff}) - \log_{10}([x_0]))])^{-a}. \quad (15)$$

The parameters x_0 , k and a were obtained by fitting the stars-only variants ($q = 0$). For each gas fraction, we then fitted a

Table 2: Comparison of stars-only (sub zero variants) simulations with Vergara et al. (2023). Our values are from single realizations; reference values include standard deviations from three realizations.

Model ID	$M_{\text{esc}} [M_{\odot}]$		$M_{\text{BH}} [M_{\odot}]$		$\epsilon_{\text{BH}} [\%]$	
	This work	Vergara+23	This work	Vergara+23	This work	Vergara+23
A ₀	28 650	28 533 ± 147	13 300	9917 ± 143	62.30	46.33 ± 2.5
B ₀	13 300	14 300 ± 687	5250	3850 ± 324	44.87	36.00 ± 3.7
C ₀	5620	5370 ± 42	1920	1660 ± 14	43.84	35.67 ± 0.5
D ₀	2540	2767 ± 111	630	610 ± 43	25.61	27.33 ± 3.1
F ₀	411	372 ± 5	88	85 ± 12	14.94	13.67 ± 2.1
H ₀	21 750	23 137 ± 1966	12 100	11 397 ± 143	42.83	42.67 ± 2.1
I ₀	4500	4253 ± 183	980	1110 ± 83	17.82	19.33 ± 1.7
J ₀	1469	1459 ± 22	435	545 ± 42	12.32	15.67 ± 1.2
K ₀	344	327 ± 8	59	59 ± 2	8.99	8.67 ± 0.5
L ₀	69 300	75 517 ± 1812	64 000	67 450 ± 2585	14.86	15.67 ± 0.5
M ₀	7170	7850 ± 283	5510	6987 ± 246	5.94	7.67 ± 0.5
N ₀	41	234 ± 44	3	71 ± 5	0.03	0.73 ± 0.0

This work: simulations run up to 5 Myr. Vergara+23: simulations of Vergara et al. (2023), run up to 10 Myr. The stellar initial conditions for each of our models correspond to the same letter in Vergara+23; e.g., our model A₀ shares the same stellar initial conditions as model A in Vergara+23.




Fig. 7: In the y-axis, top panel shows the BH efficiency (ϵ_{BH}), and bottom panel the normalized escaped mass ($M_{\text{ESC}}/M_{\text{stars}}$). The left column shows quantities as a function of M/M_{relax} , while the right column shows quantities as a function of M/M_{crit} . Symbols correspond to the different models (see legend above the figure), and colors indicate the gas fraction: blue (0%), green (5%), red (10%), purple (20%). Dashed black lines mark mass ratios equal to unity. The dash-dotted curves in the top-right panel are fits of an asymmetric sigmoid function (Eq. (15)) for each gas fraction, sharing the same shape parameters x_0 , k , a (fitted to the stars-only data) while α_q is adjusted for each gas fraction. The black dash-dotted curve is the model from Vergara et al. (2026)

single parameter α_q . The resulting parameters are $x_0 = 0.039$, $k = 0.349$, $a = -3.163$ and $\alpha_q = 26.7, 35.6, 47.8$ for $q = 0.05, 0.1, 0.2$, respectively. The fitted function shows three regimes. For $M/M_{\text{crit}} < 10^{-4}$, the efficiency is effectively 0. Between 10^{-4} and 10^5 , the efficiency increases approximately linearly with $\log_{10}(M/M_{\text{crit}})$, reaching 70-80%. For $M/M_{\text{crit}} > 10^5$, it asymptotically approaches unity. The main differences appear in the intermediate regime, where the efficiency increases with gas fraction. Compared with the fitted model of Vergara et al. (2026), our models exhibit a smoother transition between the regimes described above.

Table 3 lists the mean values for the normalized escaped mass and efficiencies, where each gas variant is normalized by its corresponding stars-only counterpart. The results are consistent

with Fig. 7. For the escaped mass, the means are within one standard deviation of unity for all gas fractions, indicating no significant difference. In particular, for the 20% gas case, the mean ratio is closest to 1 and the dispersion is the smallest among all gas fractions. The black hole mass ratios show a clearer trend. The mean ratios are 1.0840, 1.2252 and 1.4414 for 5%, 10%, and 20% gas, corresponding to increases of 8.40%, 22.52%, and 44.14% relative to the stars-only cases. Since the variance in escaped mass is negligible compared to the increases in BH mass, the efficiency ratios follow a similar trend, with mean values of 1.1256, 1.2701, and 1.4510 (increases of 12.56%, 27.01%, and 45.10%) for 5%, 10%, and 20% gas, respectively.

Table 3: Ratios of gas runs to stars-only runs. Values are means $\pm$ standard deviation over all models.

Gas fraction	$\epsilon_{\text{BH,gas}}/\epsilon_{\text{BH,0}}$	$M_{\text{BH}}/M_{\text{BH,0}}$	$M_{\text{esc}}/M_{\text{esc,0}}$
5%	1.1256 ± 0.1627	1.0840 ± 0.1589	1.0471 ± 0.0633
10%	1.2701 ± 0.2079	1.2251 ± 0.2057	1.0442 ± 0.0905
20%	1.4510 ± 0.3139	1.4414 ± 0.2132	1.0042 ± 0.0869

5. Discussion

In this work, we have performed simulations for a set of initial conditions that include coupled star-gas variants. We studied the effect of gas on SMBH seed formation in nuclear stellar clusters. Using the Astrophysical Multipurpose Software Environment (AMUSE), we coupled an N-body integrator (PH4) with an SPH code (FI) via the BRIDGE scheme. In this implementation, the interaction between gas and stars is purely gravitational. Using the initial conditions from Vergara et al. (2023) with different gas fractions, we tested the scenario proposed by Escala (2021). Although our stars-only simulations show some differences with respect to Vergara et al. (2023), the simulations with gas show an enhancement in black hole mass and efficiency, as well as an acceleration of core collapse.

5.1. The role of gas on SMBH formation

Stellar collisions in dense star clusters have been studied extensively in N-body simulations. Portegies Zwart et al. (1999) simulated young star clusters and showed that collisions are more frequent than expected from simple timescale estimates, and that they rejuvenate stars, extending their lifetimes before supernova explosions. As stars interact, massive stars tend to reach the core due to mass segregation over the half-mass relaxation timescale (Spitzer 1987). Mass segregation also implies that massive objects, such as binaries or massive stars, can return to the core after being ejected by strong encounters, as a result of DF (Spitzer 1987; Hut et al. 1992). The products of stellar collisions lose kinetic energy, causing them to sink toward the center. DF and mass segregation explain why most collisions involve a single object, and therefore make this an effective mechanism for BH formation. Numerical simulations have shown that energy equipartition is not reached under realistic initial conditions (Trenti & van der Marel 2013). This implies that massive stars remain dynamically decoupled from the lighter population, continuing to segregate to the core and enhancing the collision rate, which favors runaway growth and the formation of a massive central object. Comparing our stars-only results with those of Vergara et al. (2023), we find differences in BH mass ranging from 0% to 36% (or 0% to 6% of the initial star cluster mass), with the largest discrepancies occurring in the most collisional models (A, B, and C).

For family models A, B, C, and H, which lie in the IR, the BH masses are 13 300, 5 250, 1 920, and 12 100 $M_{\odot}$, respectively, placing them in the IMBH mass range ($\sim 10^2$ – $10^5 M_{\odot}$). Although models L and M are in the SR, their BH masses are 64 000 and 5 510 $M_{\odot}$, respectively. This is because models L and M are the most massive clusters; despite being in the SR, their large total mass still allows the formation of a massive BH. In contrast, they show lower efficiencies, meaning that the dynamics are controlled by the star cluster. If we compare models with the same initial mass, we observe that larger M/M_{crit} corresponds to higher BH mass and efficiency. For example, family models C, F, and N, all with an initial mass of $10^4 M_{\odot}$, form BHs

with masses of 1 920, 88, and 3 $M_{\odot}$, and efficiencies of 43.84%, 13.67%, and 0.03%, respectively. This suggests that more massive clusters (eventually in the CR) could form even more massive BHs, with $\epsilon_{\text{BH}} \sim 1$.

The gas can affect the cluster dynamics through two distinct mechanisms. When considered as a static background potential, it increases the stellar velocities, causing a delay in core collapse. As shown by Reinoso et al. (2020), this delay also affects the runaway growth, although for sufficiently long times the BH formation is enhanced, because the cluster maintains its compactness. On the other hand, a realistic gas can exert DF on stars through gas-star interactions (Ostriker 1999; Reinoso et al. 2020; Leigh et al. 2014). In our simulations, the stellar velocity dispersion is transonic ($\sigma \gtrsim c_s$), as shown in Figs. E.3 and E.4. In this regime, DF is expected to be particularly efficient (Ostriker 1999). While our simulations do not fully resolve the DF (see Appendix D.3), the dissipation of kinetic energy due to gas-star interactions is still present. Consistent with this, our simulations show that the efficiencies are enhanced, with increases of $12.56 \pm 16.27\%$, $27.01 \pm 20.79\%$, and $45.10 \pm 31.39\%$ relative to the stars-only simulations for gas fractions of 5%, 10%, and 20%, respectively. For the BH mass, the corresponding increases are $8.40 \pm 15.89\%$, $22.51 \pm 20.57\%$, and $44.14 \pm 21.32\%$. We also find shorter core collapse times. On average, the core collapse time is reduced to $93.77 \pm 18.36\%$, $84.48 \pm 12.32\%$ and $72.61 \pm 8.99\%$ relative to the stars-only variants for gas fractions of 5%, 10% and 20%, respectively. These results show that IMBHs can form in less massive clusters if a gaseous component is present. Although we use a modified critical mass relation compared with Vergara et al. (2023), the fitted function (Eq. (15)) describes the trend of BH efficiency ϵ_{BH} as a function of M/M_{crit} .

5.2. Implications for SMBH formation

We have shown that in our simplified models, the formation of an IMBH is possible, with gas playing an important role by enhancing the number of collisions and, consequently, the final mass of the BH. However, an important question is how massive a BH can become in more realistic NSCs. The sizes and masses of NSCs have been studied extensively (Neumayer et al. 2020), with masses ranging from 10^5 to $10^8 M_{\odot}$. Using the scaling relations from Georgiev et al. (2016), a nuclear star cluster of mass $\sim 10^7 M_{\odot}$ has a minimum effective radius of ~ 4 pc. We estimate the critical mass ratio M/M_{crit} for a cluster of equal-mass stars ($m = 1 M_{\odot}$, $r = 1 R_{\odot}$), finding $M/M_{\text{crit}} = 4.3 \times 10^{-5}$ and 6.9×10^{-5} for $t_{\text{H}} = 5$ Myr and 10 Myr, respectively. Using Eq. (15), the predicted BH efficiencies are ~ 0.014 and ~ 0.017 . Since $M/M_{\text{crit}} \sim 10^{-5}$ and $M/M_{\text{relax}} > 1$ (see Fig. 7), we can assume no mass loss, and the upper limits on the BH mass are $\sim 1.5 \times 10^5 M_{\odot}$ and $\sim 1.7 \times 10^5 M_{\odot}$, respectively. When including a gas fraction of 20% and following the same calculation, the BH efficiencies increase to ~ 0.034 and ~ 0.039 for $t_{\text{H}} = 5$ Myr and 10 Myr, respectively, yielding upper limits of $\sim 3.4 \times 10^5 M_{\odot}$ and $\sim 3.9 \times 10^5 M_{\odot}$. While the collisional and

relaxation timescales are global quantities, the core of a nuclear star cluster could reach densities high enough to collapse into a massive BH, even as the outer stars expand due to relaxation processes (Escala 2021). This suggests that NSCs should have been more compact at their formation, consistent with the observed coexistence of NSCs and SMBHs in intermediate-mass galaxies (Georgiev et al. 2016; Kritos et al. 2023; Escala 2021), and with the absence of SMBHs in low-mass NSCs. Simulations of the most compact clusters, with radius $R \leq 0.5$ pc and mass $10^7 M_\odot$, show that a million solar mass black hole can form within 100 Myr (Kritos et al. 2023). For such a system ($M = 10^7 M_\odot$, $R = 0.5$ pc) composed of stars only, the efficiencies are 0.068 and 0.077, for $t_H = 5$ Myr and 10 Myr, respectively. The escaped mass fraction is negligible, and the BH mass is estimated to be $6.7 \times 10^5 M_\odot$ and $8.9 \times 10^5 M_\odot$. For a time of 100 Myr, the mass ratios are $M/M_{\text{crit}} = 0.042$ and $M/M_{\text{relax}} = 1.080$. The corresponding efficiency is 0.114, and the escaped mass fraction is ~ 0.1 (estimated from Fig. 7; see Section 5.1). In consequence, the estimated BH mass is $1.024 \times 10^6 M_\odot$. Including a 20% of gas fraction, the estimated efficiencies are 0.126 and 0.140 for $t_H = 5$ Myr and 10 Myr, respectively. Despite the simplifications, our estimates show that in more massive NSCs it is possible to form a SMBH of $\sim 10^6 M_\odot$, with gas playing an important role (increasing the BH mass by approximately 50%). However, for realistic NSCs the predicted efficiencies are low ($\leq 5\%$), indicating that the BH would represent only a small fraction of the total NSC mass. For more compact systems ($R \leq 0.5$ pc), a SMBH can form on a timescale of 100 Myr without gas, and within 10 Myr with a 20% of gas fraction.

Several mechanisms have been proposed in the literature for the growth of IMBHs to masses in the SMBH range. Pelupessy et al. (2007) simulated halos containing IMBHs formed from Population III remnants and found that in collapsing halos with $M_{\text{halo}} \leq 10^{10} M_\odot$, accretion at the Eddington rate is not achieved due to stellar feedback. Even in the most massive halos, the growth of SMBH seeds is driven principally by BH–BH mergers. They propose that the BH mass scales with the host halo mass as $M_{\text{BH}} = 5 \times 10^6 M_\odot (M_{\text{halo}}/10^{11} M_\odot)^{0.78}$. In contrast, Shi et al. (2024) describe a scenario in which subclusters and IMBHs form within a gas cloud complex of mass $10^8 M_\odot$. The hierarchical merging of subclusters occurs on a timescale of $\lesssim 1$ Myr, and the BHs in the subclusters migrate to the center within the same period. The formation of a disk ensures conditions for large accretion rates ($\sim 1\text{--}10 M_\odot \text{yr}^{-1}$), allowing an IMBH to grow to an SMBH of $\geq 2 \times 10^6 M_\odot$ within 1 Myr. Chung et al. (2026) showed through simulations that an IMBH can form in dense stellar clusters, with masses ranging from ~ 300 to $\sim 5000 M_\odot$ via runaway collisions. The resulting BHs can grow to $\sim 6 \times 10^3 M_\odot$ on a timescale of ~ 10 Myr through Eddington-limited gas accretion and TDEs. If a steady gas supply is assumed, the BHs can reach masses of $\sim 6 \times 10^5 M_\odot$, resulting in promising SMBH seeds. These mechanisms are consistent with the formation of IMBHs in dense NSCs (such as those produced in our simulations) and suggest that gas accretion and IMBH mergers provide a viable pathway for seeds to grow to the SMBH regime.

5.3. Caveats

In this section, we summarize the simplifications made in our simulations, which should be taken into account for improvements in future work. As we described in Section 3.2, the gas is evolved using an SPH method, with self-gravity and hydrodynamics, but without radiative cooling or heating. Radiative

cooling allows the gas to lose thermal energy, which can help it remain gravitationally bound for longer periods. As shown by Silich & Tenorio-Tagle (2018), in sufficiently massive and compact clusters, radiative cooling can prevent the development of a global star cluster wind, thus suppressing gas expulsion. Moreover, recent work by Schleicher et al. (2026) suggests that in the most compact and massive young clusters detected by JWST ($M \gtrsim 6 \times 10^6 M_\odot$), supernova feedback may be insufficient to expel the gas, allowing it to be retained and potentially enhance both DF and Bondi accretion onto the central object. While our simulations do not include these dissipative processes, the findings of Schleicher et al. (2026) suggest that the gas-driven enhancement of runaway collisions we observe may be even stronger in more realistic scenarios. In addition, gas accretion could contribute to enhancing BH formation, since it increases the stellar density (Bonnell & Bate 2002; Reinoso et al. 2018). Consequently, the fraction of collisions can increase to 10%, compared with gas-free models. On the other hand, stellar winds and supernovae (SNe) contribute to the chemical enrichment of the remnant gas after star formation episodes. Although the gas is heated, it does not necessarily evaporate from the cluster. As shown by Silich & Tenorio-Tagle (2016), depending on the gas density, the remnant gas can be expelled from the cluster or remain, in which case it can contribute to the runaway collision process. Moreover, in the transonic regime where the stellar velocity dispersion is comparable to the gas sound speed, DF is particularly efficient (Ostriker 1999), contributing to mass segregation. To assess the realism of our simplifications, it is important to consider the timescales of the relevant processes. Runaway collisions begin near core collapse (t_{cc}), and a massive object is formed at approximately the global collisional timescale t_{coll} . Any process that evaporates the gas on a timescale $t \lesssim t_{\text{cc}}$ has an important impact on the runaway collision process, while for timescales $t \gtrsim t_{\text{coll}}$, the runaway is not significantly perturbed. In the intermediate regime, $t_{\text{cc}} \leq t \leq t_{\text{coll}}$, the influence may be relevant.

Some studies have investigated the timescales for gas expulsion in embedded clusters. Dinbier & Walch (2020) studied gas expulsion from embedded star clusters, where the stellar mass is one third of the total mass (i.e., a gas fraction of $q = 2$). Their results show that for clusters with total mass $M_{\text{cluster}} \lesssim 3 \times 10^3 M_\odot$, the gas expulsion timescale is $t_{\text{ge}} \gtrsim 0.9$ Myr primarily through ionizing radiation and stellar winds. This mass range corresponds to models D, F, J, and K, for which t_{coll} (stars-only) is 1.018, 10.829, 27.032, and 61.53 Myr, respectively. We note that model D, which is in the IR, would evolve with a gas component in a more realistic scenario, at least during the runaway collision process. For more massive clusters, only 10% to 30% of the total mass could become unbound, and the majority could be ionized on a timescale of ~ 1 Myr. In contrast, Calura et al. (2015) ran simulations of globular clusters with a total mass of $\sim 10^7 M_\odot$. Their hydrodynamical simulations included stellar feedback (winds and supernovae), radiative cooling and self-gravity. They found that after 3 Myr, stellar feedback removes 40% of the initial gas, and after 14 Myr, 99% is removed. Similarly, Pelupessy & Portegies Zwart (2012) showed that for relatively small embedded clusters, the gas could be expelled by different mechanisms. While high feedback coupling could remove gas on a timescale of ~ 5 Myr, low feedback coupling is not sufficient to expel the gas via stellar winds, and supernovae become the dominant mechanism (their first supernova events occur at ~ 10 Myr). More recently, Chung et al. (2026) showed that in hydrodynamical simulations of dense star cluster formation that include cooling, heating, and stellar feedback, the

gas can be retained in sufficiently compact and massive clusters, affecting the IMBH formation process via runaway collisions. Taken together, these studies suggest that gas can persist during the core collapse phase in many realistic scenarios, particularly in low-mass clusters or with low stellar feedback. However, the survival of the gas depends on the details of gravity and stellar feedback, which we have not modeled. Therefore, although it seems plausible that gas could persist during an important part of the runaway collisions and could effectively enhance the formation of an IMBH, more realistic simulations that include stellar feedback, supernovae, and radiative cooling are necessary.

Another simplification concerns the initial conditions. Stars are initialized using a Plummer distribution, with equal-mass stars in virial equilibrium. The gas is initialized using a Plummer distribution with the same spatial distribution as the stars. Although the Plummer distribution is consistent with observations (Plummer 1911) and simulations (Chung et al. 2026), it is necessary to include an IMF for the stellar mass distribution. On the other hand, cosmological simulations could provide more realistic initial conditions, including stellar and gas distributions and the remnant gas fraction. Our most massive model (L family models) has a total stellar mass of $5 \times 10^5 M_\odot$ and is therefore not able to form an SMBH. Modeling more massive systems would require a code more efficient than PH4 in AMUSE, and capable of detecting stellar collisions. Furthermore, as shown in Appendix D.3, DF is not fully resolved in our simulations, and the number of SPH particles within the gravitational radius is negligible. We therefore caution that gas DF is not fully captured. To achieve realistic DF, it would be necessary to increase the resolution of the SPH particles or to include a sub-grid model for unresolved gas DF, which could enhance the formation of an SMBH. As mentioned in Pelupessy & Portegies Zwart (2012) and Whitehead et al. (2013), small differences in initial conditions are amplified due to the chaotic nature of N-body systems. Therefore, future work should perform multiple runs with different random realizations for each set of initial conditions. Although the general results will remain, this would allow us to obtain statistics for the trends shown in Section 4.

6. Conclusions

In this work, we have studied the role of gas in the collisional runaway scenario for SMBH seed formation in nuclear stellar clusters. By performing a series of simulations with coupled N-body and SPH codes via the BRIDGE scheme, we have shown that the presence of a gaseous component, even at small fractions (5%–20% of the stellar mass), significantly affects the formation of a massive central object.

Our main findings are:

1. The gas shortens core collapse, particularly for gas fractions of 10% and 20%, with the strongest effect at 20% (reduction of $27.39\% \pm 8.99\%$ relative to gas-free simulations). The effect is less robust at 5% gas fraction ($6.23\% \pm 18.36\%$), suggesting that a minimum gas fraction is required for a significant effect.
2. The presence of gas enhances black hole formation. In our simulations, the BH mass increases by $8.40\% \pm 15.89\%$, $22.51\% \pm 20.57\%$, and $44.14\% \pm 21.32\%$ for gas fractions of 5%, 10%, and 20%, respectively, and the BH formation efficiency shows a similar trend. The enhancement is most pronounced in the models with the highest collision rates within the IR (A, B, C, H), where the density is highest and the gas can most effectively contribute to the formation of an IMBH.

3. The fitted model for BH formation efficiency shows that the effect of gas is most important in the intermediate range $10^{-4} \leq M/M_{\text{crit}} \leq 10^5$ where the efficiency increases with gas fraction. Since gas dynamical friction is not fully resolved in our simulations, this enhancement may be a conservative estimate, and higher resolution simulations could reveal an even stronger effect.

While our simulations demonstrate the potential of gas to enhance SMBH seed formation, several simplifications remain to be addressed. Our gas model is adiabatic and does not include radiative cooling, heating, stellar feedback, or supernovae. Our findings address the question of how gas affects the global instability that leads to runaway collisions and, eventually, to the formation of a SMBH. The next natural question is whether gas can remain during relevant timescales in stellar clusters. Future work should include these processes to determine whether the gas persists long enough during the core collapse phase to produce the effects observed here. Additionally, more realistic initial conditions (e.g., IMFs, cosmological initial conditions) and larger N-body systems (using more efficient codes, such as NBODY6++GPU) would allow us to confirm the robustness of our results and extend them to more massive clusters. The IMBHs formed in our simulations could serve as seeds that subsequently grow to SMBH masses via gas accretion and mergers, as suggested by recent studies (Inayoshi et al. 2022; Kritos et al. 2023; Chung et al. 2026).

We conclude that gas is a key ingredient in the collisional runaway scenario for SMBH seed formation, and its inclusion in future simulations is essential for a complete understanding of the formation of massive black holes in nuclear stellar clusters.

Acknowledgements. We gratefully acknowledge the Federal Ministry of Education and Research and the state governments of Chile for supporting this project as part of the funding of the Kultrun cluster hosted at the Departamento de Astronomía, Universidad de Concepción. B. Cuevas, A. Escala, M. C. Vergara and D. R. G. Schleicher acknowledge funding through CATA (ANID BASAL project FB21003). B. Reinoso acknowledges funding through the European Research Council via ERC Consolidator grant KETJU (no. 818930). Also D. R. G. Schleicher acknowledges the Alexander von Humboldt - Foundation, Bonn, Germany.

References

- Aarseth, S. J. & Aarseth, S. J. 2003, *Gravitational N-body simulations: tools and algorithms* (Cambridge University Press)
- Barnes, J. & Hut, P. 1986, *Nature*, 324, 446
- Bellovary, J. 2025, *The Astrophysical Journal Letters*, 984, L55
- Binney, J. & Tremaine, S. 2008, *Galactic Dynamics: Second Edition*
- Boekholt, T., Schleicher, D., Fellhauer, M., et al. 2018, *Monthly Notices of the Royal Astronomical Society*, 476, 366
- Bond, J., Arnett, W., & Carr, B. J. 1984, *Astrophysical Journal*, Part 1 (ISSN 0004-637X), vol. 280, May 15, 1984, p. 825-847. Research supported by the Science and Engineering Research Council., 280, 825
- Bonnell, I. A. & Bate, M. R. 2002, *Monthly Notices of the Royal Astronomical Society*, 336, 659
- Bromm, V. & Loeb, A. 2003, *The Astrophysical Journal*, 596, 34
- Calura, F., Few, C., Romano, D., & D’Ercole, A. 2015, *The Astrophysical Journal Letters*, 814, L14
- Chon, S., Hosokawa, T., & Omukai, K. 2021, *Monthly Notices of the Royal Astronomical Society*, 502, 700
- Chung, E., Jo, Y., Kim, J.-h., Jung, M., & Kwon, O.-k. 2026, *The Astrophysical Journal*, 1004, 250
- Das, A., Schleicher, D. R. G., Leigh, N. W. C., & Boekholt, T. C. N. 2021, *Monthly Notices of the Royal Astronomical Society*, 503, 1051
- Demircan, O. & Kahraman, G. 1991, *Bf00639097*
- Di Cintio, P., Pasquato, M., Simon-Petit, A., & Yoon, S.-J. 2022, *Astronomy & Astrophysics*, 659, A19
- Dinnbier, F. & Walch, S. 2020, *Monthly Notices of the Royal Astronomical Society*, 499, 748
- Eilers, A.-C., Simcoe, R. A., Yue, M., et al. 2023, *The Astrophysical Journal*, 950, 68

- Endsley, R., Stark, D. P., Whitler, L., et al. 2023, *Monthly Notices of the Royal Astronomical Society*, 524, 2312
- Escala, A. 2006, *The Astrophysical Journal Letters*, 648, L13
- Escala, A. 2021, *ApJ*, 908, 57
- Escala, A., Zimmermann, L., Valdebenito, S., et al. 2025, *The Astrophysical Journal*, 995, 44
- 960 Ferrarese, L., Côté, P., Dalla Bontà, E., et al. 2006, *The Astrophysical Journal Letters*, 644, L21
- Fryer, C., Woosley, S., & Heger, A. 2001, *The Astrophysical Journal*, 550, 372
- Fujii, M., Iwasawa, M., Funato, Y., & Makino, J. 2007, *PASJ*, 59, 1095
- Furtak, L. J., Labbé, I., Zitrin, A., et al. 2024, *Nature*, 628, 57
- Gardner, J. P., Mather, J. C., Abbott, R., et al. 2023, *Publications of the Astronomical Society of the Pacific*, 135, 068001
- Georgiev, I. Y., Böker, T., Leigh, N., Lützgendorf, N., & Neumayer, N. 2016, *Monthly Notices of the Royal Astronomical Society*, 457, 2122
- Gerritsen, J. P. E. & Icke, V. 1997, *A&A*, 325, 972
- Gingold, R. A. & Monaghan, J. J. 1977, *MNRAS*, 181, 375
- 970 Gültekin, K., Richstone, D. O., Gebhardt, K., et al. 2009, *The Astrophysical Journal*, 698, 198
- Gürkan, M. A., Freitag, M., & Rasio, F. A. 2004, *The Astrophysical Journal*, 604, 632
- Harikane, Y. 2025, *Astrophysics and Space Science*, 370, 85
- Häring, N. & Rix, H.-W. 2004, *The Astrophysical Journal Letters*, 604, L89
- Higham, N. J., Dennis, M. R., Glendinning, P., et al. 2015, *Princeton companion to applied mathematics* (Princeton University Press)
- Horedt, G. P. 2004, *Polytropes: applications in astrophysics and related fields* (Springer)
- 980 Hut, P., McMillan, S., Goodman, J., et al. 1992, *Publications of the Astronomical Society of the Pacific*, 104, 981
- Inayoshi, K., Haiman, Z., & Ostriker, J. P. 2016, *Monthly Notices of the Royal Astronomical Society*, 459, 3738
- Inayoshi, K., Nakatani, R., Toyouchi, D., et al. 2022, *The Astrophysical Journal*, 927, 237
- Jeon, J., Bromm, V., Liu, B., & Finkelstein, S. L. 2025, *The Astrophysical Journal*, 979, 127
- Jiang, L., McGreer, I. D., Fan, X., et al. 2015, *The Astronomical Journal*, 149, 188
- 990 Khalisi, E., Amaro-Seoane, P., & Spurzem, R. 2007, *Monthly Notices of the Royal Astronomical Society*, 374, 703
- King, A. 2015, *Monthly Notices of the Royal Astronomical Society: Letters*, 456, L109
- Kiyuna, M. 2026, *Monthly Notices of the Royal Astronomical Society*, 545, staf1935
- Kocevski, D. D., Onoue, M., Inayoshi, K., et al. 2023, *The Astrophysical Journal Letters*, 954, L4
- Kritos, K., Berti, E., & Silk, J. 2023, *Physical Review D*, 108, 083012
- Kritos, K. & Silk, J. 2026, *The Astrophysical Journal*, 1004, 226
- 1000 Kustaanheimo, P., Schinzel, A., Davenport, H., & Stiefel, E. 1965
- Labbé, I., van Dokkum, P., Nelson, E., et al. 2023, *Nature*, 616, 266
- Latif, M. & Schleicher, D. R. G. 2015, *Astronomy & Astrophysics*, 578, A118
- Latif, M. A. & Ferrara, A. 2016, *Publications of the Astronomical Society of Australia*, 33, e051
- Leigh, N. W., Mastrobuono-Battisti, A., Perets, H. B., & Böker, T. 2014, *Monthly Notices of the Royal Astronomical Society*, 441, 919
- Lodato, G. & Natarajan, P. 2006, *Monthly Notices of the Royal Astronomical Society*, 371, 1813
- Makino, J. & Aarseth, S. J. 1992, *PASJ*, 44, 141
- 1010 Marconi, A. & Hunt, L. K. 2003, *The Astrophysical Journal Letters*, 589, L21
- McMillan, S., Portegies Zwart, S., van Elteren, A., & Whitehead, A. 2012, in *Astronomical Society of the Pacific Conference Series*, Vol. 453, *Advances in Computational Astrophysics: Methods, Tools, and Outcome*, ed. R. Capuzzo-Dolcetta, M. Limongi, & A. Tornambè, 129
- Mikkola, S. & Aarseth, S. J. 1993, *Celestial Mechanics and Dynamical Astronomy*, 57, 439
- Monaghan, J. J. 1992, *ARA&A*, 30, 543
- Mortlock, D. J., Warren, S. J., Venemans, B. P., et al. 2011, *Nature*, 474, 616
- Naiman, J., Ramirez-Ruiz, E., Debuhr, J., & Ma, C.-P. 2015, *The Astrophysical Journal*, 803, 81
- 1020 Natarajan, P., Pacucci, F., Ricarte, A., et al. 2024, *The Astrophysical Journal Letters*, 960, L1
- Natarajan, P. & Treister, E. 2009, *Monthly Notices of the Royal Astronomical Society*, 393, 838
- Neumayer, N., Seth, A., & Böker, T. 2020, *The Astronomy and Astrophysics Review*, 28, 4
- Ostriker, E. C. 1999, *The Astrophysical Journal*, 513, 252
- Pelupessy, F. & Portegies Zwart, S. 2012, *Monthly Notices of the Royal Astronomical Society*, 420, 1503
- 1030 Pelupessy, F. I., Di Matteo, T., & Ciardi, B. 2007, *The Astrophysical Journal*, 665, 107
- Pelupessy, F. I., van Elteren, A., de Vries, N., et al. 2013, *A&A*, 557, A84
- Plummer, H. C. 1911, *Monthly Notices of the Royal Astronomical Society*, Vol. 71, p. 460-470, 71, 460
- Portegies Zwart, S. & McMillan, S. 2018, *Astrophysical Recipes; The art of AMUSE*
- Portegies Zwart, S., McMillan, S., Harfst, S., et al. 2009, *New A*, 14, 369
- Portegies Zwart, S., McMillan, S. L. W., van Elteren, E., Pelupessy, I., & de Vries, N. 2013, *Computer Physics Communications*, 184, 456
- Portegies Zwart, S. F., Makino, J., McMillan, S. L., & Hut, P. 1999, *Astronomy and Astrophysics*, v. 348, p. 117-126 (1999), 348, 117
- Portegies Zwart, S. F. & McMillan, S. L. 2002, *The Astrophysical Journal*, 576, 899
- Quinlan, G. D. 1996, arXiv preprint astro-ph/9606182
- Rantala, A. 2026, arXiv preprint arXiv:2604.22924
- Rasio, F. A., Freitag, M., & Gürkan, M. A. 2003, arXiv preprint astro-ph/0304038
- Regan, J. & Volonteri, M. 2024, arXiv preprint arXiv:2405.17975
- Reinoso, B., Schleicher, D., Fellhauer, M., Klessen, R., & Boekholt, T. 2018, *Astronomy & Astrophysics*, 614, A14
- 1050 Reinoso, B., Schleicher, D. R., Fellhauer, M., Leigh, N. W., & Klessen, R. S. 2020, *Astronomy & Astrophysics*, 639, A92
- Sakurai, Y., Yoshida, N., Fujii, M. S., & Hirano, S. 2017, *Monthly Notices of the Royal Astronomical Society*, 472, 1677
- Schiavi, R., Capuzzo-Dolcetta, R., Georgiev, I. Y., Arca-Sedda, M., & Mastrobuono-Battisti, A. 2021, *Monthly Notices of the Royal Astronomical Society*, 503, 594
- Schleicher, D. R., Liempi, M., Giersz, M., et al. 2026, *Astronomy & Astrophysics*, 710, A235
- Schneider, R., Valiante, R., Trinca, A., et al. 2023, *Monthly Notices of the Royal Astronomical Society*, 526, 3250
- 1060 Secunda, A., Somerville, R. S., Jiang, Y.-F., et al. 2026, *The Astrophysical Journal*, 996, 6
- Shang, C., Bryan, G. L., & Haiman, Z. 2010, *Monthly Notices of the Royal Astronomical Society*, 402, 1249
- Shi, Y., Kremer, K., & Hopkins, P. F. 2024, *The Astrophysical Journal Letters*, 969, L31
- Shlosman, I., Begelman, M. C., & Frank, J. 1990, *Nature*, 345, 679
- Silich, S. & Tenorio-Tagle, G. 2016, *Monthly Notices of the Royal Astronomical Society*, stw2879
- 1070 Silich, S. & Tenorio-Tagle, G. 2018, *Monthly Notices of the Royal Astronomical Society*, 478, 5112
- Silk, J., Begelman, M. C., Norman, C., Nusser, A., & Wyse, R. F. 2024, *The Astrophysical Journal Letters*, 961, L39
- Smith, B. D., Regan, J. A., Downes, T. P., et al. 2018, *Monthly Notices of the Royal Astronomical Society*, 480, 3762
- Spitzer, L. 1987, *Dynamical evolution of globular clusters*
- Springel, V. & Hernquist, L. 2002, *Monthly Notices of the Royal Astronomical Society*, 333, 649
- 1080 Suazo, M., Prieto, J., Escala, A., & Schleicher, D. R. 2019, *The Astrophysical Journal*, 885, 127
- Tremaine, S., Gebhardt, K., Bender, R., et al. 2002, *The Astrophysical Journal*, 574, 740
- Trenti, M. & van der Marel, R. 2013, *Monthly Notices of the Royal Astronomical Society*, 435, 3272
- Vaida, D. D. & Farber, R. J. 2026, *Frontiers in Astronomy and Space Sciences*, 13, 1779045
- van Dokkum, P., Brammer, G., Jennings, C., Pasha, I., & Baggen, J. F. 2025, *The Astrophysical Journal Letters*, 990, L48
- Vergara, M., Askar, A., Flammini Dotti, F., et al. 2026, *Astronomy & Astrophysics*, 707, A71
- 1090 Vergara, M., Schleicher, D., Boekholt, T., et al. 2021, *Astronomy & Astrophysics*, 649, A160
- Vergara, M. C., Escala, A., Schleicher, D. R. G., & Reinoso, B. 2023, *MNRAS*, 522, 4224
- Vesperini, E., McMillan, S. L., D'Ercole, A., & D'Antona, F. 2010, *The Astrophysical Journal Letters*, 713, L41
- Volonteri, M. 2010, *The Astronomy and Astrophysics Review*, 18, 279
- Wang, L., Spurzem, R., Aarseth, S., et al. 2015, *Monthly Notices of the Royal Astronomical Society*, 450, 4070
- 1100 Weisstein, E. W. 2002, <https://mathworld.wolfram.com/>
- Whitehead, A. J., McMillan, S. L., Vesperini, E., & Portegies Zwart, S. 2013, *The Astrophysical Journal*, 778, 118
- Willott, C. J., Delorme, P., Omont, A., et al. 2007, *The Astronomical Journal*, 134, 2435
- Woods, T. E., Agarwal, B., Bromm, V., et al. 2019, *Publications of the Astronomical Society of Australia*, 36, e027
- Yang, D., Hennawi, J., Guarneri, F., et al. 2026, *Astronomy & Astrophysics*, 711, A104
- Zwillinger, D. 2002, *CRC standard mathematical tables and formulae* (chapman and hall/CRC)

Appendix A: Critical mass

In this section, we present a solution of the implicit equations for the critical mass presented in Vergara et al. (2023), where virialized system is assumed. Also, we consider an external potential (i.e., gas). We begin with the derivation of the collisional timescale. For a system with N identical stars of mass m and radius r_* , velocity dispersion σ , total mass M and characteristic radius R , the collision rate is given by (Binney & Tremaine 2008):

$$\frac{1}{t_{\text{coll}}} = 16 \sqrt{\pi n \sigma r_*^2 (1 + \Theta)}, \quad (\text{A.1})$$

where Θ is the Safronov number:

$$\Theta \equiv \frac{Gm}{2\sigma^2 r_*} = 9.54 \frac{m}{M_\odot} \frac{R_\odot}{r_*} \left(\frac{100 \text{ km s}^{-1}}{\sigma} \right)^2, \quad (\text{A.2})$$

and $n = 3M/(4\pi R^3 m)$ is the number density. Defining the cross-section $\Sigma_0 = 16 \sqrt{\pi r_*^2 (1 + \Theta)}$, the collision time can be rewritten as:

$$t_{\text{coll}} = \frac{1}{n \sigma \Sigma_0}. \quad (\text{A.3})$$

We consider a stellar system in virial equilibrium with characteristic radius R . In the absence of an external potential, the velocity dispersion satisfies $\sigma = \sqrt{GM/R}$. When the system is embedded in a gravitationally bound external potential of mass M_{ext} with the same distribution (same characteristic radius R), the modified dispersion velocity yields (Reinoso et al. 2020):

$$\sigma_{\text{ext}} = \sqrt{\frac{GM}{R}} (1 + q), \quad (\text{A.4})$$

where $q = M_{\text{ext}}/M$ and $M + M_{\text{ext}} = M(1 + q)$ is the total mass. First, we replace the dispersion velocity (Eq. (A.4)) into the cross-section Σ_0 :

$$\begin{aligned} \Sigma_0 &= 16 \sqrt{\pi r_*^2 (1 + \Theta)} \\ &= 16 \sqrt{\pi r_*^2 \left(1 + 9.54 \frac{m}{M_\odot} \frac{R_\odot}{r_*} \left(\frac{v_{100}}{\sigma_{\text{ext}}} \right)^2 \right)} \\ &= 16 \sqrt{\pi r_*^2} + \frac{9.54 \times 16 \sqrt{\pi r_* m R_\odot v_{100}^2}}{M_\odot} \frac{R}{GM(1 + q)^2}. \end{aligned} \quad (\text{A.5})$$

From the relation $t_{\text{coll}} = t_H$, we place all mass-dependent terms on left side:

$$\begin{aligned} t_H = t_{\text{coll}} &= \frac{1}{n \sigma_{\text{ext}} \Sigma_0} \\ &= \frac{4\pi R^3 m}{3M} \frac{\sqrt{R}}{\sqrt{GM}(1 + q) \Sigma_0} \\ \Sigma_0 M^{3/2} &= \frac{4\pi R^3 m}{3t_H} \frac{\sqrt{R}}{\sqrt{G}(1 + q)}. \end{aligned} \quad (\text{A.6})$$

Now, we replace Eq. (A.5) into Eq. (A.6):

$$\begin{aligned} 16 \sqrt{\pi r_*^2} M^{3/2} + \frac{9.54 \times 16 \sqrt{\pi r_* m R_\odot v_{100}^2} R}{M_\odot G(1 + q)^2} M^{1/2} &= \frac{4\pi m R^{7/2}}{3G^{1/2}(1 + q)t_H} \\ M^{3/2} + \frac{9.54 m R_\odot v_{100}^2 R}{r_* G M_\odot (1 + q)^2} M^{1/2} &= \frac{\sqrt{\pi m} R^{7/2}}{12G^{1/2} r_*^2 (1 + q)t_H}. \end{aligned} \quad (\text{A.7})$$

In order to simplify, we define the variables:

$$\alpha R = A = \frac{9.54 m R_\odot v_{100}^2}{r_* G M_\odot (1 + q)^2} R \quad (\text{A.8})$$

$$\beta R^{7/2} = C = \frac{\sqrt{\pi m}}{12G^{1/2} r_*^2 (1 + q)t_H} R^{7/2}. \quad (\text{A.9})$$

Then, the equation for the mass is

$$M^{3/2} + A M^{1/2} - C = 0. \quad (\text{A.10})$$

A.1. solving for M

The critical mass can be obtained solving Eq. (A.10). To simplify, we use the change of variable $u^2 = M$ to have a cubic polynomial. Therefore, Eq. (A.10) can be rewritten as:

$$u^3 + Au - C = 0. \quad (\text{A.11})$$

This equation could have three roots, but we are only interested in real solutions. Given $p = A/3$ and $q = -C$, the discriminant of Eq. (A.11) is:

$$\Delta = 4p^3 + q^2 = \frac{4A^3}{27} + C^2. \quad (\text{A.12})$$

As $A, C > 0$ is straight that the discriminant is positive, then there is only one real solution (Zwillinger 2002):

$$u_{\text{real}} = \sqrt[3]{s_1} + \sqrt[3]{s_2} \quad (\text{A.13})$$

$$s_1 = \frac{-q + \sqrt{q^2 + 4p^3}}{2} \quad (\text{A.14})$$

$$s_2 = \frac{-q - \sqrt{q^2 + 4p^3}}{2}. \quad (\text{A.15})$$

Replacing p and q by $A/3$ and $-C$:

$$\begin{aligned} u_{\text{real}} &= \sqrt[3]{\frac{-q + \sqrt{q^2 + 4p^3}}{2}} + \sqrt[3]{\frac{-q - \sqrt{q^2 + 4p^3}}{2}} \\ &= \sqrt[3]{\frac{C}{2} + \sqrt{\frac{C^2}{4} + \frac{A^3}{27}}} + \sqrt[3]{\frac{C}{2} - \sqrt{\frac{C^2}{4} + \frac{A^3}{27}}}. \end{aligned} \quad (\text{A.16})$$

Then, the critical mass is:

$$\begin{aligned} M_{\text{crit}} &= u_{\text{real}}^2 \\ &= \left(\frac{C}{2} + \sqrt{\frac{C^2}{4} + \frac{A^3}{27}} \right)^{2/3} + \left(\frac{C}{2} - \sqrt{\frac{C^2}{4} + \frac{A^3}{27}} \right)^{2/3} + \\ &\quad 2 \sqrt[3]{\frac{C^2}{4} - \left(\frac{C^2}{4} + \frac{A^3}{27} \right)} \\ &= \left(\frac{C}{2} + \sqrt{\frac{C^2}{4} + \frac{A^3}{27}} \right)^{2/3} + \left(\frac{C}{2} - \sqrt{\frac{C^2}{4} + \frac{A^3}{27}} \right)^{2/3} - \\ &\quad \frac{2A}{3}, \end{aligned} \quad (\text{A.17})$$

or in terms of α and β :

$$\begin{aligned} M_{\text{crit}} &= \left(\frac{\beta R^{7/2}}{2} + \sqrt{\frac{\beta^2 R^7}{4} + \frac{\alpha^3 R^3}{27}} \right)^{2/3} + \\ &\quad \left(\frac{\beta R^{7/2}}{2} - \sqrt{\frac{\beta^2 R^7}{4} + \frac{\alpha^3 R^3}{27}} \right)^{2/3} - \frac{2\alpha R}{3}. \end{aligned} \quad (\text{A.18})$$

A.2. Analytical vs numerical solution

To test the analytical solution for the critical mass, we compare it with a numerical solution obtained by solving the implicit condition iteratively. Fig. A.1 shows the relative error between the numerical and analytical solutions as a function of the characteristic radius R . The relative difference between the numerical and




Fig. A.1: Relative error between the analytical solution (M_a) and the numerical solution (M_n) for the critical mass

analytical solutions is at most $\sim 1\%$ for small radii, decreasing to $\sim 0.1\%$ for larger radii. For all practical purposes, the analytical expression (Eq. (A.18)) is accurate and numerically stable for $M_{\text{crit}} \gtrsim 10 M_{\odot}$.

Appendix B: Relaxation mass

In this section we derive a characteristic mass from the two-body relaxation timescale, in the presence of an external potential. Consider a system of equal N stars of mass m , with characteristic radius R and velocity dispersion σ . A star crossing the system suffers a mean-square velocity deflection given by (Binney & Tremaine 2008):

$$\Delta v^2 \simeq 8N \left(\frac{Gm}{Rv} \right)^2 \ln \left(\frac{b_{\text{max}}}{b_{\text{min}}} \right), \quad (\text{B.1})$$

where b_{max} , b_{min} are the maximum and minimum impact parameters. Usually, $b_{\text{max}} \simeq R$, and $b_{\text{min}} = 2Gm/v^2$ is the impact parameter for a 90° deflection. The number of crossings needed for the star to change its velocity significantly is therefore

$$n_{\text{relax}} = \frac{v^2}{\Delta v^2} = \left(\frac{Rv^2}{Gm} \right)^2 \frac{1}{8N \ln(b_{\text{max}}/b_{\text{min}})}. \quad (\text{B.2})$$

As shown in Appendix A, the velocity dispersion in the presence of an external potential of mass $M_{\text{ext}} = qM$ (with the same spatial distribution) becomes $\sigma_{\text{ext}} = \sqrt{GNm/R}(1+q)$, where $M = Nm$ is the stellar mass. Setting $v = \sigma_{\text{ext}}$ we obtain $Rv^2/(Gm) = RGNm(1+q)^2/(RGm) = N(1+q)^2$ and

$$\begin{aligned} \ln \left(\frac{b_{\text{max}}}{b_{\text{min}}} \right) &= \ln \left(\frac{R}{2Gm/v^2} \right) = \ln \left(\frac{Rv^2}{2Gm} \right) \\ &= \ln \left(\frac{N(1+q)^2}{2} \right) = \ln(N(1+q)^2) - \ln(2) \\ &\simeq \ln(N(1+q)^2), \end{aligned} \quad (\text{B.3})$$

where we have dropped the constant $\ln(2)$ for simplicity. Substituting into the expression for n_{relax} yields

$$n_{\text{relax}} = \frac{N(1+q)^4}{8 \ln(N(1+q)^2)} \simeq \frac{0.1N(1+q)^4}{\ln(N(1+q)^2)}. \quad (\text{B.4})$$

Since n_{relax} is the number of crossing times required for relaxation, the relaxation time is $t_{\text{relax}} = n_{\text{relax}} t_{\text{cross}}$, where $t_{\text{cross}} = 1180 R/\sigma_{\text{ext}}$. Thus,

$$t_{\text{relax}} = n_{\text{relax}} t_{\text{cross}} = \frac{0.1N(1+q)^4}{\ln(N(1+q)^2)} t_{\text{cross}}. \quad (\text{B.5})$$

We now impose that the characteristic time t_H is shorter than the relaxation time ($t_H \leq t_{\text{relax}}$), which is the condition for the system to retain its initial properties (two-body relaxation is not yet effective). Similar to Vergara et al. (2023), we obtain a lower bound on the cluster radius:

$$\begin{aligned} t_H &\leq \frac{0.1N(1+q)^4}{\ln(N(1+q)^2)} t_{\text{cross}} = \frac{0.1N(1+q)^4}{\ln(N(1+q)^2)} \frac{R}{\sigma_{\text{ext}}} \\ &= \frac{0.1M(1+q)^4}{m \ln(M(1+q)^2/m)} \frac{R^{3/2}}{\sqrt{GM}(1+q)} \\ &= \frac{0.1M^{1/2}(1+q)^3}{\ln(M(1+q)^2/m)} \frac{R^{3/2}}{G^{1/2}m} \\ R &\geq \left(\frac{t_H m}{0.1(1+q)^3} \ln \left(\frac{M(1+q)^2}{m} \right) \right)^{2/3} \left(\frac{G}{M} \right)^{1/3}. \end{aligned} \quad (\text{B.6})$$

B.1. Solving for the relaxation mass

Inequality (B.6) can be read as a relation $R(M)$. The relaxation mass is then defined as its inverse, $M_{\text{relax}}(R) = R^{-1}(M)$. Setting the equality gives

$$\begin{aligned} R &= \left(\frac{t_H m \ln(M(1+q)^2/m)}{0.1(1+q)^3} \right)^{2/3} \left(\frac{G}{M} \right)^{1/3} \\ R^3 &= \left(\frac{t_H m \ln[M(1+q)^2/m]}{0.1(1+q)^3} \right)^2 \left(\frac{G}{M} \right) \\ M^{1/2} R^{3/2} &= \frac{mt_H \ln[M(1+q)^2/m]}{0.1(1+q)^3} G^{1/2}. \end{aligned} \quad (\text{B.7})$$

Considering the change of variable $u = M(1+q)^2/m$, Eq. (B.7) becomes

$$\begin{aligned} \frac{(um)^{1/2} R^{3/2}}{1+q} &= \frac{mt_H \ln(u)}{0.1(1+q)^3} G^{1/2} \\ u^{1/2} &= \ln(u) \frac{t_H (Gm)^{1/2}}{0.1R^{3/2}(1+q)^2}. \end{aligned} \quad (\text{B.8})$$

Defining $\lambda = t_H (Gm)^{1/2} / (0.1R^{3/2}(1+q)^2)$, Eq. (B.8) yields:

$$u^{1/2} = \lambda \ln(u), \quad (\text{B.9})$$

and introducing the change of variable $\omega = \ln(u)$

$$\begin{aligned} e^{\omega/2} &= \lambda \omega \\ -1/(2\lambda) &= (-\omega/2) e^{-\omega/2}. \end{aligned} \quad (\text{B.10})$$

The inverse of the function $f(w) = we^w$ is the Lambert function W (i.e. $w = W(y)$ if $y = we^w$) (Weisstein 2002). Hence

$$\begin{aligned} \frac{-\omega}{2} &= W \left(\frac{-1}{2\lambda} \right) \\ \omega &= -2W \left(\frac{-1}{2\lambda} \right). \end{aligned} \quad (\text{B.11})$$

Returning to $u = e^\omega$ we obtain

$$u = \exp \left[-2W \left(\frac{-1}{2\lambda} \right) \right], \quad (\text{B.12})$$

and finally the relaxation mass:

$$M_{\text{relax}} = \frac{m}{(1+q)^2} \exp \left[-2W \left(\frac{-1}{2\lambda} \right) \right]. \quad (\text{B.13})$$

Eq. (B.13) involves the Lambert function W evaluated at a negative argument $z = -1/(2\lambda)$ (since $\lambda > 0$). There are two real-valued branches, $W_0(z)$ and $W_{-1}(z)$ (Higham et al. 2015, Chapter III.17). The principal branch, $W_0(z)$, is defined for $z \in [-1/e, \infty)$. The lower branch $W_{-1}(z)$ is defined on $[-1/e, 0)$. Since $-1/(2\lambda) < 0$ and assuming $-1/e \leq -1/(2\lambda)$, both branches can be taken. From Fig. B.1, we see that for $z \in [-1/e, 0)$, the principal branch takes values in $[-1, 0]$, while $W_{-1}(z)$ takes values in $[-1, -\infty)$. Evaluating in Eq. (B.13), the




Fig. B.1: Lambert function. The black line shows the $W_0(z)$ branch, and the blue line shows the $W_{-1}(z)$ branch.

branch W_0 gives $me^2/(1+q) \geq M_{\text{relax}} > 0$, while W_{-1} gives $me^2/(1+q) \leq M_{\text{relax}} < \infty$. As we expect for the relaxation mass to be $M_{\text{relax}} \gg m$, the physically relevant solution is the lower branch. Then, the relaxation mass can be written as:

$$M_{\text{relax}} = \frac{m}{(1+q)^2} \exp \left[-2W_{-1} \left(-\frac{1}{2\lambda} \right) \right]. \quad (\text{B.14})$$

A real solution exists only when $z \geq -1/e$, i.e. $\lambda \geq e/2$. For typical star clusters ($N \gg 1$, $q \lesssim 1$, and t_H large), this condition is amply satisfied, so the expression is well-defined and yields $M_{\text{relax}} \gg m$.

Appendix C: Core Collapse Detection Method

In this section, we describe the implemented method for detecting the core collapse, using Lagrangian radii. The core collapse corresponds to the moment in which the inner Lagrangian radii reach their minimum (Di Cintio et al. 2022; Reinoso et al. 2020), followed by an expansion of the outer regions. This process is expected to occur on a timescale of $\sim 20t_{\text{relax}}$ (Quinlan 1996), and can be observed from the evolution of Lagrangian radii. However, as the most massive BH reaches a mass comparable to that enclosed within a given Lagrangian radius, the evolution of that radius can be dominated by the motion of the BH rather than by the contraction of the stellar core. To avoid this, we use the smallest Lagrangian radius that is not dominated by the most massive BH. The algorithm proceeds as follows:

1. For a given Lagrangian radius $r_i(t)$, we select data within $t \leq 40t_{\text{relax}}$.
2. We apply a Savitzky–Golay filter (implemented via the `savgol_filter` function from `scipy.signal`) with a quadratic polynomial to smooth the data using a smoothing window of size sw .
3. We define a sliding window of size $W = fN$, where f is a parameter in the range $[0, 1]$, and N is the number of data points.
4. For each window position, we perform two linear regressions: one on the first half of the data points in the window and another on the second half. We assign to the central time of the window the difference in slopes, computed as $\Delta m = m_{\text{right}} - m_{\text{left}}$.
5. After running over all the selected data, the core collapse time is identified as the time at which Δm reaches its maximum value. This corresponds to the point of maximum curvature change in the Lagrangian radius.

The window size is varied between 10% and 30% of the selected data. To preserve sufficient resolution, the smoothing window was set empirically to $ws = W/5$. The method was validated by visual inspection of the Lagrangian radii for each simulation, confirming that the identified t_{cc} consistently matches the inflection point in the $r_i(t)$ curves.

Appendix D: Code Setup and Performance

D.1. Code Parameters and Initial Conditions

Table D.1 summarizes the key parameters for the N-body code PH4 and the SPH code FI, from the AMUSE framework. Parameters not listed were left at their default values. Table D.2 shows the initial conditions of each model.

D.2. Adaptive BRIDGE Timestep

The BRIDGE and FI codes use the same timestep, which is adjusted adaptively based on the local dynamical conditions of the gas and stars. Here, t_{nb} denotes the N-body time unit, and t_{relax} is the two-body relaxation time. The algorithm proceeds as follows:

1. The minimum free-fall time of the gas is computed as:

$$t_{\text{ff}} = \sqrt{\frac{3\pi}{32G\rho_{\text{max}}}}, \quad (\text{D.1})$$

where ρ_{max} is the maximum density among SPH particles.

Table D.1: Key runtime parameters for PH4 and FI.

Code	Parameter	Value	Description
PH4	‘epsilon_squared’	$(0.05 R_\odot)^2$	Gravitational softening length squared.
	‘force_sync’	‘True’	Synchronizes forces across all particles.
	‘timestep_parameter’	0.01	Controls the accuracy of the Hermite integrator.
FI	‘eps_is_h_flag’	‘True’	Sets gravitational softening equal to the SPH smoothing length.
	‘timestep’	$1.0 t_{\text{nb}}$	Initial timestep (in N-body time units).
	‘radiation_flag’	‘False’	Disables radiative cooling and heating.
	‘use_hydro_flag’	‘True’	Enables SPH hydrodynamics.
	‘integrate_entropy_flag’	‘False’	Evolve specific internal energy (not entropy).
	‘isothermal_flag’	‘False’	Disables isothermal equation of state.
	‘star_formation_flag’	‘False’	Disables star formation.

2. A Courant-type timestep is computed from the accelerations between gas and star particles:

$$\Delta t_{\text{coup}} = \min \left(f_{\text{coup}} \sqrt{\frac{2h}{\mathbf{a}_{\text{g,s}}}}, f_{\text{coup}} \sqrt{\frac{\epsilon_{\text{nb}}}{\mathbf{a}_{\text{s,g}}}} \right), \quad (\text{D.2})$$

where h is the SPH smoothing length, ϵ_{nb} is the coupling softening length, and $\mathbf{a}_{\text{g,s}}$ and $\mathbf{a}_{\text{s,g}}$ are the accelerations of gas and star particles due to the other component.

3. The reference timescale is then defined as:

$$\alpha = \min(t_{\text{ff}}, \Delta t_{\text{coup}}). \quad (\text{D.3})$$

4. The BRIDGE timestep is adjusted so that it satisfies:

$$\frac{\alpha}{2} \leq \Delta t_{\text{BRIDGE}} \leq \alpha. \quad (\text{D.4})$$

If $\Delta t_{\text{BRIDGE}} < \frac{\alpha}{2}$, the timestep is multiplied by 2.

To maintain synchronization with the N-body integrator, the BRIDGE timestep is rounded to the nearest power of two of the N-body time (t_{nb}):

$$\Delta t_{\text{BRIDGE}} = 2^k t_{\text{nb}}, \quad (\text{D.5})$$

with k chosen such that the timestep lies within the range $[\alpha/2, \alpha]$. During the first 25 relaxation times (t_{relax}), the maximum timestep is reduced to $2^{-4} t_{\text{nb}}$ to improve temporal resolution during the core collapse phase. After this period, the timestep may grow up to the initial value of $1.0 t_{\text{nb}}$. For all simulations, $f_{\text{coup}} = 0.5$ and $\epsilon_{\text{nb}} = 5$ AU.

D.3. Dynamical friction

To verify that dynamical friction is well resolved, we check two conditions: (i) the gravitational radius should be larger than the coupling softening length (ϵ_{nb} parameter); and (ii) the number of SPH particles within the gravitational radius for a typical star should be $\gtrsim 100$. The gravitational radius is given by

$$R_g = GM/(c_s^2 + v^2), \quad (\text{D.6})$$

where M is the mass of an individual star, c_s is the gas sound speed, and v is the velocity of the star. Among all models, the gravitational radius ranges between ~ 0.1 AU and ~ 10 AU, the first condition is only marginally satisfied for the most massive stars, and is not satisfied for lower-mass stars. Furthermore,

the number of SPH particles within R_g is effectively zero for all models. We therefore conclude that dynamical friction from the gas is not fully resolved in our simulations. However, this implies that the gas-driven enhancement of core collapse and BH formation observed in our simulations may be even stronger if dynamical friction were fully resolved. Future work with higher resolution would be required to quantify this effect accurately.

D.4. Energy Conservation and Virialization

The validation of our results relies on the accuracy of energy conservation. Fig. D.1 shows, in the top panel, the error in total energy per timestep as a function of crossing time. The bottom panel shows the virial ratio, defined as the ratio of two times the total kinetic energy (including thermal energy of gas) to the absolute value of the gravitational potential energy, $2K/|U|$. The initial conditions for this test consist of a stellar system with virial radius $r_{\text{vir}} = 1$ pc, composed of 5000 stars of $1 M_\odot$ each, embedded in a gas cloud with the same virial radius, a total gas mass of $5000 M_\odot$, and containing 5000 SPH particles. We observe that during the first 50 crossing times, the system remains virialized, with an energy error of $\lesssim 10^{-4}$.




Fig. D.1: Numerical accuracy test for a system of 5000 stars of $1 M_\odot$ embedded in a gas cloud with the same mass and 5000 SPH particles. Top panel: relative error in total energy per timestep as a function of crossing time. Bottom panel: virial ratio. The system remains virialized during the first 50 crossing times, with an energy error $\lesssim 10^{-4}$.

Appendix E: Extra plots




Fig. E.1: Lagrangian radii for gas of three variants of Model A. Top panel is for A_5 , middle is for A_{10} and bottom panel is for A_{20} . Left column shows the full simulation time (5 Myr) while right column shows the evolution until 40 relaxation times.




Fig. E.2: Lagrangian radii for gas of three variants of Model N. Top panel is for N_5 , middle is for N_{10} and bottom panel is for N_{20} . Left column shows the full simulation time (5 Myr) while right column shows the evolution until 40 relaxation times.




Fig. E.3: Sound speed of gas (c_s) and velocity dispersion (σ) for family A models.




Fig. E.4: Sound speed of gas (c_s) and velocity dispersion (σ) for family N models.

Table D.2: Initial conditions for stellar clusters. R_v and M are the virial radius and total mass of the stellar cluster, r and m are the initial radius and mass of stars, N is the number of stars, q_{gas} is the gas fraction (gas mass divided by stellar mass), and σ is the velocity dispersion. For models with gas, M_{SPH} is the total gas mass, N_{SPH} is the number of SPH particles, and m_{SPH} is the mass per SPH particle, with $m_{\text{SPH}} = m/100$. For $q_{\text{gas}} = 0$, the gas-related columns are left blank.

ID	$R_v[\text{pc}]$	$M[M_\odot]$	N	$m[M_\odot]$	$r[R_\odot]$	q_{gas}	$M_{\text{SPH}}[M_\odot]$	N_{SPH}	$m_{\text{SPH}}[M_\odot]$	$\sigma[\frac{\text{km}}{\text{s}}]$
A ₀	0.005	5×10^4	10^3	50	11.7	0				146.66
A ₅	0.005	5×10^4	10^3	50	11.7	0.05	2.5×10^3	5×10^3	0.5	146.66
A ₁₀	0.005	5×10^4	10^3	50	11.7	0.1	5×10^3	10^4	0.5	146.66
A ₂₀	0.005	5×10^4	10^3	50	11.7	0.2	10^4	2×10^4	0.5	146.66
B ₀	0.005	2.5×10^4	5×10^2	50	11.7	0				103.71
B ₅	0.005	2.5×10^4	5×10^2	50	11.7	0.05	1.25×10^3	2.5×10^3	0.5	103.71
B ₁₀	0.005	2.5×10^4	5×10^2	50	11.7	0.1	2.5×10^3	5×10^3	0.5	103.71
B ₂₀	0.005	2.5×10^4	5×10^2	50	11.7	0.2	5×10^3	10^4	0.5	103.71
C ₀	0.005	10^4	10^3	10	4.7	0				65.59
C ₅	0.005	10^4	10^3	10	4.7	0.05	5×10^2	5×10^3	0.1	65.59
C ₁₀	0.005	10^4	10^3	10	4.7	0.1	10^3	10^4	0.1	65.59
C ₂₀	0.005	10^4	10^3	10	4.7	0.2	2×10^3	2×10^4	0.1	65.59
D ₀	0.005	5×10^3	5×10^2	10	4.7	0				46.38
D ₅	0.005	5×10^3	5×10^2	10	4.7	0.05	2.5×10^2	2.5×10^3	0.1	46.38
D ₁₀	0.005	5×10^3	5×10^2	10	4.7	0.1	5×10^2	5×10^3	0.1	46.38
D ₂₀	0.005	5×10^3	5×10^2	10	4.7	0.2	10^3	10^4	0.1	46.38
F ₀	0.005	10^3	10^3	1	1.0	0				20.74
F ₅	0.005	10^3	10^3	1	1.0	0.05	5×10^1	5×10^3	0.01	20.74
F ₁₀	0.005	10^3	10^3	1	1.0	0.1	10^2	10^4	0.01	20.74
F ₂₀	0.005	10^3	10^3	1	1.0	0.2	2×10^2	2×10^4	0.01	20.74
H ₀	0.01	5×10^4	5×10^3	10	4.7	0				103.71
H ₅	0.01	5×10^4	5×10^3	10	4.7	0.05	2.5×10^3	2.5×10^4	0.1	103.71
H ₁₀	0.01	5×10^4	5×10^3	10	4.7	0.1	5×10^3	5×10^4	0.1	103.71
H ₂₀	0.01	5×10^4	5×10^3	10	4.7	0.2	10^4	10^5	0.1	103.71
I ₀	0.01	10^4	10^3	10	4.7	0				46.38
I ₅	0.01	10^4	10^3	10	4.7	0.05	5×10^2	5×10^3	0.1	46.38
I ₁₀	0.01	10^4	10^3	10	4.7	0.1	10^3	10^4	0.1	46.38
I ₂₀	0.01	10^4	10^3	10	4.7	0.2	2×10^3	2×10^4	0.1	46.38
J ₀	0.01	5×10^3	5×10^3	1	1.0	0				32.79
J ₅	0.01	5×10^3	5×10^3	1	1.0	0.05	2.5×10^2	2.5×10^4	0.01	32.79
J ₁₀	0.01	5×10^3	5×10^3	1	1.0	0.1	5×10^2	5×10^4	0.01	32.79
J ₂₀	0.01	5×10^3	5×10^3	1	1.0	0.2	10^3	10^5	0.01	32.79
K ₀	0.01	10^3	10^3	1	1.0	0				14.66
K ₅	0.01	10^3	10^3	1	1.0	0.05	5×10^1	5×10^3	0.01	14.66
K ₁₀	0.01	10^3	10^3	1	1.0	0.1	10^2	10^4	0.01	14.66
K ₂₀	0.01	10^3	10^3	1	1.0	0.2	2×10^2	2×10^4	0.01	14.66
L ₀	0.1	5×10^5	10^4	50	11.7	0				103.71
L ₅	0.1	5×10^5	10^4	50	11.7	0.05	2.5×10^4	5×10^4	0.5	103.71
L ₁₀	0.1	5×10^5	10^4	50	11.7	0.1	5×10^4	10^5	0.5	103.71
L ₂₀	0.1	5×10^5	10^4	50	11.7	0.2	10^5	2×10^5	0.5	103.71
M ₀	0.1	10^5	10^4	10	4.7	0				46.38
M ₅	0.1	10^5	10^4	10	4.7	0.05	5×10^3	5×10^4	0.1	46.38
M ₁₀	0.1	10^5	10^4	10	4.7	0.1	10^4	10^5	0.1	46.38
M ₂₀	0.1	10^5	10^4	10	4.7	0.2	2×10^4	2×10^5	0.1	46.38
N ₀	0.1	10^4	10^4	1	1.0	0				14.67
N ₅	0.1	10^4	10^4	1	1.0	0.05	5×10^2	5×10^4	0.01	14.67
N ₁₀	0.1	10^4	10^4	1	1.0	0.1	10^3	10^5	0.01	14.67
N ₂₀	0.1	10^4	10^4	1	1.0	0.2	2×10^3	2×10^5	0.01	14.67

Chapter 6

Conclusions

In this thesis, we have studied the role of gas in the runaway collision process that forms SMBH seeds, within the scenario proposed by Escala (2021). We performed 48 simulations based on the initial conditions of Vergara et al. (2023), with the inclusion of a gaseous component. Our simulations couple N-body and SPH codes via the BRIDGE scheme. Our main findings are summarized as follows (see Chapter 5):

1. The core collapse time is shortened by the presence of gas. For gas fractions of 5%, 10%, and 20% the core collapse time is reduced by 6.23%, 15.52%, and 27.39% on average, respectively.
2. The BH formation efficiency is enhanced by the presence of gas. In our simulations, we find that the average increments in BH mass is $8.40 \pm 15.89\%$, $22.51 \pm 20.57\%$, and $44.14 \pm 21.32\%$ for gas fractions of 5%, 10%, and 20%, respectively. The BH formation efficiency follows the same trend, with increases of $12.56 \pm 16.27\%$, $27.01 \pm 20.79\%$, and $45.10 \pm 31.39\%$ for the same gas fractions. The enhancement is more pronounced in the models with the highest collision rates within the IR (model families A, B, C, H).
3. The fitted model for BH formation efficiency shows that the effect of gas is most important in the intermediate range $10^{-4} \leq M/M_{\text{crit}} \leq 10^5$, where ϵ_{BH} increases with both the gas fraction and M/M_{crit} , and exhibits a smoother slope than the model of Vergara et al. (2023).
4. Since gas dynamical friction is not fully resolved in our simulations, the dissipative effects are not completely captured. It is highly likely that simulations with increased resolution will produce more massive BHs. In this regards, our results are conservative.

6.1. Caveats

While our simulations demonstrate the potential of gas to enhance SMBH seed formation, several simplifications remain to be addressed. The gas is simulated using an adiabatic equation of state, evolving the energy equation instead of the entropic function (Pelupessy, 2005). Radiative cooling, heating, and stellar feedback are not included. Supernovae are also not included; however, since our simulations run for only 5 Myr, the results remain valid within this timescale. Gas accretion is not included, and gas dynamical friction is not fully resolved; consequently, the effects observed may be enhanced in simulations with higher resolution. The N-body code used does not include regularization for close encounters, then binaries can slows the global evolution. In addition, due to numerical limitations, our star cluster models are not very massive. Consequently, our simulations are limited to studying the formation of IMBHs rather than SMBHs.

Our findings address the question of how gas affects the global instability that leads to runaway collisions and, eventually, to the formation of a SMBH. The next natural question is whether gas can remain during relevant timescales in stellar clusters. Future simulations should incorporate radiative cooling, heating, and stellar feedback (including SNe) to determine whether gas persists during core collapse in more realistic scenarios, thereby validating our findings. It is also necessary to use more realistic initial conditions (e.g., IMFs, cosmological initial conditions) instead of toy models, and more efficient N-body codes to simulate larger systems.

The IMBHs formed in our simulations could serve as seeds that subsequently grow to SMBH masses via gas accretion and mergers, as suggested by recent studies (Chung et al., 2026; Inayoshi et al., 2022; Kritos et al., 2023). We conclude that gas is a key ingredient in the collisional runaway scenario for SMBH seed formation. Its inclusion in future simulations is therefore essential for a complete understanding of the formation of SMBH in NSCs.

References

- Aarseth, S. J. (1985). Direct methods for n-body simulations. In Brackbill, J. U. and Cohen, B. I., editors, *Multiple Time Scales*, pages 377–418. Academic Press.
- Abel, T., Bryan, G. L., and Norman, M. L. (2000). The formation and fragmentation of primordial molecular clouds. *The Astrophysical Journal*, 540(1):39–44.
- Alvarez, M. A., Wise, J. H., and Abel, T. (2009). Accretion onto the first stellar-mass black holes. *The Astrophysical Journal*, 701(2):L133–L137.
- Balick, B. and Brown, R. L. (1974). Intense sub-arcsecond structure in the galactic center. *The Astrophysical Journal*, 194:265–270.
- Barnes, J. and Hut, P. (1986). A hierarchical o ($n \log n$) force-calculation algorithm. *nature*, 324(6096):446–449.
- Baumgarte, T. W. and Shapiro, S. L. (1999). Evolution of rotating supermassive stars to the onset of collapse. *The Astrophysical Journal*, 526(2):941–952.
- Belczynski, K., Bulik, T., Fryer, C. L., Ruiter, A., Valsecchi, F., Vink, J. S., and Hurley, J. R. (2010). On the maximum mass of stellar black holes. *The Astrophysical Journal*, 714(2):1217–1226.
- Bianchini, P., van de Ven, G., Norris, M., Schinnerer, E., and Varri, A. (2016). A novel look at energy equipartition in globular clusters. *Monthly Notices of the Royal Astronomical Society*, 458(4):3644–3654.
- Binney, J. and Tremaine, S. (2008). *Galactic Dynamics: Second Edition*.
- Bonnell, I. A. and Bate, M. R. (2002). Accretion in stellar clusters and the collisional formation of massive stars. *Monthly Notices of the Royal Astronomical Society*, 336(2):659–669.
- Bromm, V. and Loeb, A. (2003). Formation of the first supermassive black holes. *The Astrophysical Journal*, 596(1):34–46.
- Chon, S., Hosokawa, T., and Omukai, K. (2021). Cosmological direct-collapse black hole formation sites hostile for their growth. *Monthly Notices of the Royal Astronomical Society*, 502(1):700–713.
- Chung, E., Jo, Y., Kim, J.-h., Jung, M., and Kwon, O.-k. (2026). From dense gas clouds to supermassive black hole seeds: Hybrid hydro/direct n-body simulations of runaway collision-driven

- intermediate-mass black hole formation. *The Astrophysical Journal*, 1004(2):250.
- Clark, P. C., Glover, S. C., Smith, R. J., Greif, T. H., Klessen, R. S., and Bromm, V. (2011). The formation and fragmentation of disks around primordial protostars. *Science*, 331(6020):1040–1042.
- Cohn, H. (1980). Late core collapse in star clusters and the gravothermal instability. *Astrophysical Journal, Part 1, vol. 242, Dec. 1, 1980, p. 765-771.*, 242:765–771.
- Collins, G. W. et al. (1978). The virial theorem in stellar astrophysics. *Tucson, Ariz., Pachart Publishing House (Astronomy and Astrophysics Series. Volume 7), 1978. 143 p.*
- Das, A., Schleicher, D. R., Leigh, N. W., and Boekholt, T. C. (2021). Formation of supermassive black hole seeds in nuclear star clusters via gas accretion and runaway collisions. *Monthly Notices of the Royal Astronomical Society*, 503(1):1051–1069.
- Doeleman, S., Agol, E., Backer, D., Baganoff, F., Bower, G. C., Broderick, A., Fabian, A., Fish, V., Gammie, C., Ho, P., et al. (2009). Imaging an event horizon: submm-vlbi of a super massive black hole. *arXiv preprint arXiv:0906.3899*.
- EHT Collaboration, Akiyama, K., Alberdi, A., Alef, W., Algaba, J. C., Anantua, R., Asada, K., Azulay, R., Bach, U., Baczko, A.-K., et al. (2022). First sagittarius a* event horizon telescope results. i. the shadow of the supermassive black hole in the center of the milky way. *The Astrophysical Journal Letters*, 930(2):L12.
- Escala, A. (2006). On the fueling of massive black holes and the properties of their host spheroids. *The Astrophysical Journal Letters*, 648(1):L13–L16.
- Escala, A. (2021). Observational support for massive black hole formation driven by runaway stellar collisions in galactic nuclei. *The Astrophysical Journal*, 908(1):57.
- Freitag, M., Gürkan, M. A., and Rasio, F. A. (2006a). Runaway collisions in young star clusters–ii. numerical results. *Monthly Notices of the Royal Astronomical Society*, 368(1):141–161.
- Freitag, M., Rasio, F. A., and Baumgardt, H. (2006b). Runaway collisions in young star clusters–i. methods and tests. *Monthly Notices of the Royal Astronomical Society*, 368(1):121–140.
- Fryer, C., Woosley, S., and Heger, A. (2001). Pair-instability supernovae, gravity waves, and gamma-ray transients. *The Astrophysical Journal*, 550(1):372–382.
- Fujii, M., Iwasawa, M., Funato, Y., and Makino, J. (2007). Bridge: A direct-tree hybrid-body algorithm for fully self-consistent simulations of star clusters and their parent galaxies. *Publications of the Astronomical Society of Japan*, 59(6):1095–1106.
- Garwin, L. and Lincoln, T. (2019). *A century of nature: twenty-one discoveries that changed science and the world*. University of Chicago Press.
- Gerritsen, J., Icke, V., et al. (1997). Star formation in n-body simulation, i. *Astronomy & Astrophysics*, 325.
- Ghez, A., Klein, B., Morris, M., and Becklin, E. (1999). High proper motion stars in the vicinity of sgr a*: Evidence for a supermassive black hole at the center of our galaxy. In *Observational*

- Evidence for Black Holes in the Universe: Proceedings of a Conference held in Calcutta, India, January 10–17, 1998*, pages 265–278. Springer.
- Gillessen, S., Eisenhauer, F., Fritz, T., Bartko, H., Dodds-Eden, K., Pfuhl, O., Ott, T., and Genzel, R. (2009). The orbit of the star s2 around sgr a* from very large telescope and keck data. *The Astrophysical Journal*, 707(2):L114–L117.
- Gingold, R. A. and Monaghan, J. J. (1977). Smoothed particle hydrodynamics: theory and application to non-spherical stars. *Monthly notices of the royal astronomical society*, 181(3):375–389.
- Greenstein, J. L. and Matthews, T. A. (1963). Red-shift of the unusual radio source: 3 c 48. *Nature*, 197(4872):1041–1042.
- Gürkan, M. A., Freitag, M., and Rasio, F. A. (2004). Formation of massive black holes in dense star clusters. i. mass segregation and core collapse. *The Astrophysical Journal*, 604(2):632–652.
- Heggie, D. C. and Mathieu, R. D. (1986). Standardised units and time scales. In Hut, P. and McMillan, S. L. W., editors, *The Use of Supercomputers in Stellar Dynamics*, pages 233–235, Berlin, Heidelberg. Springer Berlin Heidelberg.
- Higham, N. J., Dennis, M. R., Glendinning, P., Martin, P. A., Santosa, F., and Tanner, J. (2015). *Princeton companion to applied mathematics*. Princeton University Press.
- Horedt, G. P. (2004). *Polytropes: applications in astrophysics and related fields*. Springer.
- Hoyle, F. and Lyttleton, R. A. (1939). The evolution of the stars. In *Mathematical Proceedings of the Cambridge Philosophical Society*, volume 35, pages 592–609. Cambridge University Press.
- Inayoshi, K., Nakatani, R., Toyouchi, D., Hosokawa, T., Kuiper, R., and Onoue, M. (2022). Rapid growth of seed black holes during early bulge formation. *The Astrophysical Journal*, 927(2):237.
- Jiang, L., McGreer, I. D., Fan, X., Bian, F., Cai, Z., Clement, B., Wang, R., and Fan, Z. (2015). Discovery of eight $z > 6$ quasars in the sloan digital sky survey overlap regions. *The Astronomical Journal*, 149(6):188.
- Kritos, K., Berti, E., and Silk, J. (2023). Massive black hole assembly in nuclear star clusters. *Physical Review D*, 108(8):083012.
- Latif, M. A. and Ferrara, A. (2016). Formation of supermassive black hole seeds. *Publications of the Astronomical Society of Australia*, 33:e051.
- Lee, H. M. (1987). Dynamical effects of successive mergers on the evolution of spherical stellar systems. *The astrophysical journal*, 319:801.
- Leigh, N. W., Mastrobuono-Battisti, A., Perets, H. B., and Böker, T. (2014). Stellar dynamics in gas: the role of gas damping. *Monthly Notices of the Royal Astronomical Society*, 441(2):919–932.
- Liao, W.-T. and Turk, M. (2019). Fragmentation in a primordial accretion flow. *arXiv preprint arXiv:1911.00610*.
- Lucy, L. B. (1977). A numerical approach to the testing of the fission hypothesis. *Astronomical*

- Journal*, vol. 82, Dec. 1977, p. 1013-1024., 82:1013–1024.
- Lynden-Bell, D. (1969). Galactic nuclei as collapsed old quasars. *Nature*, 223(5207).
- Makino, J. and Aarseth, S. J. (1992). On a hermite integrator with ahmad-cohen scheme for gravitational many-body problems. *Publications of the Astronomical Society of Japan*, 44(2):141–151.
- Milosavljević, M., Couch, S. M., and Bromm, V. (2009). Accretion onto intermediate-mass black holes in dense protogalactic clouds. *The Astrophysical Journal*, 696(2):L146–L149.
- Monaghan, J. J. (1992). Smoothed particle hydrodynamics. In: *Annual review of astronomy and astrophysics*. Vol. 30 (A93-25826 09-90), p. 543-574., 30:543–574.
- Mortlock, D. J., Warren, S. J., Venemans, B. P., Patel, M., Hewett, P. C., McMahon, R. G., Simpson, C., Theuns, T., Gonzáles-Solares, E. A., Adamson, A., et al. (2011). A luminous quasar at a redshift of $z = 7.085$. *Nature*, 474(7353):616–619.
- Naiman, J., Ramirez-Ruiz, E., Debuhr, J., and Ma, C.-P. (2015). The role of nuclear star clusters in enhancing supermassive black hole feeding rates during galaxy mergers. *The Astrophysical Journal*, 803(2):81.
- Natarajan, P. and Treister, E. (2009). Is there an upper limit to black hole masses? *Monthly Notices of the Royal Astronomical Society*, 393(3):838–845.
- Nelson, R. P. and Papaloizou, J. C. (1994). Variable smoothing lengths and energy conservation in smoothed particle hydrodynamics. *Monthly Notices of the Royal Astronomical Society*, 270(1):1–20.
- O’Connor, E. and Ott, C. D. (2011). Black hole formation in failing core-collapse supernovae. *The Astrophysical Journal*, 730(2):70.
- Omukai, K., Schneider, R., and Haiman, Z. (2008). Can supermassive black holes form in metal-enriched high-redshift protogalaxies? *The Astrophysical Journal*, 686(2):801–814.
- Ostriker, E. C. (1999). Dynamical friction in a gaseous medium. *The Astrophysical Journal*, 513(1):252–258.
- O’Shea, B. W. and Norman, M. L. (2007). Population iii star formation in a λ cdm universe. i. the effect of formation redshift and environment on protostellar accretion rate. *The Astrophysical Journal*, 654(1):66–92.
- O’Shea, B. W. and Norman, M. L. (2008). Population iii star formation in a λ cdm universe. ii. effects of a photodissociating background. *The Astrophysical Journal*, 673(1):14–33.
- Pelupessy, F. I. (2005). *Numerical studies of the interstellar medium on galactic scales*. PhD thesis, éditeur non identifié.
- Pfister, H., Volonteri, M., Dubois, Y., Dotti, M., and Colpi, M. (2019). The erratic dynamical life of black hole seeds in high-redshift galaxies. *Monthly Notices of the Royal Astronomical Society*, 486(1):101–111.
- Plummer, H. C. (1911). On the problem of distribution in globular star clusters. *Monthly Notices*

- of the Royal Astronomical Society, Vol. 71, p. 460-470, 71:460–470.
- Portegies Zwart, S. and McMillan, S. (2018). *Astrophysical Recipes: the art of AMUSE*. IOP Publishing.
- Portegies Zwart, S. F., Baumgardt, H., Hut, P., Makino, J., and McMillan, S. L. (2004). Formation of massive black holes through runaway collisions in dense young star clusters. *Nature*, 428(6984):724–726.
- Portegies Zwart, S. F. and McMillan, S. L. (2002). The runaway growth of intermediate-mass black holes in dense star clusters. *The Astrophysical Journal*, 576(2):899–907.
- Rasio, F. A. (2000). Particle methods in astrophysical fluid dynamics. *Progress of Theoretical Physics Supplement*, 138:609–621.
- Rasio, F. A., Freitag, M., and Gürkan, M. A. (2003). Formation of massive black holes in dense star clusters. *arXiv preprint astro-ph/0304038*.
- Regan, J. and Volonteri, M. (2024). Massive black hole seeds. *arXiv preprint arXiv:2405.17975*.
- Reinoso, B., Klessen, R. S., Schleicher, D., Glover, S. C., and Solar, P. (2023). Formation of supermassive stars in the first star clusters. *Monthly Notices of the Royal Astronomical Society*, 521(3):3553–3569.
- Reinoso, B., Schleicher, D., Fellhauer, M., Leigh, N., and Klessen, R. (2020). The effects of a background potential in star cluster evolution: A delay in the relaxation time-scale and runaway collision processes. *Astronomy & Astrophysics*, 639:A92.
- Reinoso, B. A. (2024). *Formation of massive black hole seeds through runaway stellar collisions and gas accretion in dense stellar systems*. PhD thesis.
- Sakurai, Y., Yoshida, N., Fujii, M. S., and Hirano, S. (2017). Formation of intermediate-mass black holes through runaway collisions in the first star clusters. *Monthly Notices of the Royal Astronomical Society*, 472(2):1677–1684.
- Salmon, J. K. and Warren, M. S. (1994). Skeletons from the treecode closet. *Journal of Computational Physics*, 111(1):136–155.
- Salpeter, E. E. (1965). Accretion of interstellar matter by massive objects.
- Schmidt, M. (1963). 3c273: a star-like object with large red-shift.”. *A Century of Nature, Twenty One Discoveries That Changed Science and the World*, page 43.
- Shang, C., Bryan, G. L., and Haiman, Z. (2010). Supermassive black hole formation by direct collapse: keeping protogalactic gas h2 free in dark matter haloes with virial temperatures $t_{\text{vir}} > 104 \text{ K}$. *Monthly Notices of the Royal Astronomical Society*, 402(2):1249–1262.
- Shibata, M. and Shapiro, S. L. (2002). Collapse of a rotating supermassive star to a supermassive black hole: Fully relativistic simulations. *The Astrophysical Journal Letters*, 572(1):L39–L43.
- Shields, G. A. (1999). A brief history of active galactic nuclei. *Publications of the Astronomical Society of the Pacific*, 111(760):661–678.

- Shlosman, I., Begelman, M. C., and Frank, J. (1990). The fuelling of active galactic nuclei. *Nature*, 345(6277):679–686.
- Smith, B. D., Regan, J. A., Downes, T. P., Norman, M. L., O’Shea, B. W., and Wise, J. H. (2018). The growth of black holes from population iii remnants in the renaissance simulations. *Monthly Notices of the Royal Astronomical Society*, 480(3):3762–3773.
- Spitzer, L. (1987). *Dynamical evolution of globular clusters*.
- Springel, V. (2005). The cosmological simulation code gadget-2. *Monthly notices of the royal astronomical society*, 364(4):1105–1134.
- Springel, V. and Hernquist, L. (2002). Cosmological smoothed particle hydrodynamics simulations: the entropy equation. *Monthly Notices of the Royal Astronomical Society*, 333(3):649–664.
- Sweatman, W. L. (1993). A study of Lagrangian radii oscillations and core-wandering using N-body simulations. *MNRAS*, 261(3):497–512.
- Trenti, M. (2006). Dynamical evidence for intermediate mass black holes in old globular clusters. *arXiv preprint astro-ph/0612040*.
- Trenti, M. and van der Marel, R. (2013). No energy equipartition in globular clusters. *Monthly Notices of the Royal Astronomical Society*, 435(4):3272–3282.
- Vergara, M., Schleicher, D., Boekholt, T., Reinoso, B., Fellhauer, M., Klessen, R., and Leigh, N. (2021). Stellar collisions in flattened and rotating population iii star clusters. *Astronomy & Astrophysics*, 649:A160.
- Vergara, M., Schleicher, D., Escala, A., Reinoso, B., Flammini Dotti, F., Kamlah, A., Liempi, M., Hoyer, N., Neumayer, N., and Spurzem, R. (2024). Efficiency of black hole formation via collisions in stellar systems: Data analysis from simulations and observations. *Astronomy & Astrophysics*, 689:A34.
- Vergara, M. C., Askar, A., Kamlah, A. W., Spurzem, R., Flammini Dotti, F., Schleicher, D. R., Arca Sedda, M., Hypki, A., Giersz, M., Hurley, J., et al. (2025). Rapid formation of a very massive star ($> 50000 m_{\odot}$), and subsequently, of an imbh, from runaway collisions: Direct n-body and monte carlo simulations of dense star clusters. *Astronomy & Astrophysics*, 704:A321.
- Vergara, M. C., Escala, A., Schleicher, D., and Reinoso, B. (2023). Global instability by runaway collisions in nuclear stellar clusters: numerical tests of a route for massive black hole formation. *Monthly Notices of the Royal Astronomical Society*, 522(3):4224–4237.
- Vesperini, E., McMillan, S. L., D’Ercole, A., and D’Antona, F. (2010). Intermediate-mass black holes in early globular clusters. *The Astrophysical Journal Letters*, 713(1):L41–L44.
- Volonteri, M. (2010). Formation of supermassive black holes. *The Astronomy and Astrophysics Review*, 18(3):279–315.
- Wang, F., Yang, J., Fan, X., Hennawi, J. F., Barth, A. J., Banados, E., Bian, F., Boutsia, K., Connor, T., Davies, F. B., et al. (2021). A luminous quasar at redshift 7.642. *The Astrophysical Journal Letters*, 907(1):L1.

- Weisstein, E. W. (2002). Lambert w-function. <https://mathworld.wolfram.com/>.
- Willott, C. J., Delorme, P., Omont, A., Bergeron, J., Delfosse, X., Forveille, T., Albert, L., Reyl , C., Hill, G. J., Gully-Santiago, M., et al. (2007). Four quasars above redshift 6 discovered by the canada-france high-z quasar survey. *The Astronomical Journal*, 134(6):2435–2450.
- Woods, T., Heger, A., Whalen, D. J., Haemmerl , L., and Klessen, R. S. (2017). On the maximum mass of accreting primordial supermassive stars. *The Astrophysical Journal Letters*, 842(1):L6.
- Woods, T. E., Agarwal, B., Bromm, V., Bunker, A., Chen, K.-J., Chon, S., Ferrara, A., Glover, S. C., Haemmerle, L., Haiman, Z., et al. (2019). Titans of the early universe: The prato statement on the origin of the first supermassive black holes. *Publications of the Astronomical Society of Australia*, 36:e027.
- Yang, D., Hennawi, J., Guarneri, F., Wolf, J., Belladitta, S., Schindler, J.-T., Hughes, A., Ba ados, E., Mortlock, D., Yang, J., et al. (2026). Euclid: Discovery of 31 new quasars at $6.6 < z < 7.8$. *Astronomy & Astrophysics*, 711:A104.

Appendices

Appendix A. Derivation of Critical Masses for Collisions and Relaxation

In this section we derive two quantities relevant for the dynamical evolution of stellar clusters: the critical mass for collisions (following Escala (2021)) and the relaxation mass. For the critical mass, we first present the original derivation, then extend it to include an external potential (following Reinoso et al. (2020)), and finally generalize it by including gravitational focusing. For the relaxation mass, we derive an analogous mass scale from the two-body relaxation timescale.

A.1. Critical Mass in the Simplified Case

First, we show the derivation of the critical mass from Escala (2021). We begin with the collisional timescale (Equation (2.5)) from Section 2.1, for a star cluster of equal stars with mass m and radius r_* , total mass M and characteristic radius R :

$$t_{\text{coll}} = \frac{1}{n\Sigma_0} \sqrt{\frac{R}{GM}}, \quad (\text{A.1})$$

where $\Sigma_0 = 16\sqrt{\pi}r_*^2(1 + \Theta)$ is the effective cross-section, n is the number density, and

$$\Theta = 9.54 \frac{m}{M_\odot} \frac{R_\odot}{r_*} \left(\frac{100 \text{ km s}^{-1}}{\sigma} \right)^2, \quad (\text{A.2})$$

is the Safronov number. This derivation relies on two assumptions. First, we assume the system is virialized, so the velocity dispersion σ can be expressed as a function of the total mass M and characteristic radius R . Second, we assume $\sigma \sim 100 \text{ km s}^{-1}$, then the Safronov number depends only on the stellar mass and radius. The implication of this assumption is that the effective cross-section depends only on the geometric cross-section of the stars, effectively neglecting gravitational focusing (i.e., the dependence on the velocity dispersion). Collisions become dynamically important

when $t_{\text{coll}} \leq t_{\text{H}}$:

$$t_{\text{coll}} = \frac{1}{n\Sigma_0} \sqrt{\frac{R}{GM}} , \quad (\text{A.3})$$

$$= \frac{4\pi R^3 m}{3M\Sigma_0} \sqrt{\frac{R}{GM}} , \quad (\text{A.4})$$

$$= \frac{4\pi R^{7/2} m}{3M^{3/2} \Sigma_0 G^{1/2}} \leq t_{\text{H}} , \quad (\text{A.5})$$

where $n = 3M/(4\pi R^3 m)$ is the number density for an homogeneous system. Solving for M :

$$M^{3/2} \geq \frac{4\pi R^{7/2} m}{3t_{\text{H}} \Sigma_0 G^{1/2}} , \quad (\text{A.6})$$

$$M \geq \left(\frac{4\pi m}{3\Sigma_0 G^{1/2} t_{\text{H}}} \right)^{2/3} R^{7/3} . \quad (\text{A.7})$$

The critical mass is then:

$$M_{\text{crit}} = \left(\frac{4\pi m}{3\Sigma_0 G^{1/2} t_{\text{H}}} \right)^{2/3} R^{7/3} . \quad (\text{A.8})$$

This defines the mass limit for a virialized system of characteristic radius R , composed of equal-mass stars (mass m , radius r_*), for which the collision time equals the characteristic time ($t_{\text{coll}} = t_{\text{H}}$). If a system exceeds this mass, its collision time is shorter than t_{H} . This limit can be represented in a radius–mass diagram (Escala, 2021; Vergara et al., 2023).

A.2. Effect of an External Gravitational Potential

We now show how the critical mass changes when an external potential is considered. When no external potential is present, the virial theorem gives a relation for the time average kinetic and potential energies (K and W):

$$2K + W = 0 . \quad (\text{A.9})$$

Although K and W are time-average quantities, in virial equilibrium they fluctuate about their

mean values (for more details see Collins et al. (1978)). We can express K and W in terms of characteristic quantities:

$$K = \frac{1}{2} \langle \sum_i m_i v_i^2 \rangle = \frac{M}{2} \langle v^2 \rangle , \quad (\text{A.10})$$

$$W = -\frac{1}{2} \langle \sum_{i,j \neq i} \frac{G m_i m_j}{|r_i - r_j|} \rangle = -\frac{GM^2}{2R_{\text{vir}}} . \quad (\text{A.11})$$

Note that m_i is the mass of the i -th star, and $M = \sum_i m_i$ is the total mass of the stellar system. For a stellar system, it is standard to replace $\langle v^2 \rangle$ with the square of the velocity dispersion, σ^2 . The quantity R_{vir} is known as the virial radius (for consistency, we use R instead of R_{vir} in the following). Replacing in Equation (A.9), we have:

$$\frac{M}{2} \sigma^2 = \frac{GM^2}{2R} . \quad (\text{A.12})$$

The relation between σ and R_{vir} is then:

$$\sigma = \sqrt{\frac{GM}{R}} . \quad (\text{A.13})$$

Following Reinoso et al. (2020), if we consider an external potential of mass M_{ext} with the same spatial distribution as the stellar system, then the total mass can be rewritten as $M + M_{\text{ext}}$. Assuming the stars are in global virial equilibrium with this external potential yields:

$$\frac{M}{2} \sigma_{\text{ext}}^2 = \frac{G(M + M_{\text{ext}})^2}{2R} . \quad (\text{A.14})$$

Defining $q = M_{\text{ext}}/M$,

$$\sigma_{\text{ext}} = \sqrt{\frac{GM}{R}} (1 + q) . \quad (\text{A.15})$$

Then, for an external potential, the collision time is:

$$t_{\text{coll}} = \frac{1}{n\Sigma_0} \sqrt{\frac{R}{GM}} \frac{1}{(1+q)} , \quad (\text{A.16})$$

$$= \frac{4\pi R^3 m}{3M\Sigma_0} \sqrt{\frac{R}{GM}} \frac{1}{(1+q)} , \quad (\text{A.17})$$

$$= \frac{4\pi R^{7/2} m}{3M^{3/2} \Sigma_0 G^{1/2} (1+q)} , \quad (\text{A.18})$$

and the critical mass can be written as

$$M_{\text{crit,ext}} = \left(\frac{4\pi m}{3\Sigma_0 G^{1/2} t_{\text{H}} (1+q)} \right)^{2/3} R^{7/3} . \quad (\text{A.19})$$

A.3. Generalized Critical Mass Including Gravitational Focusing

In Section A.1, we assumed $\sigma \sim 100 \text{ km s}^{-1}$. Consequently, the Safronov number depended only on the stellar mass and radius. To account for gravitational focusing, we now derive the critical mass using the virial velocity dispersion. In the following, we use M to denote the mass of the stellar cluster. Then, the total mass of the system is $M(1+q)$ when considering an external potential (with $q = M_{\text{ext}}/M$), and M when no external potential is present. First, we expand the effective cross section:

$$\Sigma_0 = 16\sqrt{\pi} r_{\star}^2 (1 + \Theta) , \quad (\text{A.20})$$

$$= 16\sqrt{\pi} r_{\star}^2 \left(1 + 9.54 \frac{m}{M_{\odot}} \frac{R_{\odot}}{r_{\star}} \left(\frac{100 \text{ km s}^{-1}}{\sigma_{\text{ext}}} \right)^2 \right) , \quad (\text{A.21})$$

$$= 16\sqrt{\pi} r_{\star}^2 + \frac{9.54 \times 16\sqrt{\pi} r_{\star} m R_{\odot} v_{100}^2}{M_{\odot}} \frac{R}{G M (1+q)^2} . \quad (\text{A.22})$$

We have introduced σ_{ext} , and have defined $v_{100} = 100 \text{ km s}^{-1}$. From $t_{\text{coll}} = t_{\text{H}}$ in Equation (A.18), we have

$$\Sigma_0 M^{3/2} = \frac{4\pi m}{3G^{1/2} t_{\text{H}} (1+q)} R^{7/2} . \quad (\text{A.23})$$

Replacing Equation (A.22) into (A.23) yields,

$$16\sqrt{\pi}r_\star^2 M^{3/2} + \frac{9.54 \times 16\sqrt{\pi} r_\star m R_\odot v_{100}^2}{M_\odot} \frac{R M^{1/2}}{G(1+q)^2} = \frac{4\pi m}{3G^{1/2}t_H} \frac{R^{7/2}}{(1+q)} , \quad (\text{A.24})$$

$$M^{3/2} + \frac{9.54 m R_\odot v_{100}^2 R}{r_\star G M_\odot (1+q)^2} M^{1/2} = \frac{\sqrt{\pi} m}{12G^{1/2}t_H r_\star^2} \frac{R^{7/2}}{(1+q)} . \quad (\text{A.25})$$

Now, defining the coefficients

$$A = \frac{9.54 m R_\odot v_{100}^2}{r_\star G M_\odot (1+q)^2} R , \quad (\text{A.26})$$

$$B = \frac{\sqrt{\pi} m}{12G^{1/2}t_H r_\star^2 (1+q)} R^{7/2} , \quad (\text{A.27})$$

we can rewrite Equation (A.25):

$$M^{3/2} + A M^{1/2} = B . \quad (\text{A.28})$$

To continue, we introduce the change of variable $u = M^{1/2}$:

$$u^3 + Au - B = 0 . \quad (\text{A.29})$$

Note that u must be positive since $M > 0$. As A and B are positive, the function $f(u) = u^3 + Au - B$ ranges from $-B$ to ∞ . Then, $f(u)$ is strictly increasing, and there is exactly one real solution. Using Cardano's formula, the real solution for a depressed cubic equation is,

$$u_{\text{real}} = \sqrt[3]{\frac{B}{2} + \sqrt{\frac{B^2}{4} + \frac{A^3}{27}}} + \sqrt[3]{\frac{B}{2} - \sqrt{\frac{B^2}{4} + \frac{A^3}{27}}} . \quad (\text{A.30})$$

Returning to $M = u^2$, we obtain

$$M = u_{\text{real}}^2, \quad (\text{A.31})$$

$$= \left(\frac{B}{2} + \sqrt{\frac{B^2}{4} + \frac{A^3}{27}} \right)^{2/3} + \left(\frac{B}{2} - \sqrt{\frac{B^2}{4} + \frac{A^3}{27}} \right)^{2/3}, \quad (\text{A.32})$$

$$+ 2 \left(\frac{B^2}{4} - \left(\frac{B^2}{4} + \frac{A^3}{27} \right) \right)^{1/3}, \quad (\text{A.33})$$

$$= \left(\frac{B}{2} + \sqrt{\frac{B^2}{4} + \frac{A^3}{27}} \right)^{2/3} + \left(\frac{B}{2} - \sqrt{\frac{B^2}{4} + \frac{A^3}{27}} \right)^{2/3} - \frac{2A}{3}. \quad (\text{A.34})$$

Defining $\alpha = A/R$ and $\beta = B/R^{7/2}$, the critical mass is:

$$M_{\text{crit}} = \left(\frac{\beta R^{7/2}}{2} + \sqrt{\frac{\beta^2 R^7}{4} + \frac{\alpha^3 R^3}{27}} \right)^{2/3} + \left(\frac{\beta R^{7/2}}{2} - \sqrt{\frac{\beta^2 R^7}{4} + \frac{\alpha^3 R^3}{27}} \right)^{2/3} - \frac{2\alpha R}{3}. \quad (\text{A.35})$$

Equation (A.35) sets the upper mass limit for a stellar system of mass M , embedded in and in virial equilibrium with an external potential (with mass $M_{\text{ext}} = qM$), such that its collision timescale satisfies $t_{\text{coll}} = t_{\text{H}}$. In consequence, if $M > M_{\text{crit}}$, the collision time is shorter than t_{H} .

A.4. Derivation of the Relaxation Mass

In this section, we derive a similar relation for the relaxation timescale t_{relax} . In a stellar cluster, the energy of stars changes due to stellar encounters, which are typically divided into strong and weak encounters. If we consider a target star with velocity v , a strong encounter is one in which the change in velocity δv is comparable to v (i.e., $\delta v \sim v$). On the other hand, in a weak encounter, the change in velocity is small ($\delta v \ll v$). While strong encounters can produce notable effects (e.g., stellar collisions, binary formation), they are less frequent. Weak encounters, however, can accumulate over time and significantly alter the energy distribution and trajectories of stars. The relaxation timescale is the time required for a typical star's velocity to change significantly due to the cumulative effect of weak encounters. From Binney and Tremaine (2008), the deflection in the star velocity for crossing the system once is:

$$\delta v^2 \simeq 8N \left(\frac{Gm}{Rv} \right)^2 \ln(\Lambda), \quad (\text{A.36})$$

where N is the number of stars, m and v are the mass and velocity of the target star, and $\Lambda = R/b_{90}$ is the ratio of the maximum to minimum impact parameter for weak encounters. From Binney and Tremaine (2008), the minimum impact parameter is $b_{90} = 2Gm/v^2$, then $\Lambda = v^2 R/(2Gm)$. The number of system crossings required to change v significantly is

$$n_{\text{cross}} = \frac{v^2}{\delta v^2} = \frac{1}{8N \ln(\Lambda)} \left(\frac{Rv^2}{Gm} \right)^2. \quad (\text{A.37})$$

When an external potential is present, we can approximate $v^2 \sim \sigma_{\text{ext}}^2$, and n_{cross} becomes

$$n_{\text{cross}} = \frac{1}{8N \ln \left(\frac{GmN(1+q)^2}{R} \frac{R}{2Gm} \right)} \left(\frac{R}{Gm} \frac{GmN(1+q)^2}{R} \right)^2, \quad (\text{A.38})$$

$$\simeq \frac{0.1N(1+q)^4}{\ln(N(1+q)^2)}. \quad (\text{A.39})$$

In the last equation, we have used $M = Nm$, and we have neglected the $1/2$ inside $\ln(\Lambda)$. In the absence of an external potential (i.e., $q = 0$), Equation (A.39) reduces to the standard expression from Binney and Tremaine (2008). Since a typical star crosses the system in a crossing time t_{cross} , the relaxation timescale is:

$$t_{\text{relax}} = n_{\text{cross}} t_{\text{cross}} = \frac{0.1N(1+q)^4}{\ln(N(1+q)^2)} t_{\text{cross}}. \quad (\text{A.40})$$

With the crossing time estimated as $t_{\text{cross}} = R/\sigma$, and using σ_{ext} , Equation (A.40) becomes

$$t_{\text{relax}} = \frac{0.1N(1+q)^3}{\ln(N(1+q)^2)} \frac{R^{3/2}}{\sqrt{GM}}. \quad (\text{A.41})$$

Imposing $t_{\text{relax}} < t_{\text{H}}$ yields a relation between the radius R and mass M :

$$R^{3/2} \leq \frac{t_H \ln(N(1+q)^2) \sqrt{GM}}{0.1N(1+q)^3}, \quad (\text{A.42})$$

$$\leq \frac{t_H m \ln(M(1+q)^2/m) \sqrt{GM}}{0.1M(1+q)^3}, \quad (\text{A.43})$$

$$= \frac{t_H m}{0.1(1+q)^3} \ln\left(\frac{M(1+q)^2}{m}\right) \sqrt{\frac{G}{M}}, \quad (\text{A.44})$$

$$R \leq \left(\frac{t_H m}{0.1(1+q)^3} \ln\left(\frac{M(1+q)^2}{m}\right) \right)^{2/3} \left(\frac{G}{M} \right)^{1/3}. \quad (\text{A.45})$$

Equation (A.45) defines the condition for a system to significantly alter its initial conditions on a timescale $\sim t_H$. If we denote the equality case by R_{relax} , systems with $R > R_{\text{relax}}$ have a relaxation timescale longer than t_H . In a collisionless scenario, such systems are considered to be in virial equilibrium.

We now derive a mass scale from the condition in Equation (A.45). First, we rearrange the terms in (A.45) for the limit case:

$$R = \left(\frac{t_H m}{0.1(1+q)^3} \ln\left(\frac{M(1+q)^2}{m}\right) \right)^{2/3} \left(\frac{G}{M} \right)^{1/3}, \quad (\text{A.46})$$

$$R^3 M = \left(\frac{t_H m}{0.1(1+q)^3} \ln\left(\frac{M(1+q)^2}{m}\right) \right)^2 G. \quad (\text{A.47})$$

We introduce the change of variable $u \equiv M(1+q)^2/m$,

$$\frac{R^3 u m}{(1+q)^2} = \left(\frac{t_H m}{0.1(1+q)^3} \ln(u) \right)^2 G, \quad (\text{A.48})$$

$$\frac{R^{3/2} u^{1/2} m^{1/2}}{1+q} = \frac{t_H m}{0.1(1+q)^3} \ln(u) G^{1/2}, \quad (\text{A.49})$$

$$u^{1/2} = \frac{t_H \sqrt{Gm}}{0.1(1+q)^2 R^{3/2}} \ln(u). \quad (\text{A.50})$$

Defining $\lambda = (t_H \sqrt{Gm}) / (0.1(1+q)^2 R^{3/2})$:

$$u^{1/2} = \lambda \ln(u), \quad (\text{A.51})$$

and introducing another change of variable, $\omega = \ln(u)$:

$$e^{\omega/2} = \lambda \omega, \quad (\text{A.52})$$

$$\frac{1}{\lambda} = \omega e^{-\omega/2}, \quad (\text{A.53})$$

$$\frac{-1}{2\lambda} = \frac{-\omega}{2} e^{-\omega/2}. \quad (\text{A.54})$$

Equation (A.54) is solved by the Lambert W function (Weisstein, 2002). Indeed, for a function $f(w) = we^w$, its inverse is the Lambert $W(z)$ function, defined such that $W(f(w)) = w$. Hence,

$$\frac{-\omega}{2} = W\left(\frac{-1}{2\lambda}\right), \quad (\text{A.55})$$

$$\omega = -2W\left(\frac{-1}{2\lambda}\right). \quad (\text{A.56})$$

Returning to $u = e^\omega$,

$$u = \exp\left(-2W\left(\frac{-1}{2\lambda}\right)\right). \quad (\text{A.57})$$

Finally, we use $M = mu/(1+q)^2$:

$$M_{\text{relax}} = \frac{m}{(1+q)^2} \exp\left(-2W\left(\frac{-1}{2\lambda}\right)\right). \quad (\text{A.58})$$

Since $-1/2\lambda < 0$ ($\lambda > 0$), the argument of the Lambert W function is negative, and there are two real-valued branches (Higham et al., 2015, Chapter III.17). The principal branch, $W_0(z)$, is defined for $z \in [-1/e, \infty)$, and takes values $W_0(z) > -1$. In particular, for $z \in [-1/e, 0]$, it ranges from -1 to 0 . The lower branch, $W_{-1}(z)$, is defined for $z \in [-1/e, 0]$, and ranges from -1 to $-\infty$, with $\lim_{z \rightarrow 0^-} W_{-1}(z) = -\infty$. Figure A.1 shows the principal and lower branches of the Lambert function.

If we consider the principal branch $W_0(z)$, we note that the exponential in Equation (A.58) takes values between e^0 and e^2 . Consequently, the relaxation mass would be of the order of the individual stellar mass ($M_{\text{relax}} \sim m$). In contrast, for the lower branch $W_{-1}(z)$, the exponential ranges from e^2 to ∞ , yielding a relaxation mass that can reach the mass scale of the system. We therefore adopt




Figure A.1: Lambert function. The black line shows the $W_0(z)$ branch, and the blue line shows the $W_{-1}(z)$ branch.

the lower branch as the physically relevant solution:

$$M_{\text{relax}} = \frac{m}{(1+q)^2} \exp\left(-2W_{-1}\left(\frac{-1}{2\lambda}\right)\right). \quad (\text{A.59})$$

Finally, we must verify that the argument of the Lambert W function lies within the domain of the lower branch: $-1/e \leq -1/2\lambda < 0$. The right inequality is satisfied since $\lambda > 0$. The left part gives a condition over the characteristic timescale:

$$-1/e \leq -1/2\lambda , \quad (\text{A.60})$$

$$-2\lambda \leq -e , \quad (\text{A.61})$$

$$\lambda \geq e/2 , \quad (\text{A.62})$$

$$\frac{t_{\text{H}}\sqrt{Gm}}{0.1(1+q)^2R^{3/2}} \geq \frac{e}{2} , \quad (\text{A.63})$$

$$t_{\text{H}} \geq \frac{0.1e(1+q)^2R^{3/2}}{2\sqrt{Gm}} . \quad (\text{A.64})$$

Equation (A.64) implies that, for a system of characteristic radius R composed of equal-mass stars m , there exists a minimum possible relaxation time. From Equation (A.41), for fixed R and m , the number of stars that minimizes t_{relax} is $N_{\text{opt}} = e^2/(1+q)^2$. If we evaluate $t_{\text{relax}}(N_{\text{opt}})$:

$$t_{\text{relax}}(N_{\text{opt}}) = \frac{0.1N_{\text{opt}}(1+q)^3}{\ln(N_{\text{opt}}(1+q)^2)} \frac{R^{3/2}}{\sqrt{GN_{\text{opt}}m}} , \quad (\text{A.65})$$

$$= \frac{0.1e^2(1+q)^{-2}(1+q)^3}{\ln(e^2(1+q)^{-2}(1+q)^2)} \frac{R^{3/2}}{\sqrt{Ge^2(1+q)^{-2}m}} , \quad (\text{A.66})$$

$$= \frac{0.1e^2(1+q)}{2} \frac{R^{3/2}(1+q)}{\sqrt{Gme}} , \quad (\text{A.67})$$

$$= \frac{0.1e(1+q)^2R^{3/2}}{2\sqrt{Gm}} . \quad (\text{A.68})$$

Appendix B. Fourth-Order Hermite Integration Scheme

The N-body code used in this work, `ph4`, is based on a fourth-order Hermite integrator. The following subsections describe in detail the individual time step scheme (ITS) from Makino and Aarseth (1992), including the time step criterion in the N-body code.

B.1. Step 1: Particle selection

The first step is to select the next particle to be advanced. At any given time, each particle j has its own mass m_j , time t_j , time step Δt_j , coordinates $\mathbf{x}_j$ and $\mathbf{v}_j$, acceleration $\mathbf{a}_j$ and jerk $\dot{\mathbf{a}}_j$ (with $\dot{\mathbf{a}}_j = \partial \mathbf{a}_j / \partial t$). The selected particle i is the one satisfying the condition:

$$i = \arg \min_j (t_j + \Delta t_j) \quad (\text{B.1})$$

Note that always $t_j \leq t_i + \Delta t_i$ for all j . The current simulation time is then set to $t = t_i + \Delta t_i$.

B.2. Step 2: Calculate the predicted position and velocities for all particles

The positions and velocities of all particles are predicted at time t , using a Taylor expansion up to third-order (jerk):

$$\mathbf{x}_{p,j} = \mathbf{x}_j + \mathbf{v}_j(t - t_j) + \frac{\mathbf{a}_{0,j}}{2!}(t - t_j)^2 + \frac{\dot{\mathbf{a}}_{0,j}}{3!}(t - t_j)^3, \quad (\text{B.2})$$

$$\mathbf{v}_{p,j} = \mathbf{v}_j + \mathbf{a}_{0,j}(t - t_j) + \frac{\dot{\mathbf{a}}_{0,j}}{2!}(t - t_j)^2. \quad (\text{B.3})$$

The acceleration is due only to gravitational force. The expression for each particle j is:

$$\mathbf{a}_{0,j} = \sum_{k \neq j} Gm_k \frac{\mathbf{r}_{0,jk}}{(\mathbf{r}_{0,jk}^2 + \varepsilon^2)^{3/2}}, \quad (\text{B.4})$$

where ε is the softening parameter, and

$$\mathbf{r}_{0,jk} = \mathbf{r}_k - \mathbf{r}_j. \quad (\text{B.5})$$

Differentiating Eq. (B.4) with respect to time yields the jerk:

$$\dot{\mathbf{a}}_{0,j} = \frac{\partial \mathbf{a}_j}{\partial t}, \quad (\text{B.6})$$

$$= \sum_{k \neq j} Gm_k \frac{\partial}{\partial t} \frac{\mathbf{r}_{0,jk}}{(\mathbf{r}_{0,jk}^2 + \varepsilon^2)^{3/2}}, \quad (\text{B.7})$$

$$\dot{\mathbf{a}}_{0,j} = \sum_{k \neq j} Gm_k \left[\frac{\mathbf{v}_{0,jk}}{(\mathbf{r}_{0,jk}^2 + \varepsilon^2)^{3/2}} - \frac{3(\mathbf{v}_{0,jk} \cdot \mathbf{r}_{0,jk})\mathbf{r}_{0,jk}}{(\mathbf{r}_{0,jk}^2 + \varepsilon^2)^{5/2}} \right], \quad (\text{B.8})$$

where

$$\mathbf{v}_{0,jk} = \mathbf{v}_k - \mathbf{v}_j. \quad (\text{B.9})$$

B.3. Step 3: Calculate acceleration and jerk for particle i

Similar to equations (B.4) and (B.8), the acceleration and jerk are calculated using the predicted positions and velocity:

$$\mathbf{a}_{1,i} = \sum_{j \neq i} Gm_j \frac{\mathbf{r}_{ij}}{(\mathbf{r}_{ij}^2 + \epsilon^2)^{3/2}}, \quad (\text{B.10})$$

$$\dot{\mathbf{a}}_{1,i} = \sum_{j \neq i} Gm_j \left[\frac{\mathbf{v}_{ij}}{(\mathbf{r}_{ij}^2 + \epsilon^2)^{3/2}} - \frac{3(\mathbf{v}_{ij} \cdot \mathbf{r}_{ij})\mathbf{r}_{ij}}{(\mathbf{r}_{ij}^2 + \epsilon^2)^{5/2}} \right], \quad (\text{B.11})$$

$$\mathbf{r}_{ij} = \mathbf{r}_{p,j} - \mathbf{r}_{p,i}, \quad (\text{B.12})$$

$$\mathbf{v}_{ij} = \mathbf{v}_{p,j} - \mathbf{v}_{p,i}. \quad (\text{B.13})$$

In this notation, the subscript 0 denotes quantities evaluated at the beginning of the time step (t_j), while the subscript 1 denotes quantities evaluated at the end of the time step (t), using the predicted position and velocity ($\mathbf{x}_{p,i}$ and $\mathbf{v}_{p,i}$).

B.4. Step 4: Second and third time derivative of acceleration, using Hermite interpolation

To obtain a high-order correction of $\mathbf{x}$ and $\mathbf{v}$, the acceleration of particle i is modeled as a third-order polynomial in time:

$$\mathbf{a}(\tau) = \alpha\tau^3 + \beta\tau^2 + \gamma\tau + \delta, \quad (\text{B.14})$$

The successive time derivatives of $\mathbf{a}(\tau)$ are then:

$$\dot{\mathbf{a}}(\tau) = 3\alpha\tau^2 + 2\beta\tau + \gamma, \quad (\text{B.15})$$

$$\mathbf{a}^{(2)}(\tau) = 6\alpha\tau + 2\beta, \quad (\text{B.16})$$

$$\mathbf{a}^{(3)}(\tau) = 6\alpha. \quad (\text{B.17})$$

The coefficients α , β , γ , and δ are determined by using the known values of acceleration and jerk at times t_i and t . This provides a system of four equations:

$$\mathbf{a}(t_i) = \mathbf{a}_{0,i} = \alpha t_i^3 + \beta t_i^2 + \gamma t_i + \delta , \quad (\text{B.18})$$

$$\mathbf{a}(t) = \mathbf{a}_{1,i} = \alpha t^3 + \beta t^2 + \gamma t + \delta , \quad (\text{B.19})$$

$$\dot{\mathbf{a}}(t_i) = \dot{\mathbf{a}}_{0,i} = 3\alpha t_i^2 + 2\beta t_i + \gamma , \quad (\text{B.20})$$

$$\dot{\mathbf{a}}(t) = \dot{\mathbf{a}}_{1,i} = 3\alpha t^2 + 2\beta t + \gamma , \quad (\text{B.21})$$

where $t = t_i + \Delta t_i$ is the current time. As the system of equations is linearly independent, the four coefficients can be determined. However, determining α and β is enough to get the second and third time derivative of acceleration. From (B.21):

$$\gamma = \dot{\mathbf{a}}_{1,i} - 3\alpha t^2 - 2\beta t . \quad (\text{B.22})$$

Subtracting Eq. (B.20) from (B.21):

$$\beta = \frac{(\dot{\mathbf{a}}_{1,i} - \dot{\mathbf{a}}_{0,i}) - \alpha 3(t^2 - t_i^2)}{2(t - t_i)} . \quad (\text{B.23})$$

Subtracting Eq. (B.18) from (B.19), and using results from (B.22) and (B.23) gives an expression for α :

$$\alpha = \frac{(\dot{\mathbf{a}}_{1,i} + \dot{\mathbf{a}}_{0,i})(t - t_i) - 2(\mathbf{a}_{1,i} - \mathbf{a}_{0,i})}{(t - t_i)^3} . \quad (\text{B.24})$$

Replacing this result in Eq. (B.23):

$$\beta = \frac{-\dot{\mathbf{a}}_{1,i}(t + 2t_i) - \dot{\mathbf{a}}_{0,i}(2t + t_i)}{(t - t_i)^2} + \frac{3(\mathbf{a}_{1,i} - \mathbf{a}_{0,i})(t + t_i)}{(t - t_i)^3} . \quad (\text{B.25})$$

The second and third time derivatives of the acceleration at time t_i can now be evaluated. For the Eq. (B.17), and using $t - t_i = \Delta t_i$:

$$\mathbf{a}_{0,i}^{(3)} = \mathbf{a}^{(3)}(t_i) = 6\alpha , \quad (\text{B.26})$$

$$\mathbf{a}_{0,i}^{(3)} = 6 \frac{(\dot{\mathbf{a}}_{1,i} + \dot{\mathbf{a}}_{0,i})\Delta t_i - 2(\mathbf{a}_{1,i} - \mathbf{a}_{0,i})}{\Delta t_i^3} , \quad (\text{B.27})$$

$$\mathbf{a}_{0,i}^{(3)} = \frac{12(\mathbf{a}_{0,i} - \mathbf{a}_{1,i}) + 6(\dot{\mathbf{a}}_{0,i} + \dot{\mathbf{a}}_{1,i})\Delta t_i}{\Delta t_i^3} . \quad (\text{B.28})$$

Evaluating the second time derivative in t_i , and replacing the obtained for α and β :

$$\mathbf{a}_{0,i}^{(2)} = \mathbf{a}^{(2)}(t_i) = 6\alpha t_i + \beta , \quad (\text{B.29})$$

$$\mathbf{a}_{0,i}^{(2)} = \frac{-6(\mathbf{a}_{0,i} - \mathbf{a}_{1,i}) - \Delta t_i(4\dot{\mathbf{a}}_{0,i} + 2\dot{\mathbf{a}}_{1,i})}{\Delta t_i^2} . \quad (\text{B.30})$$

B.5. Step 5: Add corrections

The next step is to correct the position and velocity of particle i at time t . Using a higher-order Taylor expansion around t_i :

$$\mathbf{x}_i(t_i + \Delta t_i) = \mathbf{x}_i(t_i) + \dot{\mathbf{x}}_i(t_i)\Delta t_i + \frac{\mathbf{x}_i^{(2)}(t_i)\Delta t_i^2}{2!} + \dots + \frac{\mathbf{x}_i^{(5)}(t_i)\Delta t_i^5}{5!} , \quad (\text{B.31})$$

$$= \mathbf{x}_i + \mathbf{v}_i\Delta t_i + \frac{\mathbf{a}_{0,i}\Delta t_i^2}{2} + \frac{\dot{\mathbf{a}}_{0,i}\Delta t_i^3}{6} + \frac{\mathbf{a}_{0,i}^{(2)}\Delta t_i^4}{24} + \frac{\mathbf{a}_{0,i}^{(3)}\Delta t_i^5}{120} . \quad (\text{B.32})$$

The first four terms of this expansion correspond to the predicted position $\mathbf{x}_{p,i}$ from Eq. (B.2). Therefore, the corrected position can be written as:

$$\mathbf{x}_i(t_i + \Delta t_i) = \mathbf{x}_{p,i} + \frac{\mathbf{a}_{0,i}^{(2)}\Delta t_i^4}{24} + \frac{\mathbf{a}_{0,i}^{(3)}\Delta t_i^5}{120} . \quad (\text{B.33})$$

In a similar way, the corrected velocity is:

$$\mathbf{v}_i(t_i + \Delta t_i) = \mathbf{v}_{p,i} + \frac{\mathbf{a}_{0,i}^{(2)}\Delta t_i^3}{6} + \frac{\mathbf{a}_{0,i}^{(3)}\Delta t_i^4}{24} . \quad (\text{B.34})$$

The advantage of this scheme is that achieving high integration accuracy only requires the explicit calculation of the acceleration and jerk for all particles, and the predicted acceleration and jerk for the particle being advanced. This results in an integration scheme that is fourth-order accurate in time ($\mathcal{O}(a^{(3)})$).

B.6. Step 6: Time step update

After updating the position and velocity of particle i , a new time step Δt_i must be calculated. This step is crucial for controlling the integration error. A proven and stable criterion is given by the standard formula (Aarseth, 1985):

$$\Delta t_i = \sqrt{\eta \frac{|\mathbf{a}_{1,i}| |\mathbf{a}_{1,i}^{(2)}| + |\dot{\mathbf{a}}_{1,i}|^2}{|\dot{\mathbf{a}}_{1,i}| |\mathbf{a}_{1,i}^{(3)}| + |\mathbf{a}_{1,i}^{(2)}|^2}} . \quad (\text{B.35})$$

The values for $\mathbf{a}_{1,i}$ and $\dot{\mathbf{a}}_{1,i}$ are known from the direct calculation in Step 3. The value of $\mathbf{a}^{(3)}(t)$ is constant, since a third-order polynomial interpolation is used. Only $\mathbf{a}_{1,i}^{(2)}$ should be calculated:

$$\mathbf{a}_{1,i}^{(2)} = \mathbf{a}_{0,i}^{(2)} + \Delta t_i \mathbf{a}_{0,i}^{(3)} . \quad (\text{B.36})$$

During the initialization of the algorithm, the higher-order derivatives of the acceleration are not available, so an alternative formula for time step can be used in the startup:

$$\Delta t = \eta_s \frac{|\mathbf{a}|}{|\dot{\mathbf{a}}|} . \quad (\text{B.37})$$

The suggested value for the startup parameter is $\eta_s \sim 0.01$ (Makino and Aarseth, 1992).

Appendix C. SPH equations

In this section, we derive the hydrodynamic equations for the SPH method. We follow the derivation of Springel and Hernquist (2002). From a particle perspective, the idea of SPH is to use a discrete representation of the Lagrangian (Rasio, 2000):

$$L_{\text{SPH}} = \sum_{i=1}^N m_i \left(\frac{1}{2} \mathbf{v}_i^2 - u_i \right) . \quad (\text{C.1})$$

The sum is over all the particles (or fluid elements), described by a mass m_i , velocity $\mathbf{v}_i = d\mathbf{r}_i/dt$ and position $\mathbf{r}_i$. For Equation C.1 to be the Lagrangian of a fluid, it is necessary to have an adequate prescription to compute the density ρ_i of any particle. In SPH, the density is calculated as

$$\rho_i = \rho(\mathbf{r}_i) = \sum_j m_j W(|\mathbf{r}_i - \mathbf{r}_j|; h_i), \quad (\text{C.2})$$

where W is the smoothing kernel, and h_i is the smoothing length parameter. In particular, the kernel used in Fi is the cubic spline kernel (Monaghan, 1992; Pelulessy, 2005):

$$W(r, h) = \frac{1}{\pi h^3} \begin{cases} 1 - \frac{3}{2} \left(\frac{r}{h}\right)^2 + \frac{3}{4} \left(\frac{r}{h}\right)^3, & 0 \leq \frac{r}{h} \leq 1, \\ \frac{1}{4} \left(2 - \frac{r}{h}\right)^3, & 1 \leq \frac{r}{h} \leq 2, \\ 0, & \text{otherwise.} \end{cases} \quad (\text{C.3})$$

In some derivations, h_i is taken as a constant. However, since different particles require different resolutions, modern codes use a variable h_i . Requiring that a smoothing volume contains a fixed mass leads to $M_{\text{SPH}} = (4\pi/3)h_i^3\rho_i$, where $M_{\text{SPH}} = \bar{m}N_{\text{SPH}}$, $\bar{m}$ the mean mass among SPH particles and N_{SPH} is a fixed number of neighboring particles. The condition over h_i gives the restriction:

$$\phi_i = \frac{4\pi}{3}h_i^3(\rho_i + \rho_0) - M_{\text{SPH}} = 0. \quad (\text{C.4})$$

Here, a fixed density ρ_0 is introduced to prevent h_i from diverging in low-density regions. The constraints can be introduced into the Lagrangian via Lagrange multipliers λ_i . Then the equations of motion yield

$$\frac{d}{dt} \frac{\partial L_{\text{SPH}}}{\partial \dot{q}_i} - \frac{\partial L_{\text{SPH}}}{\partial q_i} = \sum_{j=1}^N \lambda_j \frac{\partial \phi_j}{\partial q_i}, \quad (\text{C.5})$$

where q_i is replaced by the coordinate, and $\dot{q}_i = dq_i/dt$. The advantage of this approach is that by treating h_i as independent variables in the Lagrangian, the ∇h_i terms are consistently included in the equations of motion.

C.1. Solving λ_i

If we consider a polytropic equation of state, the pressure is given by

$$P = A\rho(\mathbf{r})^\gamma, \quad (\text{C.6})$$

where γ is the polytropic index, and A is the entropic function (see Horedt (2004) for a complete derivation). In the SPH formalism, the pressure for a particle is

$$P_i = A_i \rho_i^\gamma. \quad (\text{C.7})$$

This is the entropic equation of state. Since we are interested in energy conservation, we express A_i in terms of the specific internal energy $u_i = A_i \rho_i^{\gamma-1} / (\gamma - 1)$. Then, the pressure is

$$P_i = u_i \rho_i (\gamma - 1). \quad (\text{C.8})$$

First, we solve the Lagrangian equations (Equation (C.5)) for the variable h_i . As there is no explicit dependency on $\dot{h}_i$, $\partial L_{\text{SPH}} / \partial \dot{h}_i = 0$. Then, for $q_i = h_i$ Equation (C.5) becomes:

$$-\frac{\partial}{\partial h_i} \sum_{j=1}^N m_j \left(\frac{1}{2} |\mathbf{r}_j|^2 - u_j \right) = \sum_{k=1}^N \lambda_k \frac{\partial \phi_k}{\partial h_i}, \quad (\text{C.9})$$

$$-\sum_{j=1}^N m_j \left(\mathbf{r}_j \frac{\partial \mathbf{r}_j}{\partial h_i} - \frac{\partial u_j}{\partial h_i} \right) = \sum_{k=1}^N \lambda_k \frac{\partial}{\partial h_i} \left(\frac{4\pi}{3} h_k^3 (\rho_k + \rho_0) - M_{\text{SPH}} \right), \quad (\text{C.10})$$

$$\sum_{j=1}^N m_j \frac{\partial u_j}{\partial \rho_j} \frac{\partial \rho_j}{\partial h_j} \frac{\partial h_j}{\partial h_i} = \sum_{k=1}^N \lambda_k \frac{4\pi}{3} \left(3h_k^2 (\rho_k + \rho_0) + h_k^3 \frac{\partial \rho_k}{\partial h_k} \right) \frac{\partial h_k}{\partial h_i}. \quad (\text{C.11})$$

Now, we use that $\partial u / \partial \rho = P / \rho^2$, and applying $\partial h_l / \partial h_m = \delta_{lm}$ in the summations:

$$\sum_{j=1}^N m_j \frac{P_j}{\rho_j^2} \frac{\partial \rho_j}{\partial h_j} \delta_{ji} = \sum_{k=1}^N \lambda_k \frac{4\pi}{3} \left(3h_k^2 (\rho_k + \rho_0) + h_k^3 \frac{\partial \rho_k}{\partial h_k} \right) \delta_{ki}, \quad (\text{C.12})$$

$$m_i \frac{P_i}{\rho_i^2} \frac{\partial \rho_i}{\partial h_i} = \lambda_i \frac{4\pi}{3} \left(3h_i^2 (\rho_i + \rho_0) + h_i^3 \frac{\partial \rho_i}{\partial h_i} \right), \quad (\text{C.13})$$

$$= \lambda_i \frac{4\pi h_i^3}{3} \frac{\partial \rho_i}{\partial h_i} \left(1 + \frac{3(\rho_i + \rho_0)}{h_i} \left[\frac{\partial \rho_i}{\partial h_i} \right]^{-1} \right). \quad (\text{C.14})$$

The Lagrange multipliers are then given by

$$\lambda_i = \frac{3}{4\pi} \frac{m_i}{h_i^3} \frac{P_i}{\rho_i^2} \left(1 + \frac{3(\rho_i + \rho_0)}{h_i} \left[\frac{\partial \rho_i}{\partial h_i} \right]^{-1} \right)^{-1}. \quad (\text{C.15})$$

C.2. Equation of motion for SPH particles

We now derive the equations of motion by applying Equation (C.5) to the spatial coordinates $q_i = \mathbf{r}_i$:

$$\frac{d}{dt} \frac{\partial L_{\text{SPH}}}{\partial \dot{\mathbf{r}}_i} - \frac{\partial L_{\text{SPH}}}{\partial \mathbf{r}_i} = \sum_{j=1}^N \lambda_j \frac{\partial \phi_j}{\partial \mathbf{r}_i}. \quad (\text{C.16})$$

The first term on the left can be obtained directly:

$$\frac{d}{dt} \frac{\partial L_{\text{SPH}}}{\partial \dot{\mathbf{r}}_i} = \frac{d}{dt} \sum_{j=1}^N m_j \mathbf{v}_j \frac{\partial \mathbf{v}_j}{\partial \mathbf{v}_i}, \quad (\text{C.17})$$

$$= m_i \frac{d\mathbf{v}_i}{dt}. \quad (\text{C.18})$$

The second term:

$$\frac{\partial L_{\text{SPH}}}{\partial \mathbf{r}_i} = \sum_{j=1}^N m_j \left(\dot{\mathbf{r}}_j \frac{\partial \dot{\mathbf{r}}_j}{\partial \mathbf{r}_j} - \frac{\partial u_j}{\partial \mathbf{r}_i} \right), \quad (\text{C.19})$$

$$= - \sum_{j=1}^N m_j \frac{\partial u_j}{\partial \rho_j} \nabla_i \rho_j, \quad (\text{C.20})$$

$$= \sum_{j=1}^N m_j \frac{P_j}{\rho_j^2} \nabla_i \rho_j, \quad (\text{C.21})$$

where $\nabla_i = \partial / \partial \mathbf{r}_i$. For the right side of Equation (C.16):

$$\sum_{j=1}^N \lambda_j \frac{\partial \phi_j}{\partial \mathbf{r}_i} = \sum_{j=1}^N \lambda_j \frac{\partial}{\partial \mathbf{r}_i} \left(\frac{4\pi}{3} h_j^3 (\rho_j + \rho_0) - M_{\text{SPH}} \right), \quad (\text{C.22})$$

$$= \sum_{j=1}^N \lambda_j \frac{4\pi}{3} h_j^3 \nabla_i \rho_j. \quad (\text{C.23})$$

Substituting all the terms back into Equation (C.16),

$$m_i \frac{d\mathbf{v}_i}{dt} - \sum_{j=1}^N m_j \frac{P_j}{\rho_j^2} \nabla_i \rho_j = \sum_{j=1}^N \lambda_j \frac{4\pi}{3} h_j^3 \nabla_i \rho_j, \quad (\text{C.24})$$

$$m_i \frac{d\mathbf{v}_i}{dt} = \sum_{j=1}^N m_j \frac{P_j}{\rho_j^2} \left[1 + \lambda_j \frac{4\pi}{3} \frac{h_j^3}{m_j} \frac{\rho_j^2}{P_j} \right] \nabla_i \rho_j. \quad (\text{C.25})$$

Expanding the term with λ_i ,

$$\lambda_j \frac{4\pi}{3} \frac{h_j^3}{m_j} \frac{\rho_j^2}{P_j} = \left(1 + \frac{3(\rho_j + \rho_0)}{h_j} \left[\frac{\partial \rho_j}{\partial h_j} \right]^{-1} \right)^{-1}, \quad (\text{C.26})$$

$$= \left(1 + \left[\frac{h_j}{3(\rho_j + \rho_0)} \frac{\partial \rho_j}{\partial h_j} \right]^{-1} \right)^{-1}. \quad (\text{C.27})$$

Defining

$$f_j = \left(1 + \frac{h_j}{3(\rho_j + \rho_0)} \frac{\partial \rho_j}{\partial h_j} \right)^{-1}, \quad (\text{C.28})$$

then the term involving λ_j simplifies to a compact expression, as shown below:

$$\lambda_j \frac{4\pi}{3} \frac{h_j^3}{m_j} \frac{\rho_j^2}{P_j} = \left(1 + (f_j^{-1} - 1) \right)^{-1}, \quad (\text{C.29})$$

$$= \left(\frac{f_j^{-1}}{f_j^{-1} - 1} \right)^{-1} = 1 - f_j. \quad (\text{C.30})$$

Replacing back to Equation (C.25),

$$m_i \frac{d\mathbf{v}_i}{dt} = - \sum_{j=1}^N m_j \frac{P_j}{\rho_j^2} f_j \nabla_i \rho_j. \quad (\text{C.31})$$

Using the notation $W_{ij}(h) = W(|\mathbf{r}_i - \mathbf{r}_j|, h)$, then we can write (Springel and Hernquist, 2002)

$$\nabla_i \rho_j = m_i \nabla_i W_{ij}(h_j) + \delta_{ij} \sum_{k=1}^N m_k \nabla_i W_{ki}(h_i) . \quad (\text{C.32})$$

Back into Equation (C.30),

$$m_i \frac{d\mathbf{v}_i}{dt} = - \sum_{j=1}^N m_j \frac{P_j}{\rho_j^2} f_j \left(m_i \nabla_i W_{ij}(h_j) + \delta_{ij} \sum_{k=1}^N m_k \nabla_i W_{ki}(h_i) \right) , \quad (\text{C.33})$$

$$= - \sum_{j=1}^N m_j m_i \frac{P_j}{\rho_j^2} f_j \nabla_i W_{ij}(h_j) - m_i \frac{P_i}{\rho_i^2} f_i \sum_{k=1}^N m_k \nabla_i W_{ki}(h_i) , \quad (\text{C.34})$$

$$\frac{d\mathbf{v}_i}{dt} = - \sum_{j=1}^N m_j \frac{P_j}{\rho_j^2} f_j \nabla_i W_{ij}(h_j) - \frac{P_i}{\rho_i^2} f_i \sum_{k=1}^N m_k \nabla_i W_{ki}(h_i) . \quad (\text{C.35})$$

Renaming index k to j in the second summation, and using the symmetry of the kernel, which implies $\nabla_i W_{ki}(h_i) = \nabla_i W_{ik}(h_i)$, we obtain the final form of the equation of motion:

$$\frac{d\mathbf{v}_i}{dt} = - \sum_{j=1}^N m_j \left[\frac{P_j}{\rho_j^2} f_j \nabla_i W_{ij}(h_j) + \frac{P_i}{\rho_i^2} f_i \nabla_i W_{ij}(h_i) \right] . \quad (\text{C.36})$$

C.3. Equation of energy for SPH particles

We now derive the evolution equation for the specific internal energy. The total energy of the SPH particles is given by:

$$E = \sum_{i=1}^N m_i u_i + \sum_{j=1}^N m_j \frac{\mathbf{v}_j^2}{2} , \quad (\text{C.37})$$

and under the condition of energy conservation:

$$\frac{dE}{dt} = 0 , \quad (\text{C.38})$$

$$= \sum_{i=1}^N m_i \frac{du_i}{dt} + \sum_{j=1}^N m_j \mathbf{v}_j \frac{d\mathbf{v}_j}{dt} . \quad (\text{C.39})$$

Now, we use Equation (C.36) on the right side:

$$\sum_{i=1}^N m_i \frac{du_i}{dt} = - \sum_{j=1}^N m_j \mathbf{v}_j \left(- \sum_{k=1}^N m_k \left[\frac{P_k}{\rho_k^2} f_k \nabla_j W_{jk}(h_k) + \frac{P_j}{\rho_j^2} f_j \nabla_j W_{jk}(h_j) \right] \right) , \quad (\text{C.40})$$

$$= \sum_{j,k} m_j m_k \mathbf{v}_j \frac{P_k}{\rho_k^2} f_k \nabla_j W_{jk}(h_k) + \sum_{j,k} m_j m_k \mathbf{v}_j \frac{P_j}{\rho_j^2} f_j \nabla_j W_{jk}(h_j) . \quad (\text{C.41})$$

Since j and k are dummy indices, we can relabel them to symmetrize the expression. In the first sum we set $j \rightarrow i, k \rightarrow j$, and in the second sum we set $k \rightarrow i$, yielding

$$\sum_{i=1}^N m_i \frac{du_i}{dt} = \sum_{i,j} m_i m_j \mathbf{v}_i \frac{P_j}{\rho_j^2} f_j \nabla_i W_{ij}(h_j) + \sum_{i,j} m_j m_i \mathbf{v}_j \frac{P_j}{\rho_j^2} f_j \nabla_j W_{ji}(h_j) , \quad (\text{C.42})$$

$$= \sum_{i,j} m_i m_j \mathbf{v}_i \frac{P_j}{\rho_j^2} f_j \nabla_i W_{ij}(h_j) + \sum_{i,j} m_j m_i \mathbf{v}_j \frac{P_j}{\rho_j^2} f_j (-\nabla_i W_{ij}(h_j)) , \quad (\text{C.43})$$

$$= \sum_{i,j} m_i m_j f_j \frac{P_j}{\rho_j^2} \mathbf{v}_{ij} \cdot \nabla_i W_{ij}(h_j) , \quad (\text{C.44})$$

where we have used $\nabla_j W_{ji}(h_j) = -\nabla_i W_{ij}(h_j)$ and $\mathbf{v}_{ij} = \mathbf{v}_i - \mathbf{v}_j$. Since i, j are also dummy indices, we can exchange them, having another solution for the equation of energy:

$$\sum_{i=1}^N m_i \frac{du_i}{dt} = \sum_{i,j} m_i m_j f_j \frac{P_j}{\rho_j^2} \mathbf{v}_{ij} \cdot \nabla_i W_{ij}(h_j) , \quad (\text{C.45})$$

$$= \sum_{i,j} m_i m_j f_i \frac{P_i}{\rho_i^2} \mathbf{v}_{ji} \cdot \nabla_j W_{ji}(h_i) , \quad (\text{C.46})$$

$$= \sum_{i,j} m_i m_j f_i \frac{P_i}{\rho_i^2} \mathbf{v}_{ij} \cdot \nabla_i W_{ij}(h_i) . \quad (\text{C.47})$$

Taking the average of Equations (C.44) and (C.47) yields the symmetrized form:

$$\sum_{i=1}^N m_i \frac{du_i}{dt} = \sum_{i,j} m_i m_j \frac{1}{2} \left[f_j \frac{P_j}{\rho_j^2} \mathbf{v}_{ij} \cdot \nabla_i W_{ij}(h_j) + f_i \frac{P_i}{\rho_i^2} \mathbf{v}_{ij} \cdot \nabla_i W_{ij}(h_i) \right] , \quad (\text{C.48})$$

and the evolution equation for the internal energy is

$$\frac{du_i}{dt} = \frac{1}{2} \sum_{j=1}^N m_j \left[f_j \frac{P_j}{\rho_j^2} \mathbf{v}_{ij} \cdot \nabla_i W_{ij}(h_j) + f_i \frac{P_i}{\rho_i^2} \mathbf{v}_{ij} \cdot \nabla_i W_{ij}(h_i) \right] . \quad (\text{C.49})$$