

Rigidity Properties of Higher Rank Abelian Actions

Boris Kalinin

Dynamical Systems and Group Actions

We consider a smooth action α of a group G on a compact manifold M , $\alpha : G \rightarrow \text{Diff}(M)$.

Dynamical Systems and Group Actions

We consider a smooth action α of a group G on a compact manifold M , $\alpha : G \rightarrow \text{Diff}(M)$.

Types of the acting group G :

- 1 \mathbb{Z} or \mathbb{R} . Classical dynamical system - diffeomorphism or flow.
- 2 \mathbb{Z}^k or \mathbb{R}^k , $k \geq 2$. Higher rank abelian action.

Dynamical Systems and Group Actions

We consider a smooth action α of a group G on a compact manifold M , $\alpha : G \rightarrow \text{Diff}(M)$.

Types of the acting group G :

- 1 \mathbb{Z} or \mathbb{R} . Classical dynamical system - diffeomorphism or flow.
- 2 \mathbb{Z}^k or \mathbb{R}^k , $k \geq 2$. Higher rank abelian action.

Equivalence: a conjugacy between α and α' is a homeomorphism or diffeomorphism $h : M \rightarrow M'$ such that $h \circ \alpha(g) = \alpha'(g) \circ h$ for all $g \in G$.

$$\begin{array}{ccc} M & \xrightarrow{\alpha(g)} & M \\ h \downarrow & & \downarrow h \\ M' & \xrightarrow{\alpha'(g)} & M' \end{array}$$

Hyperbolic \mathbb{Z} actions, algebraic case

Automorphisms of tori and nilmanifolds.

$A \in SL(n, \mathbb{Z})$ gives an automorphism of $\mathbb{T}^n = \mathbb{R}^n / \mathbb{Z}^n$.

A is **hyperbolic** or **Anosov** if for any eigenvalue $|\lambda| \neq 1$.

$$\text{e.g. } A = \begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix} : \mathbb{T}^2 \rightarrow \mathbb{T}^2$$

Hyperbolic \mathbb{Z} actions, algebraic case

Automorphisms of tori and nilmanifolds.

$A \in SL(n, \mathbb{Z})$ gives an automorphism of $\mathbb{T}^n = \mathbb{R}^n / \mathbb{Z}^n$.

A is **hyperbolic** or **Anosov** if for any eigenvalue $|\lambda| \neq 1$.

$$\text{e.g. } A = \begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix} : \mathbb{T}^2 \rightarrow \mathbb{T}^2$$

N is a simply connected nilpotent Lie group;

Γ is a cocompact lattice in N ;

$A : N \rightarrow N$ is an automorphism such that $A(\Gamma) = \Gamma$.

Automorphisms of tori and nilmanifolds.

$A \in SL(n, \mathbb{Z})$ gives an automorphism of $\mathbb{T}^n = \mathbb{R}^n / \mathbb{Z}^n$.

A is **hyperbolic** or **Anosov** if for any eigenvalue $|\lambda| \neq 1$.

$$\text{e.g. } A = \begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix} : \mathbb{T}^2 \rightarrow \mathbb{T}^2$$

N is a simply connected nilpotent Lie group;

Γ is a cocompact lattice in N ;

$A : N \rightarrow N$ is an automorphism such that $A(\Gamma) = \Gamma$.

Then A projects to an automorphism of the nilmanifold N/Γ .

A is **Anosov** if $D_e A : T_e N \rightarrow T_e N$ is a hyperbolic automorphism.

Hyperbolic \mathbb{Z} actions, smooth case

A diffeomorphism f of a compact manifold M is called **Anosov** if the tangent bundle has a continuous Df -invariant splitting $TM = E^s \oplus E^u$ such that for all $n > 0$

$$\begin{aligned}\|Df^n(v)\| &\leq Ce^{-\lambda n}\|v\| \quad \text{for all } v \in E^s, \\ \|Df^{-n}(v)\| &\leq Ce^{-\lambda n}\|v\| \quad \text{for all } v \in E^u.\end{aligned}$$

e.g. any C^1 -small perturbation of an Anosov automorphism.

Hyperbolic \mathbb{Z} actions, smooth case

A diffeomorphism f of a compact manifold M is called **Anosov** if the tangent bundle has a continuous Df -invariant splitting $TM = E^s \oplus E^u$ such that for all $n > 0$

$$\begin{aligned}\|Df^n(v)\| &\leq Ce^{-\lambda n}\|v\| \quad \text{for all } v \in E^s, \\ \|Df^{-n}(v)\| &\leq Ce^{-\lambda n}\|v\| \quad \text{for all } v \in E^u.\end{aligned}$$

e.g. any C^1 -small perturbation of an Anosov automorphism.

E^s and E^u are called the **stable** and **unstable** distributions. They are tangent to the stable and unstable foliations W^s and W^u .

The leaves of W^s are C^∞ injectively immersed Euclidean spaces, but E^s usually varies only Hölder continuously transversally to W^s . Similarly for W^u and E^u .

Topological rigidity problem for hyperbolic systems

Structural stability (Local topological rigidity):

Any C^1 -small perturbation g of an Anosov diffeomorphism f is Anosov and is *topologically* conjugate to f , i.e. $g \circ h = h \circ f$.

The conjugacy h is a bi-Hölder homeomorphism, but rarely smooth.

Topological rigidity problem for hyperbolic systems

Structural stability (Local topological rigidity):

Any C^1 -small perturbation g of an Anosov diffeomorphism f is Anosov and is *topologically* conjugate to f , i.e. $g \circ h = h \circ f$.

The conjugacy h is a bi-Hölder homeomorphism, but rarely smooth.

Classification on Nilmanifolds (Global topological rigidity):

Any Anosov diffeomorphism of a nilmanifold is topologically conjugate to an Anosov automorphism (Franks, Manning).

Topological rigidity problem for hyperbolic systems

Structural stability (Local topological rigidity):

Any C^1 -small perturbation g of an Anosov diffeomorphism f is Anosov and is *topologically* conjugate to f , i.e. $g \circ h = h \circ f$.

The conjugacy h is a bi-Hölder homeomorphism, but rarely smooth.

Classification on Nilmanifolds (Global topological rigidity):

Any Anosov diffeomorphism of a nilmanifold is topologically conjugate to an Anosov automorphism (Franks, Manning).

Topological classification – open problem:

Conjecture: Any Anosov diffeomorphism is topologically conjugate to a finite factor of an Anosov automorphism of a nilmanifold.

Yes, if $\dim E^s = 1$ or $\dim E^u = 1$ (Franks, Newhouse).

Topological rigidity problem for hyperbolic systems

Structural stability (Local topological rigidity):

Any C^1 -small perturbation g of an Anosov diffeomorphism f is Anosov and is *topologically* conjugate to f , i.e. $g \circ h = h \circ f$.

The conjugacy h is a bi-Hölder homeomorphism, but rarely smooth.

Classification on Nilmanifolds (Global topological rigidity):

Any Anosov diffeomorphism of a nilmanifold is topologically conjugate to an Anosov automorphism (Franks, Manning).

Topological classification – open problem:

Conjecture: Any Anosov diffeomorphism is topologically conjugate to a finite factor of an Anosov automorphism of a nilmanifold.

Yes, if $\dim E^s = 1$ or $\dim E^u = 1$ (Franks, Newhouse).

Flows: No classification conjecture (“anomalous” examples).

Higher Rank Abelian Actions

Algebraic actions of \mathbb{Z}^k :

Generated by commuting automorphisms of tori or nilmanifolds.

Higher Rank Abelian Actions

Algebraic actions of \mathbb{Z}^k :

Generated by commuting automorphisms of tori or nilmanifolds.

Consider an irreducible hyperbolic matrix $A \in SL(n, \mathbb{Z})$.

If all eigenvalues of A are real then the centralizer of A in $SL(n, \mathbb{Z})$ contains a subgroup isomorphic to \mathbb{Z}^{n-1} (Cartan action).

Higher Rank Abelian Actions

Algebraic actions of \mathbb{Z}^k :

Generated by commuting automorphisms of tori or nilmanifolds.

Consider an irreducible hyperbolic matrix $A \in SL(n, \mathbb{Z})$.

If all eigenvalues of A are real then the centralizer of A in $SL(n, \mathbb{Z})$ contains a subgroup isomorphic to \mathbb{Z}^{n-1} (Cartan action).

More generally, one can consider \mathbb{Z}^k -actions by commuting automorphisms of a nilmanifold $M = N/\Gamma$.

Higher Rank Abelian Actions

Algebraic actions of \mathbb{Z}^k :

Generated by commuting automorphisms of tori or nilmanifolds.

Consider an irreducible hyperbolic matrix $A \in SL(n, \mathbb{Z})$.

If all eigenvalues of A are real then the centralizer of A in $SL(n, \mathbb{Z})$ contains a subgroup isomorphic to \mathbb{Z}^{n-1} (Cartan action).

More generally, one can consider \mathbb{Z}^k -actions by commuting automorphisms of a nilmanifold $M = N/\Gamma$.

Smooth actions:

$\alpha : \mathbb{Z}^k \rightarrow \text{Diff}(M)$ is called a *hyperbolic action* if for some $b \in \mathbb{Z}^k$ $\alpha(b)$ is an Anosov diffeomorphism.

Higher Rank Abelian Actions

Algebraic actions of \mathbb{Z}^k :

Generated by commuting automorphisms of tori or nilmanifolds.

Consider an irreducible hyperbolic matrix $A \in SL(n, \mathbb{Z})$.

If all eigenvalues of A are real then the centralizer of A in $SL(n, \mathbb{Z})$ contains a subgroup isomorphic to \mathbb{Z}^{n-1} (Cartan action).

More generally, one can consider \mathbb{Z}^k -actions by commuting automorphisms of a nilmanifold $M = N/\Gamma$.

Smooth actions:

$\alpha : \mathbb{Z}^k \rightarrow \text{Diff}(M)$ is called a *hyperbolic action* if for some $b \in \mathbb{Z}^k$ $\alpha(b)$ is an Anosov diffeomorphism.

$\alpha : \mathbb{R}^k \rightarrow \text{Diff}(M)$ is called a *hyperbolic action* if for some $b \in \mathbb{Z}^k$ $\alpha(b)$ is normally hyperbolic to the orbit: $TM = E_b^s \oplus E^o \oplus E_b^u$.

Local smooth rigidity of algebraic actions:

Any standard higher rank hyperbolic algebraic action is C^∞ conjugate to any C^1 -small perturbation.
(Katok and Spatzier, for most types).

Rigidity properties of higher rank actions

Local smooth rigidity of algebraic actions:

Any standard higher rank hyperbolic algebraic action is C^∞ conjugate to any C^1 -small perturbation.
(Katok and Spatzier, for most types).

Conjecture: Irreducible Anosov actions of \mathbb{Z}^k and \mathbb{R}^k , $k \geq 2$, on any compact manifold are *smoothly* conjugate to algebraic actions.

Rigidity properties of higher rank actions

Local smooth rigidity of algebraic actions:

Any standard higher rank hyperbolic algebraic action is C^∞ conjugate to any C^1 -small perturbation.
(Katok and Spatzier, for most types).

Conjecture: Irreducible Anosov actions of \mathbb{Z}^k and \mathbb{R}^k , $k \geq 2$, on any compact manifold are *smoothly* conjugate to algebraic actions.

Global smooth rigidity on tori and nilmanifolds:

Under mild dynamical assumptions.

Rigidity properties of higher rank actions

Local smooth rigidity of algebraic actions:

Any standard higher rank hyperbolic algebraic action is C^∞ conjugate to any C^1 -small perturbation.
(Katok and Spatzier, for most types).

Conjecture: Irreducible Anosov actions of \mathbb{Z}^k and \mathbb{R}^k , $k \geq 2$, on any compact manifold are *smoothly* conjugate to algebraic actions.

Global smooth rigidity on tori and nilmanifolds:

Under mild dynamical assumptions.

Classification of Anosov actions on arbitrary manifolds:

Under strong dynamical assumptions.

Rigidity properties of higher rank actions

Local smooth rigidity of algebraic actions:

Any standard higher rank hyperbolic algebraic action is C^∞ conjugate to any C^1 -small perturbation.
(Katok and Spatzier, for most types).

Conjecture: Irreducible Anosov actions of \mathbb{Z}^k and \mathbb{R}^k , $k \geq 2$, on any compact manifold are *smoothly* conjugate to algebraic actions.

Global smooth rigidity on tori and nilmanifolds:

Under mild dynamical assumptions.

Classification of Anosov actions on arbitrary manifolds:

Under strong dynamical assumptions.

Measure rigidity:

Scarcity of invariant measures.

Algebraic examples: non-invertible case

Expanding endomorphisms of the circle $\mathbb{T} = \mathbb{R}/\mathbb{Z}$:

$$E_m : \mathbb{T} \rightarrow \mathbb{T}, \quad E_m x = mx \pmod{1}, \quad m \geq 2.$$

Abundance of invariant measures for a single map E_m .

Algebraic examples: non-invertible case

Expanding endomorphisms of the circle $\mathbb{T} = \mathbb{R}/\mathbb{Z}$:

$$E_m : \mathbb{T} \rightarrow \mathbb{T}, \quad E_m x = mx \pmod{1}, \quad m \geq 2.$$

Abundance of invariant measures for a single map E_m .

Two (commuting) endomorphisms of the circle

E_2 and E_3 generate \mathbb{Z}_+^2 -action on \mathbb{T} .

Furstenberg's Conjecture:

The only ergodic invariant measures for this action are the Lebesgue measure and atomic measures on periodic orbits.

Algebraic examples: non-invertible case

Expanding endomorphisms of the circle $\mathbb{T} = \mathbb{R}/\mathbb{Z}$:

$$E_m : \mathbb{T} \rightarrow \mathbb{T}, \quad E_m x = mx \pmod{1}, \quad m \geq 2.$$

Abundance of invariant measures for a single map E_m .

Two (commuting) endomorphisms of the circle

E_2 and E_3 generate \mathbb{Z}_+^2 -action on \mathbb{T} .

Furstenberg's Conjecture:

The only ergodic invariant measures for this action are the Lebesgue measure and atomic measures on periodic orbits.

Theorem (D. Rudolph) If an invariant measures for this action has positive entropy for some element then it is Lebesgue.

Open problem. Are there nontrivial zero entropy measures?

Conjecture (Margulis, Katok and Spatzier): Scarcity of invariant measures for hyperbolic algebraic actions of \mathbb{Z}^k and \mathbb{R}^k , $k \geq 2$.

Conjecture (Margulis, Katok and Spatzier): Scarcity of invariant measures for hyperbolic algebraic actions of \mathbb{Z}^k and \mathbb{R}^k , $k \geq 2$.

Example: $\mathbb{R}^2 \cong A = \left\{ \begin{pmatrix} e^t & 0 & 0 \\ 0 & e^s & 0 \\ 0 & 0 & e^{-s-t} \end{pmatrix} \mid t, s \in \mathbb{R} \right\} \subset SL(3, \mathbb{R})$

gives \mathbb{R}^2 action on $SL(3, \mathbb{R})/SL(3, \mathbb{Z})$ by left translations.

Algebraic Measure Rigidity

Conjecture (Margulis, Katok and Spatzier): Scarcity of invariant measures for hyperbolic algebraic actions of \mathbb{Z}^k and \mathbb{R}^k , $k \geq 2$.

Example: $\mathbb{R}^2 \cong A = \left\{ \begin{pmatrix} e^t & 0 & 0 \\ 0 & e^s & 0 \\ 0 & 0 & e^{-s-t} \end{pmatrix} \mid t, s \in \mathbb{R} \right\} \subset SL(3, \mathbb{R})$

gives \mathbb{R}^2 action on $SL(3, \mathbb{R})/SL(3, \mathbb{Z})$ by left translations.

Theorem (Einsiedler, Katok, Lindenstrauss)

Haar measure is the only ergodic measure with positive entropy.

Algebraic Measure Rigidity

Conjecture (Margulis, Katok and Spatzier): Scarcity of invariant measures for hyperbolic algebraic actions of \mathbb{Z}^k and \mathbb{R}^k , $k \geq 2$.

Example: $\mathbb{R}^2 \cong A = \left\{ \begin{pmatrix} e^t & 0 & 0 \\ 0 & e^s & 0 \\ 0 & 0 & e^{-s-t} \end{pmatrix} \mid t, s \in \mathbb{R} \right\} \subset SL(3, \mathbb{R})$

gives \mathbb{R}^2 action on $SL(3, \mathbb{R})/SL(3, \mathbb{Z})$ by left translations.

Theorem (Einsiedler, Katok, Lindenstrauss)

Haar measure is the only ergodic measure with positive entropy.

Littlewood Conjecture: $\liminf (n \|na\| \|nb\|) = 0$ for any $a, b \in \mathbb{R}$

Corollary The set of exceptional pairs has Hausdorff dimension 0.

Algebraic Measure Rigidity

Hope for more general \mathbb{R}^k actions by left translations on G/Γ :

Any invariant measure is *algebraic*, i.e. Haar measure on a compact homogeneous space $Hx \subset G/\Gamma$ for a subgroup $H \subset G$.

Algebraic Measure Rigidity

Hope for more general \mathbb{R}^k actions by left translations on G/Γ :
Any invariant measure is *algebraic*, i.e. Haar measure on a compact homogeneous space $Hx \subset G/\Gamma$ for a subgroup $H \subset G$.

Theorem (B.K. and R. Spatzier)

Suppose that $Ad(b)$ is diagonalizable for all $b \in \mathbb{R}^k$ and that one-parameter subgroups act ergodically with respect to μ . Then μ is an extension of a zero entropy measure in a factor with Haar measure in the fibers.

Algebraic Measure Rigidity

Hope for more general \mathbb{R}^k actions by left translations on G/Γ :
Any invariant measure is *algebraic*, i.e. Haar measure on a compact homogeneous space $Hx \subset G/\Gamma$ for a subgroup $H \subset G$.

Theorem (B.K. and R. Spatzier)

Suppose that $\text{Ad}(b)$ is diagonalizable for all $b \in \mathbb{R}^k$ and that one-parameter subgroups act ergodically with respect to μ . Then μ is an extension of a zero entropy measure in a factor with Haar measure in the fibers.

Corollary

Any measure-preserving isomorphism between two such actions is algebraic.

Algebraic Measure Rigidity

Hope for more general \mathbb{R}^k actions by left translations on G/Γ :
Any invariant measure is *algebraic*, i.e. Haar measure on a compact homogeneous space $Hx \subset G/\Gamma$ for a subgroup $H \subset G$.

Theorem (B.K. and R. Spatzier)

Suppose that $\text{Ad}(b)$ is diagonalizable for all $b \in \mathbb{R}^k$ and that one-parameter subgroups act ergodically with respect to μ . Then μ is an extension of a zero entropy measure in a factor with Haar measure in the fibers.

Corollary

Any measure-preserving isomorphism between two such actions is algebraic.

Follows by applying the theorem to the invariant measure on the graph of the measure-preserving isomorphism.

Non-algebraic actions.

Let α_0 be a Cartan action of \mathbb{Z}^k on \mathbb{T}^{k+1} :

full centralizer in $SL(k+1, \mathbb{Z})$ of an irreducible hyperbolic matrix with real eigenvalues.

We say that a \mathbb{Z}^k -action α has *Cartan homotopy data* if its elements are homotopic to the corresponding ones of α_0 .

Non-algebraic actions.

Let α_0 be a Cartan action of \mathbb{Z}^k on \mathbb{T}^{k+1} :

full centralizer in $SL(k+1, \mathbb{Z})$ of an irreducible hyperbolic matrix with real eigenvalues.

We say that a \mathbb{Z}^k -action α has *Cartan homotopy data* if its elements are homotopic to the corresponding ones of α_0 .

Theorem (B.K. and A. Katok)

Any \mathbb{Z}^k -action α by $C^{1+\varepsilon}$ diffeomorphisms of \mathbb{T}^{k+1} , $k \geq 2$, with Cartan homotopy data preserves an ergodic absolutely continuous invariant measure μ .

Such measure μ is unique and (α, μ) is isomorphic to (α_0, λ) as a measure preserving action.

Two commuting hyperbolic automorphisms $A, B : \mathbb{T}^3 \rightarrow \mathbb{T}^3$ generate a \mathbb{Z}^2 action on \mathbb{T}^3 : (m, n) acts by $A^m B^n$.

Lyapunov exponents

Two commuting hyperbolic automorphisms $A, B : \mathbb{T}^3 \rightarrow \mathbb{T}^3$ generate a \mathbb{Z}^2 action on \mathbb{T}^3 : (m, n) acts by $A^m B^n$.

$\mathbb{R}^3 = E_1 \oplus E_2 \oplus E_3$ - splitting into common eigenspaces.

$\chi_i(m, n) = \ln |i^{th} \text{ eigenvalue of } A^m B^n|$, $i = 1, 2, 3$
the Lyapunov exponents of $A^m B^n$.

Lyapunov exponents

Two commuting hyperbolic automorphisms $A, B : \mathbb{T}^3 \rightarrow \mathbb{T}^3$ generate a \mathbb{Z}^2 action on \mathbb{T}^3 : (m, n) acts by $A^m B^n$.

$\mathbb{R}^3 = E_1 \oplus E_2 \oplus E_3$ - splitting into common eigenspaces.

$\chi_i(m, n) = \ln |i^{th} \text{ eigenvalue of } A^m B^n|$, $i = 1, 2, 3$
the Lyapunov exponents of $A^m B^n$.

Linear functionals $\chi_i : \mathbb{Z}^2 \rightarrow \mathbb{R}$ extend to \mathbb{R}^2
and are called the *Lyapunov exponents* of the action.

Lyapunov exponents

Two commuting hyperbolic automorphisms $A, B : \mathbb{T}^3 \rightarrow \mathbb{T}^3$ generate a \mathbb{Z}^2 action on \mathbb{T}^3 : (m, n) acts by $A^m B^n$.

$\mathbb{R}^3 = E_1 \oplus E_2 \oplus E_3$ - splitting into common eigenspaces.

$\chi_i(m, n) = \ln |i^{th} \text{ eigenvalue of } A^m B^n|$, $i = 1, 2, 3$
the Lyapunov exponents of $A^m B^n$.

Linear functionals $\chi_i : \mathbb{Z}^2 \rightarrow \mathbb{R}$ extend to \mathbb{R}^2
and are called the *Lyapunov exponents* of the action.

The kernels $\ker(\chi_i) \subset \mathbb{R}^k$ are called *Lyapunov hyperplanes*.
Weyl chambers are the connected components of $\mathbb{R}^k \setminus \bigcup \ker(\chi)$.

Nonuniform measure rigidity.

For non-linear smooth action Multiplicative Ergodic Theorem gives similar *measurable* splitting and the Lyapunov exponents give *asymptotic* exponential rates of expansion/contraction.

Nonuniform measure rigidity.

For non-linear smooth action Multiplicative Ergodic Theorem gives similar *measurable* splitting and the Lyapunov exponents give *asymptotic* exponential rates of expansion/contraction.

We say that Lyapunov exponents are **in general position** if they are all simple and nonzero, and if the Lyapunov hyperplanes are distinct and in general position (i.e. the dimension of any intersection of them is the minimal possible).

Nonuniform measure rigidity.

For non-linear smooth action Multiplicative Ergodic Theorem gives similar *measurable* splitting and the Lyapunov exponents give *asymptotic* exponential rates of expansion/contraction.

We say that Lyapunov exponents are **in general position** if they are all simple and nonzero, and if the Lyapunov hyperplanes are distinct and in general position (i.e. the dimension of any intersection of them is the minimal possible).

Theorem (B.K., A. Katok, F. Rodriguez Hertz)

Let μ be an ergodic invariant measure for a $C^{1+\varepsilon}$ action α of \mathbb{Z}^k , $k \geq 2$, on a $(k+1)$ -dimensional manifold M . Suppose that the Lyapunov exponents of μ are in general position and that at least one element in \mathbb{Z}^k has positive entropy with respect to μ . Then μ is absolutely continuous.

Smooth rigidity for \mathbb{Z}^k and \mathbb{R}^k actions

Two cases:

- 1 \mathbb{Z}^k actions on tori and nilmanifolds.
- 2 \mathbb{Z}^k actions on an arbitrary manifold or \mathbb{R}^k actions.

Smooth rigidity for \mathbb{Z}^k and \mathbb{R}^k actions

Two cases:

- 1 \mathbb{Z}^k actions on tori and nilmanifolds.
- 2 \mathbb{Z}^k actions on an arbitrary manifold or \mathbb{R}^k actions.

If action α is of type 1, then it is topologically conjugate to the

$$\begin{array}{ccc} M & \xrightarrow{\alpha} & M \\ \text{algebraic action } \alpha_0 : h \downarrow & & \downarrow h \\ M & \xrightarrow{\alpha_0} & M \end{array} \quad \text{Need to show: } h \text{ is smooth.}$$

Smooth rigidity for \mathbb{Z}^k and \mathbb{R}^k actions

Two cases:

- 1 \mathbb{Z}^k actions on tori and nilmanifolds.
- 2 \mathbb{Z}^k actions on an arbitrary manifold or \mathbb{R}^k actions.

If action α is of type 1, then it is topologically conjugate to the

$$\begin{array}{ccc} M & \xrightarrow{\alpha} & M \\ \text{algebraic action } \alpha_0 : h \downarrow & & \downarrow h \\ M & \xrightarrow{\alpha_0} & M \end{array} \quad \text{Need to show: } h \text{ is smooth.}$$

If action α is of type 2, then there is no apparent algebraic model α_0 and the algebraic structure has to be built from scratch.

Smooth rigidity for \mathbb{Z}^k and \mathbb{R}^k actions

Two cases:

- 1 \mathbb{Z}^k actions on tori and nilmanifolds.
- 2 \mathbb{Z}^k actions on an arbitrary manifold or \mathbb{R}^k actions.

If action α is of type 1, then it is topologically conjugate to the

$$\begin{array}{ccc} M & \xrightarrow{\alpha} & M \\ \text{algebraic action } \alpha_0 : h \downarrow & & \downarrow h \\ M & \xrightarrow{\alpha_0} & M \end{array} \quad \text{Need to show: } h \text{ is smooth.}$$

If action α is of type 2, then there is no apparent algebraic model α_0 and the algebraic structure has to be built from scratch.

Building blocks: Intersections of stable foliations of different elements of the action give dynamically natural invariant foliations. The finest such intersections are called *coarse Lyapunov foliations*.

Classification results for \mathbb{Z}^k actions on arbitrary manifold

An action is called **TNS** (totally nonsymplectic) if any pair of coarse Lyapunov foliations is contracted by some element.

Classification results for \mathbb{Z}^k actions on arbitrary manifold

An action is called **TNS** (totally nonsymplectic) if any pair of coarse Lyapunov foliations is contracted by some element.

Let α be a TNS action of \mathbb{Z}^k by Anosov diffeomorphisms.

For each coarse Lyapunov distribution E of dimension more than 1 we assume that some elements of α contracting E are 1/2-pinned:

there exist $0 < \mu < \lambda < 2\mu$ and $K > 0$ such that for any $v \in E$,

$$K^{-1}e^{-n\lambda}\|v\| \leq \|Df^n(v)\| \leq Ke^{-n\mu}\|v\|.$$

Classification results for \mathbb{Z}^k actions on arbitrary manifold

An action is called **TNS** (totally nonsymplectic) if any pair of coarse Lyapunov foliations is contracted by some element.

Let α be a TNS action of \mathbb{Z}^k by Anosov diffeomorphisms.

For each coarse Lyapunov distribution E of dimension more than 1 we assume that some elements of α contracting E are 1/2-pinned:

there exist $0 < \mu < \lambda < 2\mu$ and $K > 0$ such that for any $v \in E$,

$$K^{-1}e^{-n\lambda}\|v\| \leq \|Df^n(v)\| \leq Ke^{-n\mu}\|v\|.$$

Theorem (B.K. and V. Sadovskaya)

If any two coarse Lyapunov foliations are (topologically) jointly integrable, then a finite cover of α is C^∞ conjugate to a \mathbb{Z}^k action by affine automorphisms of \mathbb{T}^n .

Classification results for \mathbb{Z}^k actions on arbitrary manifold

An action is called **TNS** (totally nonsymplectic) if any pair of coarse Lyapunov foliations is contracted by some element.

Let α be a TNS action of \mathbb{Z}^k by Anosov diffeomorphisms. For each coarse Lyapunov distribution E of dimension more than 1 we assume that some elements of α contracting E are 1/2-pinned: there exist $0 < \mu < \lambda < 2\mu$ and $K > 0$ such that for any $v \in E$,

$$K^{-1}e^{-n\lambda}\|v\| \leq \|Df^n(v)\| \leq Ke^{-n\mu}\|v\|.$$

Theorem (B.K. and V. Sadovskaya)

If any two coarse Lyapunov foliations are (topologically) jointly integrable, then a finite cover of α is C^∞ conjugate to a \mathbb{Z}^k action by affine automorphisms of \mathbb{T}^n .

For actions with stronger pinching (uniform quasiconformality), joint integrability can be replaced by certain *non-resonance* assumption on Lyapunov exponents.

Main result for \mathbb{Z}^k actions on tori and nilmanifolds

Theorem (D. Fisher, B.K., R. Spatzier)

Let α be a C^∞ -action of \mathbb{Z}^k , $k \geq 2$, on a nilmanifold N/Γ .

Let α_0 be the corresponding algebraic action.

- (1) there is an Anosov element for α in each Weyl chamber,*
- (2) there is $\mathbb{Z}^2 \subset \mathbb{Z}^k$ such that $\alpha_0(b)$ is ergodic for all $0 \neq b \in \mathbb{Z}^2$.*

Then α is C^∞ -conjugate to α_0 .

(2) is the most general *genuine higher rank* condition.

Main result for \mathbb{Z}^k actions on tori and nilmanifolds

Theorem (D. Fisher, B.K., R. Spatzier)

Let α be a C^∞ -action of \mathbb{Z}^k , $k \geq 2$, on a nilmanifold N/Γ .

Let α_0 be the corresponding algebraic action.

(1) there is an Anosov element for α in each Weyl chamber,

(2) there is $\mathbb{Z}^2 \subset \mathbb{Z}^k$ such that $\alpha_0(b)$ is ergodic for all $0 \neq b \in \mathbb{Z}^2$.

Then α is C^∞ -conjugate to α_0 .

(2) is the most general *genuine higher rank* condition.

Main new ingredients:

- Exponential decay of correlations for Hölder functions.
- New regularity result.

Regularity result

We consider Hölder foliations with smooth leaves on a manifold M . We assume that they are strongly absolutely continuous, i.e. the volume has absolutely continuous conditional measures on the leaves with densities that are smooth along the leaves and Hölder continuous transversally.

Regularity result

We consider Hölder foliations with smooth leaves on a manifold M . We assume that they are strongly absolutely continuous, i.e. the volume has absolutely continuous conditional measures on the leaves with densities that are smooth along the leaves and Hölder continuous transversally.

Theorem (D. Fisher, B.K., R. Spatzier)

Suppose that tangent spaces of foliations $\mathcal{F}_1, \dots, \mathcal{F}_r$ span TM . Let D be a distribution defined by an L^1 function ϕ on M . If the derivatives of D along each \mathcal{F}_i extend to functionals on C^θ for any $\theta > 0$, then ϕ is C^∞ .

Conjugacy equation

Fix $\mathbf{n} \in \mathbb{Z}^k$ and denote $a = \alpha(\mathbf{n})$ and $A = \alpha_0(\mathbf{n})$.

We lift a and A to \mathbb{R}^n and write the lift of conjugacy as $\text{Id} + h$.

$$(\text{Id} + h)(a(x)) = A(\text{Id} + h)(x)$$

Conjugacy equation

Fix $\mathbf{n} \in \mathbb{Z}^k$ and denote $a = \alpha(\mathbf{n})$ and $A = \alpha_0(\mathbf{n})$.

We lift a and A to \mathbb{R}^n and write the lift of conjugacy as $\text{Id} + h$.

$$(\text{Id} + h)(a(x)) = A(\text{Id} + h)(x)$$

The conjugacy equation yields functional fixed point equation for h

$$h(x) = Q(x) + A^{-1}(h(ax)), \quad \text{where } Q = A^{-1} \circ a - \text{Id}$$

as a function from \mathbb{T}^n to \mathbb{R}^n .

Conjugacy equation

Fix $\mathbf{n} \in \mathbb{Z}^k$ and denote $a = \alpha(\mathbf{n})$ and $A = \alpha_0(\mathbf{n})$.

We lift a and A to \mathbb{R}^n and write the lift of conjugacy as $\text{Id} + h$.

$$(\text{Id} + h)(a(x)) = A(\text{Id} + h)(x)$$

The conjugacy equation yields functional fixed point equation for h

$$h(x) = Q(x) + A^{-1}(h(ax)), \quad \text{where } Q = A^{-1} \circ a - \text{Id}$$

as a function from \mathbb{T}^n to \mathbb{R}^n . Projecting to an A -invariant subspace $V \subset \mathbb{R}^n$ along its A -invariant complement we obtain an equation

$$h_V(x) = Q_V(x) + A_V^{-1}(h_V(ax)) \quad =: F_V(h_V)$$

in $C^0(\mathbb{T}^n, V)$

Conjugacy equation

Fix $\mathbf{n} \in \mathbb{Z}^k$ and denote $a = \alpha(\mathbf{n})$ and $A = \alpha_0(\mathbf{n})$.

We lift a and A to \mathbb{R}^n and write the lift of conjugacy as $\text{Id} + h$.

$$(\text{Id} + h)(a(x)) = A(\text{Id} + h)(x)$$

The conjugacy equation yields functional fixed point equation for h

$$h(x) = Q(x) + A^{-1}(h(ax)), \quad \text{where } Q = A^{-1} \circ a - \text{Id}$$

as a function from \mathbb{T}^n to \mathbb{R}^n . Projecting to an A -invariant subspace $V \subset \mathbb{R}^n$ along its A -invariant complement we obtain an equation

$$h_V(x) = Q_V(x) + A_V^{-1}(h_V(ax)) \quad =: F_V(h_V)$$

in $C^0(\mathbb{T}^n, V)$ and a formal solution

$$h_V(x) = \sum_{n=0}^{\infty} A_V^{-n} Q_V(a^n x) = F_V^n(0).$$

$$h_V(x) = \sum_{n=0}^{\infty} A_V^{-n} Q_V(a^n x).$$

The equation holds in $C^0(\mathbb{T}^n, V)$ or in the space \mathcal{D} of distributions on smooth functions.

Derivatives as distributions

$$h_V(x) = \sum_{n=0}^{\infty} A_V^{-n} Q_V(a^n x).$$

The equation holds in $C^0(\mathbb{T}^n, V)$ or in the space \mathcal{D} of distributions on smooth functions. Differentiating in \mathcal{D} along foliation \mathcal{V}' we get

$$\langle h_V^{k, \mathcal{V}'}, f \rangle = \sum_{n=0}^{\infty} \langle A^{-n} (Q_V \circ a^n)^{k, \mathcal{V}'}, f \rangle.$$

Derivatives as distributions

$$h_V(x) = \sum_{n=0}^{\infty} A_V^{-n} Q_V(a^n x).$$

The equation holds in $C^0(\mathbb{T}^n, V)$ or in the space \mathcal{D} of distributions on smooth functions. Differentiating in \mathcal{D} along foliation \mathcal{V}' we get

$$\langle h_V^{k, \mathcal{V}'}, f \rangle = \sum_{n=0}^{\infty} \langle A_V^{-n} (Q_V \circ a^n)^{k, \mathcal{V}'}, f \rangle.$$

For $a = \alpha(\mathbf{n})$ with \mathbf{n} close to the Lyapunov hyperplane of \mathcal{V}' we obtain uniform slow exponential estimates on the growth of the derivatives of a^n along \mathcal{V}' .

Derivatives as distributions

$$h_V(x) = \sum_{n=0}^{\infty} A_V^{-n} Q_V(a^n x).$$

The equation holds in $C^0(\mathbb{T}^n, V)$ or in the space \mathcal{D} of distributions on smooth functions. Differentiating in \mathcal{D} along foliation \mathcal{V}' we get

$$\langle h_V^{k, \mathcal{V}'}, f \rangle = \sum_{n=0}^{\infty} \langle A^{-n} (Q_V \circ a^n)^{k, \mathcal{V}'}, f \rangle.$$

For $a = \alpha(\mathbf{n})$ with \mathbf{n} close to the Lyapunov hyperplane of \mathcal{V}' we obtain uniform slow exponential estimates on the growth of the derivatives of a^n along \mathcal{V}' . We use exponential mixing to show the pairing can be estimated by the Hölder norm of f .

Exponential mixing for Hölder functions.

Theorem (D. Fisher, B.K., R. Spatzier)

Let α be a \mathbb{Z}^k action on \mathbb{T}^n by affine automorphisms such that $\alpha(\mathbf{n})$ is ergodic for every nonzero $\mathbf{n} \in \mathbb{Z}^k$.

Then there exist $0 < r < 1$ and $C = C(\theta)$ such that for any θ -Hölder functions f and g on \mathbb{T}^n with $\int_{\mathbb{T}^n} f = \int_{\mathbb{T}^n} g = 0$ and for any $\mathbf{n} \in \mathbb{Z}^k$

$$\langle \alpha(\mathbf{n})f, g \rangle = \int_{\mathbb{T}^n} f \circ \alpha(\mathbf{n}) \cdot g < C(\|f\|_{\theta} \|g\|_2 + \|g\|_{\theta} \|f\|_2) r^{\theta \|\mathbf{n}\|}$$

Exponential mixing for Hölder functions.

Theorem (D. Fisher, B.K., R. Spatzier)

Let α be a \mathbb{Z}^k action on \mathbb{T}^n by affine automorphisms such that $\alpha(\mathbf{n})$ is ergodic for every nonzero $\mathbf{n} \in \mathbb{Z}^k$.

Then there exist $0 < r < 1$ and $C = C(\theta)$ such that for any θ -Hölder functions f and g on \mathbb{T}^n with $\int_{\mathbb{T}^n} f = \int_{\mathbb{T}^n} g = 0$ and for any $\mathbf{n} \in \mathbb{Z}^k$

$$\langle \alpha(\mathbf{n})f, g \rangle = \int_{\mathbb{T}^n} f \circ \alpha(\mathbf{n}) \cdot g < C(\|f\|_{\theta} \|g\|_2 + \|g\|_{\theta} \|f\|_2) r^{\theta \|\mathbf{n}\|}$$

The proof uses Fejér kernel approximations for f and g , Katznelson lemma for the distance of integer lattice points from an irrational invariant subspace of an integral matrix, and uniform lower bound for maximal expansion of $\alpha(\mathbf{n})$.