

Lecture 22

MATH 0200

Trigonometric
functions
composed
with their
inverses

Arccosine
plus arcsine

Lecture 22

Inverse trigonometric identities

MATH 0200

Dr. Boris Tselikhovskiy

Outline

Lecture 22

MATH 0200

Trigonometric
functions
composed
with their
inverses

Arccosine
plus arcsine

- 1 Trigonometric functions composed with their inverses
- 2 Arccosine plus arcsine

Trigonometric functions composed with their inverses

Lecture 22

MATH 0200

Trigonometric
functions
composed
with their
inverses

Arccosine
plus arcsine

Recall that if f and f^{-1} are inverse functions, then

Trigonometric functions composed with their inverses

Lecture 22

MATH 0200

Trigonometric
functions
composed
with their
inverses

Arccosine
plus arcsine

Recall that if f and f^{-1} are inverse functions, then

- $(f \circ f^{-1})(x) = x$ for any x in the domain of f^{-1} ;

Trigonometric functions composed with their inverses

Lecture 22

MATH 0200

Trigonometric
functions
composed
with their
inverses

Arccosine
plus arcsine

Recall that if f and f^{-1} are inverse functions, then

- $(f \circ f^{-1})(x) = x$ for any x in the domain of f^{-1} ;
- $(f^{-1} \circ f)(x) = x$ for any x in the domain of f .

Trigonometric functions composed with their inverses

Lecture 22

MATH 0200

Trigonometric
functions
composed
with their
inverses

Arccosine
plus arcsine

Recall that if f and f^{-1} are inverse functions, then

- $(f \circ f^{-1})(x) = x$ for any x in the domain of f^{-1} ;
- $(f^{-1} \circ f)(x) = x$ for any x in the domain of f .

Example

Evaluate $\arccos(\cos(400^\circ))$.

Trigonometric functions composed with their inverses

Lecture 22

MATH 0200

Trigonometric
functions
composed
with their
inverses

Arccosine
plus arcsine

Recall that if f and f^{-1} are inverse functions, then

- $(f \circ f^{-1})(x) = x$ for any x in the domain of f^{-1} ;
- $(f^{-1} \circ f)(x) = x$ for any x in the domain of f .

Example

Evaluate $\arccos(\cos(400^\circ))$.

First we need to find $0 \leq \alpha \leq \pi$ (or 180°) with $\cos(\alpha) = \cos(400^\circ)$. As $\cos(400^\circ) = \cos((400 - 360)^\circ) = \cos(40^\circ)$, we get $\alpha = 40^\circ = \frac{80\pi}{360} = \frac{2\pi}{9}$ and $\arccos(\cos(400^\circ)) = \arccos(\cos(\frac{2\pi}{9})) = \frac{2\pi}{9}$.

Example

Find the inverse of $f(x) = 5 - 6 \sin(3x)$ on interval $[0, \frac{\pi}{6}]$.

Example

Find the inverse of $f(x) = 5 - 6 \sin(3x)$ on interval $[0, \frac{\pi}{6}]$.

- 1 Write $y = 5 - 6 \sin(3x)$.

Example

Find the inverse of $f(x) = 5 - 6 \sin(3x)$ on interval $[0, \frac{\pi}{6}]$.

- ① Write $y = 5 - 6 \sin(3x)$.
- ② Transform $y = 5 - 6 \sin(3x) \Leftrightarrow y - 5 = -6 \sin(3x) \Leftrightarrow \frac{y-5}{-6} = \sin(3x) \Leftrightarrow 3x = \arcsin\left(\frac{y-5}{-6}\right) \Leftrightarrow x = \frac{1}{3} \arcsin\left(\frac{y-5}{-6}\right) = \frac{1}{3} \arcsin\left(\frac{5-y}{6}\right)$.

Example

Find the inverse of $f(x) = 5 - 6 \sin(3x)$ on interval $[0, \frac{\pi}{6}]$.

- ① Write $y = 5 - 6 \sin(3x)$.
- ② Transform $y = 5 - 6 \sin(3x) \Leftrightarrow y - 5 = -6 \sin(3x) \Leftrightarrow \frac{y-5}{-6} = \sin(3x) \Leftrightarrow 3x = \arcsin\left(\frac{y-5}{-6}\right) \Leftrightarrow x = \frac{1}{3} \arcsin\left(\frac{y-5}{-6}\right) = \frac{1}{3} \arcsin\left(\frac{5-y}{6}\right)$.
- ③ $f^{-1}(x) = \frac{1}{3} \arcsin\left(\frac{5-x}{6}\right)$.

Example

Find the inverse of $f(x) = 5 - 6 \sin(3x)$ on interval $[0, \frac{\pi}{6}]$.

- ① Write $y = 5 - 6 \sin(3x)$.
- ② Transform $y = 5 - 6 \sin(3x) \Leftrightarrow y - 5 = -6 \sin(3x) \Leftrightarrow \frac{y-5}{-6} = \sin(3x) \Leftrightarrow 3x = \arcsin\left(\frac{y-5}{-6}\right) \Leftrightarrow x = \frac{1}{3} \arcsin\left(\frac{y-5}{-6}\right) = \frac{1}{3} \arcsin\left(\frac{5-y}{6}\right)$.
- ③ $f^{-1}(x) = \frac{1}{3} \arcsin\left(\frac{5-x}{6}\right)$.

Check: $f \circ f^{-1}(x) = 5 - 6 \sin\left(3 \cdot \frac{1}{3} \arcsin\left(\frac{5-x}{6}\right)\right) = 5 - 6 \sin\left(\arcsin\left(\frac{5-x}{6}\right)\right) = 5 - 6 \cdot \left(\frac{5-x}{6}\right) = 5 - (5 - x) = x \checkmark$

Example

Find the inverse of $f(x) = 5 - 6 \sin(3x)$ on interval $[0, \frac{\pi}{6}]$.

- ① Write $y = 5 - 6 \sin(3x)$.
- ② Transform $y = 5 - 6 \sin(3x) \Leftrightarrow y - 5 = -6 \sin(3x) \Leftrightarrow \frac{y-5}{-6} = \sin(3x) \Leftrightarrow 3x = \arcsin\left(\frac{y-5}{-6}\right) \Leftrightarrow x = \frac{1}{3} \arcsin\left(\frac{y-5}{-6}\right) = \frac{1}{3} \arcsin\left(\frac{5-y}{6}\right)$.
- ③ $f^{-1}(x) = \frac{1}{3} \arcsin\left(\frac{5-x}{6}\right)$.

Check: $f \circ f^{-1}(x) = 5 - 6 \sin\left(3 \cdot \frac{1}{3} \arcsin\left(\frac{5-x}{6}\right)\right) = 5 - 6 \sin\left(\arcsin\left(\frac{5-x}{6}\right)\right) = 5 - 6 \cdot \left(\frac{5-x}{6}\right) = 5 - (5 - x) = x \checkmark$
 Range of $f^{-1} = \text{domain of } f = [0, \frac{\pi}{6}]$.

Example

Find the inverse of $f(x) = 5 - 6 \sin(3x)$ on interval $[0, \frac{\pi}{6}]$.

- ① Write $y = 5 - 6 \sin(3x)$.
- ② Transform $y = 5 - 6 \sin(3x) \Leftrightarrow y - 5 = -6 \sin(3x) \Leftrightarrow \frac{y-5}{-6} = \sin(3x) \Leftrightarrow 3x = \arcsin\left(\frac{y-5}{-6}\right) \Leftrightarrow x = \frac{1}{3} \arcsin\left(\frac{y-5}{-6}\right) = \frac{1}{3} \arcsin\left(\frac{5-y}{6}\right)$.
- ③ $f^{-1}(x) = \frac{1}{3} \arcsin\left(\frac{5-x}{6}\right)$.

Check: $f \circ f^{-1}(x) = 5 - 6 \sin\left(3 \cdot \frac{1}{3} \arcsin\left(\frac{5-x}{6}\right)\right) = 5 - 6 \sin\left(\arcsin\left(\frac{5-x}{6}\right)\right) = 5 - 6 \cdot \left(\frac{5-x}{6}\right) = 5 - (5 - x) = x \checkmark$
 Range of f^{-1} = domain of $f = [0, \frac{\pi}{6}]$.

As $\sin(3x)$ attains all values between $0 = \sin(0)$ and $1 = \sin\left(\frac{\pi}{6}\right)$, we get domain of f^{-1} = range of f is $[-1, 5]$.

Example

1 Evaluate $\sin(\arccos(0.4))$.

Example

- ① Evaluate $\sin(\arccos(0.4))$.

$$\begin{aligned}\sin(\arccos(0.4)) &= \sqrt{1 - \cos^2(\arccos(0.4))} = \\ &= \sqrt{1 - 0.16} = \sqrt{0.84}.\end{aligned}$$

Example

- ① Evaluate $\sin(\arccos(0.4))$.

$$\sin(\arccos(0.4)) = \sqrt{1 - \cos^2(\arccos(0.4))} = \sqrt{1 - 0.16} = \sqrt{0.84}.$$

- ② Evaluate $\tan(\arccos(0.4))$.

Example

- ① Evaluate $\sin(\arccos(0.4))$.

$$\sin(\arccos(0.4)) = \sqrt{1 - \cos^2(\arccos(0.4))} = \sqrt{1 - 0.16} = \sqrt{0.84}.$$

- ② Evaluate $\tan(\arccos(0.4))$.

$$\tan(\arccos(0.4)) = \frac{\sin(\arccos(0.4))}{\cos(\arccos(0.4))} = \frac{\sqrt{0.84}}{0.4} \approx 2.291.$$

Arccosine plus arcsine

Lecture 22

MATH 0200

Trigonometric
functions
composed
with their
inverses

Arccosine
plus arcsine

Let a be a number between 0 and 1. Consider a right triangle with a leg of length a and hypotenuse of length 1.

Arccosine plus arcsine


Lecture 22

MATH 0200

Trigonometric
functions
composed
with their
inverses

Arccosine
plus arcsine

Let a be a number between 0 and 1. Consider a right triangle with a leg of length a and hypotenuse of length 1.



Arccosine plus arcsine


Lecture 22

MATH 0200

Trigonometric
functions
composed
with their
inverses

Arccosine
plus arcsine

Let a be a number between 0 and 1. Consider a right triangle with a leg of length a and hypotenuse of length 1.



As $\sin(\alpha) = \cos\left(\frac{\pi}{2} - \alpha\right) = a/1 = a$, we get

$$\arcsin(a) + \arccos(a) = \alpha + \left(\frac{\pi}{2} - \alpha\right) = \frac{\pi}{2}.$$

Arccosine plus arcsine


Lecture 22

MATH 0200

Trigonometric
functions
composed
with their
inverses

Arccosine
plus arcsine

Let a be a number between 0 and 1. Consider a right triangle with a leg of length a and hypotenuse of length 1.



As $\sin(\alpha) = \cos\left(\frac{\pi}{2} - \alpha\right) = a/1 = a$, we get

$$\arcsin(a) + \arccos(a) = \alpha + \left(\frac{\pi}{2} - \alpha\right) = \frac{\pi}{2}.$$

The equation $\arcsin(a) + \arccos(a) = \frac{\pi}{2}$ holds true for any $-1 \leq a \leq 1$.

Lecture 22

MATH 0200

Trigonometric
functions
composed
with their
inverses

Arccosine
plus arcsine

Question

Evaluate $\arcsin(\cos(\frac{\pi}{2} - 1))$.