

Lecture 31

MATH 0200

Zeros of
polynomials

Powers of
complex
numbers

Trigonometric
identities

Lecture 31

Complex numbers (applications)

MATH 0200

Dr. Boris Tselikhovskiy

Outline

Lecture 31

MATH 0200

Zeros of
polynomials

Powers of
complex
numbers

Trigonometric
identities

- 1 Zeros of polynomials
- 2 Powers of complex numbers
- 3 Trigonometric identities

Zeros of polynomials

Lecture 31

MATH 0200

Zeros of
polynomials

Powers of
complex
numbers

Trigonometric
identities

Example

The polynomial $P(x) = x^2 + 9$ does not have any real zeros.

Reason: $\sqrt{-9}$ is not a real number.

Zeros of polynomials

Lecture 31

MATH 0200

Zeros of
polynomials

Powers of
complex
numbers

Trigonometric
identities

Example

The polynomial $P(x) = x^2 + 9$ does not have any real zeros.

Reason: $\sqrt{-9}$ is not a real number.

However, $\sqrt{-9} = \sqrt{9i^2} = 3i$ and, therefore, the complex numbers $3i$ and $-3i$ are zeros of the polynomial $P(x)$.

Zeros of polynomials

Lecture 31

MATH 0200

Zeros of
polynomials

Powers of
complex
numbers

Trigonometric
identities

Example

The polynomial $P(x) = x^2 + 9$ does not have any real zeros.

Reason: $\sqrt{-9}$ is not a real number.

However, $\sqrt{-9} = \sqrt{9i^2} = 3i$ and, therefore, the complex numbers $3i$ and $-3i$ are zeros of the polynomial $P(x)$.

More generally, recall that the zeros of a polynomial $f(x) = ax^2 + bx + c$ are $\frac{-b+\sqrt{D}}{2a}$ and $\frac{-b-\sqrt{D}}{2a}$, where $D = b^2 - 4ac$ is the discriminant of $f(x)$. If $D < 0$, then \sqrt{D} is not a real number, so there are no real zeros of $f(x)$, but there are two complex zeros, namely, $\frac{-b+i\sqrt{-D}}{2a}$ and $\frac{-b-i\sqrt{-D}}{2a}$.

Example

Find the zeros of quadratic polynomial $P(x) = x^2 - 4x + 5$.

Example

Find the zeros of quadratic polynomial $P(x) = x^2 - 4x + 5$.

We find the discriminant

$$D = (-4)^2 - 4 \cdot 5 = 16 - 20 = -4 = (2i)^2 \text{ and the zeros}$$
$$x_1 = \frac{4+2i}{2} = 2 + i \text{ and } x_2 = \frac{4-2i}{2} = 2 - i.$$

Example

Find the zeros of quadratic polynomial $P(x) = x^2 - 4x + 5$.

We find the discriminant

$$D = (-4)^2 - 4 \cdot 5 = 16 - 20 = -4 = (2i)^2 \text{ and the zeros } x_1 = \frac{4+2i}{2} = 2 + i \text{ and } x_2 = \frac{4-2i}{2} = 2 - i.$$

Remark

Notice that the numbers $2 + i$ and $2 - i$ are conjugate. This is not a coincidence. If a complex number z is a zero of a polynomial $P(x)$ with real coefficients, then its conjugate \bar{z} is a zero of $P(x)$ as well.

Example

Find the zeros of quadratic polynomial $P(x) = x^2 - 4x + 5$.

We find the discriminant

$$D = (-4)^2 - 4 \cdot 5 = 16 - 20 = -4 = (2i)^2 \text{ and the zeros } x_1 = \frac{4+2i}{2} = 2 + i \text{ and } x_2 = \frac{4-2i}{2} = 2 - i.$$

Remark

Notice that the numbers $2 + i$ and $2 - i$ are conjugate. This is not a coincidence. If a complex number z is a zero of a polynomial $P(x)$ with real coefficients, then its conjugate \bar{z} is a zero of $P(x)$ as well.

This follows from the fact a real number is equal to its conjugate, hence, $P(\bar{z}) = 0 = \bar{0} = \overline{P(z)}$.

Powers of complex numbers

Lecture 31

MATH 0200

Zeros of
polynomials

**Powers of
complex
numbers**

Trigonometric
identities

Let $z = a + bi$ be a complex number.

Powers of complex numbers

Lecture 31

MATH 0200

Zeros of
polynomials

Powers of
complex
numbers

Trigonometric
identities

Let $z = a + bi$ be a complex number.

Question

How can we compute z^{10} ?

Powers of complex numbers

Lecture 31

MATH 0200

Zeros of
polynomials

Powers of
complex
numbers

Trigonometric
identities

Let $z = a + bi$ be a complex number.

Question

How can we compute z^{10} ?

Well, it is possible to compute $(a + bi)^{10}$ directly, but would be computationally intense. Instead, we should use the polar form of z . Recall that the magnitude of z^{10} is $|z|^{10} = (\sqrt{a^2 + b^2})^{10} = (a^2 + b^2)^5$ and the argument is $Arg(z^{10}) = 10Arg(z)$ (modulo 2π).

Example

Let $z = \frac{1}{\sqrt{2}}(\sqrt{3} + i)$ and find z^{14} .

Example

Let $z = \frac{1}{\sqrt{2}}(\sqrt{3} + i)$ and find z^{14} .

We first compute $|z| = \sqrt{\left(\frac{\sqrt{3}}{\sqrt{2}}\right)^2 + \left(\frac{1}{\sqrt{2}}\right)^2} = \sqrt{\frac{3+1}{2}} = \sqrt{2}$
and $\text{Arg}(z) = \arctan\left(\frac{1}{\sqrt{3}}\right) = \frac{\pi}{6}$. It follows that $|z^{14}| = 2^7$
and $\text{Arg}(z^{14}) = \frac{14\pi}{6} = 2\pi + \frac{2\pi}{6} = 2\pi + \frac{\pi}{3} \equiv \frac{\pi}{3} \pmod{2\pi}$.

Example

Let $z = \frac{1}{\sqrt{2}}(\sqrt{3} + i)$ and find z^{14} .

We first compute $|z| = \sqrt{\left(\frac{\sqrt{3}}{\sqrt{2}}\right)^2 + \left(\frac{1}{\sqrt{2}}\right)^2} = \sqrt{\frac{3+1}{2}} = \sqrt{2}$
and $\text{Arg}(z) = \arctan\left(\frac{1}{\sqrt{3}}\right) = \frac{\pi}{6}$. It follows that $|z^{14}| = 2^7$
and $\text{Arg}(z^{14}) = \frac{14\pi}{6} = 2\pi + \frac{2\pi}{6} = 2\pi + \frac{\pi}{3} \equiv \frac{\pi}{3} \pmod{2\pi}$.

Therefore, $z^{14} = 2^7(\cos(\frac{\pi}{3}) + i \sin(\frac{\pi}{3})) = 2^7\left(\frac{1}{2} + i\frac{\sqrt{3}}{2}\right) = 2^6(1 + \sqrt{3}i) = 64(1 + \sqrt{3}i)$.

Question

Let $z = 1 + i$.

(a) Compute $|z|$.

Question

Let $z = 1 + i$.

- (a) Compute $|z|$.
- (b) Compute $\text{Arg}(z)$.

Question

Let $z = 1 + i$.

- (a) Compute $|z|$.
- (b) Compute $\text{Arg}(z)$.
- (c) Compute $|z^{10}|$.

Question

Let $z = 1 + i$.

- (a) Compute $|z|$.
- (b) Compute $\text{Arg}(z)$.
- (c) Compute $|z^{10}|$.
- (d) Compute $\text{Arg}(z^{10})$.

Question

Let $z = 1 + i$.

- (a) Compute $|z|$.
- (b) Compute $\text{Arg}(z)$.
- (c) Compute $|z^{10}|$.
- (d) Compute $\text{Arg}(z^{10})$.

Answer:

(a) $|z| = \sqrt{1^2 + 1^2} = \sqrt{2}$.

Question

Let $z = 1 + i$.

- (a) Compute $|z|$.
- (b) Compute $\text{Arg}(z)$.
- (c) Compute $|z^{10}|$.
- (d) Compute $\text{Arg}(z^{10})$.

Answer:

- (a) $|z| = \sqrt{1^2 + 1^2} = \sqrt{2}$.
- (b) $\text{Arg}(z) = \arctan\left(\frac{1}{1}\right) = \arctan(1) = \frac{\pi}{4}$.

Question

Let $z = 1 + i$.

- (a) Compute $|z|$.
- (b) Compute $\text{Arg}(z)$.
- (c) Compute $|z^{10}|$.
- (d) Compute $\text{Arg}(z^{10})$.

Answer:

- (a) $|z| = \sqrt{1^2 + 1^2} = \sqrt{2}$.
- (b) $\text{Arg}(z) = \arctan\left(\frac{1}{1}\right) = \arctan(1) = \frac{\pi}{4}$.
- (c) $|z^{10}| = \sqrt{2}^{10} = 2^5 = 32$.

Question

Let $z = 1 + i$.

- (a) Compute $|z|$.
- (b) Compute $\text{Arg}(z)$.
- (c) Compute $|z^{10}|$.
- (d) Compute $\text{Arg}(z^{10})$.

Answer:

- (a) $|z| = \sqrt{1^2 + 1^2} = \sqrt{2}$.
- (b) $\text{Arg}(z) = \arctan\left(\frac{1}{1}\right) = \arctan(1) = \frac{\pi}{4}$.
- (c) $|z^{10}| = \sqrt{2}^{10} = 2^5 = 32$.
- (d) $\text{Arg}(z^{10}) = 10\text{Arg}(z) = \frac{10\pi}{4} = 2\pi + \frac{2\pi}{4} = 2\pi + \frac{\pi}{2} \equiv \frac{\pi}{2}$.

Question

Let $z = 1 + i$.

- (a) Compute $|z|$.
- (b) Compute $\text{Arg}(z)$.
- (c) Compute $|z^{10}|$.
- (d) Compute $\text{Arg}(z^{10})$.

Answer:

(a) $|z| = \sqrt{1^2 + 1^2} = \sqrt{2}$.

(b) $\text{Arg}(z) = \arctan\left(\frac{1}{1}\right) = \arctan(1) = \frac{\pi}{4}$.

(c) $|z^{10}| = \sqrt{2}^{10} = 2^5 = 32$.

(d) $\text{Arg}(z^{10}) = 10\text{Arg}(z) = \frac{10\pi}{4} = 2\pi + \frac{2\pi}{4} = 2\pi + \frac{\pi}{2} \equiv \frac{\pi}{2}$.

We conclude that $(1 + i)^{10} = 32i$.

Let's take a look at one more example.

Example

Evaluate i^n for a non-negative integer n .

Let's take a look at one more example.

Example

Evaluate i^n for a non-negative integer n . Notice that $|i| = 1$ and $\text{Arg}(i) = \frac{\pi}{2}$, so $|i^n| = |1|^n = 1$ and


$$\text{Arg}(i^n) = \frac{\pi n}{2} = \begin{cases} 0, n\%4 = 0 \\ \frac{\pi}{2}, n\%4 = 1 \\ \pi, n\%4 = 2 \\ \frac{3\pi}{2}, n\%4 = 3 \end{cases} \quad \text{with } i^n = \begin{cases} 1, n\%4 = 0 \\ i, n\%4 = 1 \\ -1, n\%4 = 2 \\ -i, n\%4 = 3. \end{cases}$$

Let's take a look at one more example.

Example

Evaluate i^n for a non-negative integer n . Notice that $|i| = 1$ and $\text{Arg}(i) = \frac{\pi}{2}$, so $|i^n| = |1|^n = 1$ and

$$\text{Arg}(i^n) = \frac{\pi n}{2} = \begin{cases} 0, n\%4 = 0 \\ \frac{\pi}{2}, n\%4 = 1 \\ \pi, n\%4 = 2 \\ \frac{3\pi}{2}, n\%4 = 3 \end{cases} \quad \text{with } i^n = \begin{cases} 1, n\%4 = 0 \\ i, n\%4 = 1 \\ -1, n\%4 = 2 \\ -i, n\%4 = 3. \end{cases}$$



Lecture 31

MATH 0200

Zeros of
polynomials

Powers of
complex
numbers

Trigonometric
identities

Question

$$i^{2023} = ?$$

Question

$$i^{2023} = ?$$

Answer: as $2023 \% 4 = 3$, we conclude that $i^{2023} = i^3 = -i$.

Formulas for $\cos(a + b)$ and $\sin(a + b)$ revisited

Lecture 31

MATH 0200

Zeros of
polynomials

Powers of
complex
numbers

Trigonometri
identities

Let $z = \cos(a) + i \sin(a)$ and $w = \cos(b) + i \sin(b)$ be two complex numbers. Observe that $|z| = |w| = 1$, so both z and w are on the unit circle and form angles a and b with the positive x -axis, respectively.

Formulas for $\cos(a + b)$ and $\sin(a + b)$ revisited

Lecture 31

MATH 0200

Zeros of
polynomials

Powers of
complex
numbers

Trigonometri
identities

Let $z = \cos(a) + i \sin(a)$ and $w = \cos(b) + i \sin(b)$ be two complex numbers. Observe that $|z| = |w| = 1$, so both z and w are on the unit circle and form angles a and b with the positive x -axis, respectively.

Notice that $|zw| = |z| \cdot |w| = 1 \cdot 1 = 1$, while $\text{Arg}(zw) = \text{Arg}(z) + \text{Arg}(w) = a + b$. It follows that

$$zw = \cos(a + b) + i \sin(a + b).$$

Formulas for $\cos(a + b)$ and $\sin(a + b)$ revisited

Lecture 31

MATH 0200

Zeros of
polynomials

Powers of
complex
numbers

Trigonometri
identities

Let $z = \cos(a) + i \sin(a)$ and $w = \cos(b) + i \sin(b)$ be two complex numbers. Observe that $|z| = |w| = 1$, so both z and w are on the unit circle and form angles a and b with the positive x -axis, respectively.

Notice that $|zw| = |z| \cdot |w| = 1 \cdot 1 = 1$, while $\text{Arg}(zw) = \text{Arg}(z) + \text{Arg}(w) = a + b$. It follows that

$$zw = \cos(a + b) + i \sin(a + b).$$

$$\begin{aligned} \text{Also, } zw &= (\cos(a) + i \sin(a))(\cos(b) + i \sin(b)) = \\ &= \cos(a) \cos(b) + i \cos(a) \sin(b) + i \sin(a) \cos(b) + i^2 \sin(a) \sin(b) = \\ &= (\cos(a) \cos(b) - \sin(a) \sin(b)) + i(\cos(a) \sin(b) + \sin(a) \cos(b)). \end{aligned}$$

Formulas for $\cos(a + b)$ and $\sin(a + b)$ revisited

Lecture 31
MATH 0200
Zeros of
polynomials
Powers of
complex
numbers
Trigonometri
identities

Let $z = \cos(a) + i \sin(a)$ and $w = \cos(b) + i \sin(b)$ be two complex numbers. Observe that $|z| = |w| = 1$, so both z and w are on the unit circle and form angles a and b with the positive x -axis, respectively.

Notice that $|zw| = |z| \cdot |w| = 1 \cdot 1 = 1$, while $\text{Arg}(zw) = \text{Arg}(z) + \text{Arg}(w) = a + b$. It follows that

$$zw = \cos(a + b) + i \sin(a + b).$$

$$\begin{aligned} \text{Also, } zw &= (\cos(a) + i \sin(a))(\cos(b) + i \sin(b)) = \\ &= \cos(a) \cos(b) + i \cos(a) \sin(b) + i \sin(a) \cos(b) + i^2 \sin(a) \sin(b) = \\ &= (\cos(a) \cos(b) - \sin(a) \sin(b)) + i(\cos(a) \sin(b) + \sin(a) \cos(b)). \end{aligned}$$

We have established that

- $\cos(a + b) = \cos(a) \cos(b) - \sin(a) \sin(b)$ and
- $\sin(a + b) = \cos(a) \sin(b) + \sin(a) \cos(b)$.