

Lecture 8

MATH 0200

Zeros of
quadratic
functions

Completing
the square

Quadratic
formula

Parabolas

Circles

Lecture 8

Quadratic functions

MATH 0200

Dr. Boris Tselikhovskiy

Outline

Lecture 8

MATH 0200

Zeros of
quadratic
functions

Completing
the square

Quadratic
formula

Parabolas

Circles

- 1 Zeros of quadratic functions
- 2 Completing the square
- 3 Quadratic formula
- 4 Parabolas
- 5 Circles

Zeros of quadratic functions

Lecture 8

MATH 0200

Zeros of
quadratic
functions

Completing
the square

Quadratic
formula

Parabolas

Circles

Today we will talk about quadratic functions. These are functions given by polynomials of degree 2:

Zeros of quadratic functions

Lecture 8

MATH 0200

Zeros of quadratic functions

Completing the square

Quadratic formula

Parabolas

Circles

Today we will talk about quadratic functions. These are functions given by polynomials of degree 2:

$$f(x) = ax^2 + bx + c,$$

where a, b and c are some numbers and $a \neq 0$.

Zeros of quadratic functions

Lecture 8

MATH 0200

Zeros of quadratic functions

Completing the square

Quadratic formula

Parabolas

Circles

Today we will talk about quadratic functions. These are functions given by polynomials of degree 2:

$$f(x) = ax^2 + bx + c,$$

where a, b and c are some numbers and $a \neq 0$. The first question we will address is how to find zeros of quadratic functions.

Zeros of quadratic functions

Lecture 8

MATH 0200

Zeros of quadratic functions

Completing the square

Quadratic formula

Parabolas

Circles

Today we will talk about quadratic functions. These are functions given by polynomials of degree 2:

$$f(x) = ax^2 + bx + c,$$

where a, b and c are some numbers and $a \neq 0$. The first question we will address is how to find zeros of quadratic functions.

Definition

A **zero** of a function $f(x)$ is a number d with $f(d) = 0$ (the points where the graph of $f(x)$ intersects the x -axis).

Example

Let's take a look at the function $f(x) = x^2 - 4$ and find its zeros. We need to solve the equation $f(x) = 0$:

Example


Let's take a look at the function $f(x) = x^2 - 4$ and find its zeros. We need to solve the equation $f(x) = 0$:

$$x^2 - 4 = 0 \Leftrightarrow x^2 = 4 \Leftrightarrow x = \pm\sqrt{4} \Leftrightarrow x = -2 \text{ or } x = 2.$$

Example

Let's take a look at the function $f(x) = x^2 - 4$ and find its zeros. We need to solve the equation $f(x) = 0$:

$$x^2 - 4 = 0 \Leftrightarrow x^2 = 4 \Leftrightarrow x = \pm\sqrt{4} \Leftrightarrow x = -2 \text{ or } x = 2.$$



Completing the square

Lecture 8

MATH 0200

Zeros of
quadratic
functions

Completing
the square

Quadratic
formula

Parabolas

Circles

Let $f(x) = x^2 + bx + c$ and consider a 2-step procedure for completing the square.

Completing the square

Lecture 8

MATH 0200

Zeros of
quadratic
functions

Completing
the square

Quadratic
formula

Parabolas

Circles

Let $f(x) = x^2 + bx + c$ and consider a 2-step procedure for completing the square.

Step 1. Rewrite bx , the linear term of $f(x)$ as

$$bx = 2 \cdot \text{panda} \cdot x \left(\text{for } \text{panda} = \frac{b}{2} \right):$$

$$f(x) = x^2 + 2 \cdot \text{panda} \cdot x + c.$$

Completing the square

Lecture 8

MATH 0200

Zeros of quadratic functions

Completing the square

Quadratic formula

Parabolas

Circles

Let $f(x) = x^2 + bx + c$ and consider a 2-step procedure for completing the square.

Step 1. Rewrite bx , the linear term of $f(x)$ as

$$bx = 2 \cdot \text{panda} \cdot x \left(\text{for } \text{panda} = \frac{b}{2} \right):$$

$$f(x) = x^2 + 2 \cdot \text{panda} \cdot x + c.$$

Step 2. Add and subtract panda^2 :

$$\begin{aligned} f(x) &= x^2 + 2 \cdot \text{panda} \cdot x + \text{panda}^2 - \text{panda}^2 + c = \\ &= \left(x + \text{panda} \right)^2 - \text{panda}^2 + c. \end{aligned}$$

Example

Find the zeros of quadratic function $f(x) = x^2 - 2x - 8$.

Example


Find the zeros of quadratic function $f(x) = x^2 - 2x - 8$.

- 1 We write $-2x = 2 \cdot (-1) \cdot x$ and
 $f(x) = x^2 + 2 \cdot (-1) \cdot x - 8$.

Example

Find the zeros of quadratic function $f(x) = x^2 - 2x - 8$.


- ① We write $-2x = 2 \cdot (-1) \cdot x$ and
 $f(x) = x^2 + 2 \cdot (-1) \cdot x - 8$.

- ② As  $= -1$, we get
 $f(x) = x^2 + 2 \cdot (-1) \cdot x - 8 = (x + (-1))^2 - (-1)^2 - 8$.

Example

Find the zeros of quadratic function $f(x) = x^2 - 2x - 8$.

- ① We write $-2x = 2 \cdot (-1) \cdot x$ and
$$f(x) = x^2 + 2 \cdot (-1) \cdot x - 8.$$

- ② As  $= -1$, we get


$$f(x) = x^2 + 2 \cdot (-1) \cdot x - 8 = (x + (-1))^2 - (-1)^2 - 8.$$

We compute $(x - 1)^2 - 1 - 8 = 0 \Leftrightarrow (x - 1)^2 = 9 \Leftrightarrow x - 1 = \pm\sqrt{9} = \pm 3 \Leftrightarrow x = \pm 3 + 1 \Leftrightarrow x = 4$ or $x = -2$.

Example

Find the zeros of quadratic function $f(x) = x^2 - 2x - 8$.

- ① We write $-2x = 2 \cdot (-1) \cdot x$ and
$$f(x) = x^2 + 2 \cdot (-1) \cdot x - 8.$$

- ② As  $= -1$, we get

$$f(x) = x^2 + 2 \cdot (-1) \cdot x - 8 = (x + (-1))^2 - (-1)^2 - 8.$$


We compute $(x - 1)^2 - 1 - 8 = 0 \Leftrightarrow (x - 1)^2 = 9 \Leftrightarrow x - 1 = \pm\sqrt{9} = \pm 3 \Leftrightarrow x = \pm 3 + 1 \Leftrightarrow x = 4$ or $x = -2$.

Check:

Example

Find the zeros of quadratic function $f(x) = x^2 - 2x - 8$.

- ① We write $-2x = 2 \cdot (-1) \cdot x$ and
$$f(x) = x^2 + 2 \cdot (-1) \cdot x - 8.$$

- ② As  $= -1$, we get

$$f(x) = x^2 + 2 \cdot (-1) \cdot x - 8 = (x + (-1))^2 - (-1)^2 - 8.$$

We compute $(x - 1)^2 - 1 - 8 = 0 \Leftrightarrow (x - 1)^2 = 9 \Leftrightarrow x - 1 = \pm\sqrt{9} = \pm 3 \Leftrightarrow x = \pm 3 + 1 \Leftrightarrow x = 4$ or $x = -2$.


Check:

- $f(-2) = (-2)^2 - 2 \cdot (-2) - 8 = 4 + 4 - 8 = 0 \quad \checkmark$

Example

Find the zeros of quadratic function $f(x) = x^2 - 2x - 8$.

- ① We write $-2x = 2 \cdot (-1) \cdot x$ and
$$f(x) = x^2 + 2 \cdot (-1) \cdot x - 8.$$

- ② As  $= -1$, we get

$$f(x) = x^2 + 2 \cdot (-1) \cdot x - 8 = (x + (-1))^2 - (-1)^2 - 8.$$

We compute $(x - 1)^2 - 1 - 8 = 0 \Leftrightarrow (x - 1)^2 = 9 \Leftrightarrow x - 1 = \pm\sqrt{9} = \pm 3 \Leftrightarrow x = \pm 3 + 1 \Leftrightarrow x = 4$ or $x = -2$.

Check:

- $f(-2) = (-2)^2 - 2 \cdot (-2) - 8 = 4 + 4 - 8 = 0 \quad \checkmark$
- $f(4) = 4^2 - 2 \cdot 4 - 8 = 16 - 8 - 8 = 0 \quad \checkmark$

Quadratic formula

Lecture 8

MATH 0200

Zeros of
quadratic
functions

Completing
the square

Quadratic
formula

Parabolas

Circles

Definition

Let $f(x) = ax^2 + bx + c$ be a quadratic function. The number $D = b^2 - 4ac$ is called the **discriminant** of f .

Quadratic formula

Lecture 8

MATH 0200

Zeros of
quadratic
functions

Completing
the square

Quadratic
formula

Parabolas

Circles

Definition

Let $f(x) = ax^2 + bx + c$ be a quadratic function. The number $D = b^2 - 4ac$ is called the **discriminant** of f .

Now we present the formula for zeros of $f(x)$.

Quadratic formula

Lecture 8

MATH 0200

Zeros of
quadratic
functions

Completing
the square

Quadratic
formula

Parabolas

Circles

Definition

Let $f(x) = ax^2 + bx + c$ be a quadratic function. The number $D = b^2 - 4ac$ is called the **discriminant** of f .

Now we present the formula for zeros of $f(x)$.

- 1 If $D < 0$, then $f(x)$ has no real zeros.

Quadratic formula

Lecture 8

MATH 0200

Zeros of
quadratic
functions

Completing
the square

Quadratic
formula

Parabolas

Circles

Definition

Let $f(x) = ax^2 + bx + c$ be a quadratic function. The number $D = b^2 - 4ac$ is called the **discriminant** of f .

Now we present the formula for zeros of $f(x)$.

- 1 If $D < 0$, then $f(x)$ has no real zeros.
- 2 If $D = 0$, then $f(x)$ has a unique zero $x = -\frac{b}{2a}$.

Quadratic formula

Lecture 8

MATH 0200

Zeros of
quadratic
functions

Completing
the square

Quadratic
formula

Parabolas

Circles

Definition

Let $f(x) = ax^2 + bx + c$ be a quadratic function. The number $D = b^2 - 4ac$ is called the **discriminant** of f .

Now we present the formula for zeros of $f(x)$.

- 1 If $D < 0$, then $f(x)$ has no real zeros.
- 2 If $D = 0$, then $f(x)$ has a unique zero $x = -\frac{b}{2a}$.
- 3 If $D > 0$, then $f(x)$ has two zeros $x = \frac{-b+\sqrt{D}}{2a}$ and $x = \frac{-b-\sqrt{D}}{2a}$.

Quadratic formula

Lecture 8

MATH 0200

Zeros of
quadratic
functions

Completing
the square

Quadratic
formula

Parabolas

Circles

Definition

Let $f(x) = ax^2 + bx + c$ be a quadratic function. The number $D = b^2 - 4ac$ is called the **discriminant** of f .

Now we present the formula for zeros of $f(x)$.

- 1 If $D < 0$, then $f(x)$ has no real zeros.
- 2 If $D = 0$, then $f(x)$ has a unique zero $x = -\frac{b}{2a}$.
- 3 If $D > 0$, then $f(x)$ has two zeros $x = \frac{-b + \sqrt{D}}{2a}$ and $x = \frac{-b - \sqrt{D}}{2a}$.

Example

We find the zeros of $f(x) = 3x^2 - 18x - 21$. The discriminant of $f(x)$ is $D = 18^2 - 4 \cdot 3 \cdot (-21) = 324 + 252 = 576 = 24^2$, and the roots are $(18 + 24)/6 = 7$ and $(18 - 24)/6 = -1$.

Parabola

Lecture 8

MATH 0200

Zeros of
quadratic
functions

Completing
the square

Quadratic
formula

Parabolas

Circles

The graph of a quadratic function $f(x) = ax^2 + bx + c$ is a parabola. It has a **vertex**, point $(-\frac{b}{2a}, f(-\frac{b}{2a}))$, and a **directrix** (line of symmetry), vertical line $x = -\frac{b}{2a}$.

Parabola

Lecture 8

MATH 0200

Zeros of
quadratic
functions


Completing
the square

Quadratic
formula

Parabolas

Circles

The graph of a quadratic function $f(x) = ax^2 + bx + c$ is a parabola. It has a **vertex**, point $(-\frac{b}{2a}, f(-\frac{b}{2a}))$, and a **directrix** (line of symmetry), vertical line $x = -\frac{b}{2a}$.



Lecture 8

MATH 0200

Zeros of
quadratic
functions

Completing
the square

Quadratic
formula

Parabolas

Circles

Question

Find the coordinates of the vertex of parabola given by equation $f(x) = -0.5x^2 + 7x - 4$.

Properties

Lecture 8

MATH 0200

Zeros of
quadratic
functions

Completing
the square

Quadratic
formula

Parabolas

Circles

Let $f(x) = ax^2 + bx + c$ be a quadratic function.

Properties

Lecture 8

MATH 0200

Zeros of
quadratic
functions

Completing
the square

Quadratic
formula

Parabolas

Circles

Let $f(x) = ax^2 + bx + c$ be a quadratic function.

- 1 The graph of $f(x)$ is symmetric with respect to its directrix (vertical line $x = -\frac{b}{2a}$).

Properties

Lecture 8

MATH 0200

Zeros of
quadratic
functions

Completing
the square

Quadratic
formula

Parabolas

Circles

Let $f(x) = ax^2 + bx + c$ be a quadratic function.

- 1 The graph of $f(x)$ is symmetric with respect to its directrix (vertical line $x = -\frac{b}{2a}$).
- 2 If $a > 0$, then $f(x)$ attains its (global) minimal value at the vertex. This value is $f\left(-\frac{b}{2a}\right)$.

Properties

Lecture 8

MATH 0200

Zeros of
quadratic
functions

Completing
the square


Quadratic
formula

Parabolas

Circles

Let $f(x) = ax^2 + bx + c$ be a quadratic function.

- 1 The graph of $f(x)$ is symmetric with respect to its directrix (vertical line $x = -\frac{b}{2a}$).
- 2 If $a > 0$, then $f(x)$ attains its (global) minimal value at the vertex. This value is $f\left(-\frac{b}{2a}\right)$.



Properties

Lecture 8

MATH 0200

Zeros of
quadratic
functions

Completing
the square


Quadratic
formula

Parabolas

Circles

Let $f(x) = ax^2 + bx + c$ be a quadratic function.

- 1 The graph of $f(x)$ is symmetric with respect to its directrix (vertical line $x = -\frac{b}{2a}$).
- 2 If $a > 0$, then $f(x)$ attains its (global) minimal value at the vertex. This value is $f\left(-\frac{b}{2a}\right)$.



- 3 If $a < 0$, then $f(x)$ attains its (global) maximal value at the vertex. This value is $f\left(-\frac{b}{2a}\right)$.

Properties

Lecture 8

MATH 0200

Zeros of
quadratic
functions

Completing
the square


Quadratic
formula

Parabolas


Circles

Let $f(x) = ax^2 + bx + c$ be a quadratic function.

- 1 The graph of $f(x)$ is symmetric with respect to its directrix (vertical line $x = -\frac{b}{2a}$).
- 2 If $a > 0$, then $f(x)$ attains its (global) minimal value at the vertex. This value is $f\left(-\frac{b}{2a}\right)$.



- 3 If $a < 0$, then $f(x)$ attains its (global) maximal value at the vertex. This value is $f\left(-\frac{b}{2a}\right)$.



Distance between points

Lecture 8

MATH 0200

Zeros of quadratic functions

Completing the square

Quadratic formula

Parabolas

Circles

Definition

Consider two points $P = (x_1, y_1)$ and $Q = (x_2, y_2)$. The **distance** between P and Q is the length of the line segment connecting these points. It is given by the formula

$$d(P, Q) = \sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2}.$$

Distance between points

Lecture 8

MATH 0200

Zeros of
quadratic
functions

Completing
the square

Quadratic
formula


Parabolas

Circles

Definition

Consider two points $P = (x_1, y_1)$ and $Q = (x_2, y_2)$. The **distance** between P and Q is the length of the line segment connecting these points. It is given by the formula

$$d(P, Q) = \sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2}.$$



Lecture 8

MATH 0200

Zeros of
quadratic
functions

Completing
the square

Quadratic
formula

Parabolas

Circles

Question

What is the distance between the points $(-3, 1)$ and $(5, 7)$?

Definition

A **circle** is a set of all points in a plane that are at a given distance (**radius**) from a given point, the **center**.

Definition

A **circle** is a set of all points in a plane that are at a given distance (**radius**) from a given point, the **center**.


The equation of the circle of radius r centered at $\mathcal{O} = (a, b)$ is $(x - a)^2 + (y - b)^2 = r^2$.

Definition

A **circle** is a set of all points in a plane that are at a given distance (**radius**) from a given point, the **center**.

The equation of the circle of radius r centered at $\mathcal{O} = (a, b)$ is $(x - a)^2 + (y - b)^2 = r^2$.

Example



Example

Find the radius and center of the circle given by equation $x^2 - 6x + y^2 + 10y = 15$.

Example

Find the radius and center of the circle given by equation $x^2 - 6x + y^2 + 10y = 15$.

We complete the squares:

$$x^2 - 6x + y^2 + 10y = 15 \Leftrightarrow$$

$$x^2 - 2 \cdot 3x + 9 - 9 + y^2 + 2 \cdot 5y + 25 - 25 = 15 \Leftrightarrow$$

$$(x - 3)^2 + (y + 5)^2 - 34 = 15 \Leftrightarrow$$

$$(x - 3)^2 + (y + 5)^2 = 49 \Leftrightarrow$$

$$(x - 3)^2 + (y + 5)^2 = 7^2.$$

Example

Find the radius and center of the circle given by equation $x^2 - 6x + y^2 + 10y = 15$.

We complete the squares:

$$x^2 - 6x + y^2 + 10y = 15 \Leftrightarrow$$

$$x^2 - 2 \cdot 3x + 9 - 9 + y^2 + 2 \cdot 5y + 25 - 25 = 15 \Leftrightarrow$$

$$(x - 3)^2 + (y + 5)^2 - 34 = 15 \Leftrightarrow$$

$$(x - 3)^2 + (y + 5)^2 = 49 \Leftrightarrow$$

$$(x - 3)^2 + (y + 5)^2 = 7^2.$$

The center is $(3, -5)$ and the radius is equal to 7.