MATH 0200

Trigonometr in right triangles

# Lecture 19 Trigonometry in right triangles

MATH 0200

Dr. Boris Tsvelikhovskiy

# Outline

Lecture 19


MATH 0200

Trigonometry in right triangles


• Trigonometry in right triangles

MATH 0200

Trigonometry in right triangles




Trigonometr in right triangles



#### Remark

We observe that our first trigonometric identity  $\sin^2(\alpha) + \cos^2(\alpha) = 1$  is nothing else but Pythagorean theorem for triangle  $\triangle ABC$ .


Trigonometr in right triangles



#### Remark

We observe that our first trigonometric identity  $\sin^2(\alpha) + \cos^2(\alpha) = 1$  is nothing else but Pythagorean theorem for triangle  $\triangle ABC$ .

Notice that the two right triangles  $\triangle ABC$  and  $\triangle \widetilde{ABC}$  are similar.




#### Remark

We observe that our first trigonometric identity  $\sin^2(\alpha) + \cos^2(\alpha) = 1$  is nothing else but Pythagorean theorem for triangle  $\triangle ABC$ .

Notice that the two right triangles  $\triangle ABC$  and  $\triangle \widetilde{A}\widetilde{B}\widetilde{C}$  are similar. We conclude that

• 
$$\sin(\alpha) = \frac{a}{c} = \frac{\text{opposite leg}}{\text{hypothenuse}};$$





#### $^{'}$ Remark


We observe that our first trigonometric identity  $\sin^2(\alpha) + \cos^2(\alpha) = 1$  is nothing else but Pythagorean theorem for triangle  $\triangle ABC$ .

Notice that the two right triangles  $\triangle ABC$  and  $\triangle \widetilde{ABC}$  are similar. We conclude that

• 
$$\sin(\alpha) = \frac{a}{c} = \frac{\text{opposite leg}}{\text{hypothenuse}};$$

• 
$$\cos(\alpha) = \frac{b}{c} = \frac{\text{adjacent leg}}{\text{hypothenuse}};$$

Trigonometr in right triangles



#### Remark

We observe that our first trigonometric identity  $\sin^2(\alpha) + \cos^2(\alpha) = 1$  is nothing else but Pythagorean theorem for triangle  $\triangle ABC$ .

Notice that the two right triangles  $\triangle ABC$  and  $\triangle \widetilde{A}\widetilde{B}\widetilde{C}$  are similar. We conclude that

• 
$$\sin(\alpha) = \frac{a}{c} = \frac{\text{opposite leg}}{\text{hypothenuse}};$$

• 
$$\cos(\alpha) = \frac{b}{c} = \frac{\text{adjacent leg}}{\text{hypothenuse}};$$

• 
$$\tan(\alpha) = \frac{a}{b} = \frac{\text{opposite leg}}{\text{adjacent leg}};$$



MATH 0200


Trigonometry in right triangles

# Example


MATH 0200

Trigonomet in right triangles

# Example




Consider the right triangle  $\triangle ABC$  below with legs AC = 12 and BC = 5.




(a) Find the length of the hypotenuse AC.


Consider the right triangle  $\triangle ABC$  below with legs AC = 12 and BC = 5.




(a) Find the length of the hypotenuse AC. We use Pythagorean theorem to get  $c = \sqrt{5^2 + 12^2} = \sqrt{169} = 13$ .




- (a) Find the length of the hypotenuse AC. We use Pythagorean theorem to get  $c = \sqrt{5^2 + 12^2} = \sqrt{169} = 13$ .
- (b) Evaluate  $\sin(\alpha)$ .




- (a) Find the length of the hypotenuse AC. We use Pythagorean theorem to get  $c = \sqrt{5^2 + 12^2} = \sqrt{169} = 13$ .
- (b) Evaluate  $\sin(\alpha)$ . We get  $\sin(\alpha) = \frac{12}{13}$ .




- (a) Find the length of the hypotenuse AC. We use Pythagorean theorem to get  $c = \sqrt{5^2 + 12^2} = \sqrt{169} = 13$ .
- (b) Evaluate  $\sin(\alpha)$ . We get  $\sin(\alpha) = \frac{12}{13}$ .
- (c) Evaluate  $\cos(\alpha)$ .



- (a) Find the length of the hypotenuse AC. We use Pythagorean theorem to get  $c = \sqrt{5^2 + 12^2} = \sqrt{169} = 13$ .
- (b) Evaluate  $\sin(\alpha)$ . We get  $\sin(\alpha) = \frac{12}{13}$ .
- (c) Evaluate  $\cos(\alpha)$ . We get  $\cos(\alpha) = \frac{5}{13}$ .



- (a) Find the length of the hypotenuse AC. We use Pythagorean theorem to get  $c = \sqrt{5^2 + 12^2} = \sqrt{169} = 13$ .
- (b) Evaluate  $\sin(\alpha)$ . We get  $\sin(\alpha) = \frac{12}{13}$ .
- (c) Evaluate  $\cos(\alpha)$ . We get  $\cos(\alpha) = \frac{5}{13}$ .
- (d) Evaluate  $\cot(\alpha)$ .



- (a) Find the length of the hypotenuse AC. We use Pythagorean theorem to get  $c = \sqrt{5^2 + 12^2} = \sqrt{169} = 13$ .
- (b) Evaluate  $\sin(\alpha)$ . We get  $\sin(\alpha) = \frac{12}{13}$ .
- (c) Evaluate  $\cos(\alpha)$ . We get  $\cos(\alpha) = \frac{5}{13}$ .
- (d) Evaluate  $\cot(\alpha)$ . We get  $\cot(\alpha) = \frac{\cos(\alpha)}{\sin(\alpha)} = \frac{5}{13} \cdot \frac{13}{12} = \frac{5}{12}$ .

MATH 0200

Trigonometr in right triangles

#### Example

Suppose a 20-foot ladder is leaning against a wall, making a 72° angle with the ground. How high up the wall is the end of the ladder?

Suppose a 20-foot ladder is leaning against a wall, making a 72° angle with the ground. How high up the wall is the end of the ladder?



Suppose a 20-foot ladder is leaning against a wall, making a 72° angle with the ground. How high up the wall is the end of the ladder?




**Solution:**  $\sin(72^\circ) = \frac{h}{20} \Leftrightarrow h = 20\sin(72^\circ) \approx 19.021 \text{ ft.}$ 

MATH 0200

Trigonometry in right triangles Consider the right triangle depicted below.

Consider the right triangle depicted below.


#### Question



• Evaluate  $\sin(\alpha)$ .

Consider the right triangle depicted below.

# Question



- Evaluate  $\sin(\alpha)$ .
- Evaluate  $tan(\alpha)$ .