



Review

Pharmacological approaches to treating negative symptoms: A review of clinical trials

Celso Arango^a, George Garibaldi^b, Stephen R. Marder^{c,d,*}^a Child and Adolescent Psychiatry Department, Hospital General Universitario Gregorio Marañón, IISGM, School of Medicine, Universidad Complutense, CIBERSAM, Madrid, Spain^b F. Hoffmann-La Roche, Basel, Switzerland^c Semel Institute for Neuroscience, UCLA, United States^d VA Desert Pacific Mental Illness Research, Education, and Clinical Center, United States

ARTICLE INFO

Article history:

Received 12 April 2013

Received in revised form 9 July 2013

Accepted 11 July 2013

Available online 9 August 2013

Keywords:

Schizophrenia

Pharmacology

Negative symptoms

ABSTRACT

Clinical trials of pharmacological agents targeting negative symptoms in schizophrenia are reviewed. The focus is on trials that occurred in patients who were stable on an antipsychotic medication at entry to the trial. A small number of trials compared antipsychotics as monotherapy for negative symptoms. Although the data supporting amisulpride for negative symptoms is promising the trials have limitations and it is plausible that the advantages of amisulpride over placebo may result from effects on secondary negative symptoms. Among available agents, antidepressant medications may have effects in negative symptoms. Other promising agents include minocycline, glutamatergic agents, and alpha-7 nicotinic agents. More than 15 active trials are currently underway to evaluate new treatments for negative symptoms.

Published by Elsevier B.V.

1. Introduction

Negative symptoms are an important target for drug development for a number of reasons: (1) negative symptoms are relatively common with a recent study finding that nearly 58% of outpatients had at least one negative symptom (Bobes et al., 2010); (2) negative symptoms are better predictors of functioning than positive symptoms (Rabinowitz et al., 2012); and (3) there are no accepted treatments for primary negative symptoms. With the possible exception of amisulpride in some countries, antipsychotic medications are relatively ineffective for managing negative symptoms in stable patients. This lack of effectiveness is not surprising since complex disorders such as schizophrenia may include families of related disorders. In schizophrenia, patients may have impairments in a number of psychopathological domains including psychotic symptoms, cognitive impairments, and negative symptoms. As a result, it may be unrealistic to expect a single drug to treat all aspects of the disorder (Hyman and Fenton, 2003; Arango et al., 2004; Carpenter and Davis, 2012). Recent attention has focused on the development of pharmacological agents that have specific activity in treating negative symptoms that can be added to an antipsychotic medication (Arango et al., 2004; Kirkpatrick et al., 2006; Marder et al., 2011).

This review will focus on clinical trials of pharmacological agents for treating negative symptoms. We have selected trials that used specific

entry criteria for negative symptoms and excluded trials that measured negative symptom change in individuals with acute psychosis. In most cases, the trials fit recent guidelines for negative symptom trials (Marder et al., 2011). That is, negative symptoms were stable and persistent. In addition, other causes of negative symptoms such as depression, extrapyramidal side effects, and psychosis were not sufficiently severe to cause secondary negative symptoms. Studies used different criteria for assuring that persistent positive symptoms such as hallucinations and delusions were not causing negative symptoms such as emotional withdrawal and avolition. These varied from studies requiring that positive symptoms be no greater than mild to studies that permitted moderately severe positive symptoms. This review includes three sections: (1) trials of drugs that are approved for schizophrenia and other illnesses and have also been evaluated for negative symptoms; (2) newer drugs that are not approved and have been evaluated for negative symptoms; and (3) trials of agents for negative symptoms that are currently underway or that have not published results.

2. Studies with approved agents

This review includes published trials evaluating approved agents in negative symptoms of schizophrenia from 1995 to 2012. Our search found multiple published studies and meta-analyses evaluating monotherapy antipsychotics in patients with prominent or predominant negative symptoms. Most of these studies focused on negative symptom improvement in patients with acute schizophrenia and were not included. As a result, this review is limited to studies of clozapine, amisulpride and aripiprazole.

* Corresponding author at: Building 210, Room 130, West Los Angeles VA Healthcare Center, 11301 Wilshire Blvd, Los Angeles, CA 90073, United States. Tel.: +1 310 268 3647. E-mail address: marder@ucla.edu (S.R. Marder).

2.1. Clozapine

Early comparisons of clozapine with other antipsychotics in treatment resistant and acutely ill patients suggested that clozapine may be more effective for treating negative symptoms (Kane et al., 1988). Although clozapine may have been more effective when compared with haloperidol and other antipsychotics with substantial EPS liabilities, this advantage is not apparent with lower EPS agents (Essali et al., 2009). In addition, studies of a longer duration have failed to find advantages of clozapine for negative symptoms. For example, a VA Cooperative Study (Rosenheck et al., 1999) evaluated patients over a one year period and did not find that clozapine had independent effects on negative symptoms. A 29 week comparison of clozapine and haloperidol also failed to find an advantage for clozapine on negative symptoms (Kane et al., 2001).

One study (Breier et al., 1994) evaluated the effects of clozapine on negative symptoms in deficit and non-deficit patients. Deficit patients were those who demonstrated primary rather than secondary negative symptoms. Clozapine was only effective for treating negative symptoms in the non-deficit patients. Taken together, studies of clozapine suggest that it has advantages for negative symptoms that may be secondary to extrapyramidal side effects or inadequately treated positive symptoms. It does not appear to be effective for primary negative symptoms.

2.2. Amisulpride

Amisulpride, a substituted benzamide, is a selective dopamine D2 antagonist approved in multiple European countries for the treatment of positive and negative symptoms of schizophrenia. Four studies (Loo et al., 1997; Danion et al., 1999; Moller, 2001) evaluated amisulpride monotherapy compared with placebo in patients with predominant negative symptoms of schizophrenia. Patients were selected based on their high scores of the Scale for the Assessment of Negative Symptoms (SANS), low scores of the Scale for the Assessment of Positive Symptoms (SAPS) and extrapyramidal (EPS) scores. All studies used the SANS as the primary outcome parameter. Study duration varied from 6 to 26 weeks. The dose of amisulpride ranged from 50 to 300 mg.

All studies showed a significant improvement on negative symptoms compared with placebo. This improvement on negative symptoms was not accompanied by an improvement on positive symptoms in 3 out of 4 studies. In the fourth study (Danion et al., 1999), both negative and positive symptoms improved significantly compared with placebo. In all studies, placebo-treated patients improved significantly on negative symptoms compared with their baseline severity.

One study (Lecrubier et al., 2006) evaluated two doses of olanzapine (5 and 20 mg), 150 mg of amisulpride, and placebo in 244 patients with predominant negative symptoms. The 5 mg olanzapine group showed greater improvement than placebo in negative symptoms, but not the amisulpride groups. Two additional studies (Speller et al., 1997; Moller, 2000), with at least 25 patients per treatment arm, compared the efficacy of amisulpride with low doses antipsychotics (haloperidol and fluphenazine) for negative symptoms. Neither studies showed a superior efficacy of amisulpride.

2.3. Asenapine

Asenapine is approved for the treatment of schizophrenia in most countries. Trials in patients with acute exacerbations suggested that asenapine was more effective than risperidone and haloperidol for negative symptoms (Potkin et al., 2007; Kane et al., 2010a). Two randomized double-blind studies (Buchanan et al., 2012) compared the effect of asenapine with olanzapine in stable patients with predominant negative symptoms of schizophrenia. Predominant negative symptoms were defined as a PANSS (Positive and Negative Scale (Kay et al., 1987)) negative symptom subscale total of 20 or greater and a score of 4 on 3 or

more of the 7 Marder Negative Symptom Factor Scores (Marder et al., 1997). Patients with a score of 4 or greater on more than 2 items of the positive subscale of the PANSS were excluded. Patients needed to be stable for at least 5 months before screening, and have a low score on the Calgary Depression Scale of Schizophrenia (CDSS) and the abbreviated Extrapyramidal Symptoms Rating Scale (ESRS-A). Patients were excluded if they were treated with olanzapine within 5 months prior to screening and failed to benefit on negative symptoms. The prospectively defined primary efficacy variable was the change on the 16-item Negative Symptoms Scale (NSA-16 (Axelrod et al., 1993)) total at Study Week 26. Both test drugs, asenapine and olanzapine, were associated with a significant improvement on persistent negative symptoms compared with baseline. There was no significant difference between asenapine and olanzapine on the total NSA score, the primary outcome parameter.

3. Studies evaluating adjunctive therapy with available medications for the treatment of negative symptoms of schizophrenia

3.1. Antidepressant drugs

A meta-analysis (Singh et al., 2010) evaluated the efficacy of antidepressant adjunctive to antipsychotic therapy. Although there were many case reports and open label trials where antidepressants were added to an antipsychotic, only trials that used well-described criteria for negative symptoms and used double-blind methods to compare the antidepressant to a placebo were included. This resulted in the inclusion of 23 trials from 22 publications ($n = 819$ patients). In most studies, mean scores of negative symptoms were equally or more severe than those of positive symptoms. Study durations varied from 4 to 12 weeks. However, it is unclear whether patients' with depression or severe depressive symptoms were excluded. The overall standardized mean difference was modest (ES-0.33) in favor of antidepressants. Subgroup analyses revealed significant responses for fluoxetine, trazodone and ritanserin. The results supported some benefit of antidepressants in patients with prominent negative symptoms.

3.2. Minocycline

Minocycline is a tetracycline antibiotic with emerging interest for its neuroprotective properties against glutamate neurotoxicity in cell culture and in rodent models of neurodegenerative disorders. Two studies evaluated the effect of minocycline 200 mg/day in early schizophrenia for negative symptoms.

Levkovitz et al. (2010) reported results from a study evaluating the effects of minocycline. Subjects aged 18 to 35 years, within 5 years from their first episode, not receiving antipsychotic treatment for at least 6 months prior to the current episode, and the antipsychotic treatment was initiated less than 14 days prior to the study. A total of 54 patients with PANSS greater than 60 at baseline were randomized to receive minocycline (36) or placebo (18) adjunctive to atypical antipsychotic therapy. Patients treated with minocycline showed greater improvement in negative symptoms measured by the SANS than placebo patients.

Chaudhry et al. (2012) reported results from a study conducted in Brazil and Pakistan. Subjects were aged 18 to 65 years, within 5 years of diagnosis, stable on medication for 4 weeks prior to baseline. A total of 144 participating patients were randomized to receive minocycline (71) or placebo (73) adjunctive to antipsychotic treatment. Data from both participating countries are presented separately, thus making it difficult to extrapolate the overall results. However, the authors concluded that the addition of minocycline to treatment as usual early in the course of schizophrenia improves negative symptoms.

3.3. Dopamine agonists

Dopamine antagonism can produce negative symptoms in healthy volunteers and in animal models (Wise et al., 1978; Artaloytia et al., 2006). For that reason, dopamine agonists may play a role in the treatment of negative symptoms, although the evidence is mixed. Selegiline, a monoamine oxidase-B inhibitor that selectively enhances dopaminergic activity, has been shown to reduce negative symptoms of schizophrenia in open-label studies (Bodkin et al., 1996), and in double-blind, placebo-controlled studies (Bodkin et al., 2005; Amiri et al., 2008), although there are also negative findings (Jungerman et al., 1999).

Lisdexamfetamine, a D-amphetamine pro-drug was evaluated in a 10 week open label study of patients who were stabilized on an antipsychotic (Lasser et al., 2013). Lisdexamfetamine was effective for reducing negative symptoms on the PANSS. This finding clearly needs to be evaluated in a double-blind study.

Modafinil, a stimulant that modestly inhibits dopamine and norepinephrine transporters, leading to increased dopamine and norepinephrine efflux in cortical and other brain areas, has been found to improve negative symptoms in an 8-week double-blind trial when added to risperidone (Arbabi et al., 2012). Two trials evaluated a mixed group of patients with fatigue and negative symptoms where modafinil was added to typical and atypical antipsychotics (Sevy et al., 2005) or clozapine (Freudenreich et al., 2009). Neither trial found an effect on negative symptoms when compared to placebo.

There are also mixed results for the modafinil R-enantiomer armodafinil. A proof of concept trial showed that one of three doses used (200 mg/day) in a 4-week, double-blind, placebo-controlled study with 60 stable schizophrenia patients significantly reduced negative symptoms (Kane et al., 2010b). It is noteworthy that beneficial effects were found on the PANSS but not the SANS and negative symptoms were not required for study subjects. A later larger study with 285 stable schizophrenia patients with negative symptoms was unable to replicate the findings, with no difference among three different armodafinil doses (150 mg, 200 mg, or 250 mg) and placebo in a 24-week trial (Kane et al., 2012).

3.4. Cholinergic agents

Acetylcholinesterase inhibitors increase the synaptic levels of acetylcholine leading to increased stimulation of nicotinic, and muscarinic receptors. These drugs have been found to be effective for treating the cognitive symptoms of dementia and a number are approved for this indication. The effects of acetylcholinesterase inhibitors have been tested in patients with schizophrenia and cognitive impairment. In addition, a number of studies have also assessed their effects on negative symptoms. A Cochrane meta-analysis of the few studies published until 2009 suggests a benefit of an acetylcholinesterase inhibitor plus antipsychotic over antipsychotic and placebo on PANSS negative symptoms (average endpoint score (2 RCTs, $n = 31$, MD -1.69 95% CI -2.80 to -0.57)) (Singh et al., 2012). The authors mention that the many limitations of the studies included in the analysis make this weak evidence and stress the need for more well-designed, large, randomized studies.

In a 12-week, double-blind, placebo-controlled trial with 86 stable patients treated with antipsychotics, galantamine (an acetylcholinesterase inhibitor and a positive allosteric modulator at nicotinic receptors) was found to improve alogia, although there were no significant changes in global negative symptom scores (Conley et al., 2009). In another study with a very similar design, 30 stable schizophrenia patients on risperidone (4–6 mg/day) received donepezil or placebo (Akhondzadeh et al., 2008). Although there was no change in cognitive performance, donepezil-treated patients experienced an improvement in negative symptoms. This effect was not seen in 20 elderly chronic patients (mean age 70.2) in another study comparing donepezil with

placebo. The latter was a 12-week, double-blind, add-on, crossover clinical trial (Mazeh et al., 2006).

4. New pharmacological agents

4.1. Cholinergic

A partial $\alpha 7$ nicotinic receptor agonist, 3-(2,4-dimethoxybenzylidene) anabaseine (DMXB-A) was shown to improve negative symptoms as measured by the SANS in 31 stable schizophrenia patients in a double-blind, crossover, add-on study (Freedman et al., 2008). It is interesting to note that the authors developed this study to target cognitive impairment, but found more consistent effects on negative symptoms. Lieberman et al. (2013) recently reported that a partial $\alpha 7$ nicotinic receptor agonist, TC-5619, led to significant improvement in negative symptoms in a 12-week, double-blind, placebo-controlled, add-on study of 185 schizophrenia patients treated with quetiapine or risperidone monotherapy. This agent is currently being evaluated in a phase 3 trial for negative symptoms (as noted in the next section on trials in progress).

4.2. Glutamate

The “NMDA receptor hypofunction hypothesis” of schizophrenia was first documented in the 1980s (Coyle, 2012). The connection between the NMDAR and schizophrenia was based on the recognition that ketamine and phencyclidine (PCP), non-competitive antagonists of the NMDAR, had psychotomimetic effects and induce schizophrenia-like psychosis, including not only positive but also negative symptoms (Domino et al., 2004; Coyle, 2012). There is also convergent evidence that involves NMDAR-related gene expression and metabolic pathways in schizophrenia (Moghaddam and Javitt, 2012).

A number of novel pre- and postsynaptic mechanisms affecting glutamatergic synaptic transmission have emerged as viable targets for the treatment of negative symptoms in schizophrenia (Javitt, 2012). Several classes of drugs with different mechanisms of action involving the glutamatergic system have achieved promising results. Most of these drugs are metabotropic glutamate agonists, glycine transport inhibitors, drugs that act at the glycine/D-serine and redox sites of the NMDAR and pathways regulating glutamate, glycine/D-serine and glutathione synthesis/release (Javitt, 2012).

4.2.1. NMDAR glycine-site agonists

Initial studies with high doses of glycine as an add-on medication report efficacy for the treatment of negative symptoms in three small 6-week, placebo-controlled, add-on studies (Javitt et al., 1994; Heresco-Levy et al., 1996; Heresco-Levy et al., 2004). Glycine has not been shown to be efficacious when added to clozapine (Potkin et al., 1999; Evins et al., 2000). There have also been positive results for D-cycloserine. In a dose-finding study, Goff et al. (1995) found an inverted-U dose-response curve, with maximum therapeutic efficacy at 50 mg/day and loss of efficacy with doses below 50 mg/day or above 100 mg/day. Positive results have been observed in a group of 23 deficit syndrome patients treated with conventional antipsychotics (Goff et al., 1999) and inpatients who received D-cycloserine (50 mg once weekly) and who were treated with antipsychotics other than clozapine (Goff et al., 2008). On the contrary, D-cycloserine seems to worsen negative symptoms when added to clozapine (Goff et al., 1996; Goff et al., 1999) or has no effect when used at doses other than 50 mg/day (Casella et al., 1994; Rosse et al., 1996; Duncan et al., 2004). There have also been negative results with D-cycloserine used at a dose of 50 mg/day in a 6-month trial (Goff et al., 2005). D-alanine, another endogenous agonist of the NMDA-glycine site, has also been shown to improve negative symptoms in a 6-week, double-blind, placebo-controlled, add-on trial (Tsai et al., 2006).

D-serine was shown to have a beneficial effect in a 6-week, double-blind, placebo-controlled add-on trial (Tsai et al., 1998) but no effect when the same dose (30 mg/kg per day) was added to clozapine in a very similar study (Tsai et al., 1999). In a subsequent study, D-serine was superior to placebo at high doses (50 mg/day) in a 6-week, placebo-controlled, crossover trial in 11 schizophrenia patients treated with clozapine (Goff et al., 2001) and in an open-label study in 42 antipsychotic-stabilized patients at doses of 30, 60, or 120 mg/kg/day (Kantrowitz et al., 2010).

However, the two largest multicenter studies did not replicate the previous findings and no significant effect on negative symptoms was observed as compared with placebo. Buchanan et al. (2007) conducted a 16-week study (n = 157) in which neither glycine (60 mg/day) nor D-cycloserine as add-on drugs improved negative symptoms. The second multicenter study (n = 195) also did not find a positive effect for add-on D-cycloserine (2 g/day) in a 16-week study (Weiser et al., 2012).

4.2.2. Glycine transport inhibitors

Bitopertin (RG1678, Roche), a glycine transport inhibitor was found to induce beneficial effects on negative symptoms in a phase II study. Doses of 10 and 30 mg/day significantly differ from placebo in an 8-week clinical trial (Umbricht et al., 2012). There are three ongoing phase III studies with Bitopertin to assess its efficacy on negative symptoms. In a study comparing the efficacy of D-serine and sarcosine, a glycine transporter-1 (GlyT-1) antagonist, in a 6-week, double-blind, placebo-controlled, add-on trial, sarcosine treatment was better than D-serine in effect sizes for all outcome measures including negative symptoms (Lane et al., 2010).

4.3.3. Metabotropic receptors

Pomaglumedad (LY404039), a highly selective agonist for metabotropic glutamate receptor 2 (mGlu2) and metabotropic glutamate receptor 3 (mGlu3), showed efficacy for the treatment of negative symptoms in a phase II placebo-controlled clinical trial in acute schizophrenia (Patil et al., 2007). Subsequently, four pomaglumedad doses (10, 40, 80, and 160 mg/day) failed to separate from placebo in a failed trial with olanzapine as active comparator (Kinon et al., 2011) (Kinon et al., 2011). More recently a subsequent trial of pomaglumedad indicated that the drug did not separate from placebo (Hopkins, 2013).

6. Trials that have not reported results

We used the www.clinicaltrials.gov site to document trials that are still recruiting or have not reported results. Trials were located using the search terms negative symptoms and schizophrenia. The table lists the trials along with the proposed mechanism, trial duration, and the instrument used to measure negative symptoms. We only included studies that specified improvement on a negative symptom scale as a primary outcome measure.

Agent	Sponsor	Mechanism	Phase	Age	Duration (wks)	Neg symptom instrument
<i>New agents</i>						
TC5619	Targacept	Partial agonist $\alpha 7$ nicotinic receptor	2	18–60	24	SANS
RO4917838 (Bitopertin)	Hoffmann-La Roche	GlyT1 inhibitor	3	>18	24 w/ extension	PANSS neg factor
AMG 747 AZD8529 (completed)	Amgen Astra Zeneca	GlyT1 inhibitor mGlu2/3 receptor PAM	2	18–60	12 4	NSA-16
ORG 25935 (completed)	Schering-Plough	GlyT1 inhibitor	2	18–55	12	SANS
D-serine	Sheba Medical Center Yale	NMDA co-agonist	2	18–65	16	SANS
LY2140023 (completed)	Lilly	mGlu2/3 agonist	2	18–65	12 17	SANS NSA-16
<i>New indications</i>						
Lisdexamfetamine	Shire	Dextroamphetamine prodrug	3	18–65	12	NSA-16

(continued on next page)

There are also some promising results with N-Acetylcysteine (NAC), an activator of the metabotropic receptor mGluR2/3 and the precursor of cysteine, and an indirect antioxidant as a rate-limiting factor in the synthesis of glutathione (GSH). NAC was found to improve core symptoms of schizophrenia, including negative symptoms, as an add-on treatment in a double-blind controlled trial (Berk et al., 2008).

Pregnenolone, a neurosteroid that may exert its effects through positive modulation of NMDA receptors, has been shown to improve negative symptoms as compared with placebo in a small (n = 21) 8-week, double-blind trial in stable patients receiving atypical antipsychotics (Marx et al., 2009). However, a later larger study could not replicate this finding (Ritsner et al., 2010). Two double-blind, placebo-controlled, add-on studies have demonstrated that dehydroepiandrosterone (DHEA) improves negative symptoms at doses between 100 and 150 mg/day (Strous et al., 2003; Strous et al., 2007). However, there are also two negative studies with DHEA (Ritsner et al., 2006; Ritsner et al., 2010).

5. Other approaches

There are a number of other treatment strategies that have been tried in proof of concept studies. Adjunctive intranasal oxytocin for 3 weeks was determined to not be better than placebo for negative symptoms in a small group of 28 subjects (Lee et al., 2013). Ginkgo biloba has also been shown to reduce negative symptoms as compared with placebo in haloperidol treated patients with schizophrenia (Zhang et al., 2001) (Zhang et al., 2001). Recently, 2 mg of folic acid and 400 μ g of vitamin B12 showed efficacy for negative symptoms in a 16-week, randomized, double-blind, placebo-controlled clinical trial when genotype was taken into account (Roffman et al., 2013).

Serotonin 5HT₃ receptor antagonists including ondansetron, granisetron, and tropisetron have also been studied for negative symptoms. Ondansetron was evaluated in two studies as an adjunctive treatment. Both found statistically significant improvements in negative symptoms (Bennett and Vila, 2010). Granisetron and tropisetron were evaluated in separate trials as an adjunctive agent to risperidone. Both studies found improvement in negative symptoms (Khodaei-Ardakani et al., 2013; Noroozian et al., 2013). Tropisetron – which is also an agonist at alpha 7 nicotinic receptors – was also found to improve cognition (Shiina et al., 2010). These trials are certainly interesting, but each has a very modest sample size.

(continued)

Agent	Sponsor	Mechanism	Phase	Age	Duration (wks)	Neg symptom instrument
Rasagiline	U of Maryland	MAO type B inhibitor	4	18–64	12	SANS
Oxytocin	U of Maryland		2	18–64	8	SANS
Galantamine	U of Maryland	Cholinesterase inhibitor, allosteric nicotinic modulator	2	18–64	8	SANS
Citalopram	Imperial College London	SSRI	4	18–75	52	Heinrich's Quality of Life Scale
Pregnenolone	Veterans Affairs and Cornell	Neuroactive steroid	2	21–65	8	SANS
Citalopram and reboxetine	Sant Joan de Deu and CIBERSAM (Spain)	SSRI and NRI	4	18–65	24	PANSS negative
Levodopa/carbidopa	Centre for Addiction and Mental Health – Toronto	Dopamine	2	18–55	8	SANS
<i>Devices</i>						
Intermittent theta burst	Hôpital le Vinatier (France)			18–50	12	SANS
Deep transcranial magnetic stimulation	Shalvata Mental Health Center			18–65	8	SANS
Transcranial direct current stimulation	University of Sao Paulo		2	18–59	4	PANSS

The table demonstrates that negative symptoms in schizophrenia are a very active area for treatment discovery. There are a number of interesting patterns. The most common mechanisms being studied for new entities are alpha 7 nicotinic and glutamatergic. A review of trials for cognition in schizophrenia shows that this is also a very common target. Studies of cognition using these mechanisms commonly have negative symptoms as a secondary outcome measure. These trials are not included in this table. Among studies using devices, there are a number of promising approaches. As would be expected, trials of newer agents are almost always industry sponsored, whereas studies of approved compounds for a new indication are mostly coming from academic sites.

Although the SANS remains the most common scale for measuring negative symptoms, a number of recently initiated studies are utilizing the NSA-16 and PANSS factor scores. Most of the larger industry trials are 12 to 24 weeks. The relative merits of the different measures are discussed in Marder et al. in this issue.

7. Discussion

This review indicates that clinicians who are treating negative symptoms in schizophrenia have a number of current options for treating negative symptoms in schizophrenia. These options include selecting amisulpride as an antipsychotic (where it is available), or adding one of a number of adjunctive or co-medications to an antipsychotic. Among promising adjunctive medications, there is some modest evidence suggesting that adding an antidepressant may have some benefit. The evidence for other co-medications such as minocycline, modafinil, armodafinil, and galantamine is more limited. A number of promising drugs are currently in different stages of development and it is possible that in the near future more effective adjunctive drugs will be available. At this time, drugs that act at the NMDA and alpha 7 nicotinic receptors are receiving the most attention.

The studies evaluating antipsychotic monotherapy treatment have some methodological limitations. The most serious issue is the ability to demonstrate a direct effect on negative symptoms that is not mediated through alleviation of positive, mood symptoms or antipsychotic-induced motor effects. Studies to evaluate the effect of amisulpride on negative symptoms recruited patients with predominant negative symptoms and minimal positive and mood symptoms. The direct effect observed with amisulpride on primary negative symptoms is the most convincing. However, the impact of this improvement on patient functioning was not investigated. In addition, the study population that received placebo was not receiving an antipsychotic. As a result, the improvement in negative symptoms on amisulpride could have resulted from improvement in low level positive symptoms.

Studies evaluating the effect of therapies adjunctive to antipsychotics also have important limitations. First, these studies enrolled very heterogeneous patient populations using different criteria for persistent negative symptoms. Although some antidepressants have shown an effect on negative symptoms, it is unclear whether this is a direct effect on negative symptoms or if it is mediated through an improvement of mood symptoms. Finally, the effect of minocycline on negative symptoms early in the disease is inconclusive, and clearly needs to be replicated in larger trials.

Role of funding source

There was no funding source for the work described in this article.

Contributors

Drs. Marder, Arango, and Garibaldi each contributed to the literature review and the conclusions in this article.

Conflict of interests

Dr. Arango has been a consultant to or has received honoraria or grants from Abbot, AMGEN, AstraZeneca, Bristol-Myers Squibb, Caja Navarra, CIBERSAM, Fundación Alicia Koplowitz, Instituto de Salud Carlos III, Janssen Cilag, Lundbeck, Merck, Ministerio de Ciencia e Innovación, Ministerio de Sanidad, Ministerio de Economía y Competitividad, Mutua Madrileña, Otsuka, Pfizer, Roche, Servier, Takeda and Schering Plough.

Stephen R. Marder has consulted for Abbott, Boehringer Ingelheim, Bristol Meyers Squibb, Genentech, Roche, Otsuka, Pfizer, EnVivo, Targacept, Lundbeck, Shire, and Jazz. He had research support from Glaxo Smith Kline, Novartis, Sunovion, Psychogenics, and Amgen.

George Garibaldi is a full time employee of F. Hoffmann-La Roche, Ltd.

Acknowledgments

None.

References

- Akhondzadeh, S., Gerami, M., Noroozian, M., Karamghadiri, N., Ghoreishi, A., Abbasi, S.H., Rezazadeh, S.A., 2008. A 12-week, double-blind, placebo-controlled trial of donepezil adjunctive treatment to risperidone in chronic and stable schizophrenia. *Prog. Neuropsychopharmacol. Biol. Psychiatry* 32, 1810–1815.
- Amiri, A., Noorbala, A.A., Nejatisafa, A.A., Ghoreishi, A., Derakhshan, M.K., Khodaei-Ardakani, M.R., Hajiazim, M., Raznahan, M., Akhondzadeh, S., 2008. Efficacy of selegiline add on therapy to risperidone in the treatment of the negative symptoms of schizophrenia: a double-blind randomized placebo-controlled study. *Hum. psychopharmacol.* 23, 79–86.
- Arango, C., Buchanan, R.W., Kirkpatrick, B., Carpenter, W.T., 2004. The deficit syndrome in schizophrenia: implications for the treatment of negative symptoms. *Eur. Psychiatry* 19, 21–26.
- Arbabi, M., Bagheri, M., Rezaei, F., Ahmadi-Abhari, S.A., Tabrizi, M., Khalighi-Sigaroudi, F., Akhondzadeh, S., 2012. A placebo-controlled study of the modafinil added to risperidone in chronic schizophrenia. *Psychopharmacology* 220, 591–598.
- Artaloytia, J.F., Arango, C., Lahti, A., Sanz, J., Pascual, A., Cubero, P., Prieto, D., Palomo, T., 2006. Negative signs and symptoms secondary to antipsychotics: a double-blind, randomized trial of a single dose of placebo, haloperidol, and risperidone in healthy volunteers. *Am. J. Psychiatry* 163, 488–493.
- Axelrod, B.N., Goldman, R.S., Alphs, L.D., 1993. Validation of the 16-item Negative Symptom Assessment. *J. Psychiatr. Res.* 27, 253–258.

- Bennett, A.C., Vila, T.M., 2010. The role of ondansetron in the treatment of schizophrenia. *Ann. Pharmacother.* 44, 1301–1306.
- Berk, M., Copolov, D., Dean, O., Lu, K., Jeavons, S., Schapkaite, I., Anderson-Hunt, M., Judd, F., Katz, F., Katz, P., Ording-Jespersen, S., Little, J., Conus, P., Cuenod, M., Do, K.Q., Bush, A.I., 2008. N-acetyl cysteine as a glutathione precursor for schizophrenia—a double-blind, randomized, placebo-controlled trial. *Biol. Psychiatry* 64, 361–368.
- Bobes, J., Arango, C., Garcia-Garcia, M., Rejas, J., 2010. Prevalence of negative symptoms in outpatients with schizophrenia spectrum disorders treated with antipsychotics in routine clinical practice: findings from the CLAMORS study. *J. Clin. Psychiatry* 71, 280–286.
- Bodkin, J.A., Cohen, B.M., Salomon, M.S., Cannon, S.E., Zornberg, G.L., Cole, J.O., 1996. Treatment of negative symptoms in schizophrenia and schizoaffective disorder by selegiline augmentation of antipsychotic medication. A pilot study examining the role of dopamine. *J. Nerv. Ment. Dis.* 184, 295–301.
- Bodkin, J.A., Siris, S.G., Bermanzohn, P.C., Hennen, J., Cole, J.O., 2005. Double-blind, placebo-controlled, multicenter trial of selegiline augmentation of antipsychotic medication to treat negative symptoms in outpatients with schizophrenia. *Am. J. Psychiatry* 162, 388–390.
- Breier, A., Buchanan, R.W., Kirkpatrick, B., Davis, O.R., Irish, D., Summerfelt, A., Carpenter Jr., W.T., 1994. Effects of clozapine on positive and negative symptoms in outpatients with schizophrenia. *Am. J. Psychiatry* 151, 20–26.
- Buchanan, R.W., Javitt, D.C., Marder, S.R., Schooler, N.R., Gold, J.M., McMahon, R.P., Heresco-Levy, U., Carpenter, W.T., 2007. The Cognitive and Negative Symptoms in Schizophrenia Trial (CONSIST): the efficacy of glutamatergic agents for negative symptoms and cognitive impairments. *Am. J. Psychiatry* 164, 1593–1602.
- Buchanan, R.W., Panagides, J., Zhao, J., Phiri, P., den Hollander, W., Ha, X., Kouassi, A., Alphas, L., Schooler, N., Szege, A., Cazorla, P., 2012. Aripiprazole versus olanzapine in people with persistent negative symptoms of schizophrenia. *J. Clin. Psychopharmacol.* 32, 36–45.
- Carpenter Jr., W.T., Davis, J.M., 2012. Another view of the history of antipsychotic drug discovery and development. *Mol. Psychiatry* 17, 1168–1173.
- Cascella, N.G., Macciardi, F., Cavallini, C., Smeraldi, E., 1994. d-cycloserine adjuvant therapy to conventional neuroleptic treatment in schizophrenia: an open-label study. *J. Neural Transm.* 95, 105–111 (General section).
- Chaudhry, I.B., Hallak, J., Husain, N., Minhas, F., Stirling, J., Richardson, P., Dursun, S., Dunn, G., Deakin, B., 2012. Minoxidil benefits negative symptoms in early schizophrenia: a randomised double-blind placebo-controlled clinical trial in patients on standard treatment. *J. Psychopharmacol.* 26, 1185–1193.
- Conley, R.R., Boggs, D.L., Kelly, D.L., McMahon, R.P., Dickinson, D., Feldman, S., Ball, M.P., Buchanan, R.W., 2009. The effects of galantamine on psychopathology in chronic stable schizophrenia. *Clin. Neuropharmacol.* 32, 69–74.
- Coyle, J.T., 2012. NMDA receptor and schizophrenia: a brief history. *Schizophr. Bull.* 38, 920–926.
- Danion, J.M., Rein, W., Fleurot, O., 1999. Improvement of schizophrenic patients with primary negative symptoms treated with amisulpride. Amisulpride Study Group. *Am. J. Psychiatry* 156, 610–616.
- Domino, E.F., Mirzoyan, D., Tsukada, H., 2004. N-methyl-D-aspartate antagonists as drug models of schizophrenia: a surprising link to tobacco smoking. *Prog. Neuropsychopharmacol. Biol. Psychiatry* 28, 801–811.
- Duncan, E.J., Szilagyi, S., Schwartz, M.P., Bugarski-Kirola, D., Kunzova, A., Negi, S., Stephanides, M., Efferen, T.R., Angrist, B., Peselow, E., Corwin, J., Gonzenbach, S., Rotrosen, J.P., 2004. Effects of D-cycloserine on negative symptoms in schizophrenia. *Schizophr. Res.* 71, 239–248.
- Essali, A., Al-Haj Haasan, N., Li, C., Rathbone, J., 2009. Clozapine versus typical neuroleptic medication for schizophrenia. *Cochrane Database Syst. Rev.* 1–219 (CD000059).
- Evins, A.E., Fitzgerald, S.M., Wine, L., Rosselli, R., Goff, D.C., 2000. Placebo-controlled trial of glycine added to clozapine in schizophrenia. *Am. J. Psychiatry* 157, 826–828.
- Freedman, R., Olincy, A., Buchanan, R.W., Harris, J.G., Gold, J.M., Johnson, L., Allensworth, D., Guzman-Bonilla, A., Clement, B., Ball, M.P., Kutnick, J., Pender, V., Martin, L.F., Stevens, K.E., Wagner, B.D., Zerbe, G.O., Soti, F., Kem, W.R., 2008. Initial phase 2 trial of a nicotinic agonist in schizophrenia. *Am. J. Psychiatry* 165, 1040–1047.
- Freudenreich, O., Henderson, D.C., Macklin, E.A., Evins, A.E., Fan, X., Cather, C., Walsh, J.P., Goff, D.C., 2009. Modafinil for clozapine-treated schizophrenia patients: a double-blind, placebo-controlled pilot trial. *J. Clin. Psychiatry* 70, 1674–1680.
- Goff, D.C., Tsai, G., Manoach, D.S., Coyle, J.T., 1995. Dose-finding trial of D-cycloserine added to neuroleptics for negative symptoms in schizophrenia. *Am. J. Psychiatry* 152, 1213–1215.
- Goff, D.C., Tsai, G., Manoach, D.S., Flood, J., Darby, D.G., Coyle, J.T., 1996. D-cycloserine added to clozapine for patients with schizophrenia. *Am. J. Psychiatry* 153, 1628–1630.
- Goff, D.C., Tsai, G., Levitt, J., Amico, E., Manoach, D., Schoenfeld, D.A., Hayden, D.L., McCarley, R., Coyle, J.T., 1999. A placebo-controlled trial of D-cycloserine added to conventional neuroleptics in patients with schizophrenia. *Arch. Gen. Psychiatry* 56, 21–27.
- Goff, D.C., Freudenreich, O., Evins, A.E., 2001. Augmentation strategies in the treatment of schizophrenia. *CNS Spectr.* 6 (904), 907–911.
- Goff, D.C., Herz, L., Posever, T., Shih, V., Tsai, G., Henderson, D.C., Freudenreich, O., Evins, A.E., Yovel, I., Zhang, H., Schoenfeld, D., 2005. A six-month, placebo-controlled trial of D-cycloserine co-administered with conventional antipsychotics in schizophrenia patients. *Psychopharmacology* 179, 144–150.
- Goff, D.C., Cather, C., Gottlieb, J.D., Evins, A.E., Walsh, J., Raeke, L., Otto, M.W., Schoenfeld, D., Green, M.F., 2008. Once-weekly D-cycloserine effects on negative symptoms and cognition in schizophrenia: an exploratory study. *Schizophr. Res.* 106, 320–327.
- Heresco-Levy, U., Silipo, G., Javitt, D.C., 1996. Glycinergic augmentation of NMDA receptor-mediated neurotransmission in the treatment of schizophrenia. *Psychopharmacol. Bull.* 32, 731–740.
- Heresco-Levy, U., Ermilov, M., Lichtenberg, P., Bar, G., Javitt, D.C., 2004. High-dose glycine added to olanzapine and risperidone for the treatment of schizophrenia. *Biol. Psychiatry* 55, 165–171.
- Hopkins, C.R., 2013. Is there a path forward for mGlu(2) positive allosteric modulators for the treatment of schizophrenia? *ACS Chem. Neurosci.* 4, 211–213.
- Hyman, S.E., Fenton, W.S., 2003. Medicine. What are the right targets for psychopharmacology? *Science* 299, 350–351.
- Javitt, D.C., 2012. Twenty-five years of glutamate in schizophrenia: are we there yet? *Schizophr. Bull.* 38, 911–913.
- Javitt, D.C., Zylberman, I., Zukin, S.R., Heresco-Levy, U., Lindenmayer, J.P., 1994. Amelioration of negative symptoms in schizophrenia by glycine. *Am. J. Psychiatry* 151, 1234–1236.
- Jungerman, T., Rabinowitz, D., Klein, E., 1999. Deprenyl augmentation for treating negative symptoms of schizophrenia: a double-blind, controlled study. *J. Clin. Psychopharmacol.* 19, 522–525.
- Kane, J., Honigfeld, G., Singer, J., Meltzer, H., 1988. Clozapine for the treatment-resistant schizophrenic. A double-blind comparison with chlorpromazine. *Arch. Gen. Psychiatry* 45, 789–796.
- Kane, J.M., Marder, S.R., Schooler, N.R., Wirshing, W.C., Umbricht, D., Baker, R.W., Wirshing, D.A., Safferman, A., Ganguli, R., McMeniman, M., Borenstein, M., 2001. Clozapine and haloperidol in moderately refractory schizophrenia: a 6-month randomized and double-blind comparison. *Arch. Gen. Psychiatry* 58, 965–972.
- Kane, J.M., Cohen, M., Zhao, J., Alphas, L., Panagides, J., 2010a. Efficacy and safety of aripiprazole in a placebo- and haloperidol-controlled trial in patients with acute exacerbation of schizophrenia. *J. Clin. Psychopharmacol.* 30, 106–115.
- Kane, J.M., D'Souza, D.C., Patkar, A.A., Youakim, J.M., Tiller, J.M., Yang, R., Keefe, R.S., 2010b. Armodafinil as adjunctive therapy in adults with cognitive deficits associated with schizophrenia: a 4-week, double-blind, placebo-controlled study. *The Journal of clinical psychiatry* 71, 1475–1481.
- Kane, J.M., Yang, R., Youakim, J.M., 2012. Adjunctive armodafinil for negative symptoms in adults with schizophrenia: a double-blind, placebo-controlled study. *Schizophr. Res.* 135, 116–122.
- Kantrowitz, J.T., Malhotra, A.K., Cornblatt, B., Silipo, G., Balla, A., Suckow, R.F., D'Souza, C., Saksa, J., Woods, S.W., Javitt, D.C., 2010. High dose D-serine in the treatment of schizophrenia. *Schizophr. Res.* 121, 125–130.
- Kay, S.R., Fiszbein, A., Opler, L.A., 1987. The positive and negative syndrome scale (PANSS) for schizophrenia. *Schizophr. Bull.* 13, 261–276.
- Khodaei-Ardakani, M.R., Seddighi, S., Modabbernia, A., Rezaei, F., Salehi, B., Ashrafi, M., Shams-Alizadeh, N., Mohammad-Karimi, M., Esfandiari, G.R., Hajiaghah, R., Akhondzadeh, S., 2013. Granisetron as an add-on to risperidone for treatment of negative symptoms in patients with stable schizophrenia: randomized double-blind placebo-controlled study. *J. Psychiatr. Res.* 47, 472–478.
- Kinon, B.J., Zhang, L., Millen, B.A., Osuntokun, O.O., Williams, J.E., Kollack-Walker, S., Jackson, K., Kryzhanovskaya, L., Jarkova, N., Group, H.S., 2011. A multicenter, inpatient, phase 2, double-blind, placebo-controlled dose-ranging study of LY2140023 monohydrate in patients with DSM-IV schizophrenia. *J. Clin. Psychopharmacol.* 31, 349–355.
- Kirkpatrick, B., Fenton, W.S., Carpenter Jr., W.T., Marder, S.R., 2006. The NIMH-MATRICS consensus statement on negative symptoms. *Schizophr. Bull.* 32, 214–219.
- Lane, H.Y., Lin, C.H., Huang, Y.J., Liao, C.H., Chang, Y.C., Tsai, G.E., 2010. A randomized, double-blind, placebo-controlled comparison study of sarcosine (N-methylglycine) and D-serine add-on treatment for schizophrenia. *Int. J. Neuropsychopharmacol.* 13, 451–460.
- Lasser, R.A., Dirks, B., Nasrallah, H., Kirsch, C., Gao, J., Pucci, M.L., Knesevich, M.A., Lindenmayer, J.P., 2013. Adjunctive lisdexamfetamine dimesylate therapy in adult outpatients with predominant negative symptoms of schizophrenia: open-label and randomized-withdrawal phases. *Neuropsychopharmacology*.
- Leclercq, Y., Quintin, P., Bouhassira, M., Perrin, E., Lancron, S., 2006. The treatment of negative symptoms and deficit states of chronic schizophrenia: olanzapine compared to amisulpride and placebo in a 6-month double-blind controlled clinical trial. *Acta Psychiatr. Scand.* 114, 319–327.
- Lee, M.R., Wehring, H.J., McMahon, R.P., Linthicum, J., Cascella, N., Liu, F., Bellack, A., Buchanan, R.W., Strauss, G.P., Contoreggi, C., Kelly, D.L., 2013. Effects of adjunctive intranasal oxytocin on olfactory identification and clinical symptoms in schizophrenia: results from a randomized double blind placebo controlled pilot study. *Schizophr. Res.* 145, 110–115.
- Levkovitz, Y., Mendlovich, S., Riwkes, S., Braw, Y., Levkovitch-Verbin, H., Gal, G., Fennig, S., Treves, I., Kron, S., 2010. A double-blind, randomized study of minocycline for the treatment of negative and cognitive symptoms in early-phase schizophrenia. *The Journal of clinical psychiatry* 71, 138–149.
- Lieberman, J.A., Dunbar, G., Segreti, A.C., Girgis, R.R., Seoane, F., Beaver, J.S., Duan, N., Hosford, D.A., 2013. A randomized exploratory trial of an alpha-7 nicotinic receptor agonist (TC-5619) for cognitive enhancement in schizophrenia. *Neuropsychopharmacology* 38, 968–975.
- Loo, H., Poirier-Litre, M.F., Theron, M., Rein, W., Fleurot, O., 1997. Amisulpride versus placebo in the medium-term treatment of the negative symptoms of schizophrenia. *Br. J. Psychiatry* 170, 18–22.
- Marder, S.R., Davis, J.M., Chouinard, G., 1997. The effects of risperidone on the five dimensions of schizophrenia derived by factor analysis: combined results of the North American trials. *J. Clin. Psychiatry* 58, 538–546.
- Marder, S.R., Daniel, D.G., Alphas, L., Awad, A.G., Keefe, R.S., 2011. Methodological issues in negative symptom trials. *Schizophr. Bull.* 37, 250–254.
- Marx, C.E., Keefe, R.S., Buchanan, R.W., Hamer, R.M., Kilts, J.D., Bradford, D.W., Strauss, J.L., Naylor, J.C., Payne, V.M., Lieberman, J.A., Savitz, A.J., Leimone, L.A., Dunn, L., Porcu, P., Morrow, A.L., Shampine, L.J., 2009. Proof-of-concept trial with the neurosteroid pregnenolone targeting cognitive and negative symptoms in schizophrenia. *Neuropsychopharmacology* 34, 1885–1903.
- Mazeh, D., Zemishlani, H., Barak, Y., Mirecki, I., Paleacu, D., 2006. Donepezil for negative signs in elderly patients with schizophrenia: an add-on, double-blind, crossover, placebo-controlled study. *Int. Psychogeriatr.* 18, 429–436.

- Moghaddam, B., Javitt, D., 2012. From revolution to evolution: the glutamate hypothesis of schizophrenia and its implication for treatment. *Neuropsychopharmacology* 37, 4–15.
- Moller, H.J., 2000. Amisulpride: a review of its efficacy in schizophrenia. *Acta Psychiatr. Scand. Suppl.* 400, 17–22.
- Moller, H.J., 2001. Amisulpride: efficacy in the management of chronic patients with predominant negative symptoms of schizophrenia. *Eur. Arch. Psychiatry Clin. Neurosci.* 251, 217–224.
- Noroozian, M., Ghasemi, S., Hosseini, S.M., Modabbarnia, A., Khodaie-Ardakani, M.R., Mirshafiee, O., Farokhnia, M., Tajdini, M., Rezaei, F., Salehi, B., Ashrafi, M., Yekehtaz, H., Tabrizi, M., Akhondzadeh, S., 2013. A placebo-controlled study of tropisetron added to risperidone for the treatment of negative symptoms in chronic and stable schizophrenia. *Psychopharmacology* 228, 595–602.
- Patil, S.T., Zhang, L., Martenyi, F., Lowe, S.L., Jackson, K.A., Andreev, B.V., Avedisova, A.S., Bardenstein, L.M., Gurovich, I.Y., Morozova, M.A., Mosolov, S.N., Neznanov, N.G., Reznik, A.M., Smulevich, A.B., Tochilov, V.A., Johnson, B.G., Monn, J.A., Schoepp, D.D., 2007. Activation of mGlu2/3 receptors as a new approach to treat schizophrenia: a randomized phase 2 clinical trial. *Nat. Med.* 13, 1102–1107.
- Potkin, S.G., Jin, Y., Bunney, B.G., Costa, J., Gulasekaram, B., 1999. Effect of clozapine and adjunctive high-dose glycine in treatment-resistant schizophrenia. *Am. J. Psychiatry* 156, 145–147.
- Potkin, S.G., Cohen, M., Panagides, J., 2007. Efficacy and tolerability of asenapine in acute schizophrenia: a placebo- and risperidone-controlled trial. *J. Clin. Psychiatry* 68, 1492–1500.
- Rabinowitz, J., Levine, S.Z., Garibaldi, G., Bugarski-Kirola, D., Berardo, C.G., Kapur, S., 2012. Negative symptoms have greater impact on functioning than positive symptoms in schizophrenia: analysis of CATIE data. *Schizophr. Res.* 137, 147–150.
- Ritsner, M.S., Gibel, A., Ratner, Y., Tsinovoy, G., Strous, R.D., 2006. Improvement of sustained attention and visual and movement skills, but not clinical symptoms, after dehydroepiandrosterone augmentation in schizophrenia: a randomized, double-blind, placebo-controlled, crossover trial. *J. Clin. Psychopharmacol.* 26, 495–499.
- Ritsner, M.S., Gibel, A., Shleifer, T., Boguslavsky, I., Zayed, A., Maayan, R., Weizman, A., Lerner, V., 2010. Pregnenolone and dehydroepiandrosterone as an adjunctive treatment in schizophrenia and schizoaffective disorder: an 8-week, double-blind, randomized, controlled, 2-center, parallel-group trial. *The Journal of clinical psychiatry* 71, 1351–1362.
- Roffman, J.L., Lamberti, J.S., Achtyes, E., Macklin, E.A., Galendez, G.C., Raeke, L.H., Silverstein, N.J., Smoller, J.W., Hill, M., Goff, D.C., 2013. Randomized multicenter investigation of folate plus vitamin B12 supplementation in schizophrenia. *JAMA psychiatry* 1–9.
- Rosenheck, R., Dunn, L., Peszke, M., Cramer, J., Xu, W., Thomas, J., Charney, D., 1999. Impact of clozapine on negative symptoms and on the deficit syndrome in refractory schizophrenia. Department of Veterans Affairs Cooperative Study Group on Clozapine in Refractory Schizophrenia. *Am. J. Psychiatry* 156, 88–93.
- Rosse, R.B., Fay-McCarthy, M., Kendrick, K., Davis, R.E., Deutsch, S.I., 1996. D-cycloserine adjunct therapy to molindone in the treatment of schizophrenia. *Clin. Neuropharmacol.* 19, 444–450.
- Sevy, S., Rosenthal, M.H., Alvir, J., Meyer, S., Visweswarajah, H., Gunduz-Bruce, H., Schooler, N.R., 2005. Double-blind, placebo-controlled study of modafinil for fatigue and cognition in schizophrenia patients treated with psychotropic medications. *The Journal of clinical psychiatry* 66, 839–843.
- Shiina, A., Shirayama, Y., Niitsu, T., Hashimoto, T., Yoshida, T., Hasegawa, T., Haraguchi, T., Kanahara, N., Shiraishi, T., Fujisaki, M., Fukami, G., Nakazato, M., Iyo, M., Hashimoto, K., 2010. A randomised, double-blind, placebo-controlled trial of tropisetron in patients with schizophrenia. *Ann. Gen. psychiatry* 9, 27.
- Singh, S.P., Singh, V., Kar, N., Chan, K., 2010. Efficacy of antidepressants in treating the negative symptoms of chronic schizophrenia: meta-analysis. *Br. J. Psychiatry* 197, 174–179.
- Singh, J., Kour, K., Jayaram, M.B., 2012. Acetylcholinesterase inhibitors for schizophrenia. *Cochrane Database Syst. Rev.* 1 (CD007967).
- Speller, J.C., Barnes, T.R., Curson, D.A., Pantelis, C., Alberts, J.L., 1997. One-year, low-dose neuroleptic study of in-patients with chronic schizophrenia characterised by persistent negative symptoms. Amisulpride v. haloperidol. *Br. J. Psychiatry* 171, 564–568.
- Strous, R.D., Maayan, R., Lapidus, R., Stryjer, R., Lustig, M., Kotler, M., Weizman, A., 2003. Dehydroepiandrosterone augmentation in the management of negative, depressive, and anxiety symptoms in schizophrenia. *Arch. Gen. Psychiatry* 60, 133–141.
- Strous, R.D., Stryjer, R., Maayan, R., Gal, G., Viglin, D., Katz, E., Eisner, D., Weizman, A., 2007. Analysis of clinical symptomatology, extrapyramidal symptoms and neurocognitive dysfunction following dehydroepiandrosterone (DHEA) administration in olanzapine treated schizophrenia patients: a randomized, double-blind placebo controlled trial. *Psychoneuroendocrinology* 32, 96–105.
- Tsai, G., Yang, P., Chung, L.C., Lange, N., Coyle, J.T., 1998. D-serine added to antipsychotics for the treatment of schizophrenia. *Biol. Psychiatry* 44, 1081–1089.
- Tsai, G.E., Yang, P., Chung, L.C., Tsai, I.C., Tsai, C.W., Coyle, J.T., 1999. D-serine added to clozapine for the treatment of schizophrenia. *Am. J. Psychiatry* 156, 1822–1825.
- Tsai, G.E., Yang, P., Chang, Y.C., Chong, M.Y., 2006. D-alanine added to antipsychotics for the treatment of schizophrenia. *Biol. Psychiatry* 59, 230–234.
- Umbricht, D., Lentz, E., Lalonde, J., Santarelli, L.A., 2012. Post-hoc analysis of the negative symptom factor score in a proof-of-concept study with the glycine reuptake inhibitor (GRI), bitopertin, in schizophrenia. *Eur. Neuropsychopharmacol.* 22, 311.
- Weiser, M., Heresco-Levy, U., Davidson, M., Javitt, D.C., Werbeloff, N., Gershon, A.A., Abramovich, Y., Amital, D., Doron, A., Konas, S., Levkovitz, Y., Liba, D., Teitelbaum, A., Mashiach, M., Zimmerman, Y., 2012. A multicenter, add-on randomized controlled trial of low-dose d-serine for negative and cognitive symptoms of schizophrenia. *The Journal of clinical psychiatry* 73, e728–e734.
- Wise, R.A., Spindler, J., deWit, H., Gerberg, G.J., 1978. Neuroleptic-induced “anhedonia” in rats: pimozide blocks reward quality of food. *Science* 201, 262–264.
- Zhang, X.Y., Zhou, D.F., Zhang, P.Y., Wu, G.Y., Su, J.M., Cao, L.Y., 2001. A double-blind, placebo-controlled trial of extract of Ginkgo biloba added to haloperidol in treatment-resistant patients with schizophrenia. *The Journal of clinical psychiatry* 62, 878–883.