

# Sigma-1 Receptor Agonists as Therapeutic Drugs for Cognitive Impairment in Neuropsychiatric Diseases

Tomihisa Niitsu<sup>1</sup>, Masaomi Iyo<sup>2,3</sup> and Kenji Hashimoto<sup>3,\*</sup>

<sup>1</sup>Research Center for Child Mental Development, Chiba University Graduate School of Medicine, 1-8-1 Inohana, Chiba 260-8670, Japan; <sup>2</sup>Department of Psychiatry, Chiba University Graduate School of Medicine, 1-8-1 Inohana, Chiba 260-8670, Japan; <sup>3</sup>Division of Clinical Neuroscience, Chiba University Center for Forensic Mental Health, 1-8-1 Inohana, Chiba 260-8670, Japan

**Abstract:** Cognitive impairment is a core feature of patients with neuropsychiatric diseases such as schizophrenia and psychotic depression. The drugs currently used to treat cognitive impairment have significant limitations, ensuring that the search for more effective therapies remains active. Endoplasmic reticulum protein sigma-1 receptors are unique binding sites in the brain that exert a potent effect on multiple neurotransmitter systems. Accumulating evidence suggests that sigma-1 receptors play a role in both the pathophysiology of neuropsychiatric diseases, and the mechanistic action of some therapeutic drugs, such as the selective serotonin reuptake inhibitors (SSRIs), donepezil and neurosteroids. Among SSRIs, fluvoxamine, a potent sigma-1 receptor agonist, has the highest affinity at sigma-1 receptors. Sigma-1 receptor agonists greatly potentiate nerve-growth factor (NGF)-induced neurite outgrowth in PC12 cells, an effect that is antagonized by treatment with the selective sigma-1 receptor antagonist NE-100. Furthermore, phencyclidine (PCP)-induced cognitive impairment, associated with animal models of schizophrenia is significantly improved by sub-chronic administration of sigma-1 receptor agonists such as fluvoxamine, SA4503 (cutamesine) and donepezil. This effect is antagonized by co-administration of NE-100. A positron emission tomography (PET) study using the specific sigma-1 receptor ligand [<sup>11</sup>C]SA4503 demonstrates that fluvoxamine and donepezil bind to sigma-1 receptors in the healthy human brain. In clinical studies, some sigma-1 receptor agonists, including fluvoxamine, donepezil and neurosteroids, improve cognitive impairment and clinical symptoms in neuropsychiatric diseases. In this article, we review the recent findings on sigma-1 receptor agonists as potential therapeutic drugs for the treatment of cognitive impairment in schizophrenia and psychotic depression.

**Keywords:** Sigma-1 receptor, Cognition, Schizophrenia, Psychotic depression, Delirium.

## 1. INTRODUCTION

Cognitive impairment is a common symptom in patients with neuropsychiatric disorders such as schizophrenia and major depressive disorder. Schizophrenia is characterized by three distinct symptom clusters: positive symptoms (*e.g.*, hallucinations and delusions), negative symptoms (*e.g.*, affective flattening, alogia and avolition), and cognitive impairment (*e.g.*, severe deterioration of working memory and attention). Cognitive impairment is a core feature of schizophrenia and its presence predicts both vocational and social disabilities for patients [1-4]. While positive symptoms are greatly improved with atypical antipsychotic medication, cognitive impairment is not greatly improved by therapy [5].

Cognitive impairment is also a ubiquitous and characteristic feature of major depressive disorder [6], often persisting despite otherwise effective antidepressant therapy. In addition, cognitive impairment can also be an adverse effect ("cognitive toxicity"), as a direct consequence of some antidepressant therapy [7]. The severity of major depression also has a bearing on the extent and magnitude of cognitive, psychomotor and memory impairment [8, 9]. Given the relative commonality of this disorder, and the lack of highly effective therapy, there is still a strong need for the development of pharmacological agents to improve cognitive impairment associated with neuropsychiatric disorders [10-14].

Sigma-1 receptors, discovered in 1976 by Martin and co-workers [15], were cloned in 1996 [16], and characterized as having an endoplasmic reticulum (ER) retention signal. Sigma-1 receptors on the ER regulate Ca<sup>2+</sup> signaling via inositol 1,4,5-triphosphate (IP<sub>3</sub>) receptors on the ER [17]. Interestingly, the ER protein sigma-1 receptors are Ca<sup>2+</sup>-sensitive and are ligand-operated receptor chaperones at the mitochondrial associated ER membrane [18, 19].

The ER luminal domain of the sigma-1 receptor possesses robust chaperone activity that prevents the aggregation of a variety of proteins *in vitro*, which stabilizes the ER Ca<sup>2+</sup> channel IP<sub>3</sub> receptor *in vivo* [18, 20]. Sigma-1 receptors are predominantly expressed at the mitochondrial associated ER membrane, thereby regulating the IP<sub>3</sub> receptor-mediated Ca<sup>2+</sup> influx from the ER to the mitochondria [18]. Sigma-1 receptors modulate ATP production and bioenergetics within cells [20].

Sigma-1 receptors regulate a number of neurotransmitter systems, including the glutamatergic, dopaminergic, serotonergic, noradrenergic, and cholinergic systems. Several lines of evidence implicate the role of sigma-1 receptors in the pathophysiology of neuropsychiatric disorders, such as mood disorders, anxiety disorders, and schizophrenia, and suggest that the receptor ligands may be potential therapeutic agents for these diseases [21-34]. This article reviews and discusses the role of sigma-1 receptor agonists as therapeutic agents, to improve cognitive impairment in neuropsychiatric disorders, particularly schizophrenia and major depressive disorder.

## 2. SIGMA-1 RECEPTOR AGONISTS AND NEUROPLASTICITY

Sigma-1 receptors play a role in synaptogenesis and myelination in the brain [35, 36], processes implicated in the pathology of schizophrenia [37-40] and depression [41, 39].

Tricyclic antidepressants (TCA) and selective serotonin reuptake inhibitors (SSRIs) are widely used as therapeutic drugs for major depressive disorder. Although all SSRIs share the common function of the blocking serotonin transporters, leading to elevated serotonin levels throughout the central nervous system (CNS), it is well known that their secondary pharmacology is heterogeneous [42-44]. We have reported that some SSRIs possess high to moderate affinity for sigma-1, but not for sigma-2 receptors Table 1 [45]. *In vitro* experiments suggest that the affinity of SSRIs for sigma-1 receptors is as follows: fluvoxamine > sertraline >

\*Address correspondence to this author at the Division of Clinical Neuroscience, Chiba University Center for Forensic Mental Health, 1-8-1 Inohana, Chiba 260-8670, Japan; Tel: +81-43-226-2517; Fax: +81-43-226-2561; E-mail: hashimoto@faculty.chiba-u.jp

**Table 1.** *In Vitro* Affinity of Various Antidepressants for Rat Sigma-1 Binding Sites (from Ref. [45])

Drug	Ki (nM)		Ki ratio
SSRIs	Sigma-1	Sigma-2	(Sigma-2/Sigma-1)
Fluvoxamine	36	8,439	234
Sertraline	57	5,297	93
S(+)-Fluoxetine	120	5,480	46
(±)-Fluoxetine	240	16,100	68
Citalopram	292	5,410	19
Paroxetine	1,893	22,870	12
Tricyclic antidepressants			
Imipramine	343	2,107	6
Desipramine	1,987	11,430	6

fluoxetine > citalopram >> paroxetine. It is highly likely that some SSRIs such as fluvoxamine, utilize sigma-1 receptors in its mode of action [45].

Antidepressants are thought to exert their effect by inducing adaptive neuroplasticity such as neurite outgrowth [46-48, 30, 49]. The prototypic sigma-1 receptor agonist (+)-pentazocine, as well as the antidepressants imipramine and fluvoxamine, enhance nerve growth factor (NGF)-induced neurite outgrowth in PC12 cells, in a concentration dependent manner [50]. The selective sigma-1 receptor antagonist, NE-100 [51], blocks these enhancements [50]. Recently, we reported that fluvoxamine, but not sertraline or paroxetine, significantly potentiates NGF-induced neurite outgrowth in PC12 cells, in a concentration dependent manner Fig. (1) [52]. Similarly, the sigma-1 receptor agonists, 1-(3,4-dimethoxyphenethyl)-4-(3-phenylpropyl) piperazine dihydrochloride (SA4503: cutamesine) [53], 4-phenyl-1-(4-phenylbutyl) piperidine (PPBP) [54-56], dehydroepiandrosterone-sulfate (DHEA-S) [57, 58], and donepezil also potentiate NGF-induced outgrowth in PC12 cells, in a concentration-dependent manner [59, 52]. This outgrowth is greatly attenuated if NE-100 is co-administered Fig. (1) [59, 52]. All these findings suggest that sigma-1 receptors play an important role in NGF-induced neurite outgrowth, and that in the brain, selected antidepressants and compounds facilitate this neurite outgrowth, *via* sigma-1 receptors. However, the precise cellular and molecular mechanisms underlying these processes are not fully understood.

### 3. SIGMA-1 RECEPTOR AGONISTS AND COGNITION

Sigma-1 receptor agonists can improve acetylcholine (an anti-muscarinic drug)-related deficits in memory and cognition in rodent models [24, 25, 27, 60]. Selective receptor agonists, including igmesine and SA4503, reverse amnesia induced by muscarinic and nicotinic receptor antagonists [61, 62]. Furthermore, sigma-1 receptor agonists, such as (+)-SKF 10,047 and SA4503, release acetylcholine in the rat brain [63, 64]. Recently, methyl (1*R*,2*S*/1*S*,2*R*)-2-[4-hydroxy-4-phenylpiperidin-1-yl)methyl]-1-(4-methylphenyl) cyclopropanecarboxylate ((±)-PPCC), a novel sigma-1 receptor agonist, has been shown to ameliorate cognitive impairment induced by selective cholinergic lesions in rats [65]. These findings suggest that sigma-1 receptor agonists can improve acetylcholine-related cognitive impairments through a mechanism of acetylcholine in the brain [27, 60].

Multiple lines of evidence suggest that aberrant glutamatergic neurotransmission *via* the *N*-methyl-D-aspartate (NMDA) receptors

may precipitate the cognitive impairment associated with schizophrenia and major depression [66-71]. The NMDA receptor antagonists, such as phencyclidine (PCP), induce schizophrenia-like symptoms including cognitive impairment in healthy subjects [66]. As a consequence, the NMDA receptor antagonists, including PCP, are widely used to generate animal models of cognitive impairment. We have shown that PCP-induced cognitive impairment in a novel object recognition test is significantly improved by subsequent sub-chronic (2-week) administration of the atypical antipsychotic drug clozapine, but not the typical antipsychotic drug haloperidol [72]. Thus, the reversal of PCP-induced cognitive impairment as measured by the novel object recognition test, may be a potential animal model for atypical antipsychotic activity, in the amelioration of schizophrenia associated cognitive impairment [72]. Furthermore, we have found that repeated administration of PCP causes a significant reduction of sigma-1 receptors in the mouse hippocampus [73, 74], suggesting that a decrease in sigma-1 receptors may precipitate PCP-induced cognitive impairment.


## 4. POTENTIAL THERAPEUTIC DRUGS FOR COGNITIVE IMPAIRMENT IN SCHIZOPHRENIA

### 4.1. Fluvoxamine as a Sigma-1 Receptor Agonist

We have reported that PCP-induced cognitive impairment is significantly improved by subsequent sub-chronic (2-week) administration of fluvoxamine (20 mg/kg/day), but not paroxetine (10 mg/kg/day) [48] or sertraline (10 or 20 mg/kg/day) [73]. The effect of fluvoxamine on PCP-induced cognitive impairment is antagonized by co-administration of NE-100. Unlike fluvoxamine, sertraline which is an SSRI with a high affinity at sigma-1 receptors, does not attenuate PCP-induced cognitive deficits in mice. In addition, sertraline does not enhance NGF-induced neurite outgrowth in PC12 cells [52]. These findings suggest that fluvoxamine and sertraline may act as an agonist and an antagonist respectively, at sigma-1 receptors [52, 30, 73]. Also, it is likely that the agonistic activity of fluvoxamine at sigma-1 receptors mediates its therapeutic effect in PCP-induced cognitive impairment in mice.

### 4.2. Donepezil as a Sigma-1 Receptor Agonist

Donepezil is the most widely prescribed drug for Alzheimer's disease. It is thought to influence cognition and function by inhibiting acetylcholinesterase (AChE) in the brain, however, it has also been reported that donepezil binds sigma receptors in the brain [75]. This sigma-1 receptor binding is thought to promote the antidepressive, anti-amnesic and neuroprotective effects of donepezil in



**Fig. (1).** Effects of SSRIs (fluvoxamine, paroxetine, sertraline) and sigma-1 receptor agonists (SA4503, PPBP, DHEA-S) on NGF-induced neurite outgrowth in PC12 cells. \* $P < 0.05$ , \*\* $P < 0.01$ , \*\*\* $P < 0.001$  as compared with control (NGF alone) group.  $^{SSS}P < 0.001$  as compared with drugs plus NE-100 group. The data are from Ref. [52] and Ref. [59].

the mouse forced swim test [76], and CO gas-induced [77] and amyloid  $\beta_{25-35}$ -induced neurotoxicity [78].

We have reported that PCP-induced cognitive impairment is significantly improved by subsequent, sub-chronic (2-week) administration of donepezil (1.0 mg/kg/day) [74]. This effect is antagonized by co-administration of the sigma-1 receptor antagonist NE-100. In contrast, PCP-induced cognitive impairment is not improved by subsequent, sub-chronic (2-week) administration of a different AChE inhibitor, physostigmine, that has no affinity to sigma-1 receptors [74]. These findings suggest that when treating mouse, PCP induced cognitive deficits, the therapeutic effect of donepezil is mediated *via* agonistic activity at sigma-1 receptors.

#### 4.3. Neurosteroids as Sigma-1 Receptor Agonists

Neurosteroids, hormones produced in central and peripheral neurons, were first reported by Baulieu [79]. In the brain, concentrations of the neurosteroids, DHEA, DHEA-S, pregnenolone (PREG) and PREG-sulfate are far greater than in the circulatory system. Furthermore, concentrations in the brain remain high after adrenalectomy and orchietomy, suggesting that these steroids do not originate from steroidogenic tissue but rather through local brain synthesis. As these neurosteroids do have moderate affinity towards sigma-1 receptors, they are considered endogenous ligands.

Neurosteroids may influence neuronal survival, neurite outgrowth and neurogenesis [80], as well as having varied effects on NMDA,  $\alpha$ -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA),  $\gamma$ -aminobutyric acid type A (GABA<sub>A</sub>), kainate, glycine, serotonin and nicotinic acetylcholine receptors [81]. In addition, several neurosteroids have affinity for the sigma-1 receptor; for example, DHEA, DHEA-S and PREG are all sigma-1 receptor agonists, while progesterone is an antagonist [21, 82, 20]. Neurosteroids therefore, appear to be highly relevant to the pathophysiology and pharmacological treatment of mood disorders, psychosis and dementia [83]. We have previously shown that DHEA-S significantly attenuates PCP-induced cognitive impairment in mice, and that these effects are antagonized by co-administration of NE-100, suggesting that DHEA-S activity is generated through sigma-1 receptors [48].

#### 5. CLINICAL REPORTS OF SIGMA-1 RECEPTOR AGONISTS IN SCHIZOPHRENIA

A previous positron emission tomography (PET) study demonstrated that in the healthy human brain, fluvoxamine (50-200 mg) and donepezil (5 or 10 mg) bind sigma-1 receptors in a dose-dependent manner, after oral administration [84, 85]. Studies on post-mortem brains from schizophrenic patients, report a greatly reduced density of sigma receptors, compared with controls [86, 87], consistent with pre-clinical reports showing a lower level of sigma-1 receptors in PCP-treated mouse brains [73, 74].

Studies show that fluvoxamine adjunctive therapy improves negative symptoms in patients with schizophrenia [88-90] Table 2. We recently, reported that fluvoxamine is effective in correcting cognitive impairment in patients with schizophrenia [91, 92]. There is also a case demonstrating the effectiveness of fluvoxamine in treating a patient with schizoaffective disorder for whom therapy with the SSRI, escitalopram, failed [93], suggesting a possible antipsychotic effect for fluvoxamine. Both pre-clinical and clinical evidence points to fluvoxamine being a promising therapeutic drug for cognitive and clinical symptoms in patients with schizophrenia. Recently, we performed a randomized double-blind trial of fluvoxamine adjunctive therapy in patients with schizophrenia and found that fluvoxamine was effective in restoring executive functions in patients [94].

Despite small sample sizes, clinical evidence suggests that the sigma-1 receptor agonists, donepezil [95-97], PREG [98, 99], and DHEA [100-104, 98, 105], could also be promising therapeutic agents for cognitive and clinical symptoms in patients with schizo-

phrenia Table 2, although these results are inconclusive [106, 107]. Further studies on sigma-1 receptor agonism using large cohorts need to be conducted.

Cognitive impairment is also a common feature in the prodromal state of psychosis [108]. Considering that sigma-1 receptors mediate neuroprotection and neuronal plasticity, it is likely that agonists such as fluvoxamine could reduce the risk of subsequent transition to schizophrenia in susceptible patients [109]. Very recently, we reported a case that fluvoxamine was effective in preventing persons at ultra-high risk of psychotic disorder from the onset of psychosis [110]. Randomized, double-blind, placebo-controlled studies of fluvoxamine in this group of patients will be necessary to determine the drug's clinical efficacy.

#### 6. POTENTIAL THERAPEUTIC DRUGS FOR MAJOR DEPRESSIVE DISORDER

In the late 1990s, it was demonstrated that sigma-1 receptor ligands have antidepressant-like action in animal models of depression. The selective sigma-1 receptor agonist, SA4503 [53], decreases immobility time in the forced swim test without any effect on open-field locomotion [111]. Interestingly, the antidepressant-like effect of SA4503 is achieved after a single dose of the drug. A Phase II study of SA4503, in patients with major depression is currently underway. The rapid antidepressant-like action of SA4503 has more recently been replicated by different agonists such as OPC-14523, igmesine (JO1784), (+)-SKF-10,047 and DHEA-S [112, 113]. Electrophysiological studies demonstrate a rapid antidepressant-like action for sigma-1 receptor agonists [114]. In addition, sigma-1 receptor knock-out mice show longer immobility times in the forced swim test, an indicator of a depression-like phenotype in mice [115]. These results point towards rapid antidepressant-like activity for sigma-1 receptor agonists [20].

The NMDA receptor antagonist ketamine exerts a rapid antidepressant action in patients with refractory major depression and bipolar depression [116, 117]. It is possible that this activity is triggered by sigma-1 receptor agonist binding [70, 71], since ketamine has moderate affinity for this receptor [118, 119].

#### 7. CLINICAL REPORTS OF SIGMA-1 RECEPTOR AGONISTS IN PSYCHOTIC DEPRESSION

Psychotic (or delusional) major depression is a severe illness typified by marked depressive symptoms and accompanied by delusions, and sometimes, by hallucinations. Patients with psychotic depression tend to experience longer episodes, psychomotor impairment, guilt, suicidal pre-occupation, and cognitive impairment [120-122]. In addition, they have a significantly higher mortality than patients with nonpsychotic major depression [121, 123, 124]. Several reports link psychotic depression with higher cortisol levels and more severe cognitive impairment, relative to non-psychotic depression sufferers [125-127].

Unfortunately for patients, psychotic depression frequently proves difficult to treat. Clinical studies demonstrate the efficacy of combined antidepressant (either a TCA or an SSRI) and atypical antipsychotic or electroconvulsive therapy (ECT) in treating psychotic depression [120, 124]. However, combinations of antipsychotics can lead to severe side effects such as extrapyramidal symptom or tardive dyskinesia [124]. Additionally, several antidepressants, *e.g.* TCAs, can produce significant cognitive impairment [128]. Several case reports have found that fluvoxamine monotherapy, in contrast to other SSRIs, is effective in treating patients with psychotic depression [129-131]. Furthermore, fluvoxamine monotherapy has a superior efficacy in alleviating both the psychotic and depressive symptoms of this disorder [132-136] Table 3, while paroxetine has a lesser effect [137].

Sigma-1 receptors have been implicated in the pathophysiology of depression and in the therapeutic action of antidepressants [18, 30]. Unlike paroxetine, with an inhibition constant ( $K_i$ ) of 1,893



**Table 2. Clinical Studies Using Sigma-1 Receptor Agonists in Schizophrenia**

Compound	Study Design	Sample Size	Dose (mg)	Positive Symptoms	Negative Symptoms	Cognitive Symptoms	Ref.
Fluvoxamine	DBT, RCT	30	100	→	↑	NA	[88]
	DBT, RCT	25	100	→	↑	NA	[89]
	DBT, RCT	53	100	→	↑	NA	[90]
	DBT, RCT	44	150	→	→	↑	[94]
Donepezil	DBT, RCT, COT	15	5	↑ (PANSS total score)		↑	[95]
	DBT, RCT, COT	13	5-10	NA	↑(depressive symptom)	NA	[96]
	DBT, RCT, COT	13	5-10	→	↑	NA	[97]
DHEA	DBT, RCT	27	100	→	↑	NA	[100]
	DBT, RCT, COT	55	200	→	→	↑	[102]
	DBT, RCT	31	150	→	↑	→	[103]
PREG	DBT, RCT	18	500	→	↑	→	[98]
PREG DHEA	DBT, RCT	44	30	↑	→	↑	[105]
			200	→	→	→	
			400	→	→	→	

DBT: Double-blind trial, RCT: Randomized controlled trial, COT: Cross-over trial, ↑: Effective, →: No change, NA: Not assessed, PANSS: The Positive and Negative Syndrome Scale.

nM, fluvoxamine is a potent sigma-1 receptor agonist with a  $K_i$  of 36 nM [45]. Fluvoxamine, but not paroxetine, binds to sigma-1 receptors in the intact human brain [84], suggesting that sigma-1 receptors are involved in fluvoxamine's mode of action. Supporting this is the evidence that fluvoxamine, but not paroxetine, improves PCP-induced cognitive impairment in mice [48] and cognitive impairment in some schizophrenics [91, 92]. As in schizophrenia, patients with psychotic depression have a greater level of cognitive impairment compared with non-psychotic depression [125-127]. In summary, the superior efficacy of fluvoxamine monotherapy in psychotic depression may be due to its sigma-1 receptor agonist property [138, 139, 30]. Again, further studies are needed to confirm the role of sigma-1 receptors in the efficacy of fluvoxamine treatment of psychotic depression.

## 8. CLINICAL REPORTS OF FLUVOXAMINE IN DELIRIUM

Delirium, a common and deleterious complication in patients, is thought to be a neurobehavioral manifestation of imbalances in the synthesis, release, and inactivation of the neurotransmitters that normally control cognitive function, behavior, and mood [140]. Recently, Furuse and Hashimoto [141-143] reported that fluvoxamine is effective in treating the delirium associated with Alzheimer's disease and patients in intensive care units, and postoperative delirium, although these reports are case reports. Given the role of sigma-1 receptors in the regulation of neurotransmitters as well as in cognition [27, 30, 20], it is likely that again, they are the target of

fluvoxamine's therapeutic action [144]. In order to confirm the role of sigma-1 receptors in the treatment of delirium, a randomized double-blind, placebo-controlled study of selective sigma-1 receptor agonists (for example, SA4503) in patients with delirium would be of interest.

## 9. CONCLUSION

As discussed here, sigma-1 receptors play a role in neuroplasticity and regulation of various neurotransmitter systems, and may be implicated in the pathophysiology of cognitive impairment in neuropsychiatric diseases. This would make sigma-1 receptor agonists attractive therapeutic drugs in the improvement of cognitive impairment, especially in schizophrenia and psychotic depression. The evolution of more selective sigma-1 receptor agonists and the analysis of their efficacy in the treatment of cognitive impairment associated with neuropsychiatric diseases are still needed.

## ABBREVIATIONS

AChE	=	Acetylcholinesterase
AMPA	=	$\alpha$ -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid
ATP	=	Adenosine triphosphate
CNS	=	Central nervous system
DHEA	=	Dehydroepiandrosterone
DHEA-S	=	Dehydroepiandrosterone-sulphate

**Table 3. Clinical Studies Using Fluvoxamine in Psychotic Depression**

Study Design	Analyzed Sample Size	Duration of Administration	Dose (mg)	Evaluation Criteria	Efficacy (%)	Ref.
Open	59	6 weeks	300	HAM-D $\leq$ 8 DDERS=0	84.2	[133]
Open	25	6/ 24 months (30 months)	300/ 200	Rate of recurrence	0/ 20	[134]
DBT, RCT	Flu+Pla: 36 Flu+Pin: 36	6 weeks	Flu: 300 Pin: 7.5	HAM-D $\leq$ 8 DDERS=0	80.0 80.5	[135]
DBT, RCT	Flu: 14 Ven: 14	6 weeks	Flu: 300 Ven: 300	HAM-D $\leq$ 8 DDERS=0	Flu: 78.6 Ven: 58.3	[136]
DBT, RCT	Flu+Pla: 13 Flu+Hal: 11 Des+Pla: 10 Des+Hal: 14	6 weeks	Flu: 300 Des: 150 Hal: 0.1mg/kg	HAM-D $\leq$ 50% DDERS=0	Flu+Pla: 69 Flu+Hal: 45 Des+Pla: 40 Des+Hal: 64	[132]

DBT: Double-blind trial, RCT: Randomized controlled trial, HAM-D: Hamilton rating scale for depression, DDERS: Dimensions of delusional experience rating scale, Flu: Fluvoxamine, Pla: Placebo, Pin: Pindolol, Ven: Venlafaxine, Des: Desipramine, Hal: Haloperidol.

- ECT = Electroconvulsive therapy [3] Green MF, Nuechterlein KH, Gold JM, *et al.* Approaching a consensus cognitive battery for clinical trials in schizophrenia: the NIMH-MATRICES conference to select cognitive domains and test criteria. *Biol Psychiatry* 2004; 56: 301-7.
- ER = Endoplasmic reticulum [4] Kurtz MM. Neurocognitive impairment across the lifespan in schizophrenia: an update. *Schizophr Res* 2005; 74: 15-26.
- GABA<sub>A</sub> =  $\gamma$ -aminobutyric acid type A [5] Keefe RS, Bilder RM, Davis SM, *et al.* Neurocognitive effects of antipsychotic medications in patients with chronic schizophrenia in the CATIE Trial. *Arch Gen Psychiatry* 2007; 64: 633-47.
- IP<sub>3</sub> = Inositol 1,4,5-triphosphate [6] Widlocher DJ. Psychomotor retardation: clinical, theoretical, and psychometric aspects. *Psychiatr Clin North Am* 1983; 6: 27-40.
- NGF = Nerve-growth factor [7] Fava M. Symptoms of fatigue and cognitive/executive dysfunction in major depressive disorder before and after antidepressant treatment. *J Clin Psychiatry* 2003; 64 (Suppl 14): 30-4.
- NMDA = N-Methyl-D-aspartate [8] Cohen RM, Weingartner H, Smallberg SA, Pickar D, Murphy DL. Effort and cognition in depression. *Arch Gen Psychiatry* 1982; 39: 593-7.
- PCP = Phencyclidine [9] Porter RJ, Gallagher P, Thompson JM, Young AH. Neurocognitive impairment in drug-free patients with major depressive disorder. *Br J Psychiatry* 2003; 182: 214-20.
- PET = Positron emission tomography [10] Hashimoto K, Koike K, Shimizu E, Iyo M.  $\alpha 7$  Nicotinic receptor agonists as potential therapeutic drugs for schizophrenia. *Curr Med Chem CNS Agents* 2005; 5: 171-84.
- PPBP = 4-phenyl-1-(4-phenylbutyl)piperidine [11] Buchanan RW, Freedman R, Javitt DC, Abi-Dargham A, Lieberman JA. Recent advances in the development of novel pharmacological agents for the treatment of cognitive impairments in schizophrenia. *Schizophr Bull* 2007; 33: 1120-30.
- ( $\pm$ )-PPCC = Methyl (1*R*,2*S*/1*S*,2*R*)-2-[4-hydroxy-4-phenylpiperidin-1-yl)methyl]-1-(4-methylphenyl) cyclopropanecarboxylate [12] Hashimoto K. Glycine transport inhibitors for the treatment of schizophrenia. *Open Med Chem J* 2010; 4: 10-9.
- PREG = pregnenolone [13] Toyohara J, Hashimoto K.  $\alpha 7$  Nicotinic Receptor Agonists: Potential Therapeutic Drugs for Treatment of Cognitive Impairments in Schizophrenia and Alzheimer's Disease. *Open Med Chem J* 2010; 4: 37-56.
- SA4503 = 1-(3,4-dimethoxyphenethyl)-4-(3-phenylpropyl)piperazine dihydrochloride [14] Ishikawa M, Hashimoto K.  $\alpha 7$  nicotinic acetylcholine receptor as a potential therapeutic target for schizophrenia. *Curr Pharm Des* 2011; 17: 121-9.
- SSRI = Selective serotonin reuptake inhibitor [15] Martin WR, Eades CG, Thompson JA, Huppler RE, Gilbert PE. The effects of morphine- and nalorphine- like drugs in the nondependent and morphine-dependent chronic spinal dog. *J Pharmacol Exp Ther* 1976; 197: 517-32.
- TCA = Tricyclic antidepressant [16] Hanner M, Moebius FF, Flandorfer A, *et al.* Purification, molecular cloning, and expression of the mammalian signal-binding site. *Proc Natl Acad Sci USA* 1996; 93: 8072-7.

# CONFLICT OF INTEREST

Dr. Hashimoto reports having received the speakers' bureau honoraria from Abbott Pharmaceuticals. Other authors report no competing interests.

# ACKNOWLEDGEMENT

We would like to thank our collaborators who worked on some papers described in this review article, and who are listed as the co-authors of our papers in the reference list. This study was supported by a grant (to K.H.) from Comprehensive Research on Disability, Health and Welfare, Health and Labour Sciences Research Grants, Japan.

# REFERENCES

- [1] Freedman R. Schizophrenia. *N Engl J Med* 2003; 349: 1738-49.
- [2] Coyle JT, Tsai G. The NMDA receptor glycine modulatory site: a therapeutic target for improving cognition and reducing negative symptoms in schizophrenia. *Psychopharmacology (Berl)* 2004; 174: 32-38.

- [17] Hayashi T, Maurice T, Su TP.  $\text{Ca}^{2+}$  signaling *via*  $\sigma_1$ -receptors: novel regulatory mechanism affecting intracellular  $\text{Ca}^{2+}$  concentration. *J Pharmacol Exp Ther* 2000; 293: 788-98.
- [18] Hayashi T, Su TP. Sigma-1 receptor chaperones at the ER-mitochondrion interface regulate  $\text{Ca}^{2+}$  signaling and cell survival. *Cell* 2007; 131: 596-610.
- [19] Hayashi T, Rizzuto R, Hajnoczky G, Su TP. MAM: more than just a housekeeper. *Trends Cell Biol* 2009; 19: 81-8.
- [20] Hayashi T, Stahl SM. The sigma-1 receptor and its role in the treatment of mood disorders. *Drugs Future* 2009; 34: 137-46.
- [21] Maurice T, Urani A, Phan VL, Romieu P. The interaction between neuroactive steroids and the sigma-1 receptor function: behavioral consequences and therapeutic opportunities. *Brain Res Brain Res Rev* 2001; 37: 116-32.
- [22] Maurice T. Improving Alzheimer's Disease-Related Cognitive Deficits with sigma1 Receptor Agonists. *Drug News Perspect* 2002; 15: 617-25.
- [23] Su TP, Hayashi T. Understanding the molecular mechanism of sigma-1 receptors: towards a hypothesis that sigma-1 receptors are intracellular amplifiers for signal transduction. *Curr Med Chem* 2003; 10: 2073-80.
- [24] Hayashi T, Su TP. Sigma-1 receptor ligands: potential in the treatment of neuropsychiatric disorders. *CNS Drugs* 2004; 18: 269-84.
- [25] Hayashi T, Su T. The sigma receptor: evolution of the concept in neuropsychopharmacology. *Curr Neuropharmacol* 2005; 3: 267-80.
- [26] Monnet FP. Sigma-1 receptor as regulator of neuronal intracellular  $\text{Ca}^{2+}$ : clinical and therapeutic relevance. *Biol Cell* 2005; 97: 873-83.
- [27] Hashimoto K, Ishiwata K. Sigma receptor ligands: possible application as therapeutic drugs and as radiopharmaceuticals. *Curr Pharm Des* 2006; 12: 3857-76.
- [28] Maurice T, Gregoire C, Espallergues J. Neuro(steroid) actions at the neuromodulatory sigma1 (sigma-1) receptor: biochemical and physiological evidences, consequences in neuroprotection. *Pharmacol Biochem Behav* 2006; 84: 581-97.
- [29] Hayashi T, Su TP. An update on the development of drugs for neuropsychiatric disorders: focusing on the sigma-1 receptor ligand. *Expert Opin Ther Targets* 2008; 12: 45-58.
- [30] Hashimoto K. Sigma-1 receptors and selective serotonin reuptake inhibitors: clinical implications of their relationship. *Cent Nerv Syst Agents Med Chem* 2009; 9: 197-204.
- [31] Maurice T, Su TP. The pharmacology of sigma-1 receptors. *Pharmacol Ther* 2009; 124: 195-206.
- [32] Ishikawa M, Hashimoto K. The role of sigma-1 receptors in the pathophysiology of neuropsychiatric diseases. *J Receptor Ligand Channel Res* 2010; 3: 25-36.
- [33] Su TP, Hayashi T, Maurice T, Buch S, Ruoho AE. The sigma-1 receptor chaperone as an inter-organelle signaling modulator. *Trends Pharmacol Sci* 2010; 33: 557-66.
- [34] Hayashi T, Tsai SY, Mori T, Fujimoto M, Su TP. Targeting ligand-operated chaperone sigma-1 receptors in the treatment of neuropsychiatric disorders. *Expert Opin Ther Targets* 2011; 15: 557-77.
- [35] Hayashi T, Su TP. Sigma-1 receptors at galactosylceramide-enriched lipid microdomains regulate oligodendrocyte differentiation. *Proc Natl Acad Sci USA* 2004; 101: 14949-54.
- [36] Tsai SY, Hayashi T, Harvey BK, *et al.* Sigma-1 receptors regulate hippocampal dendritic spine formation *via* a free radical-sensitive mechanism involving Rac1/GTP pathway. *Proc Natl Acad Sci USA* 2009; 106: 22468-73.
- [37] Harrison PJ, Law AJ. Neuregulin 1 and schizophrenia: genetics, gene expression, and neurobiology. *Biol Psychiatry* 2006; 60: 132-40.
- [38] Walterfang M, Wood SJ, Velakoulis D, Pantelis C. Neuropathological, neurogenetic and neuroimaging evidence for white matter pathology in schizophrenia. *Neurosci Biobehav Rev* 2006; 30: 918-48.
- [39] Fields RD. White matter in learning, cognition and psychiatric disorders. *Trends Neurosci* 2008; 31: 361-70.
- [40] van Spronsen M, Hoogenraad CC. Synapse pathology in psychiatric and neurologic disease. *Curr Neurol Neurosci Rep* 2010; 10: 207-14.
- [41] Nestler EJ, Barrot M, DiLeone RJ, Eisch AJ, Gold SJ, Monteggia LM. Neurobiology of depression. *Neuron* 2002; 34: 13-25.
- [42] Stahl SM. Not so selective serotonin reuptake inhibitors. *J Clin Psychiatry* 1998; 59: 343-4.
- [43] Stahl SM. Using secondary binding properties to select a not so selective serotonin reuptake inhibitor. *J Clin Psychiatry* 1998; 59: 642-3.
- [44] Nemeroff CB, Owens MJ. Pharmacologic differences among the SSRIs: focus on monoamine transporters and the HPA axis. *CNS Spectr* 2004; 9: 23-31.
- [45] Narita N, Hashimoto K, Tomitaka S, Minabe Y. Interactions of selective serotonin reuptake inhibitors with subtypes of sigma receptors in rat brain. *Eur J Pharmacol* 1996; 307: 117-9.
- [46] Duman RS, Heninger GR, Nestler EJ. A molecular and cellular theory of depression. *Arch Gen Psychiatry* 1997; 54: 597-606.
- [47] Hashimoto K, Shimizu E, Iyo M. Critical role of brain-derived neurotrophic factor in mood disorders. *Brain Res Brain Res Rev* 2004; 45: 104-14.
- [48] Hashimoto K, Fujita Y, Iyo M. Phencyclidine-induced cognitive deficits in mice are improved by subsequent subchronic administration of fluvoxamine: role of sigma-1 receptors. *Neuropsychopharmacology* 2007; 32: 514-21.
- [49] Hashimoto K. Brain-derived neurotrophic factor as a biomarker for mood disorders: an historical overview and future directions. *Psychiatry Clin Neurosci* 2010; 64: 341-57.
- [50] Takebayashi M, Hayashi T, Su TP. Nerve growth factor-induced neurite sprouting in PC12 cells involves sigma-1 receptors: implications for antidepressants. *J Pharmacol Exp Ther* 2002; 303: 1227-37.
- [51] Okuyama S, Nakazato A. NE-100: A Novel Sigma Receptor Antagonist. *CNS Drug Reviews* 1996; 2: 226-37.
- [52] Nishimura T, Ishima T, Iyo M, Hashimoto K. Potentiation of nerve growth factor-induced neurite outgrowth by fluvoxamine: role of sigma-1 receptors, IP3 receptors and cellular signaling pathways. *PLoS ONE* 2008; 3: e2558.
- [53] Matsuno K, Mita S. SA4503: A Novel Sigma1 Receptor Agonist. *CNS Drug Reviews* 1998; 4: 1-24.
- [54] Hashimoto K, Scheffel U, London ED. *In vivo* labeling of sigma receptors in mouse brain with [ $^3\text{H}$ ]-4-phenyl-1-(4-phenylbutyl) piperidine. *Synapse* 1995; 20: 85-90.
- [55] Takahashi H, Kirsch JR, Hashimoto K, London ED, Koehler RC, Traystman RJ. PPBP [4-phenyl-1-(4-phenylbutyl) piperidine], a potent sigma-receptor ligand, decreases brain injury after transient focal ischemia in cats. *Stroke* 1995; 26: 1676-82.
- [56] Takahashi H, Kirsch JR, Hashimoto K, London ED, Koehler RC, Traystman RJ. PPBP [4-phenyl-1-(4-phenylbutyl) piperidine] decreases brain injury after transient focal ischemia in rats. *Stroke* 1996; 27: 2120-3.
- [57] Noda Y, Kamei H, Kamei Y, Nagai T, Nishida M, Nabeshima T. Neurosteroids ameliorate conditioned fear stress: an association with sigma receptors. *Neuropsychopharmacology* 2000; 23: 276-84.
- [58] Maurice T, Phan VL, Urani A, Guillemain I. Differential involvement of the sigma(1) (sigma-1) receptor in the anti-amnesic effect of neuroactive steroids, as demonstrated using an *in vivo* antisense strategy in the mouse. *Br J Pharmacol* 2001; 134: 1731-41.
- [59] Ishima T, Nishimura T, Iyo M, Hashimoto K. Potentiation of nerve growth factor-induced neurite outgrowth in PC12 cells by donepezil: role of sigma-1 receptors and IP3 receptors. *Prog Neuropsychopharmacol Biol Psychiatry* 2008; 32: 1656-9.
- [60] van Waarde A, Ramakrishnan NK, Rybczynska AA, *et al.* The cholinergic system, sigma-1 receptors and cognition. *Behav Brain Res* 2011; 221: 543-54.
- [61] Earley B, Burke M, Leonard BE, Gouret CJ, Junien JL. Evidence for an anti-amnesic effect of JO 1784 in the rat: a potent and selective ligand for the sigma receptor. *Brain Res* 1991; 546: 282-6.
- [62] Matsuno K, Senda T, Matsunaga K, Mita S. Ameliorating effects of sigma receptor ligands on the impairment of passive avoidance tasks in mice: involvement in the central acetylcholinergic system. *Eur J Pharmacol* 1994; 261: 43-51.
- [63] Kobayashi T, Matsuno K, Mita S. Regional differences of the effect of sigma receptor ligands on the acetylcholine release in the rat brain. *J Neural Transm* 1996; 103: 661-9.
- [64] Kobayashi T, Matsuno K, Nakata K, Mita S. Enhancement of acetylcholine release by SA4503, a novel sigma-1 receptor agonist, in the rat brain. *J Pharmacol Exp Ther* 1996; 279: 106-13.
- [65] Antonini V, Prezzavento O, Coradazzi M, *et al.* Anti-amnesic properties of (+/-)-PPCC, a novel sigma receptor ligand, on cognitive dysfunction induced by selective cholinergic lesion in rats. *J Neurochem* 2009; 109: 744-54.

- [66] Javitt DC, Zukin SR. Recent advances in the phencyclidine model of schizophrenia. *Am J Psychiatry* 1991; 148: 1301-8.
- [67] Hashimoto K, Fukushima T, Shimizu E, *et al.* Decreased serum levels of D-serine in patients with schizophrenia: evidence in support of the N-methyl-D-aspartate receptor hypofunction hypothesis of schizophrenia. *Arch Gen Psychiatry* 2003; 60: 572-6.
- [68] Hashimoto K, Okamura N, Shimizu E, Iyo M. Glutamate Hypothesis of Schizophrenia and Approach for Possible Therapeutic Drugs. *Curr Med Chem - Central Nervous System Agents* 2004; 4: 147-54.
- [69] Hashimoto K, Shimizu E, Iyo M. Dysfunction of Glia-Neuron Communication in Pathophysiology of Schizophrenia. *Curr Psychiatry Rev* 2005; 1: 151-63.
- [70] Hashimoto K. Emerging role of glutamate in the pathophysiology of major depressive disorder. *Brain Res Rev* 2009; 61: 105-23.
- [71] Hashimoto K. Role of the mTOR signaling pathway in the rapid antidepressant action of ketamine. *Expert Rev Neurother* 2011; 11: 33-6.
- [72] Hashimoto K, Fujita Y, Shimizu E, Iyo M. Phencyclidine-induced cognitive deficits in mice are improved by subsequent subchronic administration of clozapine, but not haloperidol. *Eur J Pharmacol* 2005; 519: 114-7.
- [73] Ishima T, Fujita Y, Kohno M, *et al.* Improvement of Phencyclidine-Induced Cognitive Deficits in Mice by Subsequent Subchronic Administration of Fluvoxamine, but not Sertraline. *Open Clin Chem J* 2009; 2: 7-11.
- [74] Kunitachi S, Fujita Y, Ishima T, *et al.* Phencyclidine-induced cognitive deficits in mice are ameliorated by subsequent subchronic administration of donepezil: role of sigma-1 receptors. *Brain Res* 2009; 1279: 189-96.
- [75] Kato K, Hayako H, Ishihara Y, Marui S, Iwane M, Miyamoto M. TAK-147, an acetylcholinesterase inhibitor, increases choline acetyltransferase activity in cultured rat septal cholinergic neurons. *Neurosci Lett* 1999; 260: 5-8.
- [76] Maurice T, Meunier J, Feng B, Ieni J, Monaghan DT. Interaction with sigma-1 protein, but not N-methyl-D-aspartate receptor, is involved in the pharmacological activity of donepezil. *J Pharmacol Exp Ther* 2006; 317: 606-14.
- [77] Meunier J, Ieni J, Maurice T. Antiamnesic and neuroprotective effects of donepezil against learning impairments induced in mice by exposure to carbon monoxide gas. *J Pharmacol Exp Ther* 2006; 317: 1307-19.
- [78] Meunier J, Ieni J, Maurice T. The anti-amnesic and neuroprotective effects of donepezil against amyloid beta<sub>25-35</sub> peptide-induced toxicity in mice involve an interaction with the sigma1 receptor. *Br J Pharmacol* 2006; 149: 998-1012.
- [79] Baulieu EE. Neurosteroids: a novel function of the brain. *Psychoneuroendocrinology* 1998; 23: 963-87.
- [80] Charalampopoulos I, Remboutsika E, Margioris AN, Gravanis A. Neurosteroids as modulators of neurogenesis and neuronal survival. *Trends Endocrinol Metab* 2008; 19: 300-7.
- [81] Rupprecht R, di Michele F, Hermann B, *et al.* Neuroactive steroids: molecular mechanisms of action and implications for neuropsychopharmacology. *Brain Res Brain Res Rev* 2001; 37: 59-67.
- [82] Monnet FP, Maurice T. The sigma1 protein as a target for the non-genomic effects of neuro(steroid)s: molecular, physiological, and behavioral aspects. *J Pharmacol Sci* 2006; 100: 93-118.
- [83] Strous RD, Maayan R, Weizman A. The relevance of neurosteroids to clinical psychiatry: from the laboratory to the bedside. *Eur Neuropsychopharmacol* 2006; 16: 155-69.
- [84] Ishikawa M, Ishiwata K, Ishii K, *et al.* High occupancy of sigma-1 receptors in the human brain after single oral administration of fluvoxamine: a positron emission tomography study using [<sup>11</sup>C]SA4503. *Biol Psychiatry* 2007; 62: 878-83.
- [85] Ishikawa M, Sakata M, Ishii K, *et al.* High occupancy of sigma1 receptors in the human brain after single oral administration of donepezil: a positron emission tomography study using [<sup>11</sup>C]SA4503. *Int J Neuropsychopharmacol* 2009; 12: 1127-31.
- [86] Weissman AD, Casanova MF, Kleinman JE, London ED, De Souza EB. Selective loss of cerebral cortical sigma, but not PCP binding sites in schizophrenia. *Biol Psychiatry* 1991; 29: 41-54.
- [87] Helmeste DM, Tang SW, Bunney WE, Jr., Potkin SG, Jones EG. Decrease in sigma but no increase in striatal dopamine D4 sites in schizophrenic brains. *Eur J Pharmacol* 1996; 314: R3-5.
- [88] Silver H, Nassar A. Fluvoxamine improves negative symptoms in treated chronic schizophrenia: an add-on double-blind, placebo-controlled study. *Biol Psychiatry* 1992; 31: 698-704.
- [89] Silver H, Shmugliakov N. Augmentation with fluvoxamine but not maprotiline improves negative symptoms in treated schizophrenia: evidence for a specific serotonergic effect from a double-blind study. *J Clin Psychopharmacol* 1998; 18: 208-11.
- [90] Silver H, Barash I, Aharon N, Kaplan A, Poyurovsky M. Fluvoxamine augmentation of antipsychotics improves negative symptoms in psychotic chronic schizophrenic patients: a placebo-controlled study. *Int Clin Psychopharmacol* 2000; 15: 257-61.
- [91] Iyo M, Shirayama Y, Watanabe H, *et al.* Fluvoxamine as a sigma-1 receptor agonist improved cognitive impairments in a patient with schizophrenia. *Prog Neuropsychopharmacol Biol Psychiatry* 2008; 32: 1072-3.
- [92] Niitsu T, Shirayama Y, Fujisaki M, Hashimoto K, Iyo M. Fluvoxamine improved cognitive impairments in a patient with schizophrenia. *Prog Neuropsychopharmacol Biol Psychiatry* 2010; 34: 1345-6.
- [93] Goldman S. Possible antipsychotic effect of fluvoxamine. *CNS Spectr* 2005; 10: 774; author reply 774-5.
- [94] Niitsu T, Fujisaki M, Shiina A, *et al.* A randomized, double-blind, placebo-controlled trial of fluvoxamine in patients with schizophrenia: a preliminary study. submitted (revision).
- [95] Erickson SK, Schwarzkopf SB, Palumbo D, Badgley-Fleeman J, Smirnow AM, Light GA. Efficacy and tolerability of low-dose donepezil in schizophrenia. *Clin Neuropharmacol* 2005; 28: 179-84.
- [96] Risch SC, Horner MD, McGurk SR, *et al.* Donepezil effects on mood in patients with schizophrenia and schizoaffective disorder. *Int J Neuropsychopharmacol* 2006; 9: 603-5.
- [97] Risch SC, Horner MD, McGurk SR, *et al.* Double-blind donepezil-placebo crossover augmentation study of atypical antipsychotics in chronic, stable schizophrenia: a pilot study. *Schizophr Res* 2007; 93: 131-5.
- [98] Marx CE, Keefe RS, Buchanan RW, *et al.* Proof-of-concept trial with the neurosteroid pregnenolone targeting cognitive and negative symptoms in schizophrenia. *Neuropsychopharmacology* 2009; 34: 1885-903.
- [99] Hashimoto K. Neurosteroid pregnenolone and sigma-1 receptor function. *Clin Psychopharmacol Neurosci* 2010; 8: 115.
- [100] Strous RD, Maayan R, Lapidus R, *et al.* Dehydroepiandrosterone augmentation in the management of negative, depressive, and anxiety symptoms in schizophrenia. *Arch Gen Psychiatry* 2003; 60: 133-41.
- [101] Nachshoni T, Ebert T, Abramovitch Y, *et al.* Improvement of extrapyramidal symptoms following dehydroepiandrosterone (DHEA) administration in antipsychotic treated schizophrenia patients: a randomized, double-blind placebo controlled trial. *Schizophr Res* 2005; 79: 251-6.
- [102] Ritsner MS, Gibel A, Ratner Y, Tsinovoy G, Strous RD. Improvement of sustained attention and visual and movement skills, but not clinical symptoms, after dehydroepiandrosterone augmentation in schizophrenia: a randomized, double-blind, placebo-controlled, crossover trial. *J Clin Psychopharmacol* 2006; 26: 495-9.
- [103] Strous RD, Stryer R, Maayan R, *et al.* Analysis of clinical symptomatology, extrapyramidal symptoms and neurocognitive dysfunction following dehydroepiandrosterone (DHEA) administration in olanzapine treated schizophrenia patients: a randomized, double-blind placebo controlled trial. *Psychoneuroendocrinology* 2007; 32: 96-105.
- [104] Strous RD, Gibel A, Maayan R, Weizman A, Ritsner MS. Hormonal response to dehydroepiandrosterone administration in schizophrenia: findings from a randomized, double-blind, placebo-controlled, crossover study. *J Clin Psychopharmacol* 2008; 28: 456-9.
- [105] Ritsner MS, Gibel A, Shleifer T, *et al.* Pregnenolone and dehydroepiandrosterone as an adjunctive treatment in schizophrenia and schizoaffective disorder: an 8-week, double-blind, randomized, controlled, 2-center, parallel-group trial. *J Clin Psychiatry* 2010; 71: 1351-62.
- [106] Ribeiz SR, Bassitt DP, Arrais JA, Avila R, Steffens DC, Bottino CM. Cholinesterase inhibitors as adjunctive therapy in patients with schizophrenia and schizoaffective disorder: a review and meta-analysis of the literature. *CNS Drugs* 2010; 24: 303-17.
- [107] Ritsner MS. Pregnenolone, dehydroepiandrosterone, and schizophrenia: alterations and clinical trials. *CNS Neurosci Ther* 2010; 16: 32-44.



- [108] Brewer WJ, Francey SM, Wood SJ, *et al.* Memory impairments identified in people at ultra-high risk for psychosis who later develop first-episode psychosis. *Am J Psychiatry* 2005; 162: 71-8.
- [109] Hashimoto K. Can the sigma-1 receptor agonist fluvoxamine prevent schizophrenia? *CNS Neurol Disord Drug Targets* 2009; 8: 470-4.
- [110] Tadokoro S, Kanahara N, Kikuchi S, Hashimoto K, Iyo M. Fluvoxamine prevents onset of psychosis: a case report of a patient at ultra-high risk of psychotic disorder. *Ann. Gen. Psychiatry* 2011; 10: 26.
- [111] Matsuno K, Kobayashi T, Tanaka MK, Mita S. Sigma-1 receptor subtype is involved in the relief of behavioral despair in the mouse forced swimming test. *Eur J Pharmacol* 1996; 312: 267-71.
- [112] Tottori K, Miwa T, Uwahodo Y, *et al.* Antidepressant-like responses to the combined sigma and 5-HT<sub>1A</sub> receptor agonist OPC-14523. *Neuropharmacology* 2001; 41: 976-88.
- [113] Urani A, Roman FJ, Phan VL, Su TP, Maurice T. The antidepressant-like effect induced by sigma-1-receptor agonists and neuroactive steroids in mice submitted to the forced swimming test. *J Pharmacol Exp Ther* 2001; 298: 1269-79.
- [114] Bermack JE, Debonnel G. The role of sigma receptors in depression. *J Pharmacol Sci* 2005; 97: 317-36.
- [115] Sabino V, Cottone P, Parylak SL, Steardo L, Zorrilla EP. Sigma-1 receptor knockout mice display a depressive-like phenotype. *Behav Brain Res* 2009; 198: 472-6.
- [116] Zarate CA, Jr., Singh JB, Carlson PJ, Brutsche NE, Ameli R, Luckenbaugh DA, *et al.* A randomized trial of an *N*-methyl-D-aspartate antagonist in treatment-resistant major depression. *Arch Gen Psychiatry* 2006; 63: 856-64.
- [117] Diazgranados N, Ibrahim L, Brutsche NE, *et al.* A randomized add-on trial of an *N*-methyl-D-aspartate antagonist in treatment-resistant bipolar depression. *Arch Gen Psychiatry* 2010; 67: 793-802.
- [118] Hustveit O, Maurset A, Oye I. Interaction of the chiral forms of ketamine with opioid, phencyclidine, sigma and muscarinic receptors. *Pharmacol Toxicol* 1995; 77: 355-9.
- [119] Stahl SM. The sigma enigma: Can sigma receptors provide a novel target for disorders of mood and cognition? *J Clin Psychiatry* 2008; 69: 1673-74.
- [120] Coryell W. The treatment of psychotic depression. *J Clin Psychiatry* 1998; 59: 22-7; discussion 28-9.
- [121] Schatzberg AF. New approaches to managing psychotic depression. *J Clin Psychiatry* 2003; 64: 19-23.
- [122] Dannon PN, Lowengrub K, Gonopolski Y, Kotler M. Current and emerging somatic treatment strategies in psychotic major depression. *Expert Rev Neurother* 2006; 6: 73-80.
- [123] Vythilingam M, Chen J, Bremner JD, Mazure CM, Maciejewski PK, Nelson JC. Psychotic depression and mortality. *Am J Psychiatry* 2003; 160: 574-6.
- [124] Hamoda HM, Osser DN. The Psychopharmacology Algorithm Project at the Harvard South Shore Program: an update on psychotic depression. *Harv Rev Psychiatry* 2008; 16: 235-47.
- [125] Belanoff JK, Kalehzan M, Sund B, Fleming Ficek SK, Schatzberg AF. Cortisol activity and cognitive changes in psychotic major depression. *Am J Psychiatry* 2001; 158: 1612-16.
- [126] Fleming SK, Blasey C, Schatzberg AF. Neuropsychological correlates of psychotic features in major depressive disorders: a review and meta-analysis. *J Psychiatr Res* 2004; 38: 27-35.
- [127] Gomez RG, Fleming SH, Keller J, *et al.* The neuropsychological profile of psychotic major depression and its relation to cortisol. *Biol Psychiatry* 2006; 60: 472-8.
- [128] Hindmarch I, Hashimoto K. Cognition and depression: the effects of fluvoxamine, a sigma-1 receptor agonist, reconsidered. *Hum Psychopharmacol* 2010; 25: 193-200.
- [129] Furuse T, Hashimoto K. Fluvoxamine monotherapy for psychotic depression: the potential role of sigma-1 receptors. *Ann Gen Psychiatry* 2009; 8: 26.
- [130] Kishimoto A, Todani A, Miura J, Kitagaki T, Hashimoto K. The opposite effects of fluvoxamine and sertraline in the treatment of psychotic major depression: a case report. *Ann Gen Psychiatry* 2010; 9: 23.
- [131] Shirayama Y, Hashimoto K. A Case of Psychotic Depression Treated with Fluvoxamine Monotherapy. *Clin Psychopharmacol Neurosci* 2010; 8: 53-4.
- [132] Bellini L, Gasperini M, Gatti F, Franchini L, Smeraldi E. In: Langer SZ, Ed. Critical issues in the treatment of affective disorders. Basel, Switzerland: Karger Press. 1994; pp. 32-6.
- [133] Gatti F, Bellini L, Gasperini M, Perez J, Zanardi R, Smeraldi E. Fluvoxamine alone in the treatment of delusional depression. *Am J Psychiatry* 1996; 153: 414-6.
- [134] Zanardi R, Franchini L, Gasperini M, Smeraldi E, Perez J. Long-term treatment of psychotic (delusional) depression with fluvoxamine: an open pilot study. *Int Clin Psychopharmacol* 1997; 12: 195-7.
- [135] Zanardi R, Franchini L, Gasperini M, Lucca A, Smeraldi E, Perez J. Faster onset of action of fluvoxamine in combination with pindolol in the treatment of delusional depression: a controlled study. *J Clin Psychopharmacol* 1998; 18: 441-6.
- [136] Zanardi R, Franchini L, Serretti A, Perez J, Smeraldi E. Venlafaxine versus fluvoxamine in the treatment of delusional depression: a pilot double-blind controlled study. *J Clin Psychiatry* 2000; 61: 26-9.
- [137] Zanardi R, Franchini L, Gasperini M, Perez J, Smeraldi E. Double-blind controlled trial of sertraline versus paroxetine in the treatment of delusional depression. *Am J Psychiatry* 1996; 153: 1631-3.
- [138] Hayashi T, Su T. Understanding the role of sigma-1 receptors in psychotic depression. *Psychiatric Times* 2005; 22: 54-63.
- [139] Stahl SM. Antidepressant treatment of psychotic major depression: potential role of the sigma receptor. *CNS Spectr* 2005; 10: 319-23.
- [140] Fong TG, Tulebaev SR, Inouye SK. Delirium in elderly adults: diagnosis, prevention and treatment. *Nat Rev Neurol* 2009; 5: 210-20.
- [141] Furuse T, Hashimoto K. Sigma-1 receptor agonist fluvoxamine for delirium in patients with Alzheimer's disease. *Ann Gen Psychiatry* 2010; 9: 6.
- [142] Furuse T, Hashimoto K. Sigma-1 receptor agonist fluvoxamine for delirium in intensive care units: report of five cases. *Ann Gen Psychiatry* 2010; 9: 18.
- [143] Furuse T, Hashimoto K. Sigma-1 receptor agonist fluvoxamine for postoperative delirium in older adults: report of three cases. *Ann Gen Psychiatry* 2010; 9: 28.
- [144] Hashimoto K, Furuse T. Sigma-1 receptor agonist fluvoxamine for delirium in elderly adults. *Int J Geriatr Psychiatry* 2011; in press.

Copyright of Current Pharmaceutical Design is the property of Bentham Science Publishers Ltd. and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.