

sPlot open - An environmentally-balanced, open-access, global dataset of vegetation plots

This manuscript ([permalink](#)) was automatically generated from [fmsabatini/sPlotOpen Manuscript@9e6650b](#) on October 14, 2020.

Authors

- **Francesco Maria Sabatini**

 [0000-0002-7202-7697](#) ·  [fmsabatini](#) ·  [sPlot iDiv](#)

German Centre for Integrative Biodiversity Research (iDiv) - Halle-Jena-Leipzig. Germany; Martin-Luther-Universität Halle-Wittenberg, Institut für Biologie, Am Kirchtor 1, 06108, Halle (Saale). Germany · Funded by None

- **Jonathan Lenoir**

 [0000-0003-0638-9582](#) ·  [lenjon](#) ·  [EkoLogIt](#)

Unité de Recherche "Ecologie et Dynamique des Systèmes Anthropisés" (EDYSAN), UMR 7058 CNRS, Université de Picardie Jules Verne, 80037 Amiens Cedex 1, France · Funded by None

- **Helge Bruelheide**

 [0000-0003-3135-0356](#) ·  [HelgeBruelheide](#)

Martin-Luther-Universität Halle-Wittenberg, Institut für Biologie, Am Kirchtor 1, 06108, Halle (Saale). Germany; German Centre for Integrative Biodiversity Research (iDiv) - Halle-Jena-Leipzig. Germany · Funded by None

Abstract

Vegetation provides the foundation of life on Earth. Assessing biodiversity status and trends in plant communities is therefore critical to understand and quantify the effects of global change on ecosystems. Here, we present the largest dataset of vegetation plots (i.e. species co-occurrence or community composition data) ever released in open access. It contains information on 91,031 vegetation plots recording the cover or abundance of each plant species that occurs in a plot of a given surface area at the date of the botanical survey. Plots were derived from 103 local to regional datasets. To improve the representation of Earth's environmental conditions, plots were resampled from a larger pool of vegetation plots using an environmentally balanced sampling design. Each vegetation plot comes with information on community-weighted means and variances of key plant functional traits. Our open-access dataset can be used to explore global patterns of diversity at the plant community level, as ground truthing data in remote sensing applications or as a baseline for biodiversity monitoring.

References
